МАТЕМАТИЧКИ ВЕСНИК
58 (2006), 7–11

UDK 517.546
оригинални научни рад
research paper

HARMONIC FUNCTIONS STARLIKE OF THE COMPLEX ORDER

Sibel Yalçin and Metin Öztürk

Abstract. The main purpose of this paper is to introduce a class $TS_H^*(\gamma)$ ($\gamma \in \mathbf{C} \setminus \{0\}$) of functions which are harmonic in the unit disc. We give necessary and sufficient conditions for the functions to be in $TS_H^*(\gamma)$.

1. Introduction

A continuous complex-valued function $f = u+iv$ defined in a simply connected domain $\mathcal{D}$ is said to be harmonic in $\mathcal{D}$ if both u and v are real harmonic in $\mathcal{D}$. In any simply connected domain we can write $f = h + \bar{g}$, where h and g are analytic in $\mathcal{D}$. A necessary and sufficient condition for f to be locally univalent and sense preserving in $\mathcal{D}$ is that $|h'(z)| > |g'(z)|$ in $\mathcal{D}$.

Denote by S_H the class of functions $f = h+\bar{g}$ that are harmonic univalent and sense preserving in the unit disc $U = \{z : |z| < 1\}$ for which $f(0) = f_z(0) - 1 = 0$. Then for $f = h + \bar{g} \in S_H$ we may express analytic functions h and g as

$$h(z) = z + \sum_{n=2}^{\infty} a_n z^n, \qquad g(z) = \sum_{n=1}^{\infty} b_n z^n, \quad |b_1| < 1. \tag{1}$$

In 1984 Clunie and Sheil-Small [2] investigated the class S_H as well as its geometric subclasses and obtained some coefficient bounds. Since then, there have been several related papers on S_H and its subclasses.

Let TS_H denote the family of functions $f = h+\bar{g}$ that are harmonic in U with the normalization

$$h(z) = z - \sum_{n=2}^{\infty} a_n z^n, \qquad g(z) = \sum_{n=1}^{\infty} b_n z^n, \quad a_n, b_n \geqslant 0, \;\; b_1 < 1. \tag{2}$$

We let $TS_H^*(\gamma)$ denote the subclass of TS_H consisting of functions $f = h+\bar{g} \in TS_H$ that satisfy the condition

$$\operatorname{Re}\left\{1 + \frac{1}{\gamma}\left(\frac{zh'(z) - \overline{zg'(z)}}{h(z) + \overline{g(z)}} - 1\right)\right\} > 0, \qquad \gamma \in \mathbf{C} \setminus \{0\}. \tag{3}$$

AMS Subject Classification: 30 C 45, 30 C 50, 31 A 05
Keywords and phrases: Harmonic functions, starlike functions.

We further let $OS_H^*(\gamma)$ denote the subclass of TS_H consisting of functions $f = h + \bar{g} \in TS_H$ that satisfy the condition

$$\sum_{n=1}^{\infty} [2(n-1+|\gamma|)a_n + (n+1+|n+1-2\gamma|)b_n] \leqslant 4|\gamma|. \tag{4}$$

Denote by $PS_H^*(\gamma)$ the subclass of TS_H consisting of functions $f = h + \bar{g} \in TS_H$ that satisfy the condition

$$\sum_{n=1}^{\infty} \left[(n-1)\frac{\mathrm{Re}(\gamma)}{|\gamma|} + |\gamma|\right] a_n + \left[(n+1)\frac{\mathrm{Re}(\gamma)}{|\gamma|} - |\gamma|\right] b_n \leqslant 2|\gamma|. \tag{5}$$

Recently, Avcı and Zlotkiewicz [1], Jahangiri [3], Silverman [4], and Silverman and Silvia [5] studied the harmonic starlike functions. Avcı and Slotkiewicz [1] proved that the coefficient condition

$$\sum_{n=2}^{\infty} n(|a_n| + |b_n|) \leqslant 1, \qquad b_1 = 0$$

is sufficient for functions $f = h + \bar{g}$ to be harmonic starlike. Silverman 4 proved that this coefficient condition is also necessary if $b_1 = 0$ and if a_n and b_n in (1) are negative. Jahangiri [3] proved that if $f = h + \bar{g}$ is given by (1) and if

$$\sum_{n=2}^{\infty} \frac{n-\alpha}{1-\alpha}|a_n| + \sum_{n=1}^{\infty} \frac{n+\alpha}{1-\alpha}|b_n| \leqslant 1, \qquad 0 \leqslant \alpha < 1,$$

then f is a harmonic, univalent and starlike function of order α in U. This condition proved to be also necessary if h and g are of the form (2). The case when $\alpha = 0$ is given in [5] and for $\alpha = b_1 = 0$ see [4].

In this paper, we give an answer to the conjecture that $TS_H^*(\gamma) = OS_H^*(\gamma)$.

2. Main results

THEOREM 2.1. *$OS_H^*(\gamma) \subset TS_H^*(\gamma)$.*

Proof. Let $f \in OS_H^*(\gamma)$. According to the condition (2) we only need to show that if (4) holds then

$$\mathrm{Re}\left\{\frac{(\gamma-1)(h(z)+\overline{g(z)}) + zh'(z) - \overline{zg'(z)}}{\gamma(h(z)+\overline{g(z)})}\right\} > 0,$$

where $\gamma \in \mathbf{C} \setminus \{0\}$. Using the fact that $\mathrm{Re}\, w > 0$ if and only if $|1+w| > |1-w|$, it suffices to show that

$$|(2\gamma-1)(h(z)+\overline{g(z)}) + zh'(z) - \overline{zg'(z)}| - |h(z)+\overline{g(z)} - zh'(z) + \overline{zg'(z)}| > 0. \tag{6}$$

Substituting for $h(z)$ and $g(z)$ in (6) yields

$$|(2\gamma-1)(h(z)+\overline{g(z)}) + zh'(z) - \overline{zg'(z)}| - |h(z)+\overline{g(z)} - zh'(z) + \overline{zg'(z)}|$$

$$= \left|2\gamma z - \sum_{n=2}^{\infty}(2\gamma - 1 + n)a_n z^n - \sum_{n=1}^{\infty}(n + 1 - 2\gamma)b_n \bar{z}^n\right| -$$
$$- \left|\sum_{n=2}^{\infty}(n-1)a_n z^n + \sum_{n=1}^{\infty}(n+1)b_n \bar{z}^n\right|$$
$$\geqslant 2|\gamma||z| - \sum_{n=2}^{\infty}2(n - 1 + |\gamma|)a_n|z|^n - \sum_{n=1}^{\infty}(n + 1 + |n + 1 - 2\gamma|)b_n|z|^n$$
$$> 2|\gamma| - \left(\sum_{n=2}^{\infty}2(n - 1 + |\gamma|)a_n + \sum_{n=1}^{\infty}(n + 1 + |n + 1 - 2\gamma|)b_n\right) \geqslant 0.$$

The functions

$$f(z) = z - \sum_{n=2}^{\infty}\frac{|\gamma|}{n - 1 + |\gamma|}x_n z^n + \sum_{n=1}^{\infty}\frac{2|\gamma|}{n + 1 + |n + 1 - 2\gamma|}y_n \bar{z}^n \tag{7}$$

where x_n, y_n are non-negative and

$$\sum_{n=2}^{\infty}x_n + \sum_{n=1}^{\infty}y_n = 1,$$

show that the coefficient bound given by (4) is sharp. The functions of the form (7) are in $TS_H^*(\gamma)$ because

$$\sum_{n=2}^{\infty}2(n - 1 + |\gamma|)a_n + \sum_{n=1}^{\infty}(n + 1 + |n + 1 - 2\gamma|)b_n$$
$$= 2|\gamma|\left(1 + \sum_{n=2}^{\infty}x_n + \sum_{n=1}^{\infty}y_n\right) = 4|\gamma|. \quad \blacksquare$$

THEOREM 2.2. $TS_H^*(\gamma) \subset PS_H^*(\gamma)$.

Proof. Let $f \in TS_H^*(\gamma)$. From (3) we have

$$\operatorname{Re}\left\{\frac{1}{\gamma}\left(\frac{-\sum_{n=2}^{\infty}(n-1)a_n z^n - \sum_{n=1}^{\infty}(n+1)b_n \bar{z}^n}{1 - \sum_{n=2}^{\infty}a_n z^n + \sum_{n=1}^{\infty}b_n \bar{z}^n}\right)\right\} > -1.$$

If we choose z on the real axis and $z \to 1^-$ we get

$$\frac{\sum_{n=2}^{\infty}(n-1)a_n + \sum_{n=1}^{\infty}(n+1)b_n}{1 - \sum_{n=2}^{\infty}a_n + \sum_{n=1}^{\infty}b_n}\operatorname{Re}\left(\frac{1}{\gamma}\right) \leqslant 1,$$

whence

$$\frac{\sum_{n=2}^{\infty}(n-1)a_n + \sum_{n=1}^{\infty}(n+1)b_n}{1 - \sum_{n=2}^{\infty}a_n + \sum_{n=1}^{\infty}b_n}\frac{\operatorname{Re}(\gamma)}{|\gamma|^2} \leqslant 1,$$

and so

$$\sum_{n=2}^{\infty}(n-1)a_n + \sum_{n=1}^{\infty}(n+1)b_n \leqslant \frac{|\gamma|^2}{\operatorname{Re}(\gamma)}\left(1 - \sum_{n=2}^{\infty}a_n + \sum_{n=1}^{\infty}b_n\right)$$

which is equivalent to (5). Thus $f \in PS_H^*(\gamma)$. $\blacksquare$

THEOREM 2.3. *If $\gamma \in (0,1]$, then $OS_H^*(\gamma) = TS_H^*(\gamma) = PS_H^*(\gamma)$.*

Proof. If $\gamma \in (0,1]$, then the inequalities (4) and (5) are equivalent; hence $OS_H^*(\gamma) = PS_H^*(\gamma)$. By using Theorem 2.1 and Theorem 2.2, from this assertion we obtain the conclusion of the present theorem. ∎

THEOREM 2.4. *If $-1/2 > \mathrm{Re}(\gamma) \leqslant 0$ or $\gamma \in (3/2, +\infty)$, then*

$$PS_H^*(\gamma) \not\subseteq TS_H^*(\gamma).$$

Proof. Let

$$f(z) = z - z^2. \tag{8}$$

Then $f \in PS_H^*(\gamma)$ when $\gamma \in \mathbf{C} \setminus \{0\}$ and $\mathrm{Re}(\gamma) < 0$, because

$$\sum_{n=1}^{\infty} \left[(n-1)\frac{\mathrm{Re}(\gamma)}{|\gamma|} + |\gamma|\right] a_n + \left[(n+1)\frac{\mathrm{Re}(\gamma)}{|\gamma|} - |\gamma|\right] b_n$$
$$= |\gamma| \cdot 1 + \frac{\mathrm{Re}(\gamma)}{|\gamma|} + |\gamma| = 2|\gamma| + \frac{\mathrm{Re}(\gamma)}{|\gamma|} \leqslant 2|\gamma|.$$

Now let $r = \mathrm{Re}(\gamma) < 0$ and let s be a negative real number such that $1+2r(1-s) > 0$. If we choose $z = \dfrac{\gamma(1-s)}{1+\gamma(1-s)}$, then $z \in U$ and for f given by (8) we have

$$1 + \frac{1}{\gamma}\left(\frac{zh'(z) - \overline{zg'(z)}}{h(z) + \overline{g(z)}} - 1\right) = s < 0,$$

hence $f \notin TS_H^*(\gamma)$.

Similarly, let

$$f(z) = z + \bar{z}^2. \tag{9}$$

Then $f \in PS_H^*(\gamma)$ when $\gamma \in (3/2, +\infty)$, because

$$\sum_{n=1}^{\infty} \left[(n-1)\frac{\mathrm{Re}(\gamma)}{|\gamma|} + |\gamma|\right] a_n + \left[(n+1)\frac{\mathrm{Re}(\gamma)}{|\gamma|} - |\gamma|\right] b_n$$
$$= |\gamma| \cdot 1 + \left(3\frac{\mathrm{Re}(\gamma)}{|\gamma|} - |\gamma|\right) \cdot 1 = 3\frac{\mathrm{Re}(\gamma)}{|\gamma|} \leqslant 2|\gamma|.$$

Now let $\gamma \in (3/2, +\infty)$ and let s be a negative real number such that $\gamma+\gamma(s-1) < 0$. If we choose $z = -\dfrac{\gamma(s-1)}{3+\gamma(s-1)}$, then $z \in U$ and for f given by (9) we have

$$1 + \frac{1}{\gamma}\left(\frac{zh'(z) - \overline{zg'(z)}}{h(z) + \overline{g(z)}} - 1\right) = s < 0,$$

hence $f \notin TS_H^*(\gamma)$. ∎

THEOREM 2.5. *If* $\gamma \in (-\infty, -1) \cup (-1/2, 0)$, *then*

$$TS_H^*(\gamma) \nsubseteq OS_H^*(\gamma).$$

Proof. Let $\gamma \in (-\infty, -1)$; we verify that the functions

$$f_\lambda(z) = z - \lambda z^2 \tag{10}$$

belong to $TS_H^*(\gamma)$ for $\lambda > \frac{\gamma}{1+\gamma}$ and that $f \notin OS_H^*(\gamma)$.

Indeed we have

$$\sum_{n=1}^{\infty} [2(n-1+|\gamma|)a_n + (n+1+|n+1-2\gamma|)b_n] = 2|\gamma| + 2(1+|\gamma|)\lambda > 4|\gamma|,$$

because $\lambda > \frac{\gamma}{1+\gamma} > 1$.

We also have

$$\operatorname{Re}\left\{1 + \frac{1}{\gamma}\left(\frac{zh'_\lambda(z) - \overline{zg'_\lambda(z)}}{h_\lambda(z) + \overline{g_\lambda(z)}} - 1\right)\right\} = \operatorname{Re}\left\{1 + \frac{\lambda z}{\gamma(\lambda z - 1)}\right\} > 0, \quad z \in U, \tag{11}$$

for $\lambda > \frac{\gamma}{1+\gamma}$ and $\gamma < -1$, hence $f_\lambda \in TS_H^*(\gamma)$.

Let now $\gamma \in (-1/2, 0)$, and let f_λ be defined by (10), where

$$-\frac{\gamma}{1-\gamma} < \lambda < -\frac{\gamma}{1+\gamma}.$$

Then $\lambda > -\frac{\gamma}{1-\gamma}$ implies $f_\lambda \notin OS_H^*(\gamma)$ and for $\lambda < -\dfrac{\gamma}{1+\gamma}$ the inequality (11) is also verified, hence $f_\lambda \in TS_H^*(\gamma)$. ∎

ACKNOWLEDGMENTS. The authors warmly thank the referee and editors for their suggestions and criticisms which have essentially improved our original paper.

REFERENCES

[**1**] Avcı, Y. and Zlotkiewicz, E., *On harmonic univalent mappings*, Ann. Univ. Mariae Curie-Sklodowska, Sect. A, **44** (1990), 1–7.

[**2**] Clunie, J. and Sheil-Small, T., *Harmonic univalent functions*, Ann. Acad. Sci. Fenn., Ser. A I Math. **9** (1984), 3–25.

[**3**] Jahangiri, J. M., *Harmonic functions starlike in the unit disk*, J. Math. Anal. Appl., **235** (1999), 470–477.

[**4**] Silverman, H., *Harmonic univalent functions with negative coefficients*, J. Math. Anal. Appl., **220** (1998), 283–289.

[**5**] Silverman, H. and Silvia, E. M., *Subclasses of harmonic univalent functions*, N. Z. J. Math., **28** (1999), 275–284.

(received 24.09.2003, in revised form 24.04.2005)

Uludağ Üniversitesi, Fen Edebiyat Fakültesi, Matematik Bölümü, 16059, Bursa, Turkey
E-mail: skarpuz@uludag.edu.tr