New York Journal of Mathematics

New York J. Math. 22 (2016) 711–714.

On smoothly superslice knots

Daniel Ruberman

ABSTRACT. We find smoothly slice (in fact doubly slice) knots in the 3-sphere with trivial Alexander polynomial that are not superslice, answering a question posed by Livingston and Meier.

1. Introduction

A recent paper of Livingston and Meier raises an interesting question about superslice knots. Recall [3] that a knot K in S^3 is said to be superslice if there is a slice disk D for K such that the double of D along K is the unknotted 2-sphere S in S^4 . We will refer to such a disk as a superslicing disk. In particular, a superslice knot is slice and also doubly slice, that is, a slice of an unknotted 2-sphere in S^4 . Livingston and Meier ask about the converse in the smooth category.

Problem 4.6 (Livingston–Meier [10]). Find a smoothly slice knot K with $\Delta_K(t) = 1$ that is not smoothly superslice.

The corresponding question in the topological (locally flat) category is completely understood [10, 12], for a knot K with $\Delta_K(t) = 1$ is topologically superslice.

In this note we give a simple solution to Problem 4.6, making use of Taubes' proof [16] that Donaldson's diagonalization theorem [5] holds for certain noncompact manifolds. For K a knot in S^3 , we write $\Sigma_k(K)$ for a k-fold cyclic branched cover of S^3 branched along K. The same notation will be used for the corresponding branched cover along an embedded disk in B^4 or sphere in S^4 .

Theorem 1.1. Suppose that J is a knot with Alexander polynomial 1, so that $\Sigma_k(J) = \partial W$, where W is simply connected and the intersection form on W is definite and not diagonalizable. Then the knot K = J # - J is smoothly doubly slice, but is not smoothly superslice.

An unpublished argument of Akbulut says that the positive Whitehead double of the trefoil is a knot J satisfying the hypotheses of the theorem, with k = 2. The construction is given as [1, Exercise 11.4] and is also

Received June 23, 2016.

²⁰¹⁰ Mathematics Subject Classification. 57M25, 57Q60.

Key words and phrases. Superslice, doubly slice.

The author was partially supported by NSF Grant 1506328.

DANIEL RUBERMAN

documented, along with some generalizations, in the paper [4]. Hence J gives an answer to Problem 4.6. We remark that for the purposes of the argument, it doesn't matter if W is positive or negative definite, as one could replace J by -J and change all the signs.

We need a simple and presumably well-known algebraic lemma.

Lemma 1.2. Suppose that



is a pushout of groups, and that $i_1 = i_2$. Then C surjects onto B.

Proof. This follows from the universal property of pushouts; the identity map id_B satisfies $id_B \circ i_1 = id_B \circ i_2$, and hence defines a homomorphism $C \to B$ with the same image as id_B .

Applying Lemma 1.2 to the decomposition of the complement of the unknot in S^4 into two disk complements, we obtain the following useful facts. (The first of these was presumably known to Kirby and Melvin; compare [8, Addendum, p. 58], and the second is due to Gordon and Sumners [6].)

Corollary 1.3. If K is superslice and D is a superslicing disk, then

$$\pi_1(B^4 - D) \cong \mathbb{Z}$$
 and $\Delta_K(t) = 1.$

Proof. The lemma says that there is a surjection

$$\mathbb{Z} \cong \pi_1(S^4 - S) \to \pi_1(B^4 - D).$$

Hence $\pi_1(B^4 - D)$ is abelian and so must be isomorphic to \mathbb{Z} . This condition implies, using Milnor duality [13] in the infinite cyclic covering, that the homology of the infinite cyclic covering of $S^3 - K$ vanishes, which is equivalent to saying that $\Delta_K(t) = 1$.

Proof of Theorem 1.1. It is standard [15] that any knot of the form J # -J is doubly slice. In fact, it is a slice of the 1-twist spin of J, which was shown by Zeeman [17] to be unknotted.

Suppose that K is superslice and let D be a superslicing disk, so $D \cup_K D = S$, an unknotted sphere. Then $S^4 = \Sigma_k(S) = V \cup_Y V$, where we have written $Y = \Sigma_k(K)$ and $V = \Sigma_k(D)$. By Corollary 1.3, the k-fold cover of $B^4 - D$ has $\pi_1 \cong \mathbb{Z}$, so the branched cover V is simply connected.

Note that $\Sigma_k(K) = \Sigma_k(J) \# -\Sigma_k(J)$. Since $\Delta_J(t) = 1$, the same is true for $\Delta_K(t)$, moreover this implies that both $\Sigma_k(J)$ and $\Sigma_k(K)$ are homology spheres. An easy Mayer-Vietoris argument says that $V = \Sigma_k(D)$ is a homology ball; in fact Claim 1.3 implies that it is contractible. Adding a

712

3-handle to V, we obtain a simply-connected homology cobordism V' from $\Sigma_k(J)$ to itself. By hypothesis, there is a manifold W with boundary $\Sigma_k(J)$ and nondiagonalizable intersection form. Stack up infinitely many copies of V', and glue them to W to make a definite periodic-end manifold M, in the sense of Taubes [16]. Since $\pi_1(V)$ is trivial, M is *admissible* (see [16, Definition 1.3]), and Taubes shows that its intersection form (which is the same as that of W) is diagonalizable. This contradiction proves the theorem. \Box

The fact that $\pi_1(B^4 - D) \cong \mathbb{Z}$ for a superslicing disk leads to a second obstruction to supersliceness, based on the Ozsváth–Szabó *d*-invariant [14]. Recall from [11] (for degree 2 covers) and [7] in general that for a knot Kand prime p, that one denotes by $\delta_{p^n}(K)$ the *d*-invariant of a particular spin structure \mathfrak{s} on $\Sigma_{p^n}(K)$ pulled back from the 3-sphere. The fact that a p^n fold branched cover of a slicing disk is a rational homology ball implies that if K slice then $\delta_{p^n}(K) = 0$. For a non-prime-power degree k, the invariant $\delta_k(K)$ might not be defined, because $\Sigma_k(K)$ is not a rational homology sphere. (One might define such an invariant using Floer homology with twisted coefficients as in [2, 9], but there's no good reason that it would be a concordance invariant.)

Theorem 1.4. If K is superslice, then for any k, the d-invariant

 $d(\Sigma_k(K),\mathfrak{s}_0)$

is defined and vanishes.

Proof. Since by Corollary 1.3 the Alexander polynomial is trivial, so $\Sigma_k(K)$ is a homology sphere, and hence $d(\Sigma_k(K), \mathfrak{s}_0)$ is defined. (There is only the one spin structure.) As in the proof of Theorem 1.1, the branched cover $\Sigma_k(D)$ is contractible, and hence [14, Theorem 1.12], $d(\Sigma_k(K), \mathfrak{s}_0) = 0$. \Box

Sadly, we do not know any examples of a slice knot where Theorem 1.4 provides an obstruction to it being superslice. For such a knot would not be ribbon, so we would also have a counterexample to the slice-ribbon conjecture!

Acknowledgements. Thanks to Hee Jung Kim for an interesting conversation that led to this paper, and to Chuck Livingston, Paul Melvin, and Nikolai Saveliev for comments on an initial draft.

References

- [1] AKBULUT, SELMAN. 4-manifolds. Oxford University Press, to appear. Preprint http: //www.math.msu.edu/~akbulut/papers/akbulut.lec.pdf.
- BEHRENS, STEFAN; GOLLA, MARCO. Heegaard Floer correction terms, with a twist. Preprint, 2015. arXiv:1505.07401.
- [3] BRAKES, W. R. Property R and superslices. Quart. J. Math. Oxford Ser. (2) 31 (1980), no. 123, 263–281. MR0587091, Zbl 0408.57003, doi: 10.1093/qmath/31.3.263.

DANIEL RUBERMAN

- [4] COCHRAN, TIM D.; GOMPF, ROBERT E. Applications of Donaldson's theorems to classical knot concordance, homology 3-spheres and property *P. Topology* 27 (1988), no. 4, 495–512. MR0976591, Zbl 0669.57003, doi: 10.1016/0040-9383(88)90028-6.
- [5] DONALDSON, S. K. An application of gauge theory to four-dimensional topology. J. Differential Geom. 18 (1983), no. 2, 279–315. MR0710056, Zbl 0507.57010.
- [6] GORDON, C. MCA.; SUMNERS, D. W. Knotted ball pairs whose product with an interval is unknotted. *Math. Ann.* **217** (1975), no. 1, 47–52. MR0380816, Zbl 0293.57010, doi: 10.1007/BF01363239.
- JABUKA, STANISLAV. Concordance invariants from higher order covers. *Topology Appl.* 159 (2012), no. 10–11, 2694–2710. MR2923439, Zbl 1250.57011, arXiv:0809.1088, doi: 10.1016/j.topol.2012.03.014.
- [8] KIRBY, ROBION; MELVIN, PAUL. Slice knots and property R. Invent. Math. 45 (1978), no. 1, 57–59. MR0467754, Zbl 0377.55002, doi: 10.1007/BF01406223.
- [9] LEVINE, ADAM SIMON; RUBERMAN, DANIEL. Generalized Heegaard Floer correction terms. Proceedings of the Gökova Geometry-Topology Conference, 2013, 76–96, Gkova Geometry/Topology Conference (GGT), Gkova, 2014. MR3287799, Zbl 1333.57024, arXiv:1403.2464.
- [10] LIVINGSTON, CHARLES; MEIER, JEFFREY. Doubly slice knots with low crossing number. New York J. Math. 21 (2015), 1007–1026. MR3425633, Zbl 1328.57009, arXiv:1504.03368.
- [11] MANOLESCU, CIPRIAN; OWENS, BRENDAN. A concordance invariant from the Floer homology of double branched covers. *Int. Math. Res. Not. IMRN* 2007, no. 20, Art. ID rnm077, 21 pp. MR2363303, Zbl 1132.57013, arXiv:math/0508065, doi:10.1093/imrn/rnm077.
- [12] MEIER, JEFFREY. Distinguishing topologically and smoothly doubly slice knots. J. Topol. 8 (2015), no. 2, 315–351. MR3356764, Zbl 1320.57027, arXiv:1401.1161, doi:10.1112/jtopol/jtu027.
- [13] MILNOR, JOHN W. Infinite cyclic coverings. 1968 Conference on the Topology of Manifolds (Michigan State Univ., E. Lansing, Mich., 1967) pp. 115–133. Prindle, Weber & Schmidt, Boston, Mass. MR0242163, Zbl 0179.52302.
- [14] OZSVÁTH, PETER; SZABÓ, ZOLTÁN. Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary. Adv. Math. 173 (2003), no. 2, 179–261. MR1957829, Zbl 1025.57016, arXiv:math/0110170, doi:10.1016/S0001-8708(02)00030-0.
- [15] SUMNERS, D. W. Invertible knot cobordisms. Comment. Math. Helv. 46 (1971), 240–256. MR0290351, Zbl 0227.57007, doi: 10.1007/BF02566842.
- [16] TAUBES, CLIFFORD HENRY. Gauge theory on asymptotically periodic 4-manifolds. J. Differential Geom. 25 (1987), no. 3, 363–430. MR0882829, Zbl 0615.57009.
- [17] ZEEMAN, E. C. Twisting spun knots. Trans. Amer. Math. Soc. 115 (1965), 471–495.
 MR0195085, Zbl 0134.42902, doi: 10.2307/1994281.

(Daniel Ruberman) Department of Mathematics, MS 050, Brandels University, Waltham, MA 02454

ruberman@brandeis.edu

This paper is available via http://nyjm.albany.edu/j/2016/22-32.html.

714