

ON THE DEFINITION OF A QUADRATIC FORM

Svetozar Kurepa

Abstract. In the first part of this paper we give a simple proof of the following wellknown theorem [3]: If a function $q : X \rightarrow C$ satisfies the parallelogram law and the homogeneity property $q(\lambda x) = |\lambda|^2 q(x)$ ($\lambda \in C, x \in X$), then there exists a sesquilinear form $L : X \times X \rightarrow C$ such that $q(x) = L(x; x)$ ($x \in X$).

If X is a real vector space then a quadratic form on X is to be defined as a function $q : X \rightarrow R$ the complexification ($q_c(q_c(x+iy)) = q(x)+q(y); x, y \in X$) of which has the homogeneity property

$$q_c(\lambda z) = |\lambda|^2 q_c(z) \quad (\lambda \in C, z \in X_c = X \times X).$$

In the second part of this paper we continue the study of quadratic forms on modules over algebras studied in [6], [7] and [4]. We assume as in [4] that the algebra A has the identity element and that it has the regularity property: For any $t \in A$ there exists a natural number n such that $t + n$ and $t + n + 1$ are invertible in A .

1. On the definition of a quadratic form

If X is a complex vector space and $L : X \times X \rightarrow C$ a sesquilinear form, then a function

$$(1) \quad q(x) = L(x, x) \quad (x \in X)$$

has properties:

$$(2) \quad q(x + y) + q(x - y) = 2q(x) + 2q(y) \quad (x, y \in X),$$

$$(3) \quad q(\lambda x) = |\lambda|^2 q(x) \quad (\lambda \in C; x \in X).$$

Proof. I. Halperin in 1963 (The New Scottish Book) asked whether for a function $q : X \rightarrow C$ which satisfies (2) and (3) there exists a sesquilinear form such that (1) holds. The positive answer to this question was given in [3] and the proof was simplified in [5]. Here we give even simpler proof of this fact.

AMS Subject Classification (1980): Primary 15A63, Secondary 39B50, 46C10, 46K99.

Key words and phrases: Modul, Vector space, Algebra, Invertible element, Bilinear form, Sesquilinear form, Quadratic form, Complexification of a vector space, Jordan derivation, Additive function.

THEOREM 1 [3]. *If X is a complex vector space and a function $q : X \rightarrow C$ satisfies conditions (2) and (3) then a functional $L : X \times X \rightarrow C$ defined by*

$$(4) \quad L(x, y) = (q(x + y) - q(x - y))/4 + i(q(x + iy) - q(x - iy))/4 \quad (x, y \in X)$$

is sesquilinear and (1) holds true.

We need three lemmas for the proof of this theorem. Although lemmas 1, 2 are well-known we prove them here for the convenience of a reader.

LEMMA 1. *If a function $q : X \rightarrow C$ satisfies the parallelogram law (2), then a function*

$$S(x, y) = q(x + y) - q(x - y) \quad (x, y \in X)$$

is biadditivem symmetric and $4q(x) = S(x, x)$ ($x \in X$).

Proof. From (2) for $x = y = 0$ we get $q(0) = 0$; for $x = 0$ we get $q(-y) = q(y)$ i.e. q is an even function. By taking $x = y$ in (2) we get $q(2x) = 4q(x)$.

For $x, y, u \in X$ we have:

$$\begin{aligned} S(x + y, 2u) &= q(x + y + 2u) - q(x + y - 2u) \\ &= q((x + u) + (y + u)) + q((x + u) - (y + u)) \\ &\quad - q((x - u) + (y - u)) - q((x - u) - (y - u)) \\ &= (2q(x + u) + 2q(y + u)) - (2q(x - u) + 2q(y - u)) \\ &= 2S(x, u) + 2S(y, u). \end{aligned}$$

From here for $y = 0$ and $x = z$ we get $S(z, 2u) = 2S(z, u)$ which for $z = x + y$ leads to

$$S(x + y, u) = S(x, u) + S(y, u). \quad \square$$

LEMMA 1. *If $q : X \rightarrow C$ satisfies (2) and (3) then the function $L : X \times X \rightarrow C$ defined by (4) is biadditive, $q(x) = L(x, x)$ ($x \in X$) and*

$$(6) \quad L(ix, y) = iL(x, y), \quad L(x, iy) = -iL(x, y) \quad (x, y \in X).$$

Proof. Using (4) and (3) for $\lambda = i$ we have

$$\begin{aligned} 4L(ix, y) &= (q(ix + y) - q(ix - y)) + i(q(ix + iy) - q(ix - iy)) \\ &= q(x - iy) - q(x + iy) + i(q(x + y) - q(x - y)) = 4iL(x, y), \\ 4L(x, iy) &= (q(x + iy) - q(x - iy)) + i(q(x + i \cdot iy) - q(x - i \cdot iy)) \\ &= q(x + iy) - q(x - iy) + i(q(x - y) - q(x + y)) = -4iL(x, y). \quad \square \end{aligned}$$

LEMMA 3 [5]. *If $f : C \rightarrow C$ is an additive function and*

$$f(\lambda) = |\lambda|^2 f(1/\lambda) \quad (\lambda \in C, \lambda \neq 0)$$

then

$$f(\lambda) = f(1) \operatorname{Re} \lambda \quad (\lambda \in C).$$

Proof. A function $g(\lambda) = f(1) \operatorname{Re} \lambda - f(\lambda)$ is additive and

$$g(1) = 0, \quad g(\lambda) = |\lambda|^2 g(1/\lambda), \quad \lambda \neq 0.$$

If $\lambda \neq 0$ then

$$\begin{aligned} g(\lambda) &= g(1 + \lambda) = |1 + \lambda|^2 g\left(\frac{1}{1 + \lambda}\right) = |1 + \lambda|^2 g\left(1 - \frac{\lambda}{1 + \lambda}\right) = -|1 + \lambda|^2 g\left(\frac{\lambda}{1 + \lambda}\right) = \\ &= -|1 + \lambda|^2 \left|\frac{\lambda}{1 + \lambda}\right|^2 g\left(\frac{1 + \lambda}{\lambda}\right) = -|\lambda|^2 g(1 + 1/\lambda) = -|\lambda|^2 g(1/\lambda) = -g(\lambda). \end{aligned}$$

Thus $g = 0$ and (8) follows. \square

Proof of Theorem 1. For any $x, y \in X$ define

$$(9) \quad f(\lambda) = L(\lambda x, y) + L(x, \lambda y) \quad (\lambda \in C).$$

Obviously, $\lambda \rightarrow f(\lambda)$ is an additive function. By use of (3) for $\lambda = 0$ we have

$$\begin{aligned} 4L(\lambda x, y) &= q(\lambda x + y) - q(\lambda x - y) + i(q(\lambda x + iy) - q(\lambda x - iy)) \\ &= |\lambda|^2 [(q(x + y/\lambda) - q(x - y/\lambda)) + i(q(x + y/\lambda) - q(x - iy/\lambda))] \\ &= 4|\lambda|^2 L(x, y/\lambda). \end{aligned}$$

In the same way we get $L(x, \lambda y) = |\lambda|^2 L(y/\lambda, x)$.

Thus the function (9) satisfies conditions of Lemma 3 so that $f(\lambda) = f(1) \operatorname{Re} \lambda$, i.e.

$$(10) \quad L(\lambda x, y) + L(x, \lambda y) = 2L(x, y) \operatorname{Re} \lambda \quad (\lambda \in C; x, y \in X).$$

If $\lambda = it$ ($t \in \mathbf{R}$), then (10) and Lemma 2 imply

$$(11) \quad L(tx, y) = L(x, ty) \quad (t \in \mathbf{R}; x, y \in X).$$

If $\lambda = t$ ($t \in \mathbf{R}$), then (10) implies

$$L(tx, y) + L(x, ty) = 2tL(x, y)$$

which together with (11) leads to

$$(12) \quad L(tx, y) = tL(x, y) \quad (t \in \mathbf{R}; x, y \in X).$$

Now if $\lambda = \sigma + i\tau$ ($\sigma, \tau \in \mathbf{R}$) then the biadditivity of L (Lemma 1) and (12) imply:

$$\begin{aligned} L(\lambda x, y) &= L(\sigma x + i\tau x, y) = L(\sigma x, y) + L(i\tau x, y) \\ &= \sigma L(x, y) + iL(\tau x, y) = \sigma L(x, y) + i\tau L(x, y) = \lambda L(x, y). \\ L(x, \lambda y) &= L(x, \sigma y + i\tau y) = L(x, \sigma y) + L(x, i\tau y) \\ &= L(x, \sigma y) - iL(x, \tau y) = L(\sigma x, y) - iL(\tau x, y) = \\ &= \sigma L(x, y) - i\tau L(x, y) = \bar{\lambda} L(x, y). \quad \square \end{aligned}$$

The complexification X_c of a real vector space X is defined as a set $X \times X$ with algebraic operations:

$$\begin{aligned}(x, y) + (x', y') &= (x + x', y + y') \quad (x, x', y, y' \in X), \\ (\sigma + i\tau)(x, y) &= (\sigma x - \tau y, \tau x + \sigma y') \quad (\sigma, \tau \in \mathbf{R}; x, y \in X).\end{aligned}$$

We write $(x, y) = x + iy$ ($x, y \in X$).

If $B : X \times X \rightarrow \mathbf{R}$ is a bilinear form then its complexification $B_c : X_c \times X_c \rightarrow C$ is defined by

$$B_c(x + iy, x' + iy') = B(x, x') + B(y, y') + i(B(y, x') - B(x, y')).$$

If B is symmetric then

$$B_c(x + iy, x + iy) = B(x, x) + B(y, y) \quad (x, y \in X)$$

THEOREM 2. *Let X be a real vector space and $q : X \rightarrow \mathbf{R}$ any function and $B : X \times X \rightarrow \mathbf{R}$ a function defined by*

$$(13) \quad B(x, y) = (q(x + y) - q(x - y))/4 \quad (x, y \in X)$$

Then, the function B is bilinear if and only if the complexification $q_c :$

$$(14) \quad q_c(x + iy) = q(x) + q(y) \quad (x, y \in X)$$

of q has the following homogeneity property

$$(15) \quad q_c(\lambda z) = |\lambda|^2 q_c(z) \quad (\lambda \in C; z \in X_c)$$

Proof. For $\lambda = \sigma + i\tau$ ($\sigma, \tau \in \mathbf{R}$) and $z = x + iy$ ($x, y \in X$) from (15) we get:

$$(16) \quad (\sigma^2 + \tau^2)(q(x) + q(y)) = q(\sigma x - \tau y) + q(\tau x + \sigma y).$$

From (16) for $\sigma = \tau = 1$ we find that q satisfies the parallelogram law (2) so that by Lemma 1 the function B is biadditive, symmetric and $q(x) = B(x, x)$. Furthermore (14) and (2) imply

$$q_c(u + v) + q_c(u - v) = 2q_c(u) + 2q_c(v)$$

for all $u, v \in X$. Now (17), (15) and Theorem 1 imply that the functional $L : X_c \times X_c \rightarrow C$ defined by

$$L(u, v) = (q_c(u + v) - q_c(u - v))/4 + i(q_c(u + iv) - q_c(u - iv))/4$$

is sesquilinear on X_c . If $u = x$ and $v = y$ are vectors in X , then

$$L(x, y) = (q(x + y) + q(0) - q(x - y) - q(0))/4 + i(q(x) + q(y) - q(x) - q(y)) = B(x, y).$$

Thus $B(tx, y) = L(tx, y) = tL(x, y)$ holds for any $t \in \mathbf{R}$ and all $x, y \in X$. This and $B(x, y) = B(y, x)$ imply that B is bilinear. \square

Using Theorem 1 we see that one can define a quadratic form on a complex vector space X as a function $q : X \rightarrow C$ which satisfies the parallelogram law (2) and has the homogeneity property (3).

According to Theorem 2 a quadratic form on a real vector space X can be defined as a function $q : X \rightarrow \mathbf{R}$ such that its complexification q_c defined by (14) satisfies the homogeneity property (15). As it is well-known for a function $q : X \rightarrow \mathbf{R}$ defined on a real vector space which satisfies the parallelogram law (2) and the homogeneity property

$$q(tx) = t^2 q(x) \quad (t \in \mathbf{R}, x \in X)$$

in general there does not exist a bilinear form $B : X \times X \rightarrow R$ such that $q(x) = B(x, x)$ ($x \in X$) (See: [2], [1]).

Remark 1. If $f, F : \mathbf{R} \rightarrow \mathbf{R}$ are additive functions such that

$$(18) \quad f(t) = t^2 f(1/t), \quad F(t) = -t^2 F(1/t), \quad (t \in \mathbf{R}, t \neq 0)$$

then $f(t) = f(1)t$ and $F(ts) = tF(s) + sF(t)$, i.e. f is continuous and F is a derivation on \mathbf{R} , hence F is not continuous unless $F = 0$.

On the other hand if $f, F : C \rightarrow C$ are additive and if

$$(19) \quad f(\lambda) = |\lambda|^2 f(1/\lambda), \quad F(\lambda) = -|\lambda|^2 F(1/\lambda) \quad (\lambda \in C, \lambda \neq 0)$$

then $f(\lambda) = f(1) \operatorname{Re} \lambda$ and $F(\lambda) = F(i) \operatorname{Im} \lambda$. In this case both functions f and F are continuous. In fact, if $F(\lambda) = -|\lambda|^2 F(1/\lambda)$ holds for all $\lambda \in C$, $\lambda \neq 0$, then a function $f_1(\lambda) = F(i\lambda)$ satisfies the condition

$$f_1(\lambda) = |\lambda|^2 f_1(1/\lambda) \quad (\lambda \in C, \lambda \neq 0).$$

By (18) we are given essentially different conditions on functions f and F while conditions given by (19) can be transformed one to another.

2. Quadratic forms on modules over algebras

By X and X' we denote complex vector spaces and by A complex algebra with unit 1. We assume that the algebra A has the following regularity property (R):

For any $t \in A$ there exists a natural number n such that $t + n$ and $t + n + 1$ are invertible elements in A .

Furthermore we assume that X is a left modul over A and that X' is left and right modul over A .

THEOREM 3. *Let A , X and X' be as above. If $q : X \rightarrow X'$ is a quadratic form i.e.*

$$(1) \quad q(x + y) + q(x - y) = 2q(x) + 2q(y) \quad (x, y \in X)$$

and if q satisfies the homogeneity condition

$$(2) \quad q(tx) = tq(x)t \quad (t \in A, x \in X)$$

then the function $M : X \times X \rightarrow X$ defined by

$$(3) \quad M(x, y) = (q(x + y) - q(x - y))/8 - i(q(x + iy) - q(x - iy))/8 \quad (x, y \in X)$$

is biadditive, symmetric,

$$q(x) = M(x, x), \quad M(ix, y) = iM(x, y) \quad (x, y \in X)$$

and

$$(4) \quad M(tx, y) + M(x, ty) = tM(x, y) + M(x, y)t \quad (t \in A; x, y \in X).$$

Furthermore, the function

$$(5) \quad h(t; x, y) = (M(tx, y) - M(x, ty))/2 \quad (t \in A; x, y \in X)$$

is a Jordan derivation on A , i.e.

$$h(t \circ s; x, y) = t \circ h(s; x, y) + h(t; x, y) \circ s$$

holds true, where

$$t \circ s = ts + st \quad (t, s \in A).$$

The proof of Theorem 3 is obtained by using the following two lemmas.

LEMMA 4. (See Lemma 1 in [4]). *If an additive function $g : A \rightarrow X'$ for each invertible element $t \in A$ satisfies the condition*

$$(7) \quad g(t) = tg(t^{-1})t$$

then

$$(8) \quad g(t) = (tg(1) + g(1)t)/2 \quad (t \in A).$$

LEMMA 5. *If an additive function $h : A \rightarrow X'$ for each invertible element $t \in A$ satisfies the condition*

$$(7) \quad h(t) = -th(t^{-1})t$$

then h is a Jordan derivation on A , i.e.

$$h(t \circ s) = h(t) \circ s + t \circ h(s) \quad (t, s \in A).$$

Proof of Lemma 5. For $t \in A$ we take a natural number n such that $t + n$ and $t + n + 1$ are invertible in A . By applying the function h on the identity

$$(t + n)^{-1} - (t + n + 1)^{-1} = (t^2 + 2nt + t + n^2 + n)^{-1}$$

and by using (9) we get

$$\begin{aligned} & -(t + n)^{-1} \cdot h(t + n) \cdot (t + n)^{-1} + (t + n + 1)^{-1} \cdot h(t + n + 1) \cdot (t + n + 1)^{-1} = \\ & = -(t^2 + 2nt + t + n^2 + n)^{-1} \cdot h(t^2 + 2nt + t + n^2 + n) \cdot (t^2 + 2nt + t + n^2 + n). \end{aligned}$$

Multiply the last relation from the left and from the right by $(t+n)(t+n+1)$ to get:

$$h(t^2 + 2nt + tn^2 + n) = (t+n+1) \cdot h(t+n) \cdot (t+n+1) - (t+n) \cdot h(t+n) \cdot (t+n)$$

from which by using $h(1) = 0$ we get

$$(11) \quad h(t^2) = th(t) + h(t)t \quad (t \in A).$$

If in (11) we replace t by $t+s$ we get (10). \square

Proof of Theorem 3. Since q is quadratic, the function M defined by (3) is biadditive and $q(x) = M(x, x)$. By using (2) it is easy to find

$$M(ix, y) = iM(x, y), \quad M(x, y) = M(y, x)$$

and

$$(12) \quad M(tx, y) = tM(x, t^{-1}y)t.$$

for any invertible element $t \in A$ and for all $x, y \in X$. If $x, y \in X$ are fixed, then the function

$$g(t) = M(tx, y) + M(x, ty) \quad (t \in A)$$

satisfies all conditions of Lemma 4 so that (8) and $g(1) = 2M(x, y)$ imply (4).

By using (2) for the function h defined by (5) we find

$$h(t; x, y) - th(t^{-1}; x, y)t$$

for any invertible $t \in A$. By applying Lemma 5 we get (6). \square

Remark 2. If X and X' are real vector spaces and A is a real algebra with the regularity property (R), then for a quadratic form which has the homogeneity property (2) the function

$$B(x, y) = (q(x+y) - q(x-y))/4 \quad (x, y \in X)$$

is biadditive, symmetric, $B(x, x) = q(x)$ ($x \in X$),

$$B(tx, y) + B(x, ty) = tB(x, y) + B(x, y) \quad (t \in A; x, y \in X)$$

and the function

$$h(t; x, y) = (B(tx, y) - B(x, ty))/4 \quad (t \in A; x, y \in X)$$

is a Jordan derivation, i.e. h has the property (6). The proof of these fact follows the proof of Theorem 3.

REFERENCES

- [1] A. M. Gleason, *The definition of quadratic form*, Amer. Math. Monthly **73** (1966), 1049–1056.

- [2] S. Kurepa, *The Cauchy functional equation and scalar product in vector spaces*, Glasnik Mat. Fiz. Astr. **19** (1964), 23–36.
- [3] S. Kurepa, *Quadratic and sesquilinear functionals*, Glasnik Mat. Fiz. Astr. **20** (1965), 79–92.
- [4] S. Kurepa, *On quadratic forms*, to appear in Aequationes Math.
- [5] P. Vrbová, *Quadratic functionals and bilinear forms*, Časopis Pest. Mat. **98** (1973), 159–161.
- [6] J. Vukan, *A result concerning additive functions in hermitian Banach *-algebras*, Proc. Amer. Math. Soc. **91** (1984), 367–372.
- [7] J. Vukman, *Some results concerning the Cauchy functional equation in certain Banach algebras*, Bull. Austral. Math. Soc. **31** (1985), 137–144.

Matematički odjel
Prirodoslovni-matematički fakultet
41000 Zagreb
Jugoslavija

(Received 20 01 1987)