

## THE DUAL STEINER FORMULA FOR CONVEX COMPACTA

Ljuban Dedić

**Abstract.** The classical Steiner formula deals with the volume of  $K + rD_n$ , where  $K$  is a convex compact set in  $\mathbf{R}^n$ ,  $r \geq 0$ , and  $D_n$  the Euclidean disc in  $\mathbf{R}^n$ . We compute the volume of  $(K + rD_n)^*$ , where  $K^*$  means the dual of  $K$ . We also represent some Steiner functionals by integrals and prove some inequalities.

### 1. Introduction and notations

To specify notations we give here some elementary properties of convex compacta in  $\mathbf{R}^n$ . For details see [1].

Let  $K(\mathbf{R}^n)$  be the set of all convex compacta in  $\mathbf{R}^n$ . If  $K \in K(\mathbf{R}^n)$  we introduce the function  $h_K : \mathbf{R}^n \rightarrow \mathbf{R}$  defined by

$$h_K(x) = \max_{y \in K} (x|y), \quad x \in \mathbf{R}^n$$

where  $(x|y) = x_1y_1 + \dots + x_ny_n$  is the standard scalar product in  $\mathbf{R}^n$ . The function  $h_K$  has the following properties:

- 1)  $h_K(tx) = th_K(x)$ ,  $t \geq 0$ ,  $x \in \mathbf{R}^n$ .
- 2)  $h_K(x + y) \leq h_K(x) + h_K(y)$ ,  $x, y \in \mathbf{R}^n$ .
- 3) If  $0 \in K$  then  $h_K \geq 0$ .
- 4) If  $0 \in K^\circ = \text{int}(K)$  then:  $h_K = 0$  iff  $x = 0$ .

If  $0 \in K^\circ$ , we define the dual  $K^*$  of  $K$  by  $K^* = \{x \in \mathbf{R}^n; h_K(x) \leq 1\}$ ; then  $K^* \in K(\mathbf{R}^n)$ ,  $0 \in K^*$ ,  $(K^*)^* = K$  and  $K = \{x \in \mathbf{R}^n; h_{K^*}(x) \leq 1\}$ . If  $K_1, \dots, K_m \in K(\mathbf{R}^n)$  and  $\lambda_1, \dots, \lambda_m \in \mathbf{R}_+$  then  $K = \lambda_1K_1 + \dots + \lambda_mK_m$  is called the Minkowski sum of  $K_1, \dots, K_m$  with coefficients  $\lambda_1, \dots, \lambda_m$  and for  $K$  we have  $h_K = \lambda_1h_{K_1} + \dots + \lambda_mh_{K_m}$ .

$K(\mathbf{R}^n)$  is a complete metric space with Hausdorff metric  $d$  defined by

$$d(K_1, K_2) = \max_{x \in S_n} |h_{K_1}(x) - h_{K_2}(x)|$$

where  $S_n = \{x \in \mathbf{R}^n; \|x\| = 1\}$ ,  $\|x\|^2 = (x|x)$ . Let  $D_n = \{x \in \mathbf{R}^n; \|x\| \leq 1\}$  be the Euclidean disc in  $\mathbf{R}^n$ . The classical Steiner formula says that

$$V(K + rD_n) = \sum_{m=0}^n \binom{n}{m} W_m(K) r^m, \quad r \geq 0, K \in K(\mathbf{R}^n)$$

where  $V(K)$  is the volume of  $K$ , i.e. the Lebesgue measure of  $K$ .

The functions  $K \rightarrow W_m(K)$ ,  $m = 0, \dots, n$ , defined by the Steiner formula, are called the Steiner functionals. They are continuous, invariant under isometries, monotonic, additive and they have many other nice properties. For the proof of these assertions see [1].

## 2. Some auxiliaries

2.1. *Definition.* Let  $H(n)$  be the set of all continuous functions  $\varphi : \mathbf{R}^n \rightarrow [0, \infty)$  such that

- 1)  $\varphi(tx) = t\varphi(x)$ ,  $t \geq 0$ ,  $x \in \mathbf{R}^n$ .
- 2)  $\varphi$  is differentiable a.e. and  $\varphi'(x) \neq 0$  a.e.

For  $\varphi \in H(n)$  we introduce the sets

$$S_\varphi = \{x \in \mathbf{R}^n; \varphi(x) = 1\}, \quad D_\varphi = \{x \in \mathbf{R}^n; \varphi(x) \leq 1\}.$$

By differentiating 1) with respect to  $t$  and  $x$  we obtain

- a)  $\varphi'(x)x = \varphi(x)$ , a.e.,  $x \in \mathbf{R}^n$ ;
- b)  $\varphi'(tx) = \varphi'(x)$ , a.e.,  $t \geq 0$ ,  $x \in \mathbf{R}^n$ .

Here we need some integral formulae involving  $\varphi \in H(n)$ . We collect all such formulae in the following theorem. For the proof of the theorem see [2].

2.2. **THEOREM.** *Let  $\varphi, \psi \in H(n)$ . Then*

$$1) \int_{\mathbf{R}^n} f(x) dx = \int_0^\infty \int_{S_\varphi} f(tx) t^{n-1} \frac{dt dh(x)}{\|\varphi'(x)\|}$$

where  $h$  is the Hausdorff  $(n-1)$ -measure in  $\mathbf{R}^n$  and  $f \in L_1(\mathbf{R}^n)$ ;

$$2) \int_{\mathbf{R}^n} f(x) \|\varphi'(x)\| dx = \int_0^\infty \int_{S_\varphi} f(tx) t^{n-1} dt dh(x);$$

$$3) \int_{S_\varphi} \frac{dh(x)}{\|\varphi'(x)\|} = nV(D_\varphi);$$

$$4) \int_{D_\varphi} \|\varphi'(x)\| dx = \frac{1}{n} h(S_\varphi);$$

$$5) \int_{S_\psi} f dh = \int_{S_\varphi} f\left(\frac{x}{\psi(x)}\right) \frac{1}{\psi(x)^n} \cdot \frac{\|\psi'(x)\|}{\|\varphi'(x)\|} dh(x)$$

where  $f \in L_1(S_\psi, h)$ ;

$$6) \int_{D_\psi} f(x) dx = \int_{D_\varphi} f\left(\frac{\varphi(x)}{\psi(x)}x\right) \frac{\varphi(x)^n}{\psi(x)^n} dx$$

where  $f \in L_1(D_\psi)$ .

2.3. *Note.* Let  $K \in K(\mathbf{R}^n)$  and  $0 \in K^\circ$ . Then by Rademacher's theorem (see [3, Theorem 3.1.6, p. 234]) we have  $h_K \in H(n)$ ,  $h_{K^*} \in H(n)$  and  $K = D_{h_K}$  and  $K^* = D_{h_{K^*}}$  and we can apply the theorem above to  $K$  and  $K^*$ .

### 3. The dual Steiner formula

3.1. THEOREM. Let  $K \in K(\mathbf{R}^n)$  and  $0 \in K^\circ$ . Then

- 1)  $V(K^*) = \frac{1}{n} \int_{S_n} \frac{dh(x)}{h_K(x)^n}$ ;
- 2)  $V((K + rD_n)^*) = \frac{1}{n} \int_{S_n} \frac{dh(x)}{(h_K(x) + r)^n}$ ,  $r \geq 0$ ;
- 3)  $W_1(K^*) = \frac{1}{n} \int_{S_n} \frac{\|h'_K(x)\|}{h_K(x)^n} dh(x)$ ;
- 4)  $W_1((K + rD_n)^*) = \frac{1}{n} \int_{S_n} \frac{\|h'_K(x) + rx\|}{(h_K(x) + r)^n} dh(x)$ ,  $r \geq 0$ .

*Proof:* 1) We apply theorem 2.2. 6) to  $\psi = h_K$ ,  $\varphi(x) = \|x\|$  and  $f = 1$ . Hence

$$V(D_\psi) = V(D_{h_K}) = V(K^*) = \int_{D_n} \frac{\|x\|^n dx}{h_K(x)^n}$$

and now by theorem 2.2. 1) we have

$$V(K^*) = \int_0^1 \int_{S_n} \frac{t^{n-1}}{h_K(x)^n} dt dh(x) = \frac{1}{n} \int_{S_n} \frac{dh(x)}{h_K(x)^n}.$$

2) Because of

$$h_{K+rD_n}(x) = h_K(x) + r\|x\|$$

the formula follows from 1).

3) We apply theorem 2.2. 5) to  $\psi = h_K$  and  $f = 1$ . Hence

$$h(S_\psi) = nW_1(D_\psi) = nW_1(K^*) = \int_{S_n} \frac{\|h'_K(x)\|}{h_K(x)^n} dh(x).$$

4) Follows from 3) as above.

3.2. COROLLARY. Let  $K \in K(\mathbf{R}^n)$  and  $0 \in K^\circ$ . Then

- 1)  $V((K + rD_n)^*) \leq \frac{\alpha(n)}{(\beta + r)^n}$ ,  $r \geq 0$ ,

where  $\alpha(n) = V(D_n)$ ,  $\beta = \min_{x \in S_n} h_K(x)$ .

- 2)  $V(K) \leq \frac{1}{n} \int_{S_n} h_K(x)^n dh(x)$ .

$$3) W_1((K + rD_n)^*) \leq \frac{\alpha(n)}{(\beta + r)^n} (\gamma^2 + r^2 + 2r\|h_K\|)^{1/2}, \quad r \geq 0,$$

where  $\beta = \min_{x \in S_n} h_K(x)$ ,  $\|h_K\| = \max_{x \in S_n} h_K(x)$ ,  $\gamma = \max_{x \in S_n} \|h'_K(x)\|$ .

*Proof:* 1) follows from theorem 3.1 2).

2) Because of  $h_K(x)h_{K^*}(y) \geq (x|y)$  we have

$$1 \leq h_K(x)h_{K^*}(x), \quad x \in S_n$$

and the inequality follows from theorem 3.1. 1).

$$3) \|h'_K(x) + rx\|^2 = \|h'_K(x)\|^2 + r^2 + 2rh_K(x). \text{ Apply now theorem 3.1 4).}$$

3.3. COROLLARY (Dual Steiner formula). *Let  $K \in K(\mathbf{R}^n)$  and  $0 \in K^\circ$ . Then*

$$V((K + rD_n)^*) = \sum_{m=0}^{\infty} \frac{1}{n} \binom{n+m-1}{m} (-1)^m r^m \int_{S_n} \frac{dh(x)}{h_K(x)^{n+m}}$$

and the series converges for  $0 \leq r < \min_{x \in S_n} h_K(x)$ .

*Proof.* We apply theorem 3.1. 2) by expanding the integrand in a power series by using the formula

$$(1 - t)^{-n} = \sum_{m=0}^{\infty} \binom{n+m-1}{m} t^m, \quad |t| < 1.$$

Hence, we have

$$\frac{1}{(h_K(x) + r)^n} = \sum_{m=0}^{\infty} \binom{n+m-1}{m} (-1)^m r^m \frac{1}{h_K(x)^{n+m}}, \quad 0 \leq r < \min_{x \in S_n} h_K(x)$$

and the formula follows.

3.4. COROLLARY. *Let  $K \in K(\mathbf{R}^n)$  and  $0 \in K^\circ$ . Then*

$$V((D_n + rK)^*) = \sum_{m=0}^{\infty} \frac{1}{n} \binom{n+m-1}{m} (-1)^m r^m \int_{S_n} h_K^m dh, \quad 0 \leq r < 1/\|h_K\|.$$

*Proof.* Put  $1/r$  in place of  $r$  in 3.1 2). Then

$$V((D_n + rK)^*) = \frac{1}{n} \int_{S_n} \frac{dh(x)}{(1 + rh_K(x))^n}, \quad r \geq 0.$$

The formula now follows as above.

3.5. THEOREM. *Let  $K \in K(\mathbf{R}^n)$ ,  $0 \in K^\circ$  and*

$$z(K) = \int_K x \, dx.$$

Then we have

$$1) z((K + rD_n)^*) = \frac{1}{n+1} \int_{S_n} \frac{x dh(x)}{(h_K(x) + r)^{n+1}}, \quad r \geq 0;$$

$$2) z((K + rD_n)^*) = \sum_{m=0}^{\infty} \frac{1}{n+1} \binom{n+m}{m} (-1)^m r^m \int_{S_n} \frac{x dh(x)}{h_K(x)^{n+m+1}}$$

and the series converges for  $0 \leq r < \min_{x \in S_n} h_K(x)$ .

*Proof.* Let us apply theorem 2.2. 6) to  $\psi = h_K$ ,  $\varphi(x) = \|x\|$  and  $f(x) = x$ . Then we have

$$\begin{aligned} z(K^*) &= \int_{K^*} x dx = \int_{D_\psi} x dx = \int_{D_n} \frac{\|x\|^{n+1}}{h_K(x)^{n+1}} x dx \\ &= \int_0^1 \int_{S_n} \frac{t^n x}{h_K(x)^{n+1}} dt dh(x) = \frac{1}{n+1} \int_{S_n} \frac{x dh(x)}{h_K(x)^{n+1}}, \end{aligned}$$

Now 1) and 2) follow by the same argument as in 3.1, 3.2 and 3.3.

3.6. Using Steiner's vector formula (see [1]):

$$z(K + rD_n) = \sum_{m=0}^n \binom{n}{m} q_m(K) r^m, \quad r \geq 0$$

to introduce Steiner vectors  $q_m(K)$ ,  $m = 0, \dots, n$ , one can apply the theory above to study the Steiner vectors of  $K$ .

The formulae in theorem 3.5 we call dual Steiner vector formulae. Steiner functionals can be used to evaluate some complicated integrals in  $\mathbf{R}^n$ . Let us give an example.

3.7. PROPOSITION. Let  $K \in K(\mathbf{R}^n)$ . Then

$$\int_{\mathbf{R}^n} f(d(x, K)) dx = f(0)V(K) + \sum_{s=0}^{n-1} n \binom{n-1}{s} W_{s+1}(K) \int_0^\infty f(t) t^s dt$$

where  $d(x, K) = \min_{y \in K} \|x - y\|$ ,  $f(d(\cdot, K)) \in L_1(\mathbf{R}^n)$ .

*Proof.* Let us use the distribution function of  $d(\cdot, K)$  i.e.

$$V(\{x \in \mathbf{R}^n; d(x, K) \leq r\}) = V(K + rD_n)$$

Hence, we have

$$\begin{aligned} \int_{\mathbf{R}^n} f(d(x, K)) dx &= f(0)V(K) + \int_0^\infty f(t) dV(K + tD_n) \\ &= f(0)V(K) + \int_0^\infty f(t) n W_1(K + tD_n) dt \end{aligned}$$

and the formula follows.

3.8. COROLLARY (Wills formula, 1973).

$$\int_{\mathbf{R}^n} \exp(-\pi d(x, K)^2) dx = \sum_{m=0}^n \binom{n}{m} \frac{1}{\alpha(m)} W_m(K), \quad K \in K(\mathbf{R}^n).$$

*Proof.* Put  $f(t) = \exp(-\pi t^2)$  in 3.7.

3.9. COROLLARY.

$$\int_{K+rD_n} d(x, K)^\alpha dx = \sum_{s=0}^{n-1} n \binom{n-1}{s} W_{s+1}(K) \frac{r^{\alpha+s+1}}{\alpha+s+1}, \quad \alpha > 0, K \in K(\mathbf{R}^n).$$

*Proof.* Put  $f(t) = t^\alpha \chi_{[0,r]}(t)$  in 3.7.

The technique above can be applied to prove many inequalities. Let us give some examples.

3.10. THEOREM. Let  $K_1, K_2 \in K(\mathbf{R}^n)$ ,  $0 \in K_1^\circ$ ,  $0 \in K_2^\circ$ , and  $\lambda \in [0, 1]$ . Then

$$V((\lambda K_1 + (1 - \lambda)K_2)^*) \leq V(K_1^*)^\lambda V(K_2^*)^{1-\lambda}.$$

*Proof.* By theorem 2.2. 1) we have

$$n!V(K_1^*) = \int_{\mathbf{R}^n} \exp(-h_{K_1}(x)) dx$$

and also

$$n!V((\lambda K_1 + (1 - \lambda)K_2)^*) = \int_{\mathbf{R}^n} \exp(-\lambda h_{K_1}(x) - (1 - \lambda)h_{K_2}(x)) dx.$$

If  $\lambda = 0$  or  $\lambda = 1$  the inequality is trivial. Hence, we can assume that  $0 < \lambda < 1$ . Let us apply Hölder's inequality for

$$p = \frac{1}{\lambda}, \quad q = \frac{1}{1-\lambda}, \quad \frac{1}{p} + \frac{1}{q} = \lambda + 1 - \lambda = 1.$$

Then we have

$$\begin{aligned} & n!V((\lambda K_1 + (1 - \lambda)K_2)^*) \\ & \leq \left( \int_{\mathbf{R}^n} \exp(-h_{K_1}(x)) dx \right)^\lambda \cdot \left( \int_{\mathbf{R}^n} \exp(-h_{K_2}(x)) dx \right)^{1-\lambda} \\ & = (n!V(K_1^*))^\lambda \cdot (n!V(K_2^*))^{1-\lambda} = n!V(K_1^*)^\lambda V(K_2^*)^{1-\lambda}. \end{aligned}$$

3.11 COROLLARY.

1)  $V((K_1 + K_2)^*) \leq 2^{-n} V(K_1^*)^{1/2} V(K_2^*)^{1/2}$ ,  $0 \in K_1^\circ$ ,  $0 \in K_2^\circ$ .

$$2) V((K + rD_n)^*)^2 \leq \frac{\alpha(n)}{4^n \cdot r^n} V(K^*), \quad 0 \in K^\circ, \quad r > 0.$$

*Proof.* 1) Put  $\lambda = 1/2$  in 3.10.

2) follows from 1).

3.12. THEOREM. Let  $K_1, K_2 \in K(\mathbf{R}^n)$ ,  $0 \in K_1^\circ$ ,  $0 \in K_2^\circ$ . Then

$$W_1((K_1 + K_2)^*) \leq W_1(K_1^*) + W_1(K_2^*).$$

*Proof.* By theorem 2.2. 2) applied to  $f(t) = e^{-t}$ ,  $\varphi = h_{K_1}$  we have

$$n!W_1(K_1^*) = \int_{\mathbf{R}^n} \exp(-h_{K_1}(x)) \|h'_{K_1}(x)\| dx.$$

Hence, we have

$$\begin{aligned} n!W_1((K_1 + K_2)^*) &= \int_{\mathbf{R}^n} \exp(-h_{K_1}(x) - h_{K_2}(x)) \|h'_{K_1}(x) + h'_{K_2}(x)\| dx \\ &\leq \int_{\mathbf{R}^n} \exp(-h_{K_1}(x) - h_{K_2}(x)) \|h'_{K_1}(x)\| dx \\ &\quad + \int_{\mathbf{R}^n} \exp(-h_{K_1}(x) - h_{K_2}(x)) \|h'_{K_2}(x)\| dx \\ &\leq \int_{\mathbf{R}^n} \exp(-h_{K_1}(x)) \|h'_{K_1}(x)\| dx + \int_{\mathbf{R}^n} \exp(-h_{K_2}(x)) \|h'_{K_2}(x)\| dx \\ &= n!W_1(K_1^*) + n!W_1(K_2^*). \end{aligned}$$

#### REFERENCES

- [1] K. Leichtweiss, *Konvexe Mengen*, Nauka, Moscow, 1985 (in Russian).
- [2] L. Dedić, *On Haar measure on  $SL(n, R)$* , Publ. Inst. Math. Belgrade **47 (61)** (1990), 56–60.
- [3] H. Federer, *Geometric measure theory*, Nauka, Moscow, 1987 (in Russian).

Sveučilište u Splitu  
Teslina 12/3  
58000 Split

(Received 06 02 1991)