

GENERALIZED HERMITE POLYNOMIALS

Gospava B. Đorđević

Abstract. We consider a new generalization of the classical Hermite polynomials and prove the basic characteristics of such polynomials $h_{n,m}^\lambda(x)$ (the generating function, an explicit representation, some recurrence relations, and the corresponding differential equation). For $m = 2$, the polynomial $h_{n,m}^\lambda(x)$ reduces to $H_n(x, \lambda)/n!$, where $H_n(x, \lambda)$ is the Hermite polynomial with a parameter. For $\lambda = 1$, $h_{n,2}^1(x) = H_n(x)/n!$, where $H_n(x)$ is the classical Hermite polynomial. Taking $\lambda = 1$ and $n = mN + q$, where $N = [n/m]$ and $0 \leq q \leq m-1$, we introduce the polynomials $P_N^{(m,q)}(t)$ by $h_{n,m}^1(x) = (2x)^q P_N^{(m,q)}((2x)^m)$, and prove that they satisfy an $(m+1)$ -term linear recurrence relation.

1. Polynomials $h_{n,m}^\lambda(x)$. At the beginning, we define polynomials $h_{n,m}^\lambda(x)$ in the following manner.

Definition 1.1. The polynomials $h_{n,m}^\lambda(x)$, $\lambda \in R^+$, $n, m \in N$, are defined by the generating function

$$F(x, t) = e^{\lambda(2xt-t^m)} = \sum_{n=0}^{\infty} h_{n,m}^\lambda(x) t^n. \quad (1.1)$$

From above we get

$$F(x, t) = e^{\lambda(2xt-t^m)} = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{[n/m]} (-1)^k \frac{\lambda^n (2x)^{n-mk}}{\lambda^{(m-1)k} k! (n-mk)!} \right) t^n.$$

Thus, we obtain the following explicit representation

$$h_{n,m}^\lambda(x) = \lambda^n \sum_{k=0}^{[n/m]} (-1)^k \frac{(2x)^{n-mk}}{\lambda^{(m-1)k} k! (n-mk)!}. \quad (1.2)$$

Starting from (1.1) we can prove the following theorem.

THEOREM 1.1. *The polynomials $h_{n,m}^\lambda(x)$ satisfy the three-term recurrence relation*

$$nh_{n,m}^\lambda(x) = \lambda(2xh_{n-1,m}^\lambda(x) - mh_{n-m,m}^\lambda(x)), \quad n \geq m \quad (1.3)$$

with initial values : $h_{n,m}^\lambda(x) = (2\lambda x)^n/n!$, $0 \leq n \leq m-1$.

Now, we prove the following theorem :

THEOREM 1.2. *The polynomials $h_{n,m}^\lambda(x)$ satisfy the following relations:*

$$2nh_{n,m}^\lambda(x) = (2x)Dh_{n,m}^\lambda(x) - mDh_{n+1-m,m}^\lambda(x); \quad (1.4)$$

$$D^k h_{n,m}^\lambda(x) = (2\lambda)^k h_{n-k,m}^\lambda(x); \quad (1.5)$$

$$\frac{(2x)^n}{n!} = \sum_{k=0}^{[n/m]} \frac{1}{k!} h_{n-mk,m}^\lambda(x) \quad (m \geq 2); \quad (1.6)$$

$$u^n h_{n,m}^l\left(\frac{x}{u}\right) = \sum_{k=0}^{[n/m]} \frac{(1-u^m)^k}{k!} h_{n-mk,m}^l(x); \quad (1.7)$$

$$h_{n,m}^l(x+y) = \sum_{k=0}^n \frac{(2y)^k}{k!} h_{n-k,m}^l(x), \quad (1.8)$$

where $D = d/dx$ is the differentiation operator.

Proof. Differentiating (1.1) with respect to x and t we find the next equalities:

$$(i) \quad \partial F(x, t)/\partial x = 2\lambda t e^{\lambda(2xt-t^m)}, \quad (ii) \quad \partial F(x, t)/\partial t = \lambda(2x - mt^{m-1})e^{\lambda(2xt-t^m)}.$$

Combining these equalities we obtain (1.4).

Differentiating the polynomials $h_{n,m}^\lambda(x)$ given by (1.2) k -times we get (1.5).

The generating function (1.1) for $\lambda = 1$ reduces to

$$e^{2xt-t^m} = \sum_{n=0}^{\infty} h_{n,m}^l(x)t^n, \quad \text{i.e. to } e^{2xt} = e^{t^m} \sum_{n=0}^{\infty} h_{n,m}^\lambda(x)t^n.$$

Developing both sides of the last equality in t , we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(2x)^n}{n!} t^n &= \left(\sum_{n=0}^{\infty} \frac{t^{mn}}{n!} \right) \left(\sum_{n=0}^{\infty} h_{n,m}^l(x)t^n \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^{[n/m]} \frac{1}{k!} h_{n-mk,m}^l(x) \right) t^n. \end{aligned}$$

Now, comparing coefficients of t^n in the last equality we get (1.6).

Starting from $e^{2xt-t^m u^m} = e^{2xt-t^m} \circ e^{t^m - u^m t^m}$, we get (1.7).

Finally, from the equality $e^{2(x+y)t-t^m} = e^{2xt-t^m} \circ e^{2yt}$, we get

$$\sum_{n=0}^{\infty} t^n \sum_{k=0}^{[n/m]} (-1)^k \frac{(2x+2y)^{n-mk}}{k!(n-mk)!} =$$

$$\left(\sum_{n=0}^{\infty} t^n \sum_{k=0}^n (-1)^k \frac{(2x)^{n-k} t^{(m-1)k}}{k!(n-k)!} \right) \left(\sum_{n=0}^{\infty} \frac{t^n (2y)^n}{n!} \right),$$

wherefrom, after some calculations, we obtain (1.8).

COROLLARY 1.1. *For $m = 2$ and $\lambda = 1$ the equalities (1.4)–(1.8) reduce to the corresponding relations for the classical Hermite polynomials.*

At the end of this section we prove that the polynomials $h_{n,m}^\lambda(x)$ have an interesting property.

THEOREM 1.3. *The polynomial $h_{n,m}^\lambda(x)$ is a particular solution of linear homogeneous equation of m -th order given by*

$$L_n(y) = y^{(m)} - 2^m m^{-1} \lambda^{m-1} (xy - ny) = 0. \quad (1.9)$$

Proof. Using (1.5) and the recurrence relation (1.3) we get

$$\begin{aligned} L_n[h_{n,m}^\lambda(x)] &= (2\lambda)^m h_{n-m,m}^\lambda(x) - 2^m m^{-1} \lambda^{m-1} x (2\lambda) h_{n-1,m}^\lambda(x) \\ &\quad + 2^m m^{-1} \lambda^{m-1} n h_{n,m}^\lambda(x) \\ &= 2^m m^{-1} \lambda^{m-1} (n h_{n,m}^\lambda(x) - 2\lambda x h_{n-1,m}^\lambda(x) + m \lambda h_{n-m,m}^\lambda(x)) = 0. \end{aligned}$$

2. Polynomials $P_N^{m,q}(t)$. In this section we introduce a class of polynomials $\{P_N^{m,q}(t)\}_{N=0}^{\infty}$. Let us suppose that $n = mN + q$, where $N = [n/m]$ and $0 \leq q \leq m - 1$. Starting from (1.2) and taking $\lambda = 1$, we have

$$\begin{aligned} h_{n,m}^l(x) &= (2x)^q \sum_{k=0}^N (-1)^k \frac{(2x)^{mN-mk}}{k!(mN+q-mk)!} \\ &= (2x)^q \sum_{k=0}^N (-1)^k \frac{((2x)^m)^{N-k}}{k!(q+m(N-k))!} \\ &= (2x)^q P_N^{(m,q)}(t), \text{ where } t = (2x)^m. \end{aligned}$$

In this way we come to

$$P_N^{m,q}(t) = \sum_{k=0}^N (-1)^k \frac{t^{N-k}}{k!(q+m(k+1))!}. \quad (2.1)$$

In fact, the polynomials $P_N^{m,q}(t)$ depend on two parameters: $m \in N$ and $q \in \{0, 1, \dots, m-1\}$.

Using (1.3) for $\lambda = 1$, i.e., $nh_{n,m}^l(x) = 2xh_{n-1,m}^l(x) - mh_{n-m,m}^l(x)$, where $n \geq m \geq 1$, we can prove the following theorem:

THEOREM 2.1 *The polynomials $P_N^{m,q}(t)$ satisfy the next recurrence relations:*

$$\begin{aligned} (mN+q)P_N^{(m,q)}(t) &= P_N^{(m,q-1)}(t) - mP_{N-1}^{(m,q)}(t), \quad \text{for } 1 \leq q \leq m-1, \\ mNP_N^{(m,0)}(t) &= tP_{N-1}^{(m,m-1)}(t) - mP_{N-1}^{(m,0)}(t), \quad \text{for } q = 0. \end{aligned}$$

It is interesting to find a recurrence relation for the polynomials $P_N^{(m,q)}(t)$ where the parameters m and q are fixed.

Using the same method as in [3] we can prove the following result:

THEOREM 2.2. *The polynomials $P_N^{(m,q)}(t)$ satisfy an $(m+1)$ -term recurrence relation of the form*

$$\sum_{i=0}^m A_{i,N,q} P_{N+1-i}^{(m,q)}(t) = B_{N,q} t P_N^{(m,q)}(t),$$

where $B_{N,q}$ and $A_{i,N,q}$ ($i = 0, 1, \dots, m$) are constants depending only on N , m and q .

According to the explicit representation of polynomials $P_N^{(m,q)}(t)$ given by (2.1), we get:

PROPOSITION 2.3 *The polynomials $P_N^{(m,q)}(t)$ have no negative real zeros.*

REFERENCES

- [1] G. B. Đorđević, *Contributions to the theory of polynomials defined by recurrence relations*, Ph. D. Thesis, Univ. of Niš, 1989. (in Serbian).
- [2] G. V. Milovanović and G. Đorđević, *On some properties of Humbert's polynomials*, Fibonacci Quart. **25**(1987), 356-360.
- [3] G. V. Milovanović and G. Đorđević, *On some properties of Humbert's polynomials*, II, Facta Univ. Ser. Math. Inform. **6**(1991), 23-30.
- [4] H. Srivastava, *A note on a generating function for the generalized Hermite polynomials*, Nederl. Akad. Wetensch. Proc. Ser. A **79**(1976), 457-461.
- [5] A. K. Varma, *A new characterization of Hermite polynomials*, Acta Math. Hungar. **49**(1-2), (1987), 169-172.

Tehnološki fakultet
Univerziteta u Nišu
16000 Leskovac
Jugoslavija

(Received 23 08 1991)
(Revised 23 03 1993)