

## LOGICS WITH TWO TYPES OF INTEGRAL OPERATORS

Radosav S. Đorđević

**Abstract.** We prove completeness theorems for absolutely continuous and singular biprobability models of a logic with integrals. Also in both cases, we prove the finite compactness theorem for a set of sentences of the form  $\tau \in [r, s]$ .

We assume throughout the paper that  $\mathcal{A}$  is a countable admissible set with  $\omega \in \mathcal{A}$ . In [2], Keisler introduced a logic  $L_{\mathcal{A}f}$  which has an integral operator which builds terms with bound variables. In our case two types of integral operators  $\int_1 \dots dx$  and  $\int_2 \dots dx$  are allowed.

A biprobability model for  $L_{\mathcal{A}f_1 f_2}$  logic is a model  $\mathfrak{A} = \langle A, R_i, c_j, \mu_1, \mu_2 \rangle_{i \in I, j \in J}$ , where  $\langle A, R_i, c_j \rangle$  is a first-order model without operations and  $\mu_1, \mu_2$  are probability measures on  $A$ . We shall see a difference in semantics for  $L_{\mathcal{A}f_1 f_2}^a$  and  $L_{\mathcal{A}f_1 f_2}^s$  by means of the following definition.

*Definition 1.* (a) An absolutely continuous biprobability model for  $L_{\mathcal{A}f_1 f_2}^a$  is a biprobability model  $\mathfrak{A}$  such that  $\mu_1$  is absolutely continuous with respect to  $\mu_2$ , i. e.  $\mu_1 \ll \mu_2$ .

(b) A singular biprobability model for  $L_{\mathcal{A}f_1 f_2}^s$  is a biprobability model  $\mathfrak{A}$  such that  $\mu_1$  is singular with respect to  $\mu_2$ , i. e.  $\mu_1 \perp \mu_2$ .  $\square$

In both cases, quantifiers are interpreted by

$$\left( \int_k \tau(x, \vec{a}) dx \right)^{\mathfrak{A}} = \int \tau(b, \vec{a})^{\mathfrak{A}} d\mu_k(b) \quad \text{for } k = 1, 2,$$

where  $\tau(x, \vec{y})$  is a term and  $\vec{a} \in A^n$ .

Diagonal products  $\mu_k^{(n)}$ , which are the corresponding restrictions of completions of  $\mu_k^n$ 's ( $k = 1, 2$ ) to  $\sigma$ -algebras generated by the measurable rectangles and the diagonal sets  $\{ \vec{x} \in A^n : x_i = x_j \}$ , can be replaced by sequences of probability measures on  $A^n$ 's which satisfy the Fubini theorem. That generalization of a probability structure is relevant for us.

---

*Mathematics Subject Classification* (1991): Primary 03C70

This research was supported by Government of Serbia grant number 0401A, through Matematički institut.

*Definition 2.* A graded biprobability model for  $L_{\mathcal{A}f_1f_2}$  is a model  $\mathfrak{A} = \langle A, R_i, c_j, \mu_n^1, \mu_n^2 \rangle_{i \in I, j \in J, n \geq 1}$  such that:

- (1) Each  $\mu_n^k$  is a countably additive probability measure on  $A^n$ .
- (2) Each  $n$ -ary relation  $R_i$  is  $\mu_n^k$ -measurable and the identity relation is  $\mu_2^k$ -measurable.
- (3)  $\mu_n^k \times \mu_m^k \subseteq \mu_{n+m}^k$ .
- (4) Each  $\mu_n^k$  is preserved under permutation of  $\{1, 2, \dots, n\}$ .
- (5)  $\langle \mu_n^k : n \in \mathbb{N} \rangle$  has the Fubini property: If  $B$  is  $\mu_{m+n}^k$ -measurable, then
  - (a) For each  $\vec{x} \in A^m$ , the section  $B_{\vec{x}} = \{\vec{y} : B(\vec{x}, \vec{y})\}$  is  $\mu_n^k$ -measurable.
  - (b) The function  $f(\vec{x}) = \mu_n^k(B_{\vec{x}})$  is  $\mu_m^k$ -measurable.
  - (c)  $\int f(\vec{x}) d\mu_m^k = \mu_{m+n}^k(B)$ .  $\square$

*Definition 3.* (a) A graded biprobability model for  $L_{\mathcal{A}f_1f_2}^a$  is a graded biprobability model  $\mathfrak{A}$  such that  $\mu_n^1 \ll \mu_n^2$  for each  $n \in \mathbb{N}$ .

(b) A graded biprobability model for  $L_{\mathcal{A}f_1f_2}^s$  is a graded biprobability model  $\mathfrak{A}$  such that  $\mu_n^1 \perp \mu_n^2$  for each  $n \in \mathbb{N}$ .  $\square$

**1. The logic  $L_{\mathcal{A}f_1f_2}^a$ .** Axioms and rules of inference for  $L_{\mathcal{A}f_1f_2}^a$  are those for  $L_{\mathcal{A}f}$ , as listed in [3] with both  $f_1$  and  $f_2$  playing the role of  $f$ , together with the following axioms:

(A<sub>1</sub>) *Axioms of continuity of integral operators:* ( $i, j = 1, 2$ )

$$(a) \bigwedge_n \bigvee_m \bigvee_k \int_i \mathbb{F}_k \left( \int_j \tau(\vec{x}, \vec{y}) d\vec{x} \right) d\vec{y} < \frac{1}{n},$$

$$\text{where } F_k(s) = \begin{cases} 1, & \text{if } r - 1/m + 1/k \leq s \leq r - 2/k \\ 0, & \text{if } s \leq r - 1/m \text{ or } s \geq r - 1/k \\ \text{linear,} & \text{for other cases} \end{cases}$$

is a continuous real function such that  $F_k \upharpoonright \mathbb{Q} \in \mathcal{A}$ .

$$(b) \bigwedge_n \bigvee_m \bigvee_k \int_i \mathbb{G}_k \left( \int_j \tau(\vec{x}, \vec{y}) d\vec{x} \right) d\vec{y} < \frac{1}{n},$$

$$\text{where } G_k(s) = \begin{cases} 1, & \text{if } r + 2/k \leq s \leq r + 1/m - 1/k \\ 0, & \text{if } s \leq r + 1/k \text{ or } s \geq r + 1/m \\ \text{linear,} & \text{for other cases.} \end{cases}$$

(A<sub>2</sub>) *Axiom of absolute continuity:*

$$\bigwedge_{\varepsilon \in \mathbb{Q}_+} \bigvee_{\delta \in \mathbb{Q}_+} \bigwedge_n \bigwedge_{\tau \in T_n} (|\int_2 \tau(\vec{x}) d\vec{x}| < \delta \implies |\int_1 \tau(\vec{x}) d\vec{x}| < \varepsilon),$$

where  $T = \bigcup_n T_n$ ,  $T_n$  is a set of terms with  $n$  free variables and  $T, T_n \in \mathcal{A}$ .

$$(A_3) \int_1 \left( \int_2 \tau dy \right) dx = \int_2 \left( \int_1 \tau dx \right) dy.$$

Now we introduce two sorts of auxiliary models.

*Definition 4.* (a) A weak model for  $L_{f_1f_2}^a$  is a model  $\langle \mathfrak{A}, I_1, I_2 \rangle$  where  $\mathfrak{A}$  is a first-order model and  $I_k$  is what may be called an  $\mathcal{A}$ -Daniell integral on  $A$ , that is,  $I_k$  is a positive linear real function on the set of terms with at most one free

variable  $x$  and parameters from  $A$ , i. e.

$$\begin{aligned} I_k(r) &= r, & k &= 1, 2 \\ I_k(r \cdot \sigma + s \cdot \tau) &= r \cdot I_k(\sigma) + s \cdot I_k(\tau), \\ \text{if } \tau(b, \vec{a})^{\mathfrak{A}} &\geq 0 \text{ for all } b \in A, \text{ then } I_k(\tau(x, \vec{a})) &\geq 0. \end{aligned}$$

(b) A middle model for  $L_{\mathcal{A} f_1 f_2}^{\alpha}$  is a weak model  $\mathfrak{A}$  such that for each  $\varepsilon > 0$  there is  $\delta > 0$  such that for each term  $\tau(x, \vec{y})$  and  $\vec{a} \in A^n$ , if  $|I_2(\tau(x, \vec{a}))| < \delta$  then  $|I_1(\tau(x, \vec{a}))| < \varepsilon$ .  $\square$

In both cases, for  $\tau$  a term, define  $\tau^{\mathfrak{A}}$  inductively as for biprobability models, except that at the integral step, we define

$$(\int_k \tau(x, \vec{a}) dx)^{\mathfrak{A}} = I_k(\tau(x, \vec{a})) .$$

LEMMA 1. (Middle Completeness Theorem for  $L_{\mathcal{A} f_1 f_2}^{\alpha}$ ) *Let  $T$  be a set of sentences of  $L_{\mathcal{A} f_1 f_2}^{\alpha}$  such that  $T$  is  $\Sigma_1$ -definable over  $\mathcal{A}$ . Then  $T$  is consistent with the axioms of this logic iff it has a middle model in which each theorem of  $L_{\mathcal{A} f_1 f_2}^{\alpha}$  is true.*

*Proof.* The soundness is easy to prove because all the axioms represent known properties of integrals (the Generalized Radon-Nikodym Theorem and the Fubini Theorem prove that each function  $\tau(x, y)^{\mathfrak{A}}: A \times A \rightarrow \mathbb{R}$  is compatible with absolutely continuous measures  $\mu_1$  and  $\mu_2$ , i. e.

$$\iint \tau(x, y)^{\mathfrak{A}} d\mu_1(x) d\mu_2(y) = \iint \tau(x, y)^{\mathfrak{A}} d\mu_2(y) d\mu_1(x) .$$

A Henkin argument is used to construct a weak model  $\langle \mathfrak{A}, I_1, I_2 \rangle$  of  $T$  in which each theorem of  $L_{\mathcal{A} f_1 f_2}^{\alpha}$  is true. Let  $K = L \cup C$  be the language introduced in this construction, where  $C$  is a set of new constant symbols and  $C \in \mathcal{A}$ . We wish the axiom  $A_2$  to hold for all the terms and that is done by the following construction (see [9]).

Let  $K'$  be a language with four kinds of variables:  $X, Y, Z, \dots$  are variables for sets,  $x, y, z, \dots$  are variables for urelements,  $r, s, t, \dots$  are variables for reals from  $[0, 1] \cap \mathcal{A}$ , and  $U, V, W, \dots$  are variables for functions  $A^n \mapsto \mathbb{R}$ ,  $n \geq 0$ . Predicates are:  $E_n^s(\vec{x}, X)$  for sets,  $n \geq 1$ ;  $E_{n+1}^t(\vec{x}, r, U)$  for terms,  $n \geq 0$ ;  $I_k(U, r)$  for  $U: A^0 \rightarrow \mathbb{R}$  or  $U: A^1 \rightarrow \mathbb{R}$ ,  $k = 1, 2$ ; and  $\leq$  for reals. Function symbols are  $f, g, h, \dots$  for each continuous real function  $F: \mathbb{R}^n \rightarrow \mathbb{R}$  such that  $F \upharpoonright \mathbb{Q}^n \in \mathcal{A}$ . Constant symbols are:  $X_{\varphi}$  for each formula  $\varphi$ ;  $U_{\tau}$  for each term  $\tau$ ; and  $\bar{r}$  for each real number  $r \in [0, 1] \cap \mathcal{A}$ .

Let  $S$  be the following theory of  $K'_{\mathcal{A}}$ :

1. *Axioms of validity:*

- 1.1  $(\forall X) \bigwedge_{n < m} \neg(\exists \vec{x}, \vec{y})(E_m^s(\vec{x}, \vec{y}, X) \wedge E_n^s(\vec{x}, X)), \text{ where } \{\vec{x}\} \cap \{\vec{y}\} = \emptyset;$
- 1.2  $(\forall U) \bigwedge_{n < m} \neg(\exists \vec{x}, \vec{y}, r, s)(E_{m+1}^t(\vec{x}, \vec{y}, r, U) \wedge E_{n+1}^t(\vec{x}, s, U));$
- 1.3  $(\forall U)(\forall \vec{x}, r, s)((E_{n+1}^t(\vec{x}, r, U) \wedge E_{n+1}^t(\vec{x}, s, U)) \implies r = s);$

## 2. Axioms of extensionality:

- 2.1  $(\forall \vec{x})(E_n^s(\vec{x}, X) \iff E_n^s(\vec{x}, Y)) \iff X = Y;$   
 2.2  $(\forall \vec{x}, r)(E_{n+1}^t(\vec{x}, r, U) \iff E_{n+1}^t(\vec{x}, r, V)) \iff U = V;$

## 3. Axioms of terms:

- 3.1  $(\forall \vec{x})(E_{n+1}^t(\vec{x}, 0, U_\tau) \vee E_{n+1}^t(\vec{x}, 1, U_\tau))$  if  $\tau$  is  $\mathbf{1}(R(\vec{x}))$ ;  
 3.2  $(\forall x, y)(E_{2+1}^t(x, y, 0, U_\tau) \vee E_{2+1}^t(x, y, 1, U_\tau))$  if  $\tau$  is  $\mathbf{1}(x = y)$ ;  
 3.3  $E_{0+1}^t(\vec{r}, U_\tau)$  if  $\tau$  is  $r$ ;  
 3.4  $(\forall \vec{x}, r)(E_{n+1}^t(\vec{x}, r, U_\tau) \iff (\exists \vec{s})(\bigwedge_{i=1}^k E_{n+1}^t(\vec{x}, s_i, U_{\tau_i}) \wedge$   
 $\wedge f(s_1, \dots, s_k) = r))$  if  $\tau$  is  $\mathbb{F}(\tau_1, \dots, \tau_k)$ ;  
 3.5  $(\forall \vec{x}, r)(E_{n+1}^t(\vec{x}, r, U_\tau) \iff (\exists V)((\forall y, s)(E_{1+1}^t(y, s, V) \iff$   
 $\iff E_{n+1+1}^t(\vec{x}, y, s, U_\sigma)) \wedge I_k(V, r))$  if  $\tau$  is  $\int_k \sigma(\vec{v}, v_0) dv_0$ ,  $k = 1, 2$ ;

## 4. Axioms of satisfaction:

- 4.1  $(\forall \vec{x})(E_n^s(\vec{x}, X_\varphi) \iff (\exists r \geq 0)E_{n+1}^t(\vec{x}, r, U_\tau))$  if  $\varphi$  is  $\tau \geq 0$ ;  
 4.2  $(\forall \vec{x})(E_n^s(\vec{x}, X_{\neg\varphi}) \iff \neg E_n^s(\vec{x}, X_\varphi))$ ;  
 4.3  $(\forall \vec{x})(E_n^s(\vec{x}, X_{\wedge\Phi}) \iff \bigwedge_{\varphi \in \Phi} E_n^s(\vec{x}, X_\varphi))$ ;

## 5. Axioms of integral operators:

- 5.1  $(\forall U)((\bigwedge_{n \geq 2} \neg(\exists \vec{x}, r)E_{n+1}^t(\vec{x}, r, U)) \iff (\exists_1 s)I_k(U, s))$ ,  $k = 1, 2$ ;  
 5.2  $(\forall r)I_k(U_r, r)$ ;  
 5.3  $(\forall U, V, r, s)I_k(r \cdot U + s \cdot V) = r \cdot I_k(U) + s \cdot I_k(V)$ , where  $I_k(U) = r$  iff  $I_k(U, r)$ ;  
 5.4  $(\forall U)((\forall x)(\exists r \geq 0)E_{1+1}^t(x, r, U) \implies (\exists s \geq 0)I_k(U, s))$ ;

## 6. Axiom of absolute continuity:

$$(\forall \varepsilon > 0)(\exists \delta > 0)(\forall U)(|I_2(U)| < \delta \implies |I_1(U)| < \varepsilon);$$

## 7. Axioms for an Archimedean field;

8. Transformations of axioms of  $K_{\mathcal{A} f_1 f_2}^a$ :

$$(\forall \vec{x})E_n^s(\vec{x}, X_\varphi), \text{ where } \varphi \text{ is an axiom of this logic};$$

9. Axioms of realizability of all sentences  $\varphi$  of  $T$  :

$$(\forall x_0)E_1^s(x_0, X_\varphi).$$

A weak model  $\langle \mathfrak{A}, I_1, I_2 \rangle$  for  $K_{\mathcal{A} f_1 f_2}^a$  can be transformed to a standard model  $\mathfrak{B}$  for  $K_{\mathcal{A}}'$  by taking:  $B_\varphi^{\mathfrak{B}} = \{ \vec{a} \in A^n : \mathfrak{A} \models \varphi[\vec{a}] \}$ ,  $U_\tau^{\mathfrak{B}}(\vec{a}) = \tau^{\mathfrak{A}}(\vec{a})$  for  $\vec{a} \in A^n$  and  $I_k^{\mathfrak{B}}(U_\tau^{\mathfrak{B}}) = I_k(\tau)$  for each term  $\tau$  with at most one free variable. By the Barwise Compactness Theorem (see [1]), it can be shown that  $S$  has a standard model  $\mathfrak{D}$ , because  $S$  is  $\Sigma$ -definable over  $\mathfrak{A}$  and  $A_2$  holds in  $\mathfrak{A}$ .  $\mathfrak{D}$  can be transformed to a middle model  $\mathfrak{C}$  of  $T$  by taking:

$$R^{\mathfrak{C}} = \{ \vec{x} \in D^n : E_n^s(\vec{x}, X_{\mathbf{1}(R(\vec{x})=1)}) \} \text{ and}$$

$$I_k^{\mathfrak{C}}(\tau(x, \vec{a})) = I_k^{\mathfrak{D}}(U_{\tau(x, \vec{a})}) \text{ for } \vec{a} \in D^n \text{ and } k = 1, 2.$$

This completes the proof of the Middle Completeness Theorem.  $\square$

In order to construct an absolutely continuous biprobability model, we need the following lemma.

LEMMA 2. (Loeb [4]) *In an  $\omega_1$ -saturated nonstandard universe, let  $M$  be an internal vector lattice of functions from an internal set  $A$  into  ${}^*\mathbb{R}$  (the set of hyperreal numbers), and let  $I$  be an internal positive linear functional on  $M$ , such that  $\mathbf{1} \in M$  and  $I(\mathbf{1}) = 1$ . Then there is a complete probability measure  $\mu$  on  $A$  such that for each finitely bounded  $\varphi \in M$ , the standard part of  $\varphi$  is integrable with respect to  $\mu$  and its integral is equal to the standard part of  $I(\varphi)$ .*

THEOREM 1. (Completeness Theorem for  $L_{\mathcal{A} f_1 f_2}^\alpha$ ) *Let  $T$  be a set of sentences of  $L_{\mathcal{A} f_1 f_2}^\alpha$  such that  $T$  is  $\Sigma_1$  on  $\mathcal{A}$  and consistent. Then there is an absolutely continuous biprobability model of  $T$ .*

*Proof.* Let  $\langle \mathfrak{A}, I_1, I_2 \rangle$  be a middle model of  $T$  in which each theorem of  $L_{\mathcal{A} f_1 f_2}^\alpha$  is true. The Daniell integral construction from Lemma 2 produces probability measures  $\mu_1, \mu_2$  on  ${}^*A$  such that for each  $*$ -term  $\tau(x)$ , the standard part of  ${}^*I_k(\tau)$  is the integral  $\int \text{st}(\tau(b)^{\mathfrak{A}}) d\mu_k(b)$  (we define measures  $\mu_n^k$  on  ${}^*A^n$  by using iterated integrals). The absolute continuity in the middle model  $\mathfrak{A}$  implies the absolute continuity for all measurable sets. Also, using axiom  $A_3$ , it can be shown that  $\mu_n^1 \ll \mu_n^2$  for each  $n \in \mathbb{N}$ . This graded biprobability model  $\hat{\mathfrak{A}} = \langle {}^*\mathfrak{A}, \mu_n^1, \mu_n^2 \rangle$  can be used to produce an absolutely continuous biprobability model of  $T$  (see [3]).  $\square$

We can look only for a part of  $L_{\mathcal{A} f_1 f_2}^\alpha$  which satisfies the finite compactness property, because this logic cannot satisfy the full compactness (for example, each finite subset of  $T = \{ \int_1 \mathbf{1}(R(x)) dx > 0 \} \cup \{ \int_1 \mathbf{1}(R(x)) dx \leq \frac{1}{n} : n \in \mathbb{N} \}$ , where  $R$  is a unary predicate, has a probability model, but not  $T$  itself).

THEOREM 2. *Let  $T$  be a set of sentences of  $L_{\mathcal{A} f_1 f_2}^\alpha$  of the form  $\tau \in [r, s]$ . If every finite subset of  $T$  has a graded biprobability model, then  $T$  has a graded biprobability model.*

*Proof.* Let us suppose that each finite subset  $\Psi \subseteq T$  has a model  $\mathfrak{A}_\Psi$ . By Lemma 1 we can suppose that  $\mathfrak{A}_\Psi$  is a middle model. Take an ultraproduct  ${}^*\mathfrak{A}$  such that, for each  $\varphi \in T$ , almost every  $\mathfrak{A}_\Psi$  satisfies  $\varphi$ . Then form a graded biprobability model  $\hat{\mathfrak{A}}$  from  ${}^*\mathfrak{A}$  by the Daniell integral construction (Lemma 2). It can be shown by induction that every sentence of  $L_{\mathcal{A} f_1 f_2}^\alpha$  of the form  $\tau \in [r, s]$  which is true in almost all  $\mathfrak{A}_\Psi$  holds in  $\hat{\mathfrak{A}}$ , too. The absolute continuity condition can be expressed in the middle model by the first-order sentence

$$(\forall \varepsilon > 0)(\exists \delta > 0)(\forall U)(|I_2(U)| < \delta \implies |I_1(U)| < \varepsilon) .$$

By Los's Theorem and Loeb construction the sentence

$$(\forall \varepsilon > 0)(\exists \delta > 0)(\forall X)(\mu_2(X) < \delta \implies \mu_1(X) < \varepsilon)$$

holds in  $\hat{\mathfrak{A}}$ .  $\square$

**2. The logic  $L_{\mathcal{A}f_1f_2}^s$ .** Axioms and rules of inference for the logic  $L_{\mathcal{A}f_1f_2}^s$  are those of  $L_{\mathcal{A}f}$  (with both  $\int_1$  and  $\int_2$  in place of  $\int$ , see [3]) together with the axioms of continuity  $A_1$  and

$$(A_4) \text{ Axiom of singularity:}$$

$$\bigvee_k \int_i \mathbb{H}_k \left( \int_1 \mathbf{1}(x=y) dy, \int_2 \mathbf{1}(x=y) dy \right) dx = 0, \quad i = 1, 2,$$

$$\text{where } H_k(s, t) = \begin{cases} 1, & \text{if } s \geq \frac{2}{k} \text{ and } t \geq \frac{2}{k} \\ 0, & \text{if } s \leq \frac{1}{k} \text{ or } t \leq \frac{1}{k} \\ \text{linear,} & \text{for other cases} \end{cases}.$$

**THEOREM 3.** (Completeness Theorem for  $L_{\mathcal{A}f_1f_2}^s$ ) *A theory  $T$  of  $L_{\mathcal{A}f_1f_2}^s$  is consistent iff  $T$  has a singular biprobability model.*

*Proof.* The proof of soundness is easy. Let  $\langle \mathfrak{A}, I_1, I_2 \rangle$  be a weak model of  $T$  in which each theorem of  $L_{\mathcal{A}f_1f_2}^s$  is true. Let  $\mathcal{F} = \{ B \subseteq A : \chi_B \in \text{dom}(I_1) = \text{dom}(I_2) \}$  be an algebra of subsets of  $A$ , where  $\chi_B(x) = \begin{cases} 1, & \text{if } x \in B \\ 0, & \text{if } x \notin B \end{cases}$ . Define finitely additive probability measures  $\nu_1, \nu_2$  on  $\mathcal{F}$  by  $\nu_k(B) = I_k(\chi_B)$ ,  $B \in \mathcal{F}$  and  $k = 1, 2$ .

Then, for  $a \in A$ , the singleton  $\{a\}$  belongs to  $\mathcal{F}$  because  $\chi_{\{a\}} = (x=a)^{\mathfrak{A}}$ , the set  $B = \{a \in A : \nu_1\{a\} > 0 \ \& \ \nu_2\{a\} > 0\}$  belongs to  $\mathcal{F}$  and  $\nu_k(B) = 0$   $k = 1, 2$  by  $A_4$ .

By construction from [7], the measures  $\nu_1, \nu_2$  can be extended so that  $\nu_1 \subseteq \bar{\nu}_1$ ,  $\nu_2 \subseteq \bar{\nu}_2$  and the measures  $\bar{\nu}_1, \bar{\nu}_2$  are singular. Then construct a middle biprobability model  $\langle \mathfrak{A}, \bar{I}_1, \bar{I}_2 \rangle$  of  $T$  by

$$\text{dom}(\bar{I}_k) = \text{dom}(I_k) \cup \{ \chi_C : C \in \bar{\mathcal{F}} \setminus \mathcal{F} \} \quad \text{and} \quad \bar{I}_k(\chi_C) = \bar{\nu}_k(C),$$

for each  $C$  from the extension  $\bar{\mathcal{F}}$  of  $\mathcal{F}$ .

By Loeb's construction (Lemma 2) and the construction of the biprobability model from a graded biprobability model (see [3]), the singularity of finitely additive measures in the middle model will be preserved in the biprobability model.  $\square$

Finally, we prove Finite Compactness Theorem for the singular case.

**THEOREM 4.** *Let  $T$  be a set of sentences of  $L_{\mathcal{A}f_1f_2}^s$  of the form  $\tau \in [r, s]$ . If every finite subset of  $T$  has a graded biprobability model, then  $T$  has a graded biprobability model.*

*Proof.* As in Theorem 2, our proof is based on the ultraproduct and Daniell integral construction. Now, we can suppose that  $\mathfrak{A}_\Psi$  is a weak model for each finite subset  $\Psi \subseteq T$ . Let  $\bar{\mathfrak{A}}_\Psi$  be a middle model as in Theorem 3. Take an ultraproducts  ${}^*\mathfrak{A} = \prod \bar{\mathfrak{A}}_\Psi$  such that, for each  $\varphi \in T$ , almost every  $\bar{\mathfrak{A}}_\Psi$  satisfies  $\varphi$ . The condition of singularity can be express in the middle model by the first-order sentence  $(\exists f)(\bar{I}_1(f) = 1 \wedge \bar{I}_2 = 0)$ . By Los's Theorem and Loeb's construction the sentence  $(\exists X)(\mu_1(X) = 1 \wedge \mu_2(X) = 0)$  holds in  $\bar{\mathfrak{A}}$ .  $\square$

## REFERENCES

- [1] J. Barwise, *Admissible Sets and Structures*, Springer-Verlag, Berlin, 1975.
- [2] H.J. Keisler, *Hyperfinite model theory*; in Logic Colloquium '76, North-Holland, Amsterdam, 1977, pp.5-110.
- [3] H.J. Keisler, *Probability quantifiers*; in Model Theoretic Languages (J. Barwise and S. Feferman, eds.), Springer-Verlag, Berlin, 1985, pp. 509-556.
- [4] P.A. Loeb, *A functional approach to nonstandard measure theory*, Contemporary Math. **26** (1984), 251-261.
- [5] J. Los, E. Marczewski, *Extension of measures*, Fund. Math. **36** (1949), 267-276.
- [6] M.D. Rašković, *Completeness theorem for biprobability models*, J. Symbolic Logic **51**, (1986), 586-590.
- [7] M.D. Rašković, *Completeness theorem for singular biprobability models*, Proc. Amer. Math. Soc. **102** (1988), 389-391.
- [8] M.D. Rašković, R.S. Đorđević, *Finite compactness theorem for biprobability logics*, Math. Balkanica **5** (1991), 12-14.
- [9] R.S. Đorđević, *Analytic completeness theorem for absolutely continuous biprobability models*, Zeitschr. Math. Logik Grundlagen Math. **38** (1992), 241-246.

Prirodno-matematički fakultet  
Radoja Domanovića 12  
34001 Kragujevac, p.p. 60  
Jugoslavija

(Received 16 11 1992)