

ON THE FOURTH MOMENT OF THE RIEMANN ZETA-FUNCTION

Aleksandar Ivić

Dedicated to the memory of Professor Đuro Kurepa

Abstract. Atkinson proved in 1941 that $\int_0^\infty e^{-t/T} |\zeta(1/2 + it)|^4 dt = TQ_4(\log T) + O(T^c)$ with $c = 8/9 + \varepsilon$, where $Q_4(y)$ is a suitable polynomial in y of degree four. We improve Atkinson's result by showing that $c = 1/2$ is possible, and we provide explicit expressions for all the coefficients of $Q_4(y)$ and the closely related polynomial $P_4(y)$.

1. Introduction

In recent years there has been much progress with problems involving the function $E_2(T)$ (see [7], [8], [9], [10], [12], [13], [15]). This important function, which represents the error term in the asymptotic formula for the fourth moment of the Riemann zeta-function $\zeta(s)$ on the so-called "critical line" $\text{Res} = \frac{1}{2}$, is defined by the relation

$$\int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^4 dt = T \sum_{j=0}^4 a_j \log^j T + E_2(T). \quad (1.1)$$

In 1926 Ingham [4] proved that $a_4 = 1/(2\pi^2)$. Much later in 1979 Heath-Brown [3] proved that $E_2(T) \ll T^{7/8+\varepsilon}$ ($f \ll g$ and $f = O(g)$ both mean that $|f(x)| \leq Cg(x)$ for $x \geq x_0$, $C > 0$ and $g(x) > 0$), and calculated

$$a_3 = 2(4\gamma - 1 - \log(2\pi) - 12\zeta'(2)\pi^{-2})\pi^{-2}, \quad (1.2)$$

where as usual $\gamma = 0.577215\dots$ is Euler's constant. The constants a_2 , a_1 and a_0 are more complicated, and were not stated explicitly in [3]. Heath-Brown's bound for $E_2(T)$ was improved to

$$E_2(T) = O(T^{2/3} \log^C T) \quad (C > 0) \quad (1.3)$$

in [9] by Motohashi and the author (see also [7]), where it was also proved that

$$E_2(T) = \Omega(T^{\frac{1}{2}}), \quad (1.4)$$

and as usual $f = \Omega(g)$, $g > 0$ means that $\lim_{x \rightarrow \infty} f(x)/g(x) \neq 0$. Therein it was also shown that

$$\int_0^T E_2(t) dt = O(T^{3/2}). \quad (1.5)$$

Recently Motohashi [15] improved (1.4) to $E_2(T) = \Omega_{\pm}(T^{\frac{1}{2}})$, and in [10] Motohashi and the author established that, with some $C > 0$,

$$\int_0^T E_2^2(t) dt = O(T^2 \log^C T). \quad (1.6)$$

In general, one can define the error term function $E_k(T)$ by the relation

$$\int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt = TP_{k^2}(\log T) + E_k(T), \quad (1.7)$$

where $k \geq 1$ is a fixed integer, and $P_{k^2}(y)$ is a suitable polynomial in y of degree k^2 . Apart from the classical case $k = 1$ (see [6] and [7] for an extensive discussion) and $k = 2$, our knowledge about the general $E_k(T)$ (see Ch. 4 of [7]) is very modest. At present it is not known $E_k(T) = o(T)$ as $T \rightarrow \infty$ for any $k \geq 3$, and in fact it is not clear how to define properly the coefficients of $P_{k^2}(y)$ for $k \geq 3$.

Instead at (1.7) one may look at the related formula of the Laplace transform (see Ch. 7 of Titchmarsh [16])

$$\int_0^{\infty} e^{-\delta t} \zeta \left(\frac{1}{2} + it \right) \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt \quad (\delta \rightarrow 0+, \quad k \geq 1 \text{ an integer}), \quad (1.8)$$

since Laplace transforms of many functions are easier to handle than the original functions. Kober [11] proved that

$$\int_0^{\infty} e^{-2\delta t} \left| \zeta \left(\frac{1}{2} + it \right) \right|^2 dt = \frac{\gamma - \log(4\pi\delta)}{2\sin\delta} + \sum_{n=0}^N c_n \delta^n + O(\delta^{N+1}) \quad (1.9)$$

for $\delta \rightarrow 0+$, any fixed integer $N \geq 1$ and suitable constants c_n . This is much sharper than the corresponding asymptotic formula (1.7) when $k = 1$. Atkinson [1] obtained

$$\begin{aligned} & \int_0^{\infty} e^{-\delta t} \left| \zeta \left(\frac{1}{2} + it \right) \right|^4 dt \\ &= \frac{1}{\delta} (A \log^4 \delta^{-1} + B \log^3 \delta^{-1} + C \log^2 \delta^{-1} + D \log \delta^{-1} + E) + L_2(\delta^{-1}) \end{aligned} \quad (1.10)$$

as $\delta \rightarrow 0+$ with

$$A = 1/(2\pi^2), \quad B = \pi^{-2}(2\log(2\pi) - 6\gamma + 24\zeta'(2)\pi^{-2}) \tag{1.11}$$

and $L_2(1/\delta) \ll (1/\delta)^{13/14+\varepsilon}$ for any given $\varepsilon > 0$. Atkinson's proof used bounds for the function $E(x, r)$ in (4.6). In his work it was indicated on p. 185 how a better bound for $E(x, r)$, which depends on bounds for Kloosterman sums, will lead to the better bound

$$L_2\left(\frac{1}{\delta}\right) \ll \left(\frac{1}{\delta}\right)^{8/9+\varepsilon} \quad (\delta \rightarrow 0+). \tag{1.12}$$

In analogy with (1.7) we define (writing in (1.8) $T = 1/\delta$ with $T \rightarrow \infty$) the function $L_k(T)$ by the relation

$$\int_0^\infty e^{-t/T} \left| \zeta\left(\frac{1}{2} + it\right) \right|^{2k} dt = TQ_{k^2}(\log T) + L_k(T), \tag{1.13}$$

where $k \geq 1$ is a fixed integer, $Q_{k^2}(y)$ is a suitable polynomial in y of degree k^2 , and one should have $L_k(T) = o(T)$ as $T \rightarrow \infty$. At present (analogously to $E_k(T) = o(T)$) the relation $L_k(T) = o(T)$ is not known to hold for $k \geq 3$.

2. Statement of results

A comparison of the asymptotic formulas (1.1) and (1.10) shows that $a_4 = A = 1/(2\pi^2)$, but that $a_3 \neq B$. One actually has

$$B = a_3 + 2\pi^{-2}(1 - \gamma), \tag{2.1}$$

the reason for which will become clear later. Recent advances concerning $E_2(T)$ make it possible to improve (1.12), and we have

THEOREM 1. *If L_2 is defined by (1.10), then as $T \rightarrow \infty$ one has*

$$L_2(T) = O(T^{1/2}). \tag{2.2}$$

This result gives a substantial improvement over Atkinson's exponent $8/9 + \varepsilon$ in (1.12). The exponent $1/2$ in (2.2) is the limit of the method of proof, which is based on the use of (1.5). One can prove, more generally, the following

THEOREM 2. *Suppose $k \geq 2$ is a fixed integer and*

$$\int_0^T E_k(t) dt = O(T^{c_k}) \tag{2.3}$$

holds for some $c_k > 0$. If L_k is defined by (1.13), then as $T \rightarrow \infty$

$$L_k(T) = O(T^{c_k-1}). \tag{2.4}$$

Moreover the coefficients of $Q_{k^2}(y)$ can be expressed as linear combinations of the coefficients of $P_{k^2}(y)$, defined by (1.7).

The remaining aim of this paper is to provide explicit expressions for the coefficients a_2 , a_1 and a_0 in (1.1) that were not stated explicitly by Heath-Brown [3]. From the nature of the problem it is clear that the expressions for these coefficients will be more complicated than the expression (1.2) for a_3 . For this reason they will not be stated here as a theorem, but will be dealt with in section 4, where all the appropriate notation will be introduced. From the proof of Theorem 2 in section 3 it will be clear that we can evaluate explicitly C , D and E in (1.10) as linear combinations of the a'_j 's. Conversely, if we know explicitly the coefficients of $Q_{k^2}(y)$, then it is not difficult to see that the coefficients of $P_{k^2}(y)$ can be written as linear combinations of the coefficients of $Q_{k^2}(y)$.

3. The Laplace transform of the $2k$ -th moment

It is enough to prove Theorem 2, since Theorem 1 is its consequence because, by (1.5), we have $c_2 = 3/2$ in (2.3) for $k = 2$. Let

$$I_k(T) = \int_0^T \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt = TP_{k^2}(\log T) + E_k(T)$$

with

$$P_{k^2}(y) = \sum_{j=0}^{k^2} a_j y^j, \quad a_j = a_j(k). \quad (3.1)$$

Then integration by parts gives

$$\begin{aligned} \int_0^\infty e^{-t/T} \left| \zeta \left(\frac{1}{2} + it \right) \right|^{2k} dt &= T^{-1} \int_0^\infty e^{-t/T} I_k(t) dt \\ &= T^{-1} \int_0^\infty e^{-t/T} t P_{k^2}(\log t) dt + T^{-1} \int_0^\infty e^{-t/T} E_k(t) dt \\ &= T \int_0^\infty e^{-x} x P_{k^2}(\log x + \log T) dx + T^{-2} \int_0^\infty e^{-t/T} \left(\int_0^t E_k(y) dy \right) dt = I' + I'', \end{aligned} \quad (3.2)$$

say. Inserting (3.1) in the expression for I' we obtain

$$\begin{aligned} I' &= T \int_0^\infty e^{-x} x \sum_{j=0}^{k^2} a_j (\log x + \log T)^j dx \\ &= T \int_0^\infty e^{-x} x \sum_{j=0}^{k^2} a_j \sum_{i=0}^j \binom{j}{i} \log^i T \cdot \log^{j-i} x \cdot dx \end{aligned}$$

$$= T \sum_{j=0}^{k^2} a_j \sum_{i=0}^j \binom{j}{i} \log^i T \left(\int_0^\infty e^{-x} x \cdot \log^{j-i} x \cdot dx \right).$$

But $\Gamma^{(k)}(z) = \int_0^\infty e^{-t} t^{z-1} (\log t)^k dt$, for $\text{Re} z > 0$ and $k \geq 0$ which gives

$$I' = T \sum_{j=0}^{k^2} a_j \sum_{i=0}^j \binom{j}{i} \Gamma^{(j-i)}(2) \cdot \log^i T = T \sum_{i=0}^{k^2} b_i \log^i T \tag{3.3}$$

with

$$b_i = b_i(k) = \sum_{j=i}^{k^2} \binom{j}{i} a_j \Gamma^{(j-i)}(2) \quad (i = 0, 1, \dots, k^2), \tag{3.4}$$

so that the coefficients of Q_{k^2} are linear combinations of the coefficients of P_{k^2} . By using (2.3) we obtain

$$\begin{aligned} I'' &= T^{-2} \int_0^\infty e^{-t/T} \left(\int_0^t E_k(y) dy \right) dt \ll T^{-2} \int_0^\infty e^{-t/T} t^{c_k} dt \\ &= T^{c_k-1} \Gamma(c_k + 1) \ll T^{c_k-1}, \end{aligned} \tag{3.5}$$

so that Theorem 2 follows from (1.13) and (3.2)–(3.5). Note that in the particular case $k = 2$ (3.4) yields

$$\begin{aligned} A &= b_4 = a_4 = 1/(2\pi^2), \quad B = b_3 = 4a_4\Gamma'(2) + a_3\Gamma(2), \\ C &= b_2 = 6a_4\Gamma''(2) + 3a_3\Gamma'(2) + a_2\Gamma(2), \\ D &= b_1 = 4a_4\Gamma^{(3)}(2) + 3a_3\Gamma''(2) + 2a_2\Gamma'(2) + a_1\Gamma(2), \\ E &= b_0 = a_4\Gamma^{(4)}(2) + a_3\Gamma^{(3)}(2) + a_2\Gamma''(2) + a_1\Gamma'(2) + a_0\Gamma(2). \end{aligned} \tag{3.6}$$

Since $\gamma = -\int_0^\infty e^{-x} \log x dx = -\Gamma'(1)$, it follows by an integration by parts that $-\gamma = -\int_0^\infty x(e^{-x} \log x)' dx = \Gamma'(2) - 1$. Thus $\Gamma'(2) = 1 - \gamma$, and (3.6) yields $B = b_3 = 4 \cdot \frac{1}{2\pi^2}(1 - \gamma) + a_3$, which is (2.1). Conversely, if the b'_j s are known, then (3.4) is a system of $k^2 + 1$ linear equations in $k^2 + 1$ unknowns a_0, a_1, \dots, a_{k^2} with a triangular determinant whose value is $(\Gamma(2))^{k^2+1} = 1$, so that the a'_j s can be uniquely expressed as linear combinations of the b'_j s.

4. The coefficients of the main term in the fourth moment formula

There are several ways to obtain explicitly the coefficients a_j in (1.1). This can be achieved by following the proofs of the fourth moment formula (see Ch. 4 of [7] or [13]). Here we shall follow the method of Heath-Brown [3], who showed that the main term $TP_4(\log T)$ in (1.1) consists of two parts: the part coming from the “diagonal” terms, and the part coming from the “non-diagonal” terms of a sum involving the number of divisors function $d(n)$.

The diagonal terms furnish an expression of the form $TR_4(\log T)$, and the non-diagonal terms the expression $TQ_2(\log T)$. Here $R_4(x)$ and $Q_2(x)$ denote suitable polynomials of degree four and two, respectively. It will turn out that the coefficients of $Q_2(x)$ have a more complex form than those of $R_4(x)$. Thus we shall have

$$P_4(x) = R_4(x) + Q_2(x). \quad (4.1)$$

As on p. 403 of [3] the diagonal terms make a contribution which is equal to

$$\begin{aligned} 2 \sum_{m \leq T/(2\pi)} d^2(m)m^{-1}(T - 2\pi m) \\ = \frac{1}{2\pi i} \left\{ \int_{1-i\infty}^{1+i\infty} 4\pi \zeta^4(s+1) \zeta^{-1}(2s+2) \left(\frac{T}{2\pi}\right)^{s+1} \frac{ds}{s(s+1)} \right\}. \end{aligned}$$

The term $TR_4(\log T)$ will be the residue of the simple pole (of the integrand in curly brackets) at $s = 0$. Near $s = 0$ we have the expansions

$$\zeta^{-1}(2s+2) = \zeta^{-1}(2) - \frac{2s\zeta'(2)}{\zeta^2(2)} + c_2s^2 + c_3s^3 + \dots$$

with

$$\begin{aligned} c_k &= \frac{2^k}{k!} \sum_{n=1}^{\infty} \mu(n)(-\log n)^k n^{-2} = \frac{1}{k!} \left\{ \frac{d^k}{ds^k} (\zeta^{-1}(2s+2)) \right\} \Big|_{s=0}, \\ \frac{1}{s+1} &= 1 - s + s^2 - s^3 + \dots, \\ \left(\frac{T}{2\pi}\right)^{s+1} &= \frac{T}{2\pi} \left\{ 1 + s \log\left(\frac{T}{2\pi}\right) + \frac{s^2}{2!} \log^2\left(\frac{T}{2\pi}\right) + \frac{s^3}{3!} \log^3\left(\frac{T}{2\pi}\right) + \dots \right\}, \\ \zeta^4(s+1) &= \frac{1}{s^4} + \frac{4\gamma}{s^3} + \frac{b_{-2}}{s^2} + \frac{b_{-1}}{s} + b_0 + b_1s + b_2s^2 + \dots \end{aligned} \quad (4.2)$$

The coefficients b_k ($k \geq -2$) may be found from the relation

$$\zeta^4(s+1) = \left(\frac{1}{s} + \gamma_0 + \gamma_1s + \gamma_2s^2 + \gamma_3s^3 + \dots \right)^4, \quad (4.3)$$

where one has (see Theorem 1.3 of [6])

$$\gamma_0 = \gamma, \gamma_k = \frac{(-1)^k}{k!} \lim_{N \rightarrow \infty} \left(\sum_{m \leq N} \frac{\log^k m}{m} - \frac{\log^{k+1} N}{k+1} \right).$$

Israilov [5] calculated

$$\gamma_1 = 0.072815846\dots, \quad \gamma_2 = -0.004845182\dots, \quad \gamma_3 = -0.000342305\dots,$$

and Euler's constant $\gamma = \gamma_0$ is of course known with much greater accuracy,

$$\gamma = 0.5772\ 15664\ 90153\ 28606\ 06512\dots$$

From (4.2) and (4.3) we obtain by comparing the coefficients

$$\begin{aligned} b_{-2} &= 4\gamma_1 + 6\gamma^2, & b_{-1} &= 4\gamma_2 + 12\gamma\gamma_1 + 4\gamma^3, \\ b_0 &= 4\gamma_3 + 12\gamma\gamma_2 + 6\gamma_1^2 + 12\gamma^2\gamma_1 + \gamma^4. \end{aligned}$$

Since the residue is the coefficient of s^{-1} , we obtain

$$\begin{aligned} TR_4(\log T) & \\ &= \frac{2T}{\zeta(2)} \left\{ \frac{1}{24} \log^4 \left(\frac{T}{2\pi} \right) + \frac{a}{6} \log^3 \left(\frac{T}{2\pi} \right) + \frac{b}{2} \log^2 \left(\frac{T}{2\pi} \right) + c \log \left(\frac{T}{2\pi} \right) + d \right\} \end{aligned} \quad (4.4)$$

with

$$\begin{aligned} a &= 4\gamma - 1 - \frac{2\zeta'(2)}{\zeta(2)}, & b &= 1 + \frac{2\zeta'(2)}{\zeta(2)} + c_2\zeta(2) + 4\gamma \left(-1 - \frac{2\zeta'(2)}{\zeta(2)} \right) + b_{-2}, \\ c &= -1 - \frac{2\zeta'(2)}{\zeta(2)} + (c_3 - c_2)\zeta(2) + 4\gamma \left(1 + c_2\zeta(2) + \frac{2\zeta'(2)}{\zeta(2)} \right) \\ &\quad + b_{-2} \left(-1 - \frac{2\zeta'(2)}{\zeta(2)} \right) + b_{-1}, \\ d &= 1 + \frac{2\zeta'(2)}{\zeta(2)} + (c_2 - c_3 + c_4)\zeta(2) + 4\gamma \left(-1 - \frac{2\zeta'(2)}{\zeta(2)} + (c_3 - c_2)\zeta(2) \right) \\ &\quad + b_{-2} \left(1 + \frac{2\zeta'(2)}{\zeta(2)} + c_2\zeta(2) \right) + b_{-1} \left(-1 - \frac{2\zeta'(2)}{\zeta(2)} \right) + b_0. \end{aligned} \quad (4.5)$$

Expanding $\log^j(T/2\pi) = (\log T - \log 2\pi)^j$ by the binomial theorem we obtain ($TR_4(\log T) = g_1(T)$ in Heath-Brown's notation)

$$\begin{aligned} TR_4(\log T) &= \frac{T}{\pi^2} \left\{ \frac{1}{2} \log^4 T + (2a - 2 \log(2\pi)) \log^3 T + (3 \log^2(2\pi) - 6a \log(2\pi) \right. \\ &\quad + 6b) \log^2 T + (-2 \log^3(2\pi) + 6a \log^2(2\pi) - 12b \log(2\pi) + 12c) \log T \\ &\quad \left. + \left(\frac{1}{2} \log^4(2\pi) - 2a \log^3(2\pi) + 6b \log^2(2\pi) - 12c \log(2\pi) + 12d \right) \right\}. \end{aligned}$$

The coefficients a_4 and a_3 in (1.1) are the coefficients of $\log^4 T$ and $\log^3 T$, respectively, and so by the first formula in (4.5) they are

$$a_4 = \frac{1}{2\pi^2}, \quad a_3 = \frac{2a - 2 \log(2\pi)}{\pi^2} = \frac{8\gamma - 2 - 24\zeta'(2)\pi^{-2} - 2 \log(2\pi)}{\pi^2}.$$

These are the same values that were obtained by Ingham and Heath-Brown.

For the remaining part $TQ_2(\log T)$ in the main term one has (see p. 404 of [3]) that it is the main term in the asymptotic formula for

$$f_0(T) \sim 2\text{Re} \left\{ \sum_{r=1}^R (ir)^{-1} S_r \right\} \quad (R \rightarrow \infty),$$

where $S_r \sim \int_0^{T/(2\pi)} m'(x, r) e^{iTr/x} dx$. Here $m(x, r)$ stands for the main term in the asymptotic formula for the so-called binary additive divisor problem (see Motohashi [14] for an extensive discussion), namely

$$\sum_{n \leq x} d(n)d(n+r) = m(x, r) + E(x, r), \quad m(x, r) = x \sum_{j=0}^2 c_{3-j}(r) \log^j x. \quad (4.6)$$

We have $m'(x, r) = d_0(r) \log^2 x + d_1(r) \log x + d_2(r)$ with

$$d_0(r) = c_1(r), \quad d_1(r) = c_2(r) + 2c_1(r), \quad d_2(r) = c_2(r) + c_3(r). \quad (4.7)$$

In Theorem 2 of [3] Heath-Brown evaluated the constants $c_i(r)$ in (4.6), but his expressions are cumbersome. We find it more expedient to use the expressions given by Balakrishnan and Sengupta [2], and from these expressions and (4.7) we obtain, with the notation

$$\sigma_z(n) = \sum_{d|n} d^z, \quad \sigma'_z(n) = \frac{d}{dz}(\sigma_z(n)) = \sum_{d|n} d^z \log d, \quad \sigma''_z(n) = \sum_{d|n} d^z \log^2 d,$$

the following:

$$\begin{aligned} d_0(r) &= \frac{\sigma_{-1}(r)}{\zeta(2)}, \quad d_1(r) = d_0(r) \left\{ 4\gamma - 4\frac{\zeta'(2)}{\zeta} - 4\frac{\sigma'_{-1}(r)}{\sigma_{-1}} \right\}, \\ d_2(r) &= d_0(r) \left\{ 4\gamma - 1 - 4\frac{\zeta'(2)}{\zeta} - 4\frac{\sigma'_{-1}(r)}{\sigma_{-1}} - 4\frac{\zeta''(2)}{\zeta} - 4\left(\frac{\zeta'(2)}{\zeta}\right)^2 \right. \\ &\quad \left. + 4\frac{\sigma''_{-1}(r)}{\sigma_{-1}} - 4\left(\frac{\sigma'_{-1}(r)}{\sigma_{-1}}\right)^2 + \left(2\gamma - 1 - 2\frac{\zeta'(2)}{\zeta} - 2\frac{\sigma'_{-1}(r)}{\sigma_{-1}}\right)^2 \right\}. \end{aligned} \quad (4.8)$$

Further, after a change of variable, we have

$$\begin{aligned} S_r &\sim \int_0^{T/(2\pi)} \{d_0(r) \log^2 x + d_1(r) \log x + d_2(r)\} e^{iTr/x} dx \\ &= Tr \int_{2\pi r}^{\infty} \{d_0(r) \log^2 \left(\frac{Tr}{y}\right) + d_1(r) \log \left(\frac{Tr}{y}\right) + d_2(r)\} e^{iy} \frac{dy}{y^2}. \end{aligned}$$

Therefore it follows that

$$TQ_2(\log T) = 2\operatorname{Re} \left\{ \sum_{r=1}^{\infty} (ir)^{-1} S_r \right\} = T(e_0 \log^2 T + e_1 \log T + e_2)$$

with

$$\begin{aligned} e_0 &= 2 \sum_{r=1}^{\infty} d_0(r) \int_{2\pi r}^{\infty} \frac{\sin x}{x^2} dx, \\ e_1 &= \sum_{r=1}^{\infty} \left\{ d_0(r) \int_{2\pi r}^{\infty} (2 \log r - 2 \log x) \frac{\sin x}{x^2} dx + d_1(r) \int_{2\pi r}^{\infty} \frac{\sin x}{x^2} dx \right\}, \end{aligned} \quad (4.9)$$

$$e_2 = \sum_{r=1}^{\infty} \int_{2\pi r}^{\infty} \left\{ d_0(r) \log^2 x - (2d_0(r) \log r + d_1(r)) \log x + d_0(r) \log^2 r + d_1(r) \log r + d_2(r) \right\} \frac{\sin x}{x^2} dx.$$

If, for $\text{Re } a < 1$ and $x > 0$ we introduce the standard notation

$$C(x, a) = \int_x^{\infty} t^{a-1} \cos t \cdot dt, \quad S(x, a) = \int_x^{\infty} t^{a-1} \sin t \cdot dt,$$

then we have

$$\frac{\partial S(x, a)}{\partial a} = \int_x^{\infty} t^{a-1} \log t \cdot \sin t \cdot dt, \quad \frac{\partial^2 S(x, a)}{\partial a^2} = \int_x^{\infty} t^{a-1} \log^2 t \cdot \sin t \cdot dt,$$

and

$$e^{-\frac{1}{2}\pi ia} \Gamma(a, ix) = C(x, a) - iS(x, a), \quad S(x, a) = -\text{Im}\{e^{-\frac{1}{2}\pi ia} \Gamma(a, ix)\},$$

$$\Gamma(a, x) = \Gamma(a) - \gamma(a, x) = \int_x^{\infty} e^{-t} t^{a-1} dt,$$

where $\gamma(a, x) = \int_0^x e^{-t} t^{a-1} dt$ is the incomplete gamma-function. With this notation we have

$$e_0 = 2 \sum_{r=1}^{\infty} d_0(r) S(2\pi r, -1),$$

$$e_1 = 2 \sum_{r=1}^{\infty} \left\{ (2d_0(r) \log r + d_1(r)) S(2\pi r, -1) - 2d_0(r) \frac{\partial S}{\partial a}(2\pi r, -1) \right\}, \quad (4.10)$$

$$e_2 = 2 \sum_{r=1}^{\infty} \left\{ (d_0(r) \frac{\partial^2 S}{\partial a^2}(2\pi r, -1) - (2d_0(r) \log r + d_1(r)) \frac{\partial S}{\partial a}(2\pi r, -1) + (d_0(r) \log^2 r + d_1(r) \log r + d_2(r)) S(2\pi r, -1) \right\}.$$

Hence finally from (4.1) and the above expressions we obtain

THEOREM 3. *For the coefficients a_j in (1.1) we have*

$$a_4 = \frac{1}{2\pi^2}, \quad a_3 = 2(4\gamma - 1 - \log(2\pi) - 12\zeta'(2)\pi^{-2})\pi^{-2},$$

$$a_2 = (3 \log^2(2\pi) - 6a \log(2\pi) + 6b)\pi^{-2} + e_0,$$

$$a_1 = (-2 \log^3(2\pi) + 6a \log^2(2\pi) - 12b \log(2\pi) + 12c)\pi^{-2} + e_1,$$

$$a_0 = \left(\frac{1}{2} \log^4(2\pi) - 2a \log^3(2\pi) + 6b \log^2(2\pi) - 12c \log(2\pi) + 12d \right) \pi^{-2} + e_2,$$

where a, b, c, d are given by (4.5) and e_0, e_1, e_2 by (4.10).

REFERENCES

- [1] F.V. Atkinson, *The mean value of the zeta-function on the critical line*, Proc. London Math. Soc. (2) **47** (1941), 174–200.
- [2] U. Balakrishnan and J. Sengupta, *On the sum $\sum d(n)\sigma(n+N)$* , Manuscr. Math. **67** (1990), 367–378.
- [3] D.R. Heath-Brown, *The fourth power moment of the Riemann zeta-function*, Proc. London Math. Soc. (3) **38** (1979), 385–422.
- [4] A.E. Ingham, *Mean-value theorems in the theory of the Riemann zeta-function*, Proc. London Math. Soc. (2) **27** (1926), 273–300.
- [5] M.I. Israilov, *On the Laurent expansion of the Riemann zeta-function (Russian)*, Trudy Mat. Inst. AN SSSR **158** (1981), 98–104.
- [6] A. Ivić, *The Riemann zeta-function*, John Wiley & Sons, New York, 1985.
- [7] A. Ivić, *The mean values of the Riemann zeta-function*, Lecture Notes 82, Tata Institute, Bombay, 1991 (distr. by Springer Verlag, Berlin etc.).
- [8] A. Ivić and Y. Motohashi, *A note on the mean value of the zeta and L-functions VII*, Proc. Japan Acad. Ser. A **66** (1990), 150–152.
- [9] A. Ivić and Y. Motohashi, *On the fourth power moment of the Riemann zeta-function*, J. Number Theory **51** (1995), 16–45.
- [10] A. Ivić and Y. Motohashi, *The mean square of the error term for the fourth power moment of the zeta-function*, Proc. London Math. Soc. (3) **69** (1994), 309–329.
- [11] H. Kober, *Eine Mittelwertformel der Riemannschen Zetafunktion*, Compositio Math. **3** (1936), 174–189.
- [12] Y. Motohashi, *The fourth power mean of the Riemann zeta-function*, in “Proc. Conf. Analytic Number Theory (Amalfi, 1989)”, eds. E. Bombieri et al., Università di Salerno, Salerno, 1992, 325–344.
- [13] Y. Motohashi, *An explicit formula for the fourth power mean of the Riemann zeta-function*, Acta Math. **170** (1993), 181–220.
- [14] Y. Motohashi, *The binary additive divisor problem*, Ann. Scien. Ecole Norm. Sup. 4 série **27** (1994), 529–572.
- [15] Y. Motohashi, *A relation between the Riemann zeta-function and the hyperbolic Laplacian*, to appear in Ann. Scuola Norm. Sup. Pisa classe di Scienze.
- [16] E.C. Titchmarsh, *The theory of the Riemann zeta-function (2nd ed.)*, Clarendon Press, Oxford, 1986.

Katedra Matematike RGF-a
 Universiteta u Beogradu
 Dušina 7
 11000 Beograd
 Serbia (Yugoslavia)

(Received 24 01 1995)