PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 72(86) (2002), 49–54

ON GAPS BETWEEN BOUNDED OPERATORS

Dragana Cvetković

Communicated by Miroljub Jevtić

Abstract. We consider the distance between an arbitrary bounded operator on a Banach space X and the null operator. The distance is computed in terms of various gaps. Thus, we generalize the Habibi's result from [4]

1. Introduction

In [4] Habibi considered the spherical gap between a linear operator and the null operator on finite dimensional Hilbert spaces. We extend (see also [1]) his result to various gaps on arbitrary Banach spaces.

Let X, Y be arbitrary Banach spaces. We denote by $\mathcal{G}(X)$, and $\mathcal{B}(X,Y)$, respectively, the set of all closed subspaces of X and the set of all bounded linear operators from X to Y. The closed unit sphere of a Banach space X is denoted by $S(X)$. If x is a vector of a Banach space X and M is a subset of X we put $\operatorname{dist}(x,M) = \inf_{m\in M} \|x-m\|$. If M, N are subspaces of X, the set of all invertible operators C on X such that $C(M) = N$ is denoted by $\mathcal{B}(X; M, N)^{-1}$. Let $X \times Y$ be the space with the norm

$$\|(x,y)\| = (\|x\|^p + \|y\|^p)^{1/p}, \quad x \in X,\ y \in Y,\ p \geqslant 1.$$

Let $M, N \in \mathcal{G}(X)$. The spherical gap between M and N is defined by [2]

$$\widetilde{\Theta}(M,N) = \max\left\{\widetilde{\Theta}_0(M,N), \widetilde{\Theta}_0(N,M)\right\},$$

where

$$\widetilde{\Theta}_0(M,N) = \sup_{m\in S(M)} d(m, S(N)).$$

2000 *Mathematics Subject Classification.* Primary: 46B20; Secondary 15A60.
Key words and phrases. spherical gap; geometric gap.

The geometric gap between M and N is defined by [6]

$$\Theta(M,N) = \max\{\Theta_0(M,N), \Theta_0(N,M)\},$$

where

$$\Theta_0(M,N) = \sup_{m\in S(M)} d(m,N).$$

If $A, B \in \mathcal{B}(X,Y)$, then their graphs $G(A), G(B)$ are closed subspaces of $X\times Y$, so the gap between the operators A and B is defined as the gap between their graphs, i.e.,

$$\widetilde{\Theta}(A,B) = \widetilde{\Theta}(G(A),G(B)) \text{ and } \Theta(A,B) = \Theta(G(A),G(B)).$$

Also,

$$\widetilde{\Theta}_0(A,B) = \widetilde{\Theta}_0(G(A),G(B)) \quad \text{and} \quad \Theta_0(A,B) = \Theta_0(G(A),G(B)).$$

Obviously,

$$\widetilde{\Theta}(A,B) = \max\left\{\widetilde{\Theta}_0(A,B), \widetilde{\Theta}_0(B,A)\right\},$$

$$\Theta(A,B) = \max\{\Theta_0(A,B), \Theta_0(B,A)\}.$$

In [4] Habibi considered the spherical gap between operators on finite dimensional Hilbert spaces, assuming $p = 2$. We shall consider spherical and geometric gap of bounded operators on Banach spaces, where $p \geqslant 1$ is arbitrary. Thus, a generalization of Habibi's result is obtained.

2. Results

First we compute the spherical gap between a bounded operator and the null operator.

THEOREM 2.1. *If X, Y are Banach spaces and $A \in \mathcal{B}(X,Y)$, then*

$$\widetilde{\Theta}(A) = \widetilde{\Theta}(A,0) = \left(\left(1 - \frac{1}{(1+\|A\|^p)^{1/p}}\right)^p + \frac{\|A\|^p}{1+\|A\|^p}\right)^{1/p}.$$

Proof. Since

$$G(A) = \{(x,Ax) : x \in X\} \text{ and } G(0) = \{(x,0) : x \in X\},$$

it follows that

$$\begin{aligned}\widetilde{\Theta}_0(A,0) &= \sup_{(y,Ay)\in S(G(A))} \inf_{(x,0)\in S(G(0))} \|(x,0)-(y,Ay)\| \\ &= \sup_{\substack{y\in X \\ \|y\|^p+\|Ay\|^p=1}} \inf_{\substack{x\in X \\ \|x\|=1}} \left(\|x-y\|^p + \|Ay\|^p\right)^{1/p}.\end{aligned}$$

Consider the function

$$f(x,y) = (\|x-y\|^p + \|Ay\|^p)^{1/p}.$$

Suppose that $x \in X$, $y \in Y$ satisfy

$$\|x\| = 1, \ \|y\|^p + \|Ay\|^p = 1. \tag{1}$$

Then

$$f(x,y) \geqslant ((\|x\| - \|y\|)^p + \|Ay\|^p)^{1/p} = ((1-\|y\|)^p + \|Ay\|^p)^{1/p}.$$

Hence,

$$\inf_{\substack{x\in X\\ \|x\|=1}} f(x,y) \geqslant ((1-\|y\|)^p + \|Ay\|^p)^{1/p}. \tag{2}$$

Also,

$$\inf_{\substack{x\in X\\ \|x\|=1}} f(x,y) \leqslant f\left(\frac{y}{\|y\|}, y\right) = ((1-\|y\|)^p + \|Ay\|^p)^{1/p}. \tag{3}$$

So, from (2) and (3) it follows that

$$\inf_{\substack{x\in X\\ \|x\|=1}} f(x,y) = ((1-\|y\|)^p + \|Ay\|^p)^{1/p}. \tag{4}$$

According to the condition (1) we have

$$\|y\| \geqslant \frac{1}{(1+\|A\|^p)^{1/p}}.$$

Since the function $\phi(t) = (1-t)^p + 1 - t^p$ is decreasing for $0 \leqslant t \leqslant 1$, we have

$$\begin{aligned} \widetilde{\Theta}_0(A,0) &= \sup_{\substack{y\in X\\ \|y\|^p+\|Ay\|^p=1}} ((1-\|y\|)^p + 1 - \|y\|^p)^{1/p} \\ &= \left((1 - \frac{1}{(1+\|A\|^p)^{1/p}})^p + \frac{\|A\|^p}{1+\|A\|^p}\right)^{1/p}. \end{aligned} \tag{5}$$

Let $y = cx$, where $c = \frac{1}{(1+\|Ax\|^p)^{1/p}}$. It is obvious that y satisfy the condition (1) and $f(x,y) = \left(\left(1 - \frac{1}{(1+\|Ax\|^p)^{1/p}}\right)^p + \frac{\|Ax\|^p}{1+\|Ax\|^p}\right)^{1/p}$.

So,

$$\inf_{\substack{y\in Y\\ \|y\|^p+\|Ay\|^p=1}} f(x,y) \leqslant \left(\left(1 - \frac{1}{(1+\|Ax\|^p)^{1/p}}\right)^p + \frac{\|Ax\|^p}{1+\|Ax\|^p}\right)^{1/p}. \tag{6}$$

The function $\psi(t) = (1 - \frac{1}{(1+t)^{1/p}})^p + \frac{t}{1+t}$ is increasing, so from (6) it follows that

$$\widetilde{\Theta}_0(0,A) \leqslant \sup_{\substack{x\in X\\ \|x\|=1}} \left(\left(1 - \frac{1}{(1+\|Ax\|^p)^{1/p}}\right)^p + \frac{\|Ax\|^p}{1+\|Ax\|^p}\right)^{1/p}$$

$$= \left(\left(1 - \frac{1}{(1+\|A\|^p)^{1/p}}\right)^p + \frac{\|A\|^p}{1+\|A\|^p}\right)^{1/p}. \tag{7}$$

Finally, from (5) and (7) we have that

$$\widetilde{\Theta}(A) = \max\left\{\widetilde{\Theta}_0(A,0), \widetilde{\Theta}_0(0,A)\right\} = \widetilde{\Theta}_0(A,0)$$

$$= \left(\left(1 - \frac{1}{(1+\|A\|^p)^{1/p}}\right)^p + \frac{\|A\|^p}{1+\|A\|^p}\right)^{1/p} \quad \square$$

From the proof of Theorem 2.1 it follows that

$$\widetilde{\Theta}_0(0,A) \leqslant \widetilde{\Theta}_0(A,0) = \left((1 - \frac{1}{(1+\|A\|^p)^{1/p}})^p + \frac{\|A\|^p}{1+\|A\|^p}\right)^{1/p}.$$

Notice that Habibi's main result in [4] is a special case of our Theorem 2.1 for $p = 2$.

Now, we consider the geometric gap.

THEOREM 2.2. *If X, Y are Banach spaces and $A \in \mathcal{B}(X,Y)$, then*

$$\Theta_0(A,0) = \frac{\|A\|}{(1+\|A\|^p)^{1/p}}.$$

Proof. It follows that

$$\Theta_0(A,0) = \sup_{\substack{y\in X\\ \|y\|^p+\|Ay\|^p=1}} \inf_{x\in X} (\|x-y\|^p + \|Ay\|^p)^{1/p}$$

$$= \sup_{\substack{y\in X\\ \|y\|^p+\|Ay\|^p=1}} \|Ay\| = \sup_{z\in X} \frac{\|Az\|}{(\|z\|^p + \|Az\|^p)^{1/p}} = \frac{\|A\|}{(1+\|A\|^p)^{1/p}} \quad \square$$

In the case $p = 2$ we can obtain the following Habibi's result.

THEOREM 2.3. *If X and Y are Banach spaces, $A \in \mathcal{B}(X,Y)$ and $p = 2$, then the following holds:*

$$\Theta(A,0) = \frac{\|A\|}{(1+\|A\|^2)^{1/2}}.$$

Proof. By Theorem 2.2 it is sufficient to prove

$$\Theta_0(0,A) \leqslant \frac{\|A\|}{(1+\|A\|^2)^{1/2}}.$$

Actually, this is a Habibi's result from [3]. For reader's convenience, we give a complete proof. Consider the function $f(x,y) = (\|x-y\|^2 + \|Ay\|^2)^{1/2}$. Let $x \in S(X)$ and $y = \frac{x}{1+\|Ax\|^2}$. Then we have

$$\operatorname{dist}((x,0),G(A)) \leqslant f(x,y) = \frac{\|Ax\|}{(1+\|Ax\|^2)^{1/2}}.$$

Since the function $\mu(t) = \frac{t}{1+t}$ is increasing, it follows that

$$\begin{aligned}\Theta_0(0,A) &= \sup_{\|x\|=1} \operatorname{dist}((x,0),G(A)) \leqslant \sup_{\|x\|=1} \frac{\|Ax\|}{(1+\|Ax\|^2)^{1/2}} \\ &= \frac{\|A\|}{(1+\|A\|^2)^{1/2}}.\end{aligned}$$

□

Finally, we can prove the following result.

THEOREM 2.4. *Let X, Y be Banach spaces and $X \times Y$ is the space with the norm*

$$\|(x,y)\| = (\|x\|^2 + \|y\|^2)^{1/2}, \quad x \in X,\ y \in Y.$$

If $A, B \in B(X,Y)$, then

$$\Theta(A,B) \leqslant 2(1+\min\{\|A\|,\|B\|\}^2)\frac{\|A-B\|}{(1+\|A-B\|^2)^{1/2}}.$$

Proof. From [5],Theorem 2.17, page 204, it follows that

$$\Theta(A,B) = \Theta((A-B)+B, 0+B) \leqslant 2(1+\|B\|^2)\Theta(A-B,0),$$

and

$$\Theta(A,B) = \Theta(0+A,(B-A)+A) \leqslant 2(1+\|A\|^2)\Theta(B-A,0).$$

Now, by Theorem 2.3 we have that

$$\Theta(A,B) \leqslant 2(1+\|B\|^2)\frac{\|A-B\|}{(1+\|A-B\|^2)^{1/2}},$$

and

$$\Theta(A,B) \leqslant 2(1+\|A\|^2)\frac{\|A-B\|}{(1+\|A-B\|^2)^{1/2}},$$

i.e.,

$$\Theta(A,B) \leqslant 2(1+\min\{\|A\|,\|B\|\}^2)\frac{\|A-B\|}{(1+\|A-B\|^2)^{1/2}}. \quad \square$$

The author wishes to thank referee for suggestions which greatly improved several proofs.

References

[1] D. Cvetković, *Gaps between Banach spaces(in Serbian), Master Thesis*, Faculty of Science and Mathematics, University of Nis (2000), 1–81.

[2] I. C.Gohberg and A. S. Markus, *Two theorems on the opening between subspaces of a Banach space*, Uspekhi Mat. Nauk. **14 (5)** (1959), 135–140 (Russian).

[3] J. F. Habibi, *The gap of the graph of a matrix*, Linear Algebra Appl. **186** (1993), 55–57.

[4] J. F. Habibi, *The spherical gap of the graph of a linear transformation*, Linear Algebra Appl. **212** (1994), 501–503.

[5] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin-Heidelberg-New York, 1976.

[6] M. G. Krein, M. A. Krasnoselskii and D. P. Milman, *On the defect numbers of linear operators in a Banach space and on some geometric questions*, Sbornik Trudov Inst. Mat. AN Ukrainian SSR **11** (1948), 97–112 (Russian).

Prirodno-atematički fakultet
18000 Niš
Serbia and Montenegro

dragana@pmf.pmf.ni.ac.yu

(Received 17 01 2001)
(Revised 28 02 2002)