

## HÖLDER SPACES OF QUASICONFORMAL MAPPINGS

Leonid V. Kovalev

ABSTRACT. We prove that a  $K$ -quasiconformal mapping belongs to the little Hölder space  $c^{0,1/K}$  if and only if its local modulus of continuity has an appropriate order of vanishing at every point. No such characterization is possible for Hölder spaces with exponent greater than  $1/K$ .

### 1. Introduction

Let  $\Omega$  denote a domain in  $\mathbb{C}$ , and let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous complex-valued function. Given  $E \subset \Omega$ , define the modulus of continuity of  $f|_E$  by

$$\omega_f(E, \delta) = \sup\{|f(z_1) - f(z_2)| : z_1, z_2 \in E, |z_1 - z_2| \leq \delta\}.$$

For  $0 < \alpha < 1$  we consider the Hölder space

$$C^{0,\alpha}(E) = \{f : E \rightarrow \mathbb{C} : \sup_{\delta > 0} \delta^{-\alpha} \omega_f(E, \delta) < \infty\},$$

with the seminorm

$$\|f\|_{E,\alpha} = \limsup_{\delta \rightarrow 0} \delta^{-\alpha} \omega_f(E, \delta).$$

This seminorm vanishes on the little Hölder space

$$c^{0,\alpha}(E) = \{f \in C^{0,\alpha}(E) : \|f\|_{E,\alpha} = 0\}.$$

Furthermore, define  $C_{\text{loc}}^{0,\alpha}(\Omega) = \bigcap_{E \Subset \Omega} C^{0,\alpha}(E)$  and similarly for  $c_{\text{loc}}^{0,\alpha}(\Omega)$ .

We can also consider the local modulus of continuity at a point  $z \in \Omega$ :

$$\omega_f(z, \delta) = \sup\{|f(\zeta) - f(z)| : \zeta \in \Omega, |\zeta - z| \leq \delta\}.$$

If  $U$  is a neighborhood of  $z$  in  $\Omega$ , then  $\omega_f(z, \delta) \leq \omega_f(U, \delta)$  for all sufficiently small  $\delta > 0$ . In particular,

$$(1.1) \quad \limsup_{\delta \rightarrow 0} \delta^{-\alpha} \omega_f(z, \delta) \leq \|f\|_{U,\alpha}.$$

---

2000 *Mathematics Subject Classification*: Primary 30C62; Secondary 26B35.

*Key words and phrases*: Quasiconformal mappings, Hölder spaces, linear dilatation, modulus of continuity.

Inequality (1.1) provides a simple necessary condition for a continuous mapping  $f : \Omega \rightarrow \mathbb{C}$  to be in the class  $c_{\text{loc}}^{0,\alpha}(\Omega)$ ; namely,

$$(1.2) \quad f \in c_{\text{loc}}^{0,\alpha}(\Omega) \implies \limsup_{\delta \rightarrow 0} \delta^{-\alpha} \omega_f(z, \delta) = 0 \quad \forall z \in \Omega.$$

This condition can be helpful because it is often easier to estimate  $\omega_f(z, \delta)$  for  $z \in \Omega$  than to estimate  $\omega_f(E, \delta)$  for all  $E \Subset \Omega$ . Unfortunately, the implication in (1.2) cannot be reversed in general.

The present paper deals with the following question: is the reverse implication in (1.2) true under the additional assumption that  $f$  is a  $K$ -quasiconformal mapping from  $\Omega$  to  $\mathbb{C}$ ? It is well-known that under this assumption  $f$  belongs to  $C_{\text{loc}}^{0,1/K}(\Omega)$  [1, 3, 7], but not necessarily to  $c_{\text{loc}}^{0,1/K}(\Omega)$  (for example,  $f(z) = |z|^{1/K-1}z$  is  $K$ -quasiconformal in  $\mathbb{C}$ , but  $f \notin c_{\text{loc}}^{0,1/K}(\mathbb{C})$ ). Therefore, our question is nontrivial only when  $1/K \leq \alpha < 1$ .

The answer turns out to be affirmative in the case  $\alpha = 1/K$  (Theorem 2.1) and negative in the case  $1/K < \alpha < 1$  (Proposition 2.1).

## 2. Main results

We start by showing that in general one cannot determine the degree of Hölder continuity of a quasiconformal mapping from its local behavior. More precisely, the following proposition exhibits a  $K$ -quasiconformal mapping which has linear local modulus of continuity at every point, yet does not belong to  $c_{\text{loc}}^{0,\alpha}(\Omega)$  with  $\alpha$  arbitrarily close to  $1/K$ . We use notations  $\mathbb{D}(a, r) = \{z \in \mathbb{C} : |z - a| < r\}$  and  $\mathbb{D} = \mathbb{D}(0, 1)$ .

**PROPOSITION 2.1.** *Given  $K > 1$  and  $1/K < \alpha < 1$ , there exists a  $K$ -quasiconformal automorphism  $f : \mathbb{D} \rightarrow \mathbb{D}$  such that  $f \notin c_{\text{loc}}^{0,\alpha}(\mathbb{D})$ , but*

$$(2.1) \quad \limsup_{\zeta \rightarrow z} \frac{|f(\zeta) - f(z)|}{|\zeta - z|} < \infty$$

for every  $z \in \mathbb{D}$ .

**PROOF.** Choose  $\varepsilon > 0$  so that  $(\alpha - \varepsilon)/(1 - \varepsilon) = 1/K$ . Consider two sequences of open disks  $D_n = \mathbb{D}(2^{-n}, 2^{-(n+2)})$  and  $D'_n = \mathbb{D}(2^{-n}, 2^{-(n+2)/\varepsilon})$ ,  $n \geq 1$ . We will define  $f$  separately on  $D'_n$ ,  $D_n \setminus D'_n$  and  $\mathbb{D} \setminus \bigcup_{n=1}^{\infty} D_n$ . Each disk  $D'_n$  is stretched under  $f$  by the factor of  $2^{(1-\alpha)(n+2)/\varepsilon}$ :

$$f(2^{-n} + re^{i\varphi}) = 2^{-n} + 2^{(1-\alpha)(n+2)/\varepsilon} r e^{i\varphi}, \quad 0 \leq r < 2^{-(n+2)/\varepsilon}, \quad \varphi \in \mathbb{R}.$$

Thus  $f(D'_n)$  is a disk that is concentric with  $D_n$  and has the radius  $2^{-\alpha(n+2)/\varepsilon} < 2^{-(n+2)}$ . Hence  $f(D'_n) \subset D_n$ . Next,  $f$  maps the annulus  $D_n \setminus D'_n$  onto  $D_n \setminus f(D'_n)$  by means of the “extremal  $K$ -quasiconformal stretch mapping” (cf. [5, p.63]).

$$f(2^{-n} + re^{i\varphi}) = 2^{-n} + 2^{(n+2)(1/K-1)} r^{1/K} e^{i\varphi}, \quad 2^{-(n+2)/\varepsilon} \leq r < 2^{-n-2}, \quad \varphi \in \mathbb{R}.$$

Finally, let  $f(z) = z$  for  $z \notin \bigcup_{n=1}^{\infty} D_n$ . It is easy to see that  $f$  is continuous and thus  $K$ -quasiconformal in  $\mathbb{D}$ . It is also evident that  $f$  is locally Lipschitz in  $\mathbb{D} \setminus \{0\}$ ,

which implies that (2.1) holds for  $z \in \mathbb{D} \setminus \{0\}$ . To verify (2.1) for  $z = 0$ , observe that  $f$  maps each disk  $D_n$  onto itself. Hence for every  $\zeta \in D_n$  we have

$$\frac{|f(\zeta)|}{|\zeta|} \leq \frac{2^{-n} + 2^{-n-2}}{2^{-n} - 2^{-n-2}} = \frac{5}{3}.$$

Thus (2.1) holds for all  $z \in \mathbb{D}$ .

Now let  $a_n = 2^{-n} + 2^{-(n+2)/\varepsilon}$  and  $b_n = 2^{-n}$ ,  $n \geq 1$ . By the definition of  $f$  we have

$$\begin{aligned} f(a_n) &= 2^{-n} + 2^{(n+2)(1/K-1)} \left( 2^{-(n+2)/\varepsilon} \right)^{1/K} = 2^{-n} + 2^{(n+2)((\varepsilon-1)/\varepsilon K-1)} \\ &= 2^{-n} + 2^{-\alpha(n+2)/\varepsilon} \end{aligned}$$

and  $f(b_n) = 2^{-n}$ . Since

$$\frac{|f(a_n) - f(b_n)|}{|a_n - b_n|^\alpha} = \frac{2^{-\alpha(n+2)/\varepsilon}}{2^{-\alpha(n+2)/\varepsilon}} = 1,$$

it follows that for every  $r > 0$  the mapping  $f$  fails to be in  $c^{0,\alpha}(\mathbb{D}(0,r))$ .  $\square$

Surprisingly, the situation is different for the critical Hölder exponent  $1/K$ . According to the following theorem, one can determine if a  $K$ -quasiconformal mapping belongs to  $c^{0,1/K}$  just by looking at its local modulus of continuity. Its proof uses some ideas from [4].

**THEOREM 2.1.** *Let  $f : \Omega \rightarrow \mathbb{C}$  be a  $K$ -quasiconformal mapping, and let  $E$  be a compact subset of  $\Omega$ . Then  $f \in c^{0,1/K}(E)$  if and only if for every  $z \in E$*

$$(2.2) \quad \lim_{\substack{\zeta \rightarrow z \\ \zeta \in E}} \frac{|f(\zeta) - f(z)|}{|\zeta - z|^{1/K}} = 0.$$

**PROOF.** If  $f \in c^{0,1/K}(E)$ , then (2.2) follows immediately from the definition of  $c^{0,1/K}(E)$ . Conversely, suppose that  $f \notin c^{0,1/K}(E)$ , i.e.  $\|f\|_{E,1/K} > 0$ . Our goal is to prove that (2.2) fails for some  $z \in E$ .

By the definition of  $\|f\|_{E,1/K}$  there exists a sequence  $\delta_j \rightarrow 0$  and points  $a_j, b_j \in E$  such that  $|a_j - b_j| = \delta_j$  and

$$(2.3) \quad |f(a_j) - f(b_j)| = \|f\|_{E,1/K} \delta_j^{1/K} (1 + o(1)), \quad j \rightarrow \infty.$$

Without loss of generality we may assume that  $a_j \rightarrow 0 \in E$ ,  $\overline{\mathbb{D}}(a_j, \delta_j) \subset \mathbb{D}$  for every  $j$ ,  $\overline{\mathbb{D}} \subset \Omega$ , and  $f(0) = 0$ . Since  $f$  is continuous in  $\overline{\mathbb{D}}$ , the domain  $\Omega' = f(\overline{\mathbb{D}})$  is bounded. Let  $R = \text{diam } \Omega'$  be its diameter.

The set  $F_j = f(\overline{\mathbb{D}}(a_j, \delta_j))$  is connected and its diameter is controlled by (2.3). We are going to use this information to estimate its capacity from below. On the other hand, the quasiconformality of  $f$  will lead to an upper bound for the capacity of  $F_j$ . Comparison of the two estimates will show that  $f$  satisfies the hypotheses of [2, Thm.1], which in turn implies that (2.2) fails for  $z = 0$ .

Let us begin by defining the conformal capacity of a compact set  $E$  with respect to a domain  $\Omega \supset E$ .

$$(2.4) \quad \text{cap}(\Omega, E) = \inf \left\{ \int_{\Omega} |\nabla u(z)|^2 d\mathcal{L}^2(z) : u \in C_0^\infty(\Omega) \text{ and } u \geq 1 \text{ on } E \right\},$$

where  $\mathcal{L}^2$  is the 2-dimensional Lebesgue measure. Since  $\Omega' \subset \mathbb{D}(f(a_j), R)$ , it follows from (2.4) that  $\text{cap}(\Omega', F_j) \geq \text{cap}(\mathbb{D}(f(a_j), R), F_j)$ . Observe that  $\mathbb{D}(f(a_j), R) \setminus F_j$  is a doubly-connected domain. There is another well-known conformal invariant associated with such objects, namely, the ring module [7, 5.49]. It can be defined as follows:  $M(\mathbb{D}(f(a_j), R) \setminus F_j) = \log(r_2/r_1)$  if  $\mathbb{D}(f(a_j), R) \setminus F_j$  is conformally equivalent to the circular ring  $\{z : r_1 < |z| < r_2\}$ . The relation between capacity and module is given by

$$\text{cap}(\mathbb{D}(f(a_j), R), F_j) = \frac{2\pi}{M(\mathbb{D}(f(a_j), R) \setminus F_j)}$$

(compare [7, 7.8] with [7, 5.49]).

Since  $F_j$  is connected and contains both  $f(a_j)$  and  $f(b_j)$ , the Grötzsch module theorem [5, p.54] and the estimate (2.10) in [5, p.61] imply

$$M(\mathbb{D}(f(a_j), R) \setminus F_j) \leq \log(4R/|f(a_j) - f(b_j)|).$$

Hence

$$(2.5) \quad \text{cap}(\Omega', F_j) \geq \text{cap}(\mathbb{D}(f(a_j), R), F_j) \geq \frac{2\pi}{\log(4R/|f(a_j) - f(b_j)|)}.$$

Now plug (2.3) into (2.5) to obtain

$$\begin{aligned} \text{cap}(\Omega', F_j) &\geq \frac{2\pi}{\log(4R/\|f\|_{E,1/K}) + K^{-1} \log(1/\delta_j) + o(1)} \\ &= \frac{2\pi K}{\log(1/\delta_j)} \left( 1 + K \frac{\log(4R/\|f\|_{E,1/K})}{\log(1/\delta_j)} + o\left(\frac{1}{\log(1/\delta_j)}\right) \right)^{-1} \\ &= \frac{2\pi K}{\log(1/\delta_j)} \left( 1 - K \frac{\log(4R/\|f\|_{E,1/K})}{\log(1/\delta_j)} + o\left(\frac{1}{\log(1/\delta_j)}\right) \right). \end{aligned}$$

Let  $C = 2\pi K^2 \log(4R/\|f\|_{E,1/K}) + 1$ ; then for all sufficiently large  $j$  we have

$$(2.6) \quad \text{cap}(\Omega', F_j) \geq \frac{2\pi K}{\log(1/\delta_j)} - \frac{C}{(\log(1/\delta_j))^2}.$$

To obtain an upper bound for  $\text{cap}(\Omega', F_j)$ , we proceed as follows. Let  $g : \Omega' \rightarrow \mathbb{D}$  be the inverse of  $f$  and define

$$u(w) = \frac{\log^+ \{(1 - |a_j|)/|g(w) - a_j|\}}{\log\{(1 - |a_j|)/\delta_j\}}$$

for  $w \in \Omega'$ . (Here  $\log^+ t = \max\{\log t, 0\}$ .) It is easy to see that the function  $u$  is Hölder continuous in  $\Omega' \setminus F_j$ ,  $\min\{u, 1\} \in W_0^{1,2}(\Omega')$ , and  $u|_{F_j} \geq 1$ . Therefore,

$$(2.7) \quad \begin{aligned} \text{cap}(\Omega', F_j) &\leq \int_{\Omega' \setminus F_j} |\nabla u(w)|^2 d\mathcal{L}^2(w) \\ &\leq (\log\{(1 - |a_j|)/\delta_j\})^{-2} \int_{\Omega' \setminus F_j} |\nabla \log |g(w) - a_j||^2 d\mathcal{L}^2(w). \end{aligned}$$

At the points where  $\log |g - a_j|$  is differentiable, its gradient can be written in terms of the complex differential operators  $\partial$  and  $\bar{\partial}$ .

$$\begin{aligned} |\nabla \log |g - a_j||^2 &= 4|\partial \log |g - a_j||^2 = |\partial \log(g - a_j) + \partial \log \overline{(g - a_j)}|^2 \\ &= \left| \frac{\partial g}{g - a_j} + \overline{\left( \frac{\bar{\partial} g}{g - a_j} \right)} \right|^2. \end{aligned}$$

Since  $\partial g(w)|_{w=f(z)} = \overline{\partial f(z)} J_f(z)^{-1}$  and  $\bar{\partial} g(w)|_{w=f(z)} = -\bar{\partial} f(z) J_f(z)^{-1}$ , we can express the last integral in (2.7) in terms of the complex dilatation  $\mu = \bar{\partial} f / \partial f$ . Indeed, using notation  $\varphi_j = \arg(z - a_j)$ , we have

$$\begin{aligned} &\int_{\Omega' \setminus F_j} |\nabla \log |g(w) - a_j||^2 d\mathcal{L}^2(w) \\ &= \int_{\mathbb{D} \setminus \bar{\mathbb{D}}(a_j, \delta_j)} \left| \frac{\overline{\partial f(z)}}{(z - a_j) J_f(z)} - \overline{\left( \frac{\bar{\partial} f(z)}{(z - a_j) J_f(z)} \right)} \right|^2 J_f(z) d\mathcal{L}^2(z) \\ &= \int_{\mathbb{D} \setminus \bar{\mathbb{D}}(a_j, \delta_j)} \frac{|\partial f(z) - e^{-2i\varphi_j} \bar{\partial} f(z)|^2}{|\partial f(z)|^2 - |\bar{\partial} f(z)|^2} |z - a_j|^{-2} d\mathcal{L}^2(z) \\ &= \int_{\mathbb{D} \setminus \bar{\mathbb{D}}(a_j, \delta_j)} \frac{|1 - e^{-2i\varphi_j} \mu(z)|^2}{1 - |\mu(z)|^2} |z - a_j|^{-2} d\mathcal{L}^2(z). \end{aligned}$$

This, together with (2.6) and (2.7), yields

$$(2.8) \quad \begin{aligned} &\int_{\mathbb{D} \setminus \bar{\mathbb{D}}(a_j, r_j)} \frac{|1 - e^{-2i\varphi_j} \mu(z)|^2}{1 - |\mu(z)|^2} |z - a_j|^{-2} d\mathcal{L}^2(z) \\ &\geq (\log\{(1 - |a_j|)/\delta_j\})^2 \left( \frac{2\pi K}{\log(1/\delta_j)} - \frac{C}{(\log(1/\delta_j))^2} \right) \end{aligned}$$

for large  $j$ . Since  $a_j \rightarrow 0$ , it follows that

$$(\log\{(1 - |a_j|)/\delta_j\})^2 = (\log(1/\delta_j))^2 + o(\log(1/\delta_j)), \quad j \rightarrow \infty$$

Hence the right-hand side of (2.8) is bounded from below by

$$2\pi K \log(1/\delta_j) - C + o(1), \quad j \rightarrow \infty.$$

For all sufficiently large  $j$  we have

$$\int_{\mathbb{D} \setminus \bar{\mathbb{D}}(a_j, r_j)} \frac{|1 - e^{-2i\varphi_j} \mu(z)|^2}{1 - |\mu(z)|^2} |z - a_j|^{-2} d\mathcal{L}^2(z) \geq 2\pi K \log(1/\delta_j) - C_1,$$

where  $C_1 = C + 1$ . Since

$$\int_{\mathbb{D} \setminus \overline{\mathbb{D}}(a_j, \delta_j)} \frac{d\mathcal{L}^2(z)}{|z - a_j|^2} \leq \int_{\mathbb{D}(a_j, 2) \setminus \overline{\mathbb{D}}(a_j, \delta_j)} \frac{d\mathcal{L}^2(z)}{|z - a_j|^2} = 2\pi \log(2/\delta_j),$$

it follows that

$$\int_{\mathbb{D} \setminus \overline{\mathbb{D}}(a_j, r_j)} \left( K - \frac{|1 - e^{-2i\varphi_j} \mu(z)|^2}{1 - |\mu(z)|^2} \right) |z - a_j|^{-2} d\mathcal{L}^2(z) \leq C_1 + 2\pi K \log 2.$$

Note that the integrand is non-negative because  $|\mu| \leq (K - 1)/(K + 1)$  for  $K$ -quasiconformal mappings. (See also Proposition 2.2 below.) This allows us to pass to the limit  $j \rightarrow \infty$  using Fatou's lemma, thus obtaining

$$(2.9) \quad \int_{\mathbb{D}} \left| K - \frac{|1 - e^{-2i\varphi} \mu(z)|^2}{1 - |\mu(z)|^2} \right| |z|^{-2} d\mathcal{L}^2(z) < \infty,$$

where  $\varphi = \arg z$ . By (2.9) and Proposition 2.2

$$(2.10) \quad \int_{\mathbb{D}} \left| K^{-1} - \frac{|1 + e^{-2i\varphi} \mu(z)|^2}{1 - |\mu(z)|^2} \right| |z|^{-2} d\mathcal{L}^2(z) < \infty.$$

By virtue of (2.9) and (2.10) we can apply Theorem 1 of [2] which asserts that there exists  $A > 0$  such that  $|f(z)|/|z|^{1/K} \rightarrow A$  as  $z \rightarrow 0$ . This leads to the conclusion that (2.2) does not hold at the point  $z = 0$ , because 0 is a non-isolated point of the set  $E$ .  $\square$

**PROPOSITION 2.2.** *If  $\nu \in \mathbb{C}$  and  $K \geq 1$  are such that  $|\nu| \leq (K - 1)/(K + 1)$ , then*

$$0 \leq \frac{|1 + \nu|^2}{1 - |\nu|^2} - \frac{1}{K} \leq K - \frac{|1 - \nu|^2}{1 - |\nu|^2}.$$

**PROOF.** The first inequality follows from

$$\frac{|1 + \nu|^2}{1 - |\nu|^2} \geq \frac{(1 - |\nu|)^2}{1 - |\nu|^2} = \frac{1 - |\nu|}{1 + |\nu|} \geq \frac{1}{K},$$

while the second one follows from

$$\frac{|1 + \nu|^2}{1 - |\nu|^2} + \frac{|1 - \nu|^2}{1 - |\nu|^2} = 2 \frac{1 + |\nu|^2}{1 - |\nu|^2} \leq 2 \frac{(K + 1)^2 + (K - 1)^2}{(K + 1)^2 - (K - 1)^2} = K + \frac{1}{K}.$$

$\square$

It was recently proved [4] that for a  $K$ -quasiconformal mapping  $f$  the limit  $\lim_{\zeta \rightarrow z} |f(\zeta) - f(z)|/|\zeta - z|^{1/K}$  exists at every point  $z$  in its domain of definition. At the points where this limit is positive, the linear dilatation of  $f$

$$H_f(z) = \limsup_{r \rightarrow 0} \sup_{z_1, z_2} \left\{ \frac{|f(z_1) - f(z)|}{|f(z_2) - f(z)|} : |z_1 - z| = r = |z_2 - z| \right\}.$$

is evidently equal to 1. Thus we arrive at the following corollary.

**COROLLARY 2.1.** *For a  $K$ -quasiconformal mapping  $f : \Omega \rightarrow \mathbb{C}$ , one of the following statements is true: (a)  $f \in \mathcal{C}_{\text{loc}}^{0, 1/K}(\Omega)$ ; (b)  $H_f(z) = 1$  for some  $z \in \Omega$ .*

It is likely that the following quantitative version of Theorem 2.1 is true.

CONJECTURE 2.1. *Let  $f : \Omega \rightarrow \mathbb{C}$  be a  $K$ -quasiconformal mapping, and let  $E$  be a compact subset of  $\Omega$ . Then*

$$(2.11) \quad \|f\|_{E,1/K} = \sup_{z \in E} \lim_{\substack{\zeta \rightarrow z \\ \zeta \in E}} \frac{|f(\zeta) - f(z)|}{|\zeta - z|^{1/K}}.$$

It is obvious that the right-hand side of (2.11) does not exceed  $\|f\|_{E,1/K}$ , but the reverse inequality seems much harder to prove.

### 3. Concluding remarks

As Corollary 2.1 indicates, there is a tight connection between the modulus of continuity of a quasiconformal mapping and its linear dilatation. Recall that the linear dilatation  $H_f$  of a  $K$ -quasiconformal mapping  $f$  can exceed  $K$  (see [5] or [6], where the sharp upper bound for  $H_f$  is found). On the other hand,  $H_f(z) \leq K$  if  $f$  has a non-zero derivative at  $z$  [6]. Also,  $H_f(z) = 1$  if the upper limit

$$\limsup_{\zeta \rightarrow z} \frac{|f(\zeta) - f(z)|}{|\zeta - z|^{1/K}}$$

is strictly positive [4]. This naturally leads to the following question: what is the exact value of

$$H(\alpha) = \sup \left\{ H_f(z) : f \text{ is } K\text{-qc and } \limsup_{\zeta \rightarrow z} \frac{|f(\zeta) - f(z)|}{|\zeta - z|^\alpha} > 0 \right\}$$

for  $\alpha$  between  $1/K$  and  $K$ ? The function  $H$  increases from  $H(1/K) = 1$  to  $H(K) = \lambda(K)$  (as defined in [5, II(6.4)] or [6, (11)]). Apparently, none of its intermediate values are known, although it seems likely that  $H(1) = K$ .

Note that the authors of [6] use a symmetrization argument to show that

$$(3.1) \quad \sup_f H_f(z) = \sup_f \limsup_{\zeta \rightarrow 0} \frac{|f(z + \zeta) - f(z)|}{|f(z - \zeta) - f(z)|},$$

where the supremum is taken over all  $K$ -quasiconformal mappings of the plane. The identity (3.1) is one of the crucial points in [6], and it is not clear if it still holds when the supremum on both sides is taken only over those  $K$ -quasiconformal mappings for which

$$\limsup_{\zeta \rightarrow z} \frac{|f(\zeta) - f(z)|}{|\zeta - z|^\alpha} > 0.$$

### Acknowledgements

The author is grateful to Albert Baernstein and David Opěla for several helpful discussions. The referee's comments helped to improve the paper considerably.

**References**

- [1] L. V. Ahlfors, *Lectures on quasiconformal mappings*, Van Nostrand, New York, 1966.
- [2] M. Brakalova and J. A. Jenkins, *On the local behavior of certain homeomorphisms. II*, J. Math. Sci., New York **95** (1999), no.3, 2178–2184.
- [3] T. Iwaniec and G. Martin, *Geometric function theory and nonlinear analysis*, Oxford Univ. Press, New York, 2001.
- [4] L. V. Kovalev, *Quasiregular mappings of maximal local modulus of continuity*, to appear in Ann. Acad. Sci. Fenn. Math. **29** (2004), 211–222.
- [5] O. Lehto and K. I. Virtanen, *Quasiconformal mappings in the plane*, 2nd ed. Springer-Verlag, Berlin-Heidelberg-New York, 1973.
- [6] O. Lehto, K. I. Virtanen and J. Väisälä, *Contributions to the distortion theory of quasiconformal mappings*, Ann. Acad. Sci. Fenn. Math. **273** (1959), 1–14.
- [7] M. Vuorinen, *Conformal geometry and quasiregular mappings*, Lecture Notes in Math. Vol. 1319. Springer-Verlag, Berlin-Heidelberg-New York, 1988.

Department of Mathematics  
Washington University  
St. Louis, MO 63130, USA  
lkovalev@math.wustl.edu

(Received 07 09 2003)