

A NOTE ON THE FEKETE–SZEGÖ PROBLEM FOR CLOSE-TO-CONVEX FUNCTIONS WITH RESPECT TO CONVEX FUNCTIONS

Bogumiła Kowalczyk, Adam Lecko, and H. M. Srivastava

ABSTRACT. We discuss the sharpness of the bound of the Fekete–Szegő functional for close-to-convex functions with respect to convex functions. We also briefly consider other related developments involving the Fekete–Szegő functional $|a_3 - \lambda a_2^2|$ ($0 \leq \lambda \leq 1$) as well as the corresponding Hankel determinant for the Taylor–Maclaurin coefficients $\{a_n\}_{n \in \mathbb{N} \setminus \{1\}}$ of normalized univalent functions in the open unit disk \mathbb{D} , \mathbb{N} being the set of positive integers.

1. Introduction

A classical problem in geometric function theory of complex analysis, which was settled by Fekete and Szegő [4], is to find for each $\lambda \in [0, 1]$ the maximum value of the coefficient functional $\Phi_\lambda(f)$ given by

$$(1.1) \quad \Phi_\lambda(f) := |a_3 - \lambda a_2^2|$$

over the class \mathcal{S} of univalent functions f in the open unit disk

$$\mathbb{D} := \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$$

of the following normalized form (see, for details, [5, 22, 24]):

$$(1.2) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (z \in \mathbb{D}).$$

By applying the Loewner method, Fekete and Szegő [4] proved that

$$\max_{f \in \mathcal{S}} \Phi_\lambda(f) = \begin{cases} 1 + 2 \exp\left(-\frac{2\lambda}{1-\lambda}\right) & (0 \leq \lambda < 1) \\ 1 & (\lambda = 1). \end{cases}$$

2010 *Mathematics Subject Classification*: Primary 30C45.

Key words and phrases: analytic functions, convex functions, Fekete–Szegő problem, Hankel determinant, Taylor–Maclaurin coefficients, close-to-convex functions with respect to a convex function, Carathéodory class, Schwarz functions.

Communicated by Stevan Pilipović.

For various compact subclasses \mathcal{F} of the class \mathcal{A} of all analytic functions f in \mathbb{D} of the form (1.2), as well as with λ being an arbitrary real or complex number, many authors computed

$$(1.3) \quad \max_{f \in \mathcal{F}} \Phi_\lambda(f)$$

or calculated the upper bound of (1.3) (see, e.g., [2, 8, 11, 21]).

Let \mathcal{S}^* denote the class of *starlike* functions, that is, $f \in \mathcal{S}^*$ if

$$f \in \mathcal{A} \quad \text{and} \quad \operatorname{Re} \left(\frac{zf'(z)}{f(z)} \right) > 0 \quad (z \in \mathbb{D}).$$

Given $\delta \in (-\frac{\pi}{2}, \frac{\pi}{2})$ and $g \in \mathcal{S}^*$, let $\mathcal{C}_\delta(g)$ denote the class of functions called *close-to-convex with argument δ with respect to g* , that is, the class of all functions $f \in \mathcal{A}$ such that

$$(1.4) \quad \operatorname{Re} \left(e^{i\delta} \frac{zf'(z)}{g(z)} \right) > 0 \quad (z \in \mathbb{D}).$$

We also suppose that, given $g \in \mathcal{S}^*$, $\mathcal{C}(g) := \bigcup_{g \in \mathcal{S}^*} \mathcal{C}_\delta(g)$ and that, given $\delta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, $\mathcal{C}_\delta := \bigcup_{g \in \mathcal{S}^*} \mathcal{C}_\delta(g)$. Let

$$\mathcal{C} := \bigcup_{\delta \in (-\frac{\pi}{2}, \frac{\pi}{2})} \bigcup_{g \in \mathcal{S}^*} \mathcal{C}_\delta(g)$$

denote the class of *close-to-convex* functions (see, for details, [20, pp. 184–185], [6, 10]).

For the whole class \mathcal{C} , the sharp bound of the Fekete–Szegő coefficient functional Φ_λ for $\lambda \in [0, 1]$, given by (1.1), was calculated by Koepf [13] who extended the earlier result for the class \mathcal{C}_0 and for $\lambda \in \mathbb{R}$ due to Keogh and Merkes [11], namely, it holds

$$\max_{f \in \mathcal{C}} \Phi_\lambda(f) = \max_{f \in \mathcal{C}_0} \Phi_\lambda(f) = \begin{cases} |3 - 4\lambda| & (\lambda \in (-\infty, \frac{1}{3}] \cup [1, \infty)) \\ \frac{1}{3} + \frac{4}{9\lambda} & (\lambda \in [\frac{1}{3}, \frac{2}{3}]) \\ 1 & (\lambda \in [\frac{2}{3}, 1]). \end{cases}$$

For various subclasses of the class of close-to-convex functions, the problem to estimate the coefficient functional Φ_λ is continued in several subsequent works (see, for details, [9, 12, 14–16]). Some interesting and important subclasses of the class \mathcal{C} are the classes \mathcal{C}_δ^c and \mathcal{C}^c , which are defined below.

Let \mathcal{S}^c denote the class of *convex* functions, that is, $f \in \mathcal{S}^c$ if

$$f \in \mathcal{A} \quad \text{and} \quad \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > 0 \quad (z \in \mathbb{D}).$$

Since $\mathcal{S}^c \subsetneq \mathcal{S}^*$, the class $\mathcal{C}_\delta^c := \bigcup_{g \in \mathcal{S}^c} \mathcal{C}_\delta(g)$ is a proper subclass of the class \mathcal{C}_δ and the class

$$\mathcal{C}^c := \bigcup_{\delta \in (-\frac{\pi}{2}, \frac{\pi}{2})} \bigcup_{g \in \mathcal{S}^c} \mathcal{C}_\delta(g)$$

is a proper subclass of the class \mathcal{C} .

The class \mathcal{C}_0^c was defined by Abdel-Gawad and Thomas [1]. The class \mathcal{C}^c of *close-to-convex functions with respect to convex functions* was introduced by Srivastava, Mishra and Das [23]. In both of these cited papers, the authors (Abdel-Gawad and Thomas [1] and Srivastava, Mishra and Das [23]) considered the coefficient functional Φ_λ with $\lambda \in [0, 1]$ also. In fact, in Srivastava, Mishra and Das [23] extended, for the class \mathcal{C}^c , the earlier result of Abdel-Gawad and Thomas [1] for the class \mathcal{C}_0^c . However, in each of the above-cited papers, the proof for the sharpness of the bound in (1.3) for $\lambda \in (\frac{2}{3}, 1]$ was proposed incorrectly as $5/6$.

This note is motivated essentially by the earlier papers [1] and [23]. The main purpose of our investigation here is to discuss such sharpness results for the bound in (1.3). We also provide a rather brief consideration of other related developments involving the Fekete–Szegő functional $|a_3 - \lambda a_2^2|$ ($0 \leq \lambda \leq 1$) in (1.1) as well as the corresponding Hankel determinant for the Taylor–Maclaurin coefficients $\{a_n\}_{n \in \mathbb{N} \setminus \{1\}}$ of normalized univalent functions of the form (1.2).

2. Main Observation

As we remarked in Section 1, in both of the afore cited papers [1, 23], the upper bounds of the Fekete–Szegő coefficient functional Φ_λ ($0 \leq \lambda \leq 1$) for the classes \mathcal{C}_0^c and \mathcal{C}^c , were computed. In fact, Theorems 5 and 6 of Srivastava, Mishra and Das [23] state that the following sharp inequality

$$(2.1) \quad \max_{f \in \mathcal{C}^c} \Phi_\lambda(f) \leq \frac{5}{6} \quad (\lambda \in [\frac{2}{3}, 1])$$

holds true and that this result is the same as in [1] for the class \mathcal{C}_0^c (a part of Theorem 3). However, the assertion that the extremal function, for which the equality in (2.1) is satisfied when $\lambda \in (\frac{2}{3}, 1]$, belongs to \mathcal{C}^c is incorrect. Indeed, here in this section, we note that the above-cited papers [1, 23] contain a statement to the effect that the equality in (2.1) is attained by a function $f \in \mathcal{A}$ given by

$$(2.2) \quad \frac{zf'(z)}{h(z)} = \frac{1 + \omega(z)}{1 - \omega(z)} \quad (z \in \mathbb{D}),$$

where $h \in \mathcal{S}^c$ is of the form

$$(2.3) \quad h(z) = z + \sum_{n=2}^{\infty} b_n z^n \quad (z \in \mathbb{D}; b_2 = b_3 := 1)$$

and ω is a function of the form

$$(2.4) \quad \omega(z) = \sum_{n=1}^{\infty} \beta_n z^n \quad (z \in \mathbb{D})$$

with

$$(2.5) \quad \beta_1 := \frac{2 - 3\lambda}{6\lambda} \pm i \frac{\sqrt{6\lambda - 4}}{6\lambda} \quad \text{and} \quad \beta_2 := 1 - \beta_1^2.$$

Unfortunately, however, ω is not a Schwarz function for $\lambda \in (\frac{2}{3}, 1]$. We recall here that a Schwarz function means an analytic self-mapping of \mathbb{D} with $\omega(0) := 0$. Let us

denote the class of Schwarz functions by \mathcal{B}_0 . In order to see that $\omega \notin \mathcal{B}_0$, we verify (by straightforward computation) that, for $\lambda \in (\frac{2}{3}, 1]$, the following inequality:

$$(2.6) \quad |\beta_2| \leq 1 - |\beta_1|^2$$

is false, so a necessary condition for ω to be in \mathcal{B}_0 (see, for example, [5, Vol. II, p. 78]) does not hold true. Alternatively, in order to get a contradiction, we suppose that ω with its coefficients in (2.5) is a Schwarz function. Thus, clearly, (2.6) holds true. Hence we find from (2.5) that $1 - |\beta_1|^2 \geq |\beta_2| = |1 - \beta_1^2| \geq 1 - |\beta_1|^2$. Thus we have $|1 - \beta_1^2| = 1 - |\beta_1|^2$ and, therefore, $\beta_1 = |\beta_1|$ or $\beta_1 = -|\beta_1|$. This means that β_1 is a real number, which by (2.5) is possible only for $\lambda = \frac{2}{3}$. Consequently, for $\lambda \in (\frac{2}{3}, 1]$, the function ω with its coefficients in (2.5) does not belong to \mathcal{B}_0 . So, in light of (2.2), it does not follow that f is in \mathcal{C}^c or in \mathcal{C}_0^c .

Equivalently, let

$$(2.7) \quad p(z) := \frac{1 + \omega(z)}{1 - \omega(z)} \quad (z \in \mathbb{D}),$$

where ω is as given above. Then

$$(2.8) \quad p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n \quad (z \in \mathbb{D}),$$

where, in view of (2.7), (2.4) and (2.5), we have $c_1 = 2\beta_1$ and $c_2 = 2(\beta_2 + \beta_1^2) = 2$. We observe further that, for $\lambda \in (\frac{2}{3}, 1]$, the function p does not belong to the Carathéodory class. We recall here that the Carathéodory class, denoted as \mathcal{P} , contains analytic functions p of the form (2.8) with a positive real part. In order to see that $p \notin \mathcal{P}$, we verify for $\lambda \in (\frac{2}{3}, 1]$ that the inequality $|c_2 - c_1^2/2| \leq 2 - |c_1|^2/2$, is false, which happens to be a necessary condition for p to be in the class \mathcal{P} (see, for example, [22, p. 166]).

3. Concluding remarks and further developments

By means of Theorem 3 of Abdel-Gawad and Thomas [1], Theorems 1 to 4 of Srivastava, Mishra and Das [23], and in light of our observation in Section 2, we arrive at the following result.

THEOREM 1. *Each of the following assertions holds true:*

$$(3.1) \quad \max_{f \in \mathcal{C}^c} \Phi_\lambda(f) = \max_{f \in \mathcal{C}_0^c} \Phi_\lambda(f) = \begin{cases} \frac{5}{3} - \frac{9\lambda}{4} & (\lambda \in [0, \frac{2}{9}]) \\ \frac{2}{3} + \frac{1}{9\lambda} & (\lambda \in [\frac{2}{9}, \frac{2}{3}]) \end{cases}$$

$$(3.2) \quad \max_{f \in \mathcal{C}^c} \Phi_\lambda(f) \leq \frac{5}{6} \quad (\lambda \in (\frac{2}{3}, 1]).$$

REMARK 1. The sharpness of the inequality in (3.2) for the classes \mathcal{C}^c and \mathcal{C}_0^c is an *open problem*.

We now note that, by Loewner Theorem (see, for example, [5, Vol. I, p. 1127]), the function $h \in \mathcal{S}^c$ of the form (2.3) (with $b_2 = b_3 := 1$) is uniquely determined, that is, $h(z) = \frac{z}{1-z} = \sum_{n=1}^{\infty} z^n$ ($z \in \mathbb{D}$). Then (1.4) with $g := h$ is of the form

$$(3.3) \quad \operatorname{Re}(e^{i\delta}(1-z)f'(z)) > 0 \quad (z \in \mathbb{D})$$

and defines the class $\mathcal{C}_\delta(h)$, and further the class $\mathcal{C}(h)$. For the first time, the inequality in (3.3), treated as the univalence criterion, was distinguished explicitly in [20, p. 185]. For the class $\mathcal{C}(h)$, the upper bound of the Fekete–Szegő coefficient functional Φ_λ for $\lambda \in \mathbb{R}$ was recently obtained in [14], where the following result was proven.

THEOREM 2. *It is asserted that*

$$(3.4) \quad \max_{f \in \mathcal{C}(h)} \Phi_\lambda(f) \leq \begin{cases} \left| \frac{1}{3} - \frac{1}{4}\lambda \right| + \frac{2}{3} |2 - 3\lambda| & (\lambda \in (-\infty, \frac{2}{9}] \cup [\frac{10}{9}, \infty)) \\ \frac{1}{12} \cdot \frac{(2-3\lambda)^2}{2-|2-3\lambda|} + \left| \frac{1}{3} - \frac{1}{4}\lambda \right| + \frac{2}{3} & (\lambda \in [\frac{2}{9}, \frac{10}{9}]). \end{cases}$$

For each $\lambda \in (-\infty, \frac{2}{9}] \cup [\frac{4}{3}, \infty)$, the inequality is sharp and the equality in (2) is attained by a function in $\mathcal{C}_0(h)$.

REMARK 2. For $\lambda \in (-\infty, \frac{2}{9}] \cup [\frac{4}{3}, \infty)$, we can rewrite (3.4) as the following corollary.

COROLLARY 1. *The following assertion holds true:*

$$(3.5) \quad \max_{f \in \mathcal{C}(h)} \Phi_\lambda(f) = \begin{cases} \left| \frac{5}{3} - \frac{9\lambda}{4} \right| & (\lambda \in (-\infty, \frac{2}{9}] \cup [\frac{4}{3}, \infty)) \\ \frac{2}{3} + \frac{1}{9\lambda} & (\lambda \in [\frac{2}{9}, \frac{2}{3}]). \end{cases}$$

REMARK 3. For $\lambda \in [0, \frac{2}{3}]$, the result (3.5) asserted by Corollary 3.5 coincides with (3.1). Thus, naturally, Theorem 1 and Theorem 2 yield Corollary 2 below.

COROLLARY 2. *Each of the following assertions holds true:*

$$\begin{aligned} \max_{f \in \mathcal{C}(h)} \Phi_\lambda(f) &= \max_{f \in \mathcal{C}_0^c} \Phi_\lambda(f) = \max_{f \in \mathcal{C}^c} \Phi_\lambda(f) \quad (\lambda \in [0, \frac{2}{3}]), \\ \max_{f \in \mathcal{C}(h)} \Phi_\lambda(f) &\leq \frac{9\lambda^2 - 30\lambda + 26}{6(4 - 3\lambda)} \leq \frac{5}{6} \quad (\lambda \in (\frac{2}{3}, 1]). \end{aligned}$$

REMARK 4. The maximum of Φ_λ for $\lambda \in [0, \frac{2}{3}]$, over the class \mathcal{C}^c of close-to-convex functions with respect to convex functions and over its subclass $\mathcal{C}(h)$ of close-to-convex functions with respect to convex function h , are identical.

REMARK 5. The sharpness of the inequality in (3.4) for $\lambda \in (\frac{2}{9}, \frac{4}{3})$ is an *open problem*.

REMARK 6. We reiterate the fact that the Fekete–Szegő coefficient functional $|a_3 - \lambda a_2^2|$ is well known for its rich history in geometric function theory. Its origin was in the disproof by Fekete and Szegő [4] of the 1933 conjecture of Littlewood and Paley that the coefficients of odd univalent functions are bounded by unity (see, for details, [4]). The λ -generalized Fekete–Szegő coefficient functional $|a_3 - \lambda a_2^2|$ has since received great attention, particularly in connection with many subclasses of the class \mathcal{S} of normalized analytic and univalent functions. On the other hand, in the year 1976, Noonan and Thomas [17] defined the q th Hankel determinant of

the function f in (1.2) by

$$H_q(n) = \begin{vmatrix} a_n & a_{n+1} & \cdots & a_{n+q-1} \\ a_{n+1} & a_{n+2} & \cdots & a_{n+q} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n+q-1} & a_{n+q} & \cdots & a_{n+2q-2} \end{vmatrix} \quad (n, q \in \mathbb{N}; a_1 := 1).$$

The determinant $H_q(n)$ has also been considered by several other authors. For example, Noor [18] determined the rate of growth of $H_q(n)$ as $n \rightarrow \infty$ for functions f given by (1.1) with bounded boundary. In particular, sharp upper bounds on $H_2(2)$ were obtained in the recent works [7, 18] for different classes of functions. We note, in particular, that

$$H_2(1) = \begin{vmatrix} a_1 & a_2 \\ a_2 & a_3 \end{vmatrix} = a_3 - a_2^2 \quad \text{and} \quad H_2(2) = \begin{vmatrix} a_2 & a_3 \\ a_3 & a_4 \end{vmatrix} = a_2 a_4 - a_3^2.$$

The Hankel determinant $H_2(1) = a_3 - a_2^2$ is the *classical* Fekete–Szegő coefficient functional. The upper bounds of $H_2(2)$ for some specific analytic function classes were discussed quite recently by Deniz et al. [3] (see also [19]).

References

1. H. R. Abdel-Gawad, D. K. Thomas, *A subclass of close-to-convex functions*, Publ. Inst. Math., Nouv. Sér. **49**(63) (1991), 61–66.
2. B. Bhowmik, S. Ponnusamy, K. J. Wirths, *On the Fekete–Szegő problem for concave univalent functions*, J. Math. Anal. Appl. **373** (2011), 432–438.
3. E. Deniz, M. Çağlar, H. Orhan, *Second Hankel determinant for bi-starlike and bi-convex functions of order β* , Appl. Math. Comput. **271** (2015), 301–307.
4. M. Fekete, G. Szegő, *Eine Bemerkung über ungerade schlichte Funktionen*, J. Lond. Math. Soc. **8** (1933), 85–89.
5. A. W. Goodman, *Univalent Functions*, Mariner, Tampa, Florida, 1983.
6. A. W. Goodman, E. B. Saff, *On the definition of close-to-convex function*, Int. J. Math. Math. Sci. **1** (1978), 125–132.
7. T. Hayami, S. Owa, *Generalized Hankel determinant for certain classes*, Int. J. Math. Anal. **52** (2010), 2473–2585.
8. Z. J. Jakubowski, *Sur le maximum de la fonctionnelle $|A_3 - \alpha A_2^2|$ ($0 \leq \alpha < 1$) dans la famille de fonctions F_M* , Bull. Soc. Sci. Lett. Łódź **13**(1) (1962), 1–19.
9. S. Kanas, A. Lecko, *On the Fekete–Szegő problem and the domain of convexity for a certain class of univalent functions*, Zesz. Nauk. Politech. Rzeszowskiej, Folia Sci. Univ. Tech. Resoviensis **73** (1990), 49–57.
10. W. Kaplan, *Close to convex schlicht functions*, Mich. Math. J. **1** (1952), 169–185.
11. F. R. Keogh, E. P. Merkes, *A coefficient inequality for certain classes of analytic functions*, Proc. Am. Math. Soc. **20**, (1969), 8–12.
12. Y. C. Kim, J. H. Choi, T. Sugawa, *Coefficient bounds and convolution properties for certain classes of close-to-convex functions*, Proc. Japan Acad., Ser. A **76**(6) (2000), 95–98.
13. W. Koepf, *On the Fekete–Szegő problem for close-to-convex functions*, Proc. Am. Math. Soc. **101** (1987), 89–95.
14. B. Kowalczyk, A. Lecko, *Fekete–Szegő problem for a certain subclass of close-to-convex functions*, Bull. Malays. Math. Sci. Soc. (2) **38** (2015), 1393–1410.
15. ———, *Fekete–Szegő problem for close-to-convex functions with respect to a certain convex function dependent on a real parameter*, Front. Math. China **11** (2016), 1471–1500.
16. R. R. London, *Fekete–Szegő inequalities for close-to-convex functions*, Proc. Am. Math. Soc. **117** (1993), 947–950.

17. J. W. Noonan, D. K. Thomas, *On the second Hankel determinant of areally mean p -valent functions*, Trans. Am. Math. Soc. **223** (1976), 337–346.
18. K. I. Noor, *Hankel determinant problem for the class of functions with bounded boundary rotation*, Rev. Roum. Math. Pures Appl. **28** (1983), 731–739.
19. H. Orhan, N. Magesh, J. Yamini, *Bounds for the second Hankel determinant of certain bi-univalent functions*, Turk. J. Math. **40** (2016), 679–687.
20. S. Ozaki, *On the theory of multivalent functions*, Sci. Rep. Tokyo Bunrika Daigaku, Sect. A **2** (1935), 167–188.
21. A. Pfluger, *The Fekete–Szegő inequality for complex parameter*, Complex Variables, Theory Appl. **7** (1986), 149–160.
22. Ch. Pommerenke, *Univalent Functions*, Vandenhoeck and Ruprecht, Göttingen, 1975.
23. H. M. Srivastava, A. K. Mishra, M. K. Das, *The Fekete–Szegő problem for a Subclass of Close-to-Convex Functions*, Complex Variables, Theory Appl. **44** (2001), 145–163.
24. H. M. Srivastava, S. Owa, *Current Topics in Analytic Function Theory*, World Scientific, Singapore, New Jersey, London and Hong Kong, 1992.

Faculty of Mathematics and Computer Science
Department of Complex Analysis
University of Warmia and Mazury
Olsztyn
Poland
b.kowalczyk@matman.uwm.edu.pl
alecko@matman.uwm.edu.pl

(Received 18 06 2016)

(Revised 12 11 2016)

Department of Mathematics and Statistics
University of Victoria
Victoria
Canada;
China Medical University
Taichung
Taiwan
Republic of China
harimsri@math.uvic.ca