PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 101(115) (2017), 169–182 https://doi.org/10.2298/PIM1715169K

ABSOLUTE AND UNIFORM CONVERGENCE OF SPECTRAL EXPANSION OF THE FUNCTION FROM THE CLASS $W_p^1(G)$, $p > 1$, IN EIGENFUNCTIONS OF THIRD ORDER DIFFERENTIAL OPERATOR

V. M. Kurbanov and E. B. Akhundova

Abstract. We study an ordinary differential operator of third order and absolute and uniform convergence of spectral expansion of the function from the class $W_p^1(G)$, $G = (0,1)$, $p > 1$, in eigenfunctions of the operator. Uniform convergence rate of this expansion is estimated.

1. Basic notion and formulation of results

Consider on the interval $G = (0,1)$ a formal differential operator

$$Lu = u^{(3)} + p_1(x)u^{(2)} + p_2(x)u^{(1)} + p_3(x)u$$

with summable complex valued coefficients $p_1(x) \in L_2(G)$, $p_l(x) \in L_1(G)$, $l = 2,3$.

Denote by $D(G)$ a class of functions absolutely continuous together with own derivatives to second order, inclusively on the closed interval $\bar{G} = [0,1]$. Following [**1**], under the eigenfunction of the operator L responding to the eigenvalue λ, we understand any not identically equal to zero function $u(x) \in D(G)$ satisfying almost everywhere in G the equation $Lu + \lambda u = 0$.

Let $\{u_n(x)\}_{n=1}^{\infty}$ be a complete orthonormalized in $L_2(G)$ system consisting of eigenfunctions of the operator L, and $\{\lambda_n\}_{n=1}^{\infty}$ be an appropriate system of eigenvalues, moreover $\operatorname{Re}\lambda_n = 0$ (it is supposed that the coefficients of the operator L admit the existence of such a system $\{u_n(x)\}_{n=1}^{\infty}$). Under rather smooth coefficients the existence of such systems follows from the monograph [**2**])

Denote by μ_n the number $(\mp i\lambda_n)^{1/3}$ for $\pm \operatorname{Im}\lambda_n \geqslant 0$. We will say that the function $f(x)$ belongs to $W_p^1(G), 1 \leqslant p \leqslant \infty$ if $f(x)$ is absolutely continuous on $\bar{G}$ and $f'(x) \in L_p(G)$.

Introduce a partial sum of spectral expansion of the function $f(x) \in W_p^1(G)$ in the system $\{u_n(x)\}_{n=1}^{\infty}$ by $\sigma_\nu(x,f) = \sum_{\mu_n \leqslant \nu} f_n u_n(x)$, $\nu > 0$, where $f_n =$

2010 *Mathematics Subject Classification:* 34L10; 42A20.
Key words and phrases: absolute convergence; uniform convergence; eigenfunctions; spectral expansion.
Communicated by Gradimir Milovanović.

$(f,u_n)=\int_0^1 f(x)\overline{u_n(x)}dx$. Denote $R_\nu(x,f)=f(x)-\sigma_\nu(x,f)$. We prove the following theorems.

THEOREM 1.1. *Let $p_1(x)\equiv 0$, $p_l(x)\in L_1(G)$, $l=2,3$; $f(x)\in W_p^1(G)$, $p>1$, and the following condition be fulfilled*

$$\left|f(x)\overline{u_n^{(2)}(x)}\right|_0^1\leqslant C(f)\mu_n^\alpha\|u_n\|_\infty,\quad 0\leqslant\alpha<2,\quad \mu_n\geqslant 1, \tag{1.1}$$

where $C(f)>0$ is a constant dependent on the function $f(x)$. Then the spectral expansion of the function $f(x)$ in the system $\{u_n(x)\}_{n=1}^\infty$ converges absolutely and uniformly on $\bar{G}=[0,1]$, and we have the estimation

$$\begin{aligned}\sup_{x\in\bar{G}}|R_\nu(x,f)|&\\ \leqslant \text{const}&\left\{C(f)\nu^{\alpha-2}+\nu^{-\beta}\|f'\|_p+\nu^{-1}(\|f\|_\infty+\|f'\|_1)\sum_{r=2}^{3}\nu^{2-r}\|p_r\|_1\right\},\end{aligned} \tag{1.2}$$

where $\beta=\min\left\{\frac{1}{2},\frac{1}{q}\right\}$, $\frac{1}{p}+\frac{1}{q}=1$, $\nu\geqslant 2$, const *is independent of $f(x)$, and $\|\cdot\|_p=\|\cdot\|_{L_p(G)}$.*

COROLLARY 1.1. *If in Theorem* 1.1*, the function $f(x)$ satisfies the condition $f(0)=f(1)=0$, then obviously* (1.1) *is fulfilled and ve have the estimation*

$$\sup_{x\in\bar{G}}|R_\nu(x,f)|\leqslant\text{const}\ \nu^{-\beta}\|f'\|_p,\quad \nu\geqslant 2;$$

if $C(f)=0$ or $0\leqslant\alpha<2-\beta$, then $\sup_{x\in\bar{G}}|R_\nu(x,f)|=o(\nu^{-\beta})$, $\nu\to+\infty$.

THEOREM 1.2. *Let $p_1(x)\in L_2(G)$, $p_l(x)\in L_1(G)$, $l=2,3$; $f(x)\in W_2^1(G)$ and condition* (1.1) *be fulfilled. Then the spectral expansion of the function $f(x)$ in system $\{u_n(x)\}_{n=1}^\infty$ converges absolutely and uniformly on $\bar{G}=[0,1]$, and we have the estimation*

$$\begin{aligned}\sup_{x\in\bar{G}}|R_\nu(x,f)|\leqslant\text{const}&\left\{C(f)\nu^{\alpha-2}+\nu^{-\frac{1}{2}}(\|f\,\overline{p_1}\|_2+\|f'\|_2)\right.\\ &\left.+\nu^{-1}\|f\|_\infty\sum_{r=2}^{3}\nu^{2-r}\|p_r\|_1\right\},\quad \nu\geqslant 2.\end{aligned}$$

COROLLARY 1.2. *If in Theorem* 1.2*, $C(f)=0$ or $0\leqslant\alpha<3/2$, then*

$$\sup_{x\in\bar{G}}|R_\nu(x,f)|=o\big(\nu^{-\frac{1}{2}}\big),\quad \nu\to+\infty.$$

THEOREM 1.3. *Let $p_1(x)\in L_2(G), p_l(x)\in L_1(G)$, $l=2,3$; $f(x)\in W_p^1(G)$, $1<p<2$, condition* (1.1) *be fulfilled, and the system $\{u_n(x)\}_{n=1}^\infty$ be uniformly bounded. Then the spectral expansion of the function $f(x)$ in system $\{u_n(x)\}_{n=1}^\infty$ converges absolutely and uniformly on $\bar{G}$, and we have the estimation*

$$\begin{aligned}\sup_{x\in\bar{G}}|R_\nu(x,f)|\leqslant\text{const}&\left\{C(f)\nu^{\alpha-2}+\nu^{-\frac{1}{2}}\|f\overline{p_1}\|_2\right.\\ &\left.+\nu^{-\frac{1}{q}}\|f'\|_p+\nu^{-1}\|f\|_\infty\sum_{r=2}^{3}\nu^{2-r}\|p_r\|_1\right\},\quad \mu\geqslant 2,\end{aligned}$$

where $p^{-1}+q^{-1}=1$.

COROLLARY 1.3. *If in Theorem* 1.3, $C(f)=0$ *or* $0\leqslant\alpha<2-q^{-1}$, *then*

$$\sup_{x\in\bar{G}}|R_\nu(x,f)|=o\big(\nu^{-\frac{1}{q}}\big),\quad \nu\to+\infty.$$

Note that such results for the Schrodinger operator $L_1=-\frac{d^2}{dx^2}+q(x)$ were obtained in [**3**–**7**]. In the case $f(x)\in W_1^1(G)$ the absolute and uniform convergence of spectral expansion in eigenfunctions of a third order operator was studied in [**8**].

2. Some auxiliary statements

To prove, the formulated results it is necessary to estimate the Fourier coefficients of the function $f(x)$ in the system $\{u_n(x)\}_{n=1}^{\infty}$. To this end we established the representation for the eigenfunction $u_n(x)$. Introduce the notation

$$K(z)=\sum_{j=1}^{3}\omega_j\exp(i\omega_j(\operatorname{sgn}\operatorname{Im}\lambda_n)z),$$

$$M(u_n(\xi))=\frac{1}{3}\mu_n^{-2}\sum_{r=1}^{3}p_r(\xi)u_n^{(3-r)}(\xi),$$

$$X_j^{\pm}(x)=\frac{1}{3}\mu_n^{-2}\sum_{m=0}^{2}(\pm i\mu_n)^m\omega_j^{m+1}u_n^{(2-m)}(x),$$

where $\omega_1=-1$, $\omega_2=\exp(-i\pi/3)$, $\omega_3=\exp(i\pi/3)$.

LEMMA 2.1. *For the eigenfunction* $u_n(x)$, *the following representations are valid* ($\lambda_n\neq0$, $l=\overline{0,2}$)

$$u_n^{(l)}(x+t)=\sum_{j=1}^{3}(-i\omega_j\mu_n)^lX_j^-(x)\exp(-i\omega_j\mu_nt)-\int_x^{x+t}M(u_n(\xi))K_t^{(l)}(\xi-x-t)d\xi,\quad \textit{if } \operatorname{Im}\lambda_k>0; \tag{2.1}$$

$$u_n^{(l)}(x+t)=\sum_{j=1}^{3}(i\omega_j\mu_n)^lX_j^+(x)\exp(i\omega_j\mu_nt)-\int_x^{x+t}M(u_n(\xi))K_t^{(l)}(\xi-x-t)d\xi,\quad \textit{if } \operatorname{Im}\lambda_n<0. \tag{2.2}$$

PROOF. For definiteness we consider the case $\operatorname{Im}\lambda_n<0$. Multiply each side of the equation $Lu_n(\xi)+\lambda_nu_n(\xi)=0$ by the function $K_t^{(l)}(\xi-x-t)$ and integrate the obtained equality with respect to ξ from x to $x+t$, where $x,x+t\in G$

$$\sum_{j=1}^{3}\omega_j(i\omega_j\mu_n)^l\int_x^{x+t}u_n^{(3)}(\xi)\exp(-i\omega_j\mu_n(\xi-x-t))d\xi \tag{2.3}$$

$$+\int_x^{x+t}\left\{p_1(\xi)u_n^{(2)}(\xi)+p_2(\xi)u_n^{(1)}(\xi)+p_3(\xi)u_n(\xi)\right\}$$
$$\times K_t^{(l)}(\xi-x-t)d\xi+\lambda_n\int_x^{x+t}u_n(\xi)K_t^{(l)}(\xi-x-t)d\xi=0.$$

Integrating by parts and using $\sum_{j=1}^{3}\omega_j^s=3\delta_{3s}$ (δ_{ks} is the Kronecker symbol), we transform the first expression in equality (2.3) in the following way

$$\sum_{j=1}^{3}\omega_j(i\omega_j\mu_n)^l\int_x^{x+t}u_n^{(3)}(\xi)\exp(-i\omega_j\mu_n(\xi-x-t))d\xi$$
$$=\sum_{j=1}^{3}\omega_j(i\omega_j\mu_n)^l\sum_{m=0}^{2}(-1)^m(-i\omega_j\mu_n)^m$$
$$\times\left[u_n^{(2-m)}(x+t)-u_n^{(2-m)}(x)\exp(i\omega_j\mu_n t)\right]$$
$$-\sum_{j=1}^{3}\omega_j(i\omega_j\mu_n)^l(-\omega_j\mu_n)^3\int_x^{x+t}u_n(\xi)\exp(-i\omega_j\mu_n(\xi-x-t))d\xi$$
$$=3\mu_n^2u_n^{(l)}(x+t)$$
$$-\sum_{j=1}^{3}\omega_j(i\omega_j\mu_n)^l\sum_{m=0}^{2}(i\omega_j\mu_n)^mu_n^{(2-m)}(x)\exp(i\omega_j\mu_n t)+(-i\mu_n)^3$$
$$\times\sum_{j=1}^{2}\omega_j(i\omega_j\mu_n)^l\int_x^{x+t}u_n(\xi)\exp(-i\omega_j\mu_n(\xi-x-t)d\xi.$$

Taking this into account in (2.3) and taking into attention the equality $(-i\mu_n)^3=-\lambda_n$, we get

$$3\mu_n^2u_n^{(l)}(x+t)-\sum_{j=1}^{3}(i\omega_j\mu_n)^l\left[\sum_{m=0}^{2}\omega_j^{m+1}(i\mu_n)^mu_n^{(2-m)}(x)\right]\exp(i\omega_j\mu_n t)$$
$$+\int_x^{x+t}\left\{\sum_{\tau=1}^{3}p_r(\xi)u_n^{(3-r)}(\xi)\right\}K_t^{(l)}(\xi-x-t)d\xi=0.$$

Hence we find $u_n^{(l)}(x+t)$ and get formula (2.2). Lemma 2.1 is proved. □

For $x=0$ we write formulas (2.1) and (2.2) in a more convenient form

$$\mu_n^{-l}u_n^{(l)}(t)=\sum_{j=1}^{2}X_j^-(0)(-i\omega_j)^l\exp(-i\omega_j\mu_n t)$$
$$-(-i\omega_3)B_3^-\exp(i\omega_3\mu_n(1-t))$$
$$-\sum_{j=1}^{2}(-i)^l\omega_j^{l+1}\int_0^t M(u_n(\xi))\exp(i\omega_j\mu_n(\xi-t))d\xi \tag{2.4}$$

$$+ (-i)^l \omega_3^{l+1} \int_t^1 M(u_n(\xi)) \exp(i\omega_3\mu_n(\xi - t))d\xi,$$

$\operatorname{Im}\lambda_n > 0$, $l = \overline{0,2}$, where

$$B_3^- = X_3^-(0)\exp(-i\omega_3\mu_n) - \omega_3 \int_0^1 M(u_n(\xi))\exp(-i\omega_3\mu_n(\xi-1))d\xi;$$

$$\begin{aligned}(2.5)\qquad \mu_n^{-l} u_n^{(l)}(t) = &\sum_{j=1,j\neq 2}^{2} (i\omega_j)^l X_j^+(0)\exp(i\omega_j\mu_n t)\\ &+ (i\omega_2)^l B_2^+ \exp(-i\omega_2\mu_n(1-t))\\ &- \sum_{j=1,j\neq 2}^{3} (i)^l \omega_j^{l+1} \int_0^t M(u_n(\xi))\exp(-i\omega_j\mu_n(\xi-t))d\xi\\ &+ (i)^l\omega_3^{l+1}\int_t^1 M(u_n(\xi))\exp(-i\omega_2\mu_n(\xi-t))d\xi,\end{aligned}$$

where $\operatorname{Im}\lambda_n < 0$, $l = \overline{0,2}$

$$B_2^+ = X_2^+(0)\exp(i\omega_2\mu_n) - \omega_2\int_0^1 M(u_n(\xi))\exp(-i\omega_2\mu_n(\xi-t))d\xi.$$

For the coefficients in formulas (2.4) and (2.5) we note that if the system $\{u_n(x)\}_{n=1}^{\infty}$ is a Bessel system in $L_2(G)$, then for, them the following estimations are fulfilled (see [**9**, **10**])

$$\begin{aligned}(2.6)\qquad |X_1^{\pm}(0)| \leqslant \|u_n\|_2, \qquad |X_2^{\pm}(0)| \leqslant C\|u_n\|_\infty\\ |B_2^+| \leqslant C\|u_n\|_\infty, \qquad |B_3^-| \leqslant C\|u_n\|_\infty.\end{aligned}$$

Now, using formulas (2.4) and (2.5), estimate the Fourier coefficients of the function $f(x) \in W_p^1(G), p > 1$, in system $\{u_n(x)\}_{n=1}^{\infty}$.

LEMMA 2.2. *Let the function $f(x) \in W_p^1(G), p > 1$, and the system $\{u_n(x)\}_{n=1}^{\infty}$ satisfy condition* (1.1). *Then for the Fourier coefficients f_n the estimations* ($\mu_n \geqslant 1$) *are valid*

$$\begin{aligned}(2.7)\quad |f_n| \leqslant \operatorname{const}\Bigg\{ &C(f)\mu_n^{\alpha-3}\|u_n\|_\infty + \mu_n^{-1}|(f\overline{p_1}, \mu_n^{-2}u_n^{(2)})|\\ &+ \mu_n^{-1}\left|\left(f', \mu_n^{-2}u_n^{(2)}\right)\right| + \mu_n^{-2}\left(\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\right)\|f\|_\infty\|u\|_\infty\Bigg\};\end{aligned}$$

$$\begin{aligned}(2.8)\quad |f_n| \leqslant \operatorname{const}\Bigg\{ &\Bigg[C(f)\mu_n^{\alpha-3} + \mu_n^{-1}|(f', \exp(i\omega_3\mu_n t))|\\ &+ \mu_n^{-1}|(f', \exp(-i\omega_2\mu_n(1-t)))| + (\|f\|_\infty + \|f'\|_1)\mu_n^{-2}\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\Bigg]\\ &\times \|u_n\|_\infty + \mu_n^{-1}|(f', \exp(-i\mu_n t))|\Bigg\},\end{aligned}$$

if $p_1(x) \equiv 0, \operatorname{Im}\lambda_n < 0$*;*

$$|f_n| \leqslant \text{const}\left\{\left[C(f)\mu_n^{\alpha-3} + \mu_n^{-1}|(f', \exp(-i\omega_2\mu_n t))| + \mu_n^{-1}|(f', \exp(i\omega_3\mu_n(1-t)))| + (\|f\|_\infty + \|f'\|_1)\mu_n^{-2}\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\right] \times \|u_n\|_\infty + \mu_n^{-1}|(f', \exp(i\mu_n t))|\right\}, \tag{2.9}$$

if $p_1(x) \equiv 0, \operatorname{Im}\lambda_n > 0$.

Proof. By definition of the eigenfunction $u_n(x)$, the Fourier coefficients f_n for $\mu_n \geqslant 1$ are calculated by the formula

$$\begin{aligned} f_n = (f, u_n) &= -\frac{1}{\overline{\lambda_n}}(f, Lu_n) \\ &= -\frac{1}{\overline{\lambda_n}}(f, u^{(3)}) - \frac{1}{\overline{\lambda_n}}\big(f, p_1u_n^{(2)}\big) - \frac{1}{\overline{\lambda_n}}\big(f, p_2u_n^{(1)}\big) - \frac{1}{\overline{\lambda_n}}(f, p_3u_n). \end{aligned} \tag{2.10}$$

Applying the estimation (see [**10**])

$$\big\|u_n^{(s)}\big\|_\infty \leqslant \text{const}(1+\mu_m)^{s+\frac{1}{p}}\|u_n\|_p, \quad p \geqslant 1, s = \overline{0,2} \tag{2.11}$$

we find

$$\begin{aligned} &\frac{1}{|\lambda_n|}\big|\big(f, p_2u_n^{(1)}\big)\big| + \frac{1}{|\lambda_n|}|(f, p_3u_n)| \\ &\qquad \leqslant \text{const}\,\|f\|_\infty\mu_n^{-2}\left(\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\right)\|u_n\|_\infty. \end{aligned} \tag{2.12}$$

Making integration in parts in the first summand on the right-hand side of equality (2.10) and taking into account condition (1.1), we get

$$\frac{1}{|\lambda_n|}\big|\big(f, u_n^{(3)}\big)\big| \leqslant C(f)\mu_n^{\alpha-3}\|u_n\|_\infty + \mu_n^{-3}\big|\big(f', u_n^{(2)}\big)\big|. \tag{2.13}$$

From (2.10), (2.12) and (2.13) it follows estimation (2.7).

Estimate the expression $\mu_n^{-3}\big|\big(f', u_n^{(2)}\big)\big|$ in the case $p_1(x) \equiv 0$. For that we use formulas (2.4) and (2.5) depending on the sign of $\operatorname{Im}\lambda_n$. For definiteness we consider the case $\operatorname{Im}\lambda_n < 0$ and apply formula (2.5) for $l = 2$. Then by estimations (2.6), (2.11) and

$$\begin{aligned} |M(u_n(\xi))| &\leqslant \frac{1}{3}\mu_n^{-2}\sum_{r=2}^{3}|p_r(\xi)|\big|u_n^{(3-r)}(\xi)\big| \\ &\leqslant \text{const}\,\mu_n^{-1}\left(\sum_{r=2}^{3}|p_r(\xi)|\mu_n^{2-r}\right)\|u_n\|_\infty \end{aligned}$$

we get that

$$\mu_n^{-3}\big|\big(f', u_n^{(2)}\big)\big| = \mu_n^{-1}\big|\big(f', \mu_n^{-2}u_n^{(2)}\big)\big|$$

$$\begin{aligned}
&\leqslant \mu_n^{-1}\sum_{j=1,j\neq 2}^{3}|X_j^+(0)||(f',\exp(i\omega_j\mu_n t))|\\
&\quad+\mu_n^{-1}|B_2^+||(f',\exp(-i\omega_2\mu_n(1-t)))|\\
&\quad+\mu_n^{-1}\sum_{j=1,j\neq 2}^{3}\left|\left(f',\int_0^t M(u_n(\xi))\exp(-i\omega_j\mu_n(\xi-t)d\xi)\right)\right|\\
&\quad+\mu_n^{-1}\left|\left(f',\int_t^1 M(u_n(\xi))\exp(-i\omega_2\mu_n(\xi-t)d\xi)\right)\right|\\
&\leqslant \operatorname{const}\mu_n^{-1}\big\{|(f',\exp(-i\mu_n t))|+\big[|(f',\exp(i\omega_3\mu_n t))|\\
&\qquad\qquad+|(f',\exp(-i\omega_2\mu_n(1-t)))|\big]\|u_n\|_\infty\big\}\\
&\quad+\operatorname{const}\mu_n^{-2}\|f'\|_1\left(\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\right)\|u_n\|_\infty.
\end{aligned}$$

Thus, in the case $p_1(x)\equiv 0$, $\operatorname{Im}\lambda_n<0$ the following estimation is fulfilled

$$\begin{aligned}
(2.14)\quad \mu_n^{-3}\big|\big(f',u_n^{(3)}\big)\big|&\leqslant \operatorname{const}\mu_n^{-1}\Big\{|(f',\exp(-i\mu_n t))|+\Big[|(f',\exp(i\omega_3\mu_n t))|\\
&\quad+|(f',\exp(i\omega_2\mu_n(t-1)))|+\|f'\|_1\mu_n^{-1}\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\Big]\|u_n\|_\infty\Big\}.
\end{aligned}$$

Consequently, estimation (2.8) follows from (2.10), (2.12)–(2.14). Estimation (2.9) for $p_1(x)\equiv 0$, $\operatorname{Im}\lambda_n>0$ is proved in the same way. The Lemma 2.2 is proved. □

LEMMA 2.3 (see [**10**]). *Let $p_1(x)\in L_2(G)$, $p_l(x)\in L_1(G)$, $l=2,3$ and $\{u_n(x)\}$ be an orthonormalized in $L_2(G)$ system of eigenfunctions of the operator L. Then the following estimations are fulfilled:*

$$(2.15)\qquad \sum_{\tau\leqslant\mu_n\leqslant tau+1}1\leqslant\operatorname{const},\quad \textit{for all } \tau\geqslant 0,$$

$$(2.16)\qquad \sum_{0\leqslant\mu_n\leqslant\tau}\|u_n\|_\infty^2\leqslant\operatorname{const}(1+\tau),\quad \textit{for all } \tau>0.$$

LEMMA 2.4 (see [**9**]). *If conditions of Lemma 2.3 are fulfilled, then the system $\big\{\mu_n^{-2}u_n^{(2)}(x)\big\}$, $\mu_n\geqslant 1$, is a Bessel system, i.e., for any $g(x)\in L_2(G)$ the following inequality is fulfilled*

$$\sum_{\mu_n\geqslant 1}\big|\big(f,\mu_n^{-2}u_n^{(2)}(x)\big)\big|^2\leqslant\operatorname{const}\|g\|_2^2.$$

LEMMA 2.5. *Under conditions of Lemma 2.3, the system $\{\exp(-i\mu_n t)\}$ for $\operatorname{Im}\lambda_n<0$ and the system $\{\exp(i\mu_n t)\}$ for $\operatorname{Im}\lambda_n>0$ satisfies the Riesz inequality for $1<p\leqslant 2$.*

PROOF. As these system satisfy the Bessel inequality in $L_2(G)$ (see [**11**]) subject to condition (2.15), furthermore, for any $g(x) \in L_1(G)$ it is valid

$$\left| \int_0^1 g(x)\overline{\varphi_n(x)}dx \right| \leqslant \text{const}\, \|g\|_1$$

where $\{\varphi_n(x)\}$ is any from the above mentioned systems, then by the Riesz–Torin theorem (see [**12**]), for these systems the Riesz inequality is valid, i.e.,

$$\sum_n \left| \int_0^1 g(x)\overline{\varphi_n(x)}dx \right|^q \leqslant \text{const}\, \|g\|_p^q$$

for any $g(x) \in L_p(G)$, $1 < p \leqslant 2$. Lemma 2.5 is proved. □

LEMMA 2.6. *Let the conditions of Lemma* 2.3 *be fulfilled. Then*

$$\sum_{\mu_n \geqslant \mu} \frac{\|u_n\|_\infty^2}{\mu_n^{\theta+1}} \leqslant \text{const}\, \mu^{-\theta}, \quad \theta > 0, \quad \textit{for all } \mu \geqslant 2. \tag{2.17}$$

PROOF. By estimations (2.15), (2.16) and Abel transformation for any $l \in N$

$$\begin{aligned}
\sum_{\mu \leqslant \mu_n \leqslant [\mu]+l} \frac{\|u_n\|_\infty^2}{\mu_n^{1+\theta}} &\leqslant \sum_{k=[\mu]}^{[\mu]+l} \frac{1}{k^{1+\theta}} \Bigg(\sum_{k \leqslant \mu_n < k+1} \|u_n\|_\infty^2 \Bigg) \\
&\leqslant \sum_{k=[\mu]}^{[\mu]+l-1} \Bigg(\sum_{1 \leqslant \mu_n \leqslant k+1} \|u_n\|_\infty^2 \Bigg) \Big(\frac{1}{k^{1+\theta}} - \frac{1}{(k+1)^{1+\theta}} \Big) \\
&\quad + \Bigg(\sum_{1 \leqslant \mu_n \leqslant [\mu]+l} \|u_n\|_\infty^2 \Bigg) ([\mu]+l)^{-(1+\theta)} + \Bigg(\sum_{1 \leqslant \mu_n \leqslant [\mu]-1} \|u_n\|_\infty^2 \Bigg) [\mu]^{-(1+\theta)} \\
&\leqslant \text{const} \sum_{k=[\mu]}^{[\mu]+l-1} (k+1) \frac{(k+1)^\theta (1+\theta)}{(k(k+1))^{1+\theta}} \\
&\quad + \text{const}([\mu]+l)^{-\theta} + \text{const}[\mu]^{-\theta} \leqslant \text{const} \Bigg(\sum_{k=[\mu]}^{\infty} \frac{1+\theta}{k^{1+\theta}} + [\mu]^{-\theta} \Bigg).
\end{aligned}$$

Hence, by the arbitrariness of the natural number l we get estimation (2.17). □

LEMMA 2.7. *Let the conditions of Lemma* 2.3 *be fulfilled. Then*

$$\sum_{\mu_n \geqslant \mu} \frac{\|u_n\|_\infty^p}{\mu_n^p} \leqslant \text{const}\, \mu^{1-p}, \quad 1 < p \leqslant 2, \quad \textit{for all } \mu \geqslant 2. \tag{2.18}$$

PROOF. For $p = 2$ estimation (2.18) follows from (2.17) for $\theta = 1$. Consider the case $p \neq 2$ and apply the Holder inequality for $p' = \frac{2}{p}$, $q' = \frac{2}{2-p}$:

$$\sum_{\mu_n \geqslant \mu} \frac{\|u_n\|_\infty^p}{\mu_n^p} = \sum_{\mu_n \geqslant \mu} \frac{\|u_n\|_\infty^p}{\mu_n^{p-\frac{1}{2}}} \frac{1}{\mu_n^{\frac{1}{2}}} \leqslant \Bigg(\sum_{\mu_n \geqslant \mu} \frac{\|u_n\|_\infty^2}{\mu_n^{2-\frac{1}{p}}} \Bigg)^{p/2} \Bigg(\sum_{\mu_n \geqslant \mu} \frac{1}{\mu_n^{\frac{1}{2-p}}} \Bigg)^{\frac{2-p}{2}}$$

$$\leqslant \Big(\sum_{\mu_n \geqslant \mu} \frac{\|u_n\|_\infty^2}{\mu_n^{2-\frac{1}{p}}} \Big)^{p/2} \Big(\sum_{k=[\mu]}^{\infty} \frac{1}{k^{\frac{1}{2-p}}} \Big(\sum_{k \leqslant \mu_n \leqslant k+1} 1 \Big) \Big)^{\frac{2-p}{2}}.$$

Having applied here Lemma 2.6 for $\theta = 1 - \frac{1}{p}$ and estimation (2.15), we get

$$\sum_{\mu_n \geqslant \mu} \frac{\|u_n\|_\infty^2}{\mu_n^p} \leqslant \operatorname{const} \big(\mu^{(\frac{1}{p}-1)}\big)^{p/2} [\mu]^{\frac{1-p}{2}} \leqslant \operatorname{const} \mu^{1-p}.$$

Lemma 2.7 is proved. □

LEMMA 2.8 (see [**9**]). *Let $\{\alpha_m\}_{m=0}^\infty$ be a numerical sequence with the elements $\alpha_m \geqslant 0, \beta$ be a complex number for which $\operatorname{Re}\beta > 0$. Then for the inequality*

$$\Big(\sum_{m=0}^{\infty} \alpha_m \Big| \int_G f(x) \exp(-m\beta x) dx \Big|^q \Big)^{\frac{1}{q}} \leqslant M_p \|f\|_p$$

to be fulfilled for any function $f(x) \in L_p(G)$, $1 < p \leqslant 2$, $q = \frac{p}{p-1}$, it is necessary and sufficient the existence of a constant K such that for all $N = 1, 2, \dots$ the following estimation is valid $\sum_{m=0}^N \alpha_m \leqslant KN$.

LEMMA 2.9. *If under the conditions of Lemma 2.3, we have that for each system $\big\{\|u_n\|_\infty^{\frac{2}{q}} e^{i\omega_3 \mu_n t}\big\}_{n=1}^\infty$ and $\big\{\|u_n\|_\infty^{\frac{2}{q}} e^{-i\omega_2 \mu_n (1-t)}\big\}_{n=1}^\infty$ for $\operatorname{Im}\lambda_n < 0$, for each system $\big\{\|u_n\|_\infty^{\frac{2}{q}} e^{-i\omega_2 \mu_n t}\big\}_{n=1}^\infty$ and $\big\{\|u_n\|_\infty^{\frac{2}{q}} e^{i\omega_3 \mu_n (1-t)}\big\}_{n=1}^\infty$ for $\operatorname{Im}\lambda_n < 0$ the Riesz inequality is fulfilled for $1 < p \leqslant 2$, where $p^{-1} + q^{+1} = 1$.*

PROOF. Let us consider the first one of these systems and prove for it the Riesz inequality (the remaining systems are considered similarly). Since $\mu_n \in [0, +\infty)$ and $\omega_3 = i\frac{\sqrt{3}}{2} + \frac{1}{2}$, then $i\omega_3 \mu_n t = \big(i\frac{1}{2} - \frac{\sqrt{3}}{2}\big)\mu_n t$ and $|e^{i\omega_3 \mu_n t}| = e^{-\frac{\sqrt{3}}{2}\mu_n t}$. Taking this into account, we get that for any $f(x) \in L_p(G)$

$$\begin{aligned}
\sum_{\operatorname{Im}\lambda_n < 0} \|u_n\|_\infty^2 \Big| \int_0^1 \overline{f(t)} e^{i\omega_3 \mu_n t} dt \Big|^q &\leqslant \sum_{\operatorname{Im}\lambda_n < 0} \|u_n\|_\infty^2 \Big| \int_0^1 \overline{f(t)} e^{\left(i\frac{1}{2} - \frac{\sqrt{3}}{2}\right)\mu_n t} dt \Big|^q \\
&\leqslant \sum_{n=1}^{\infty} \|u_n\|_\infty^2 \Big(\int_0^1 |f(t)| e^{-\frac{1}{2}\mu_n t} dt \Big)^q \\
&\leqslant \sum_{k=0}^{\infty} \sum_{k \leqslant \mu_n < k+1} \|u_n\|_\infty^2 \Big(\int_0^1 |f(t)| e^{-\frac{1}{2}\mu_n t} dt \Big)^q \\
&\leqslant \sum_{k=0}^{\infty} \Big(\sum_{k \leqslant \mu_n < k+1} \|u_n\|_\infty^2 \Big) \Big(\int_0^1 |f(t)| e^{-\frac{1}{2}kt} dt \Big)^q \\
&= \sum_{k=0}^{\infty} \alpha_k \Big(\int_0^1 |f(t)| e^{-\frac{1}{2}kt} dt \Big)^q,
\end{aligned}$$

where $\alpha_k = \sum_{k \leqslant \mu_n < k+1} \|u_n\|_\infty^2$.

By inequality (2.16), for any natural number N it is fulfilled:

$$\sum_{k=0}^{N}\alpha_k=\sum_{k=0}^{N}\Big(\sum_{k\leqslant\mu_n<k+1}\|u_n\|_\infty^2\Big)=\sum_{0\leqslant\mu_n<N+1}\|u_n\|_\infty^2\leqslant\mathrm{const}\cdot N.$$

Consequently, the condition of Lemma 2.8 is fulfilled. Therefore, the following inequality is valid

$$\Big\{\sum_{k=1}^{\infty}\alpha_k\Big(\int_0^1|f(t)|e^{-\frac{1}{2}kt}dt\Big)^q\Big\}^{1/q}\leqslant M(p)\|f\|_p.$$

Lemma 2.9 is proved. □

3. Proof of the basic results

Proof of Theorem 1.1. If suffices to consider the case $1<p\leqslant 2$. Prove the uniform convergence of the series $\sum_{n=1}^{\infty}|f_n||u_n(x)|$ on $\bar{G}$. To this end we partition this series in two sums: $\sum_{0\leqslant\mu_n\leqslant 2}|f_n||u_n(x)|$ and $\sum_{\mu_n>2}|f_n||u_n(x)|$. The first sum by inequality (2.16) doesn't exceed the quantity $\mathrm{const}\,\|f\|_1$. For investigating the second series we apply Lemma 2.2, i.e., estimations (2.8) and (2.9) depending on the sign of $\operatorname{Im}\lambda_n$. For that we represent the given series in the form

$$\sum_{\mu_n>2}|f_n||u_n(x)|=\sum_{n\in J_1}|f_n||u_n(x)|+\sum_{n\in J_2}|f_n||u_n(x)|=I_1+I_2,$$

where $J_1=\{n:\mu_n>2,\operatorname{Im}\lambda_n<0\}$, $J_2=\{n:\mu_n>2,\operatorname{Im}\lambda_n>0\}$. By estimation (2.8)

$$\begin{aligned}I_1=\sum_{n\in J_1}|f_n||u_n(x)|&\leqslant\mathrm{const}\sum_{n\in J_1}C(f)\mu_n^{\alpha-3}\|u_n\|_\infty^2\\&+\mathrm{const}\sum_{n\in J_1}\mu_n^{-1}|(f',e^{i\omega_3\mu_n t})|\|u_n\|_\infty^2\\&+\mathrm{const}\sum_{n\in J_1}\mu_n^{-1}|(f',e^{-i\omega_2\mu_n(1-t)})|\|u_n\|_\infty^2\\&+\mathrm{const}(\|f\|_\infty+\|f'\|_1)\sum_{n\in J_1}\mu_n^{-2}\Big(\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\Big)\|u_n\|_\infty^2\\&+\mathrm{const}\sum_{n\in J_1}\mu_n^{-1}|(f,e^{-i\mu_n t})|\|u_n\|_\infty\\&=\mathrm{const}(I_1^1+I_1^2+I_1^3+I_1^4+I_1^5).\end{aligned}$$

Estimate the series I_1^j, $j=\overline{1,5}$. By Lemma 2.6 and condition $0\leqslant\alpha<2$ we find

$$\begin{aligned}I_1^1=C(f)\sum_{n\in J_1}\mu_n^{\alpha-3}\|u_n\|_\infty^2&\leqslant C(f)\sum_{\mu_n\geqslant 2}\frac{\|u_n\|_\infty^2}{\mu_n^{1+(2-\alpha)}}\\&\leqslant\mathrm{const}\,C(f)2^{\alpha-2}<\infty.\end{aligned}\tag{3.1}$$

For estimating the series I_1^2 we apply at first the Holder inequality for the sum, and then Lemmas 2.6 and 2.9:

$$\begin{aligned} I_1^2 &= \sum_{n\in J_1} \|u_n\|_\infty^{2/p}\mu_n^{-1}\big(\|u_n\|_\infty^{2/q}|(f', e^{i\omega_3\mu_n t})|\big) \\ &\leqslant \left\{\sum_{n\in J_1}\frac{\|u_n\|^2}{\mu_n^p}\right\}^{1/p}\left\{\sum_{n\in J_1}\|u_n\|_\infty^2|(f', e^{i\omega_3\mu_n t})|^q\right\}^{1/q} \\ &\leqslant \left\{\sum_{\mu_n\geqslant 2}\frac{\|u_n\|^2}{\mu_n^p}\right\}^{1/p}\left\{\sum_{n\in J_1}\|u_n\|_\infty^2|(f', e^{i\omega_3\mu_n t})|^q\right\}^{1/q} \\ &\leqslant \text{const}\, 2^{-1/q}M(p)\|f'\|_p < \infty. \end{aligned}$$

The series I_1^3 is estimated in the same way as the series I_1^2. For estimating the series I_1^4 we apply Lemma 2.6.

$$\begin{aligned} I_1^4 &= (\|f\|_\infty + \|f'\|_1)\sum_{n\in J_1}\mu_n^{-2}\left(\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\right)\|u\|_\infty^2 \\ &\leqslant \text{const}(\|f\|_\infty + \|f'\|_1)\left(\sum_{r=2}^{3}\|p_r\|_1 2^{1-r}\right) < \infty. \end{aligned} \tag{3.2}$$

Now estimate the series I_1^5. For that we apply the Holder inequality, and then Lemmas 2.5 and 2.7:

$$\begin{aligned} I_1^5 &= \sum_{n\in J_1}\frac{\|u_n\|_\infty}{\mu_n}|(f', e^{-i\mu_n t})| \\ &\leqslant \left(\sum_{n\in J_1}\frac{\|u_n\|_\infty^p}{\mu_n^p}\right)^{1/p}\left(\sum_{n\in J_1}|(f', e^{-i\mu_n t})|^q\right)^{\frac{1}{q}} \leqslant \text{const}\,\|f'\|_p 2^{-\frac{1}{q}} < \infty. \end{aligned}$$

Thus, the series I_1 uniformly converges on $\bar{G}$. Applying estimation (2.9) for the coefficients f_n in the same way we prove the uniform convergence of the series I_2 on $\bar{G}$. Consequently, the series $\sum_{n=1}^{\infty}|f_n|\,|u_n(x)|$ uniformly converges on $\bar{G}$. By the completeness of the system $\{u_n(x)\}_{n=1}^{\infty}$ in $L_2(G)$ and continuity of the function $f(x)$ on $\bar{G}$ the series $\sum_{n=1}^{\infty}f_n u_n(x)$ uniformly converges to $f(x)$, i.e. it holds the equality

$$f(x) = \sum_{n=1}^{\infty} f_n u_n(x), \quad x\in\bar{G} \tag{3.3}$$

Now establish estimate (1.2). By equality (3.3)

$$|R_\nu(x,f)| = |f(x) - \sigma_\nu(x,f)| = \left|\sum_{\mu_n>\nu} f_n u_n(x)\right|$$

$$\leqslant \sum_{\mu_n>\nu} |f_n||u_n(x)| = \sum_{n\in A_1(\nu)} + \sum_{n\in A_2(\nu)} = T_1(\nu) + T_2(\nu),$$

where $A_1(\nu) = \{n : \mu_n \geqslant \nu, \operatorname{Im}\lambda_n < 0\}$, $A_2(\nu) = \{n : \mu_n \geqslant \nu, \operatorname{Im}\lambda_n < 0\}$. The series $T_1(\nu), T_2(\nu)$ are estimated by the scheme demonstrated by estimating the series I_1. As a result we find

$$T_j(\nu) \leqslant \text{const}\left\{C(f)\nu^{\alpha-2} + \nu^{-\frac{1}{q}}\|f'\|_p + \nu^{-1}(\|f\|_\infty + \|f'\|_1)\Big(\sum_{r=2}^{3}\|p_r\|_1\nu^{2-r}\Big)\right\},$$
$$j = 1, 2, \quad x \in \bar{G}.$$

Consequently, estimation (1.2) is valid for $1 < p \leqslant 2$. For $p > 2$ the validity of estimation (1.2) follows from the embedding $L_p(G) \subset L_2(G)$.

Theorem 1.1 is proved. □

Corollary 1.1 follows from Theorem 1.1 with regard to the inequality $\|f\|_\infty \leqslant \|f'\|_1$ that is fulfilled for the function$f(x) \in W_p^1(G)$, $f(0) = f(1) = 0$.

Proof of Theorem 1.2. In the present case it is necessary to prove the uniform convergence of the series $\sum_{\mu_n\geqslant 2}|f_n||u_n(x)|$ on $\bar{G}$. By estimation (2.7) we find

$$\begin{aligned}\sum_{\mu_n\geqslant 2}|f_n||u_n(x)| \leqslant \text{const}\Big\{&C(f)\sum_{\mu_n\geqslant 2}\mu^{\alpha-3}\|u_n\|_\infty^2 \\ &+ \sum_{\mu_n\geqslant 2}\mu_n^{-1}\left|\Big(f\,\overline{p_1}, \frac{u_n^{(2)}}{\mu_n^2}\Big)\right|\|u_n\|_\infty^2 + \sum_{\mu_n\geqslant 2}\mu_n^{-1}\|u_n\|_\infty\left|\Big(f', \frac{u_n^{(2)}}{\mu_n^2}\Big)\right| \\ &+ \|f\|_\infty\sum_{\mu_n\geqslant 2}\mu_n^{-2}\|u_n\|_\infty^2\Big(\sum_{r=2}^{3}\mu_n^{2-r}\|p_r\|_1\Big)\Big\} \\ &\leqslant \text{const}\{V_1 + V_2 + V_3 + V_4\}.\end{aligned}$$

The series V_1 and V_4 are estimated in the same way as the series I_1^1 and I_1^4. For V_1 estimation (3.1) is fulfilled, and for V_4 estimation (3.2) is fulfilled by changing the multiplier $(\|f\|_\infty + \|f'\|_1)$ by the multiplier $\|f\|_\infty$.

By estimating the series V_2 and V_3, we apply the Bessel inequality for the system $\{u_n^{(2)}(x)/\mu_n^2\}$, $\mu_n \geqslant 2$, whose validity is established in the paper [**9**], and also Lemma 2.7 for $p = 2$.

As a result, we find

$$\begin{aligned}V_2 = \sum_{\mu_n\geqslant 2}\mu_n^{-1}\|u_n\|_\infty\Big(f\,\overline{P_1}, \frac{u_n^{(2)}}{\mu_n^2}\Big) &\leqslant \Big(\sum_{\mu_n\geqslant 2}\frac{\|u_n\|_\infty^2}{\mu_n^2}\Big)^{1/2} \\ &\times\Big(\sum_{\mu_n\geqslant 2}\left|\Big(f\,\overline{p_1}, \frac{u_n^{(2)}}{\mu_n^2}\Big)\right|^2\Big)^{1/2} \leqslant \text{const}\,2^{-\frac{1}{2}}\|f\,\overline{p_1}\|_2;\end{aligned}$$

$$V_3 \leqslant \Big(\sum_{\mu_n\geqslant 2}\frac{\|u_n\|_\infty^2}{\mu_n^2}\Big)^{1/2}\Big(\sum_{\mu_n\geqslant 2}\Big|\Big(f',\frac{u_n^{(2)}}{\mu_n^2}\Big)\Big|^2\Big)^{1/2}\leqslant \operatorname{const} 2^{-\frac{1}{2}}\|f'\|_2.$$

Consequently, the series $\sum_{n=1}^{\infty} f_n u_n(x)$ converges absolutely and uniformly on $\bar{G}$, and the following equality $f(x)=\sum_{n=1}^{\infty} f_n u_n(x)$, $x\in G$, is valid. It is easy to see that for the remainder $R_\nu(x,f)$ of this series the following estimation will be valid (in the remainder the summation is conducted according to numbers n, for which $\mu_n>\nu$)

$$\sup|R_\nu(x,f)|\leqslant \operatorname{const}\Big\{C(f)\nu^{\alpha-2}+\nu^{-\frac{1}{2}}(\|f\overline{p_1}\|_2+\|f'\|_2)+\nu^{-1}\sum_{r=2}^{3}\nu^{2-r}\|p_r\|_1\Big\},$$
$$\nu\geqslant 2.$$

Theorem 1.2 is proved. □

For justification of Corollary 1.2, it suffices to take into account that the sequence of remainders of the converging series tends to zero, more exactly,

$$\sum_{\mu_n\geqslant\nu}\Big|\Big(f\,\overline{p_1},\frac{u_n^{(2)}}{\mu_n^2}\Big)\Big|^2=o(1),\quad \nu\to+\infty;$$
$$\sum_{\mu_n\geqslant\nu}\Big|\Big(f',\frac{u_n^{(2)}}{\mu_n^2}\Big)\Big|^2=o(1),\quad \nu\to+\infty.$$

Proof of Theorem 1.3. By orthonormality of the system $\{u_n(x)\}_{n=1}^{\infty}$ in $L_2(G)$, condition (2.15) is fulfilled. On the other hand,

$$1=|(u_n,u_n)|\leqslant\|u_n\|_p\|u_n\|_q\leqslant\|u_n\|_\infty\|u_n\|_q.$$

Hence we find $\|u_n\|_q^{-q}\leqslant\|u_n\|_\infty^q$. Therefore, by inequality (2.15) and uniform boundedness of the system $\{u_n(x)\}_{n=1}^{\infty}$ we find

$$\sum_{0\leqslant\mu_n\leqslant\tau}\|u_n\|_\infty^q\|u_n\|_q^{-q}\leqslant\sum_{0\leqslant\mu_n\leqslant\tau}\|u_n\|_\infty^{2q}\leqslant C\sum_{0\leqslant\mu_n\leqslant\tau}1\leqslant\operatorname{const}\tau,\quad\text{for all }\tau>0.$$

Thus, for the system $\{u_n(x)\}_{n=1}^{\infty}$, all the conditions of the sufficiency part of [**9**, Theorem 3] are fulfilled. Therefore, for the system $\{\frac{u_n(x)}{\mu_n^2}\}$, $\mu_n\geqslant 1$ the Riesz inequality is valid.

For proving Theorem 1.3, it suffices to estimate the series $\mathbf{V}_3$ (all remaining series V_1,V_2,V_4 were estimated in Theorem 1.2 without requirement of uniform boundedness of the system $\{u_n(x)\}_{n=1}^{\infty}$). At first, apply the Hölder inequality, and then the Riesz inequality and Lemma 2.7. As a result, for V_3 and its remainder we get

$$V_3=\sum_{\mu_n\geqslant 2}\mu_n^{-1}\|u_n\|_\infty\Big|\Big(f',\frac{u_n^{(2)}}{\mu_n^2}\Big)\Big|$$
$$\leqslant\Big(\sum_{\mu_n\geqslant 2}\frac{\|u_n\|_\infty^p}{\mu_n^p}\Big)^{1/p}\Big(\sum_{\mu_n\geqslant 2}\Big|\Big(f',\frac{u_n^{(2)}}{\mu_n^2}\Big)\Big|^q\Big)^{1/q}\leqslant\operatorname{const}2^{-\frac{1}{q}}\|f'\|_2;$$

$$\sum_{\mu_n\geqslant\nu}\frac{\|u_n\|_\infty}{\mu_n}\left|\left(f,\frac{u_n^{(2)}}{\mu_n^2}\right)\right|$$
$$\leqslant\left(\sum_{\mu_n\geqslant\nu}\frac{\|u_n\|_\infty^p}{\mu_n^p}\right)^{1/p}\left(\sum_{\mu_n\geqslant\nu}\left|\left(f',\frac{u_n^{(2)}}{\mu_n}\right)\right|^q\right)^{1/q}\leqslant\operatorname{const}\nu^{-\frac{1}{q}}\|f'\|_p.$$

Theorem 1.3 is proved. □

References

1. V. A. Il'in, *Necessary and sufficient conditions for spatial decompositions to be bases and to be equiconvergent with atrigonometric series I*, Differ. Uravn. **16**(5) (1980), 37l–794.
2. M. A. Naymark, *Lineyar Differential Operators*, Nauka, Moskva, 1969 (In Russian).
3. N. L. Lažetić, *On uniform convergence of spectral expansions and their derivatives arising by selfadjoint extensions of an one-dimensional Schrodinger operator*, Publ. Inst. Math., Nouv. Sér. **69(83)** (2001), 59–77.
4. ———, *On uniform convergence on closed intervals of spectral expansions and their derivatives for functions from* $W_p^{(1)}$, Mat. Vesn. **56** (2004), 91–104.
5. V. M. Kurbanov, R. A. Safarov, *On uniform convergence of orthogonal expansions in eigenfunctions of Sturm–Liouville operator*, Trans. Natl. Acad. Sci. Azerb., Ser. Phys.-Tech. Math. Sci. **24**(1) (2004), 161–167.
6. ———, *Uniform convergence of biorthogonal expansion responding to the Schrodinger operator*, Proc. Inst. Math. Mech., Natl. Acad. Sci. Azerb. **20** (2004), 63–70.
7. V. M. Kurbanov, A. T. Garayeva, *Absolute and uniform convergence of expansions in the root vector functions of the Schrodinger operator with matrix potential*, Dokl. Math. **87**(3) (2013), 304–306.
8. V. M. Kurbanov, E. B. Akhundova, *Absolute convergence of spectral expansions in eigenfunctions of third order ordinary differential operator*, Proc. Inst. Math. Mech., Natl. Acad. Sci. Azerb. **40** (2014), Spec. Iss., 264–274.
9. V. M. Kurbanov, *On an analog of the Riesz theorem and basis property of the system of root functions of differential operator in* L_p*: I, II*, Differ. Uravn. **49**(1) (2013), 7–19 (**4**, 437–449).
10. ———, *On the distribution of eigenfunctions and a criterion for the Bessel property of root functions of a differential operator I*, Differ. Uravn. **4l**(4) (2005), 464–478.
11. V. A. Il'in, *Unconditional basis property on a closed interval of systems of eigen- and associated functions of a second order differential operator*, Dokl. Akad. Nauk SSSR **273**(5) (1983), 1048–1053.
12. A. Zygmund, *Trigonometric series*, Cambridge, 1959. Russian transl.: *Trigonometricheskie ryady*, Gonti, Moskva, 1965.

Institute of Mathematics and Mechanics of NAS of Azerbaijan
Baku, Azerbaijan
q.vali@yahoo.com

(Received 09 05 2015)