

Refined restricted inversion sequences

Dongsu Kim^{*1} and Zhicong Lin^{†2}

¹Department of Mathematical Sciences, Korea Advanced Institute of Science and Technology,
Daejeon 305-701, Republic of Korea

²School of Science, Jimei University, Xiamen 361021, P.R. China & CAMP, National Institute
for Mathematical Sciences, Daejeon 305-811, Republic of Korea

Abstract. Recently, the study of patterns in inversion sequences was initiated by Corteel-Martinez-Savage-Weselcouch and Mansour-Shattuck independently. Motivated by their works and a double Eulerian equidistribution due to Foata (1977), we investigate several classical statistics on restricted inversion sequences that are either known or conjectured to be enumerated by *Catalan*, *Large Schröder*, *Euler* and *Baxter* numbers. One of the two highlights of our results is an intriguing bijection between 021-avoiding inversion sequences and (2413, 4213)-avoiding permutations, which proves a sextuple equidistribution involving double Eulerian statistics. The other one is a refinement of a conjecture due to Martinez and Savage that the cardinality of $\mathbf{I}_n(\geq, \geq, >)$ is the n -th Baxter number, which is proved via the so-called *obstinate kernel method* developed by Bousquet-Mélou.

Keywords: Inversion sequences, ascents, distinct entries, last entry, Schröder numbers, Baxter numbers

1 Introduction

For each $n \geq 1$, the set of *inversion sequences* of length n , denoted \mathbf{I}_n , is defined by $\mathbf{I}_n = \{(e_1, e_2, \dots, e_n) : 0 \leq e_i < i\}$. It serves as various kind of codings for \mathfrak{S}_n , the set of permutations of $[n] := \{1, 2, \dots, n\}$. By a *coding* of \mathfrak{S}_n , we mean a bijection from \mathfrak{S}_n to \mathbf{I}_n . For example, the map $\Theta(\pi) : \mathfrak{S}_n \rightarrow \mathbf{I}_n$ defined for $\pi = \pi_1\pi_2 \dots \pi_n \in \mathfrak{S}_n$ as

$$\Theta(\pi) = (e_1, e_2, \dots, e_n), \quad \text{where } e_i := |\{j < i : \pi_j > \pi_i\}|,$$

is a natural coding of \mathfrak{S}_n . Clearly, the sum of the entries of $\Theta(\pi)$ equals the number of *inversions* of π , i.e., the number of pairs $i < j$ such that $\pi_i > \pi_j$. This is the reason why \mathbf{I}_n is named inversion sequences here.

Pattern avoidance in permutations has already been extensively studied in the literature (see the book by Kitave [9]), while the systematic study of patterns in inversion

*dongsu.kim@kaist.ac.kr

†lin@nims.re.kr. Supported in part by the National Science Foundation of China grant 11501244.

C_n	$\mathbf{I}_n(132), \mathbf{I}_n(123)$: classical result [9]; $\mathbf{I}_n(\geq, -, \geq)$: conjectured in [12]
S_n	$\mathfrak{S}_n(2413, 3142), \mathfrak{S}_n(2413, 4213), \mathfrak{S}_n(3124, 3214)$: classical result [10] $\mathbf{I}_n(021)$: proved in [5, 11]; $\mathbf{I}_n(\geq, \neq, \geq), \mathbf{I}_n(>, -, \geq), \mathbf{I}_n(\geq, -, >)$: proved in [12]
B_n	$\mathfrak{S}_n(\underline{2413}, \underline{3142})$: classical result [4]; $\mathbf{I}_n(\geq, \geq, >)$: conjectured in [12]
E_{n+1}	Simsun permutations of $[n]$: classical result [16]; $I_n(000)$: proved in [5]

Figure 1: Sets enumerated by C_n, S_n, B_n or E_{n+1} .

sequences was initiated only recently in [5] and [11]. Since both permutations and inversion sequences will be regarded as words over $\mathbb{N} = \{0, 1, \dots\}$, their patterns can be defined in a unified way as follows.

For two words $W = w_1w_2 \cdots w_n$ and $P = p_1p_2 \cdots p_k$ ($k \leq n$) on \mathbb{N} , we say that W contains the pattern P if there exist some indices $i_1 < i_2 < \cdots < i_k$ such that the subword $W' = w_{i_1}w_{i_2} \cdots w_{i_k}$ of W is order isomorphic to P . Otherwise, W is said to avoid the pattern P . For example, the word $W = 32421$ contains the pattern 231, because the subword $w_2w_3w_5 = 241$ of W has the same relative order as 231. However, W is 101-avoiding. For a set of words \mathcal{W} , the set of words in \mathcal{W} avoiding patterns P_1, \dots, P_r is denoted by $\mathcal{W}(P_1, \dots, P_r)$. One well-known enumeration result in this area, attributed to MacMahon and Knuth (cf. [9]), is that $|\mathfrak{S}_n(123)| = C_n = |\mathfrak{S}_n(132)|$, where $C_n := \frac{1}{n+1} \binom{2n}{n}$ is the n -th Catalan number.

In [5, 11], inversion sequences avoiding patterns of length 3 are exploited, where a number of familiar combinatorial sequences, such as *large Schröder numbers* (denoted S_n) and *Euler numbers* (denoted by E_n), arise. Martinez and Savage [12] further considered a generalization of pattern avoidance to a fixed triple of binary relations (ρ_1, ρ_2, ρ_3) . For each triple of relations $(\rho_1, \rho_2, \rho_3) \in \{<, >, \leq, \geq, =, \neq, -\}^3$, they studied the set $\mathbf{I}_n(\rho_1, \rho_2, \rho_3)$ consisting of those $e \in \mathbf{I}_n$ with no $i < j < k$ such that $e_i \rho_1 e_j$, $e_j \rho_2 e_k$ and $e_i \rho_3 e_k$. Here the relation " $-$ " on a set S is all of $S \times S$, i.e., $x - y$ for all $x, y \in S$. For example, $\mathbf{I}_n(<, >, <) = \mathbf{I}_n(021)$ and $\mathbf{I}_n(\geq, -, \geq) = \mathbf{I}_n(000, 101, 110)$. In Figure 1, we summarize some of their enumeration results and conjectures, as well as corresponding classical facts in permutation patterns. Based on these results, we will investigate more connections between restricted permutations and inversion sequences by considering several classical statistics that we recall below.

For each $\pi \in \mathfrak{S}_n$ and each $e \in \mathbf{I}_n$, let

$$\text{DES}(\pi) := \{i \in [n-1] : \pi_i > \pi_{i+1}\} \quad \text{and} \quad \text{ASC}(e) := \{i \in [n-1] : e_i < e_{i+1}\}$$

be the **descent set** of π and the **ascent set** of e , respectively. Another important property

of the coding Θ is that $\text{DES}(\pi) = \text{ASC}(\Theta(\pi))$ for each $\pi \in \mathfrak{S}_n$. Thus,

$$\sum_{\pi \in \mathfrak{S}_n} t^{\text{DES}(\pi)} = \sum_{e \in \mathbf{I}_n} t^{\text{ASC}(e)}, \quad (1.1)$$

where $t^S := \prod_{i \in S} t_i$ for any set S of positive integers. Throughout this paper, we use the convention that if “ST” is a set-valued statistic, then “st” is the corresponding numerical statistic. For example, $\text{des}(\pi)$ is the cardinality of $\text{DES}(\pi)$ for each π . It is known that $A_n(t) := \sum_{\pi \in \mathfrak{S}_n} t^{\text{des}(\pi)}$ is the classical n -th Eulerian polynomial [7] and each statistic whose distribution gives $A_n(t)$ is called a Eulerian statistic. In view of (1.1), “asc” is a Eulerian statistic on inversion sequences. Let $\text{dist}(e)$ be the number of **distinct positive entries** of e . This statistic was first introduced by Dumont [6], who also showed that it is a Eulerian statistic on inversion sequences. Amazingly, Foata [7] later invented two different codings of permutations called *V-code* and *S-code* to prove the following extension of (1.1).

Theorem 1.1 (Foata 1977). *For each $\pi \in \mathfrak{S}_n$ let $\text{idcs}(\pi) := \text{des}(\pi^{-1})$ be the number of inverse descents of π . Then,*

$$\sum_{\pi \in \mathfrak{S}_n} s^{\text{idcs}(\pi)} t^{\text{DES}(\pi)} = \sum_{e \in \mathbf{I}_n} s^{\text{dist}(e)} t^{\text{ASC}(e)}. \quad (1.2)$$

Partial results regarding the statistics “asc” and “dist” on restricted inversion sequences have already been obtained in [5, 11, 12]. In particular, the ascent polynomial $S_n(t) := \sum_{e \in \mathbf{I}_n(021)} t^{\text{asc}(e)}$ was shown to be *palindromic* via a connection with some *black-white rooted binary trees* in [5]. Inspired by Foata’s result, we will consider the joint distribution of “asc” and “dist” on restricted inversion sequences and prove several restricted versions of (1.2). Another interesting statistic for $e \in \mathbf{I}_n$ is the **last entry** of e , that we denote $\text{last}(e)$. This statistic turns out to be useful in solving some real root problems in [14] and will also lead us to solve two enumeration conjectures.

The rest of this paper deals with refinements of Catalan, Schröder, Baxter and Euler numbers. Two highlights of our results are: (i) a bijection from $\mathbf{I}_n(021)$ to $\mathfrak{S}_n(2413, 4213)$ (see Section 3.2.2); (ii) a refinement of a conjecture due to Martinez and Savage [12] that asserts the cardinality of $\mathbf{I}_n(\geq, \geq, >)$ is the n -th *Baxter number* (denoted B_n), which is proved via Bousquet-Mélou’s *obstinate kernel method* (see Section 4).

2 Catalan numbers

Let (ρ_1, ρ_2, ρ_3) be a relation triple in $\{(\geq, -, \geq), (\geq, -, >), (\geq, \geq, >)\}$. We introduce the parameter $\text{cri}(e)$ for each $e \in \mathbf{I}_n(\rho_1, \rho_2, \rho_3)$, that we call the **critical value** of e , as the minimal integer c such that $(e_1, \dots, e_n, c) \in \mathbf{I}_{n+1}(\rho_1, \rho_2, \rho_3)$. Note that “cri” depends on the relation triple (ρ_1, ρ_2, ρ_3) . For example, if we consider $e = (0, 1, 0, 2, 2, 4)$ as inversion

sequence in $\mathbf{I}_6(\geq, -, >)$, then $\text{cri}(e) = 2$. However, $\text{cri}(e) = 3$ when e is considered as an inversion sequence in $\mathbf{I}_6(\geq, -, \geq)$. The reason to introduce “cri” is that if $e \in \mathbf{I}_n(\rho_1, \rho_2, \rho_3)$, then (e_1, \dots, e_n, k) is in $\mathbf{I}_{n+1}(\rho_1, \rho_2, \rho_3)$ if and only if $\text{cri}(e) \leq k \leq n$. This parameter will play an important role in our study of the *Catalan*, *Schröder* and *Baxter triangles* induced by the statistic “last”.

As a warm-up, we will first show how the critical value can be used to prove that $|\mathbf{I}_n(\geq, -, \geq)| = C_n$, which was conjectured in [12]. Let us define the refinement $C_{n,k} := |\{e \in \mathbf{I}_n(\geq, -, \geq) : \text{last}(e) = k\}|$. The following recurrence shows that the numbers $C_{n,k}$ generate the *Catalan triangle* that has already been widely studied (see [OEIS: A009766](#)).

Proposition 2.1. *For $0 \leq k \leq n - 1$, we have the three-term recurrence*

$$C_{n,k} = C_{n,k-1} + C_{n-1,k}.$$

Proof. Let $\mathfrak{C}_{n,k} := \{e \in \mathbf{I}_n(\geq, -, \geq) : \text{last}(e) = k\}$. We divide $\mathfrak{C}_{n,k}$ into the disjoint union $\mathfrak{A}_{n,k} \cup \mathfrak{B}_{n,k}$, where $\mathfrak{A}_{n,k} = \{e \in \mathfrak{C}_{n,k} : \text{cri}(e_1, e_2, \dots, e_{n-1}) = k\}$ and $\mathfrak{B}_{n,k} = \mathfrak{C}_{n,k} \setminus \mathfrak{A}_{n,k}$. Since $\text{cri}(e_1, e_2, \dots, e_{n-1}) \leq k - 1$ for $e \in \mathfrak{B}_{n,k}$, the mapping that sends $(e_1, e_2, \dots, e_{n-1}, k)$ to $(e_1, e_2, \dots, e_{n-1}, k - 1)$ is a bijection from $\mathfrak{B}_{n,k}$ to $\mathfrak{C}_{n-1,k-1}$. Therefore, the cardinality of $\mathfrak{B}_{n,k}$ is $C_{n-1,k-1}$ and so it remains to show that $|\mathfrak{A}_{n,k}| = C_{n-1,k}$. Now, we are going to construct a bijection $g : \mathfrak{A}_{n,k} \rightarrow \mathfrak{C}_{n-1,k}$, which will complete the proof of the recurrence for $C_{n,k}$. For each $e \in \mathfrak{A}_{n,k}$, there is a unique index i such that $e_i = k - 1$ and $e_{i+1} \leq k - 1$. Define $g(e)$ to be the inversion sequence obtained from e by deleting e_{n-1} , if $e_{n-1} = n - 2$, or by deleting e_i , otherwise. For example, we have $g(0, 1, 1, 3, 2) = (0, 1, 1, 2)$ while $g(0, 1, 1, 2, 2) = (0, 1, 2, 2)$. It is routine to check that g is actually a bijection. \square

Theorem 2.2. *For $n \geq 1$, we have the equidistribution*

$$\sum_{\pi \in \mathfrak{S}_n(123)} t^{\text{des}(\pi)} = \sum_{e \in \mathbf{I}_n(\geq, -, \geq)} t^{\text{dist}(e)}.$$

Proof. Let $e \in \mathbf{I}_n(\geq, -, \geq)$. If $t = \max\{i : e_i = i - 1\} < n$, then it is straightforward to show that e can be decomposed into two smaller inversion sequences: $(e_1, \dots, e_{t-1}, e_{t+1})$ in $\mathbf{I}_t(\geq, -, \geq)$ and $(e_{t+2} - t, e_{t+3} - t, \dots, e_n - t)$ in $\mathbf{I}_{n-1-t}(\geq, -, \geq)$. Using this decomposition, one can show easily that

$$\sum_{n \geq 1} x^n \sum_{e \in \mathbf{I}_n(\geq, -, \geq)} t^{\text{dist}(e)} = \frac{-1 + 2tx(1 + x - tx) + \sqrt{1 - 4tx(1 + x - tx)}}{2t^2x(tx - 1 - x)}.$$

The desired result then follows by comparing this with the o.g.f. for $\sum_{\pi \in \mathfrak{S}_n(123)} t^{\text{des}(\pi)}$ in [OEIS: A166073](#). \square

3 Schröder numbers

3.1 A new Schröder triangle

Theorem 3.1. For $n \geq 1$ and $0 \leq k \leq n - 1$, we have

$$|\{e \in \mathbf{I}_n(\geq, -, >) : \text{last}(e) = k\}| = |\{e \in \mathbf{I}_n(021) : \text{last}(e) \equiv k + 1 \pmod{n}\}|. \quad (3.1)$$

Note that this result is obviously true for $k = n - 1, n - 2, n - 3$. Let us define the Schröder triangle $S_{n,k} := |\{e \in \mathbf{I}_n(\geq, -, >) : \text{last}(e) = k\}|$. We have the following simple recurrence for $S_{n,k}$.

Lemma 3.2. For $0 \leq k \leq n - 3$, we have the four-term recurrence

$$S_{n,k} = S_{n,k-1} + 2S_{n-1,k} - S_{n-1,k-1}.$$

Proof. As in the Catalan case, we divide the set $\mathcal{S}_{n,k} := \{\mathbf{I}_n(\geq, -, >) : \text{last}(e) = k\}$ into the disjoint union $\mathcal{A}_{n,k} \cup \mathcal{B}_{n,k}$, where $\mathcal{A}_{n,k} := \{e \in \mathcal{S}_{n,k} : \text{cri}(e_1, \dots, e_{n-1}) = k\}$ and $\mathcal{B}_{n,k} = \mathcal{S}_{n,k} \setminus \mathcal{A}_{n,k}$. Clearly, there is a natural bijection from $\mathcal{B}_{n,k}$ to $\mathcal{S}_{n,k-1}$, which maps (e_1, \dots, e_{n-1}, k) to $(e_1, \dots, e_{n-1}, k - 1)$. Therefore, the cardinality of $\mathcal{B}_{n,k}$ is $S_{n,k-1}$ and so it remains to show $|\mathcal{A}_{n,k}| = 2S_{n-1,k} - S_{n-1,k-1}$, assuming $k \leq n - 3$. To do this, we further divide $\mathcal{A}_{n,k}$ into the disjoint union $\mathcal{C}_{n,k} \cup \mathcal{D}_{n,k}$, where

$$\mathcal{C}_{n,k} := \{e \in \mathcal{A}_{n,k} : e_{n-1} = n - 2, \text{cri}(e_1, \dots, e_{n-2}) = k\}$$

and $\mathcal{D}_{n,k} = \mathcal{A}_{n,k} \setminus \mathcal{C}_{n,k}$. Obviously, we have $\{(e_1, \dots, e_{n-2}, e_n) : e \in \mathcal{C}_{n,k}\} = \mathcal{A}_{n-1,k}$. Thus, $|\mathcal{C}_{n,k}| = |\mathcal{A}_{n-1,k}| = |\mathcal{S}_{n-1,k}| - |\mathcal{B}_{n-1,k}| = S_{n-1,k} - S_{n-1,k-1}$, which will end the proof once we can define a bijection from $\mathcal{D}_{n,k}$ to $\mathcal{S}_{n-1,k}$.

For each $e \in \mathcal{D}_{n,k}$, if e_i is the left-most entry that equals $\text{cri}(e) = k$, then the entries $e_i, e_{i+1}, \dots, e_{n-1}$ of e must satisfy: (i) $e_i = k$ and $e_{i+1} \leq k$; (ii) $k \leq e_{i+2} \leq e_{i+3} \leq \dots \leq e_{n-1}$, where the inequalities after the entries greater than k are strict. Now removing the right-most entry e_j , such that $e_j = k$ and $i \leq j \leq n - 1$, from e results in an inversion sequence in $\mathcal{S}_{n-1,k}$ (since $e_{n-1} \leq n - 3$) that we denote $f(e)$. For example, we have $f(0, 1, 2, 0, 2, 2) = (0, 1, 2, 0, 2)$, $f(0, 1, 0, 2, 2, 2) = (0, 1, 0, 2, 2)$ and $f(0, 1, 2, 1, 3, 2) = (0, 1, 1, 3, 2)$. We claim that the map $f : \mathcal{D}_{n,k} \rightarrow \mathcal{S}_{n-1,k}$ is a bijection. \square

Proof of Theorem 3.1. It is not hard to show that the right-hand side of (3.2) satisfies the same recurrence relation as $S_{n,k}$, which completes the proof of the theorem. \square

One may ask if there is any other interpretation of $S_{n,k}$ in terms of pattern-avoiding permutations. The following conjecture will answer this question completely, if true.




Figure 2: The outline of inversion sequence $(0, 1, 0, 1, 2, 0, 4)$.

Conjecture 3.3. Let (σ, π) be a pair of patterns of length 4. Then,

$$S_{n,k} = |\{\pi \in \mathfrak{S}_n(\sigma, \pi) : \text{last}(\pi) - 1 = k\}|$$

for any $0 \leq k < n$ if and only if (σ, π) is one of the following nine pairs:

$$(4321, 3421), (3241, 2341), (2431, 2341), (4231, 3241), \\ (4231, 2431), (4231, 3421), (2431, 3241), (3421, 2431), (3421, 3241).$$

3.2 Double Eulerian equidistributions

3.2.1 Statistics

Let $\pi \in \mathfrak{S}_n$ be a permutation. The **values of inverse descents** of π is

$$\text{VID}(\pi) := \{2 \leq i \leq n : \pi_i + 1 \text{ appears to the left of } \pi_i\},$$

which is an important set-valued extension of “ides”. The **positions of left-to-right maxima** of π is $\text{LMA}(\pi) := \{i \in [n] : \pi_i > \pi_j \text{ for all } 1 \leq j < i\}$. Similarly, we can define the **positions of left-to-right mixima** $\text{LMI}(\pi)$, the **positions of right-to-left maxima** $\text{RMA}(\pi)$ and the **positions of right-to-left minima** $\text{RMI}(\pi)$ of π .

Let $e \in \mathbf{I}_n$ be an inversion sequence. The **positions of the last occurrence of distinct positive entries** of e is $\text{DIST}(e) := \{2 \leq i \leq n : e_i \neq 0 \text{ and } e_i \neq e_j \text{ for all } j > i\}$. The **positions of zeros** in e is $\text{ZERO}(e) := \{i \in [n] : e_i = 0\}$. The **positions of the entries of e that achieve maximum** is $\text{EMA}(e) := \{i \in [n] : e_i = i - 1\}$ and the **positions of right-to-left minima** of e is $\text{RMI}(e) := \{i \in [n] : e_i < e_j \text{ for all } j > i\}$.

3.2.2 A sextuple equidistribution

Note that an inversion sequence avoids 021 if and only if its positive entries are weakly increasing, which inspires the following geometric representation.

Definition 3.4 (Outline). Recall that a *Dyck path* of length n is a lattice path in \mathbb{N}^2 from $(0,0)$ to (n,n) using the *east step* $(1,0)$ and the *north step* $(0,1)$, which does not pass above the line $y = x$. Here a Dyck path will be represented as $d_1d_2 \dots d_n$, where d_i is the height of its i -th east step. For each $e \in \mathbf{I}_n(021)$, we associate it with a two-colored Dyck path $d(e) = d_1d_2 \dots d_n$, where the red east steps indicate the positions of zero entries of e like this:

$$d_i = \begin{cases} e_i & \text{if } e_i \neq 0, \\ k & \text{if } e_i = 0 \text{ and } k = \max\{e_1, \dots, e_i\}. \end{cases}$$

For example, if $e = (0,1,0,1,2,0,4) \in \mathbf{I}_7(021)$, then $d(e)$ is the two-colored Dyck path in [Figure 2](#). The two-colored Dyck path $d(e)$ is called the *outline* of e . We introduce the *exposed positions* of e as

$$\text{EXPO}(e) := \{i : i \notin \mathcal{C}(e) \text{ and } i - d_i < j - d_j \text{ for all } j > i\},$$

where $\mathcal{C}(e) = \{i : e_i = 0 \text{ and there is } (a,b), a < i < b, \text{ satisfying } e_a = e_b \neq 0\}$. Continuing with our example, we have $\text{EXPO}(e) = \{2,7\}$.

Theorem 3.5. *There exists a bijection $\Psi : \mathbf{I}_n(021) \rightarrow \mathfrak{S}_n(2413,4213)$ such that*

$$(\text{DIST}, \text{ASC}, \text{ZERO}, \text{EMA}, \text{RMI}, \text{EXPO})e = (\text{VID}, \text{DES}, \text{LMA}, \text{LMI}, \text{RMA}, \text{RMI})\Psi(e)$$

for each $e \in \mathbf{I}_n(021)$.

The idea of constructing Ψ is to draw lines parallel to the diagonal $y = x$ in some specified order and successively label the east steps (of the outline) touched by them. The details are provided in [8]. [Theorem 3.5](#) has two interesting applications: (i) the calculation of the double Eulerian distribution $(\text{des}, \text{ides})$ on $\mathfrak{S}_n(2413,4213)$ using the natural structure of two-colored Dyck paths; (ii) an interpretation of the γ -coefficients of $S_n(t)$ in terms of 021-avoiding inversion sequences via the so-called *Foata–Strehl group action*, which implies the palindromicity and *unimodality* of $S_n(t)$.

3.2.3 Two more equidistributions

Based on calculations, Martinez and Savage [12] suspected that

$$\sum_{e \in \mathbf{I}_n(021)} t^{\text{asc}(e)} = \sum_{e \in \mathbf{I}_n(\geq, \neq, \geq)} t^{\text{asc}(e)} = \sum_{e \in \mathbf{I}_n(>, -, \geq)} s^{\text{asc}(e)}.$$

This follows from [Theorem 3.5](#), the palindromicity of $S_n(t)$ and two more multivariate equidistributions ([Theorems 3.6](#) and [3.7](#)) stated below.

First we introduce a set-valued extension of “dist” different from “DIST”:

$$\text{ROW}(e) := \{e_1, e_2, \dots, e_n\} \setminus \{0\}, \text{ for each } e \in \mathbf{I}_n.$$

Theorem 3.6. For $n \geq 1$, we have

$$\sum_{e \in \mathbf{I}_n(\geq, \neq, \geq)} s^{\text{ROW}(e)} t^{\text{ASC}(e)} u^{\text{last}(e)} = \sum_{e \in \mathbf{I}_n(>, -, \geq)} s^{\text{ROW}(e)} t^{\text{ASC}(e)} u^{\text{last}(e)}.$$

Proof. We can construct a bijection from $\mathbf{I}_n(\geq, \neq, \geq)$ to $\mathbf{I}_n(>, -, \geq)$, which preserves the triple statistics (ROW, ASC, last). Notice that $\mathbf{I}_n(\geq, \neq, \geq) = \mathbf{I}_n(\mathbf{110}, 101, 201, 210)$, while $\mathbf{I}_n(>, -, \geq) = \mathbf{I}_n(\mathbf{100}, 101, 201, 210)$. The idea is to replace iteratively occurrences of pattern 100 in an inversion sequence in $\mathbf{I}_n(\geq, \neq, \geq) \setminus \mathbf{I}_n(>, -, \geq)$ with those of patterns 110, the details of which will be omitted here. \square

Recently, Baril and Vajnovszki [1] constructed a new coding $\Phi : \mathfrak{S}_n \rightarrow \mathbf{I}_n$ satisfying

$$(\text{VID}, \text{DES}, \text{LMA}, \text{LMI}, \text{RMA})\pi = (\text{DIST}, \text{ASC}, \text{ZERO}, \text{EMA}, \text{RMI})\Phi(\pi)$$

for each $\pi \in \mathfrak{S}_n$.

Theorem 3.7. For $n \geq 1$, we have

$$\sum_{\pi \in \mathfrak{S}_n(3142, 3124)} s^{\text{VID}(\pi)} t^{\text{DES}(\pi)} = \sum_{e \in \mathbf{I}_n(\geq, \neq, \geq)} s^{\text{DIST}(e)} t^{\text{ASC}(e)}.$$

Proof. We can show that the Baril–Vajnovszki coding Φ restricts to a bijection between $\mathfrak{S}_n(3124, 3142)$ and $\mathbf{I}_n(\geq, \neq, \geq)$. The details are omitted here. \square

Remark 3.8. Interestingly, we have also been able to show that Φ restricts to a bijection between $\mathfrak{S}_n(2413, 4213)$ and $\mathbf{I}_n(021)$. This restricted Φ does not transform “RMI” to “EXPO”, while our bijection Ψ in [Theorem 3.5](#) does.

4 Baxter numbers

A permutation avoiding both *vincular patterns* (see [9] for the definition) $\underline{2413}$ and $\underline{3142}$ is called a *Baxter permutation*. It is a result of Chung et al. [4] that

$$B_n = |\mathfrak{S}_n(\underline{2413}, \underline{3142})| = \frac{1}{\binom{n+1}{1} \binom{n+1}{2}} \sum_{k=0}^{n-1} \binom{n+1}{k} \binom{n+1}{k+1} \binom{n+1}{k+2}.$$

The number B_n is known as the n -th *Baxter number*. Martinez and Savage [12] conjectured that $|\mathbf{I}_n(\geq, \geq, >)| = B_n$, which can be refined as follows.

Theorem 4.1. For $n \geq 1$, we have the equidistribution

$$\sum_{e \in \mathbf{I}_n(\geq, \geq, >)} u^{n+1-\text{cri}(e)} = \sum_{\pi \in \mathfrak{S}_n(\underline{2413}, \underline{3142})} u^{\text{lma}(\pi) + \text{rma}(\pi)}. \quad (4.1)$$

Corollary 4.2. Define the Baxter triangle as $B_{n,k} := |\{e \in \mathbf{I}_n(\geq, \geq, >) : \text{last}(e) = k\}|$. Then,

$$B_{n,k} = |\{\pi \in \mathfrak{S}_{n-1}(2413, 3142) : \text{lma}(\pi) + \text{rma}(\pi) \geq n - k\}|.$$

The rest of this section is devoted to a sketch of our proof of [Theorem 4.1](#). For each $e \in \mathbf{I}_n(\geq, \geq, >)$, introduce the parameters (p, q) of e , where $p = m + 1 - \text{cri}(e)$ and $q = n - m$ with $m = \max\{e_1, \dots, e_n\}$. After a careful discussion we can obtain the following new rewriting rule (see [\[3\]](#) for other known rewriting rules for Baxter families).

Lemma 4.3. Let $e \in \mathbf{I}_n(\geq, \geq, >)$ be an inversion sequence with parameters (p, q) . Exactly $p + q$ inversion sequences in $\mathbf{I}_{n+1}(\geq, \geq, >)$ when removing their last entries will become e , and their parameters are respectively:

$$(p-1, q+1), (p-2, q+1), \dots, (1, q+1), \\ (1, q+1), (p+1, q), (p+2, q-1), \dots, (p+q, 1).$$

The order in which the parameters are listed corresponds to the inversion sequences with last entries from c to n , where $c = n + 1 - (p + q)$.

Define the formal power series $F(t; u, v) = F(u, v) := \sum_{n,p,q \geq 1} F_{n,p,q} t^n u^p v^q$, where $F_{n,p,q}$ is the number of inversion sequences in $\mathbf{I}_n(\geq, \geq, >)$ with parameters (p, q) . We can turn the above lemma into a functional equation as follows.

Proposition 4.4. We have the following equation for $F(u, v)$:

$$\left(1 + \frac{tv}{1-u} + \frac{tv}{1-v/u}\right) F(u, v) = tuv + tuv \left(1 + \frac{1}{1-u}\right) F(1, v) + \frac{tv}{1-v/u} F(u, u). \quad (4.2)$$

Let $G(u, v) := \sum_{n \geq 1} t^n \sum_{\pi \in \mathfrak{S}_n(2413, 3142)} u^{\text{lma}(\pi)} v^{\text{rma}(\pi)}$. This formal power series $G(u, v)$ was first introduced and studied by Bousquet-Mélou [\[2\]](#). Now, [Theorem 4.1](#) is equivalent to $G(u, u) = F(u, u)$, which will be established by solving [\(4.2\)](#).

Proof of [Theorem 4.1](#). It will be convenient to set $w = v/u$ in [\(4.2\)](#). The equation then becomes

$$\left(1 + \frac{tuw}{1-u} + \frac{tuw}{1-w}\right) F(u, wu) = tu^2w + tu^2w \left(1 + \frac{1}{1-u}\right) F(1, wu) + \frac{tuw}{1-w} F(u, u).$$

Further setting $u = 1 + x$ and $w = 1 + y$ in the above equation yields

$$\frac{xy - t(1+x)(1+y)(x+y)}{t(1+x)(1+y)} F(1+x, (1+x)(1+y)) \\ = xy(1+x) - (1-x^2)yF(1, (1+x)(1+y)) - \tilde{F}(x), \quad (4.3)$$

where $\tilde{F}(x) := xF(1+x, 1+x)$. We call the numerator $K(x, y)$ of the coefficient of $F(1+x, (1+x)(1+y))$ the *kernel* of the above equation:

$$K(x, y) = xy - t(1+x)(1+y)(x+y).$$

We are going to apply the so-called *kernel method* (cf. [2]) to this equation.

As a polynomial in y , the kernel has two roots:

$$Y(x) = \frac{1 - t(1+x)(1+\bar{x}) - \sqrt{1 - 2t(1+x)(1+\bar{x}) - t^2(1-x^2)(1-\bar{x}^2)}}{2t(1+\bar{x})},$$

$$Y'(x) = \frac{1 - t(1+x)(1+\bar{x}) + \sqrt{1 - 2t(1+x)(1+\bar{x}) - t^2(1-x^2)(1-\bar{x}^2)}}{2t(1+\bar{x})},$$

where $\bar{x} = 1/x$. Only the first root can be substituted for y in (4.3), because the term $F(1+x, (1+x)(1+Y'))$ is not a well-defined power series in t (the Taylor expansion of Y' in t does not exist).

Now, we will adopt the *obstinate kernel method* that was invented by Bousquet-Mélou [2, Section 2.2] for producing all the pairs (x, y) that can be legally substituted in (4.3): those are the pairs $(x, Y), (\bar{x}Y, Y), (\bar{x}Y, \bar{x})$ and their dual $(Y, x), (Y, \bar{x}Y), (\bar{x}, \bar{x}Y)$, thanks to the symmetry of the kernel $K(x, y)$. Substituting these 6 pairs into (4.3) and after some manipulations, we obtain

$$\begin{cases} (x - xY^2)\tilde{F}(x) - (Y - x^2Y)\tilde{F}(Y) = (Y - Y^3)(x^2 + x^3) - (x - x^3)(Y^2 + Y^3), \\ (Y\bar{x} - Y^3\bar{x})\tilde{F}(Y\bar{x}) - (Y - Y^3\bar{x}^2)\tilde{F}(Y) = (Y - Y^3)(Y^2\bar{x}^2 + Y^3\bar{x}^3) - (Y\bar{x} - Y^3\bar{x}^3)(Y^2 + Y^3), \\ (Y\bar{x} - Y\bar{x}^3)\tilde{F}(Y\bar{x}) - (\bar{x} - Y^2\bar{x}^3)\tilde{F}(\bar{x}) = (\bar{x} - \bar{x}^3)(Y^2\bar{x}^2 + Y^3\bar{x}^3) - (Y\bar{x} - Y^3\bar{x}^3)(\bar{x}^2 + \bar{x}^3). \end{cases}$$

By eliminating $\tilde{F}(Y)$ and $\tilde{F}(Y\bar{x})$, we get a relation between $\tilde{F}(x)$ and $\tilde{F}(\bar{x})$:

$$\tilde{F}(x) + \tilde{F}(\bar{x}) = \frac{Y(1+x)(x^4 - 2Yx^3 + 2Y^2x - 2Y + 1)}{x^2(Y-1)(Y-x)}. \quad (4.4)$$

But $\tilde{F}(x) = xF(1+x, 1+x)$ is a formal power series in t with coefficients in $x\mathbb{N}[x]$, while $\tilde{F}(\bar{x})$ is a formal power series in t with coefficients in $\bar{x}\mathbb{N}[\bar{x}]$. Therefore, the positive part in x of the right hand side of (4.4) is exactly $\tilde{F}(x)$.

On the other hand, it has been showed in [2, Corollary 3] that if we let $\tilde{G}(x) := xG(1+x, 1+x)$, then

$$\frac{x - 2t(1+x)^2}{t(1+x)^2} \tilde{G}(x) = x^2 - 2R(x), \quad (4.5)$$

where $R(x) = xG(1+x, 1)$. Combining with the relation between $R(x)$ and $R(\bar{x})$ proved in [2, Eq. (8)]:

$$R(x) + R(\bar{x}) = \bar{x}^2Y(1+x^3 - xY),$$

we have

$$\begin{aligned} \tilde{G}(x) + \tilde{G}(\bar{x}) &= \frac{t(1+x)^2}{x-2t(1+x)^2} (x^2 + \bar{x}^2 - 2(R(x) + R(\bar{x}))) \\ &= \frac{t(1+x)^2}{x-2t(1+x)^2} (x^2 + \bar{x}^2 - 2\bar{x}^2Y(1+x^3-xY)). \end{aligned}$$

To check that $\frac{t(1+x)^2}{x-2t(1+x)^2} (x^2 + \bar{x}^2 - 2\bar{x}^2Y(1+x^3-xY))$ equals the right hand side of (4.4) is routine by Maple, which proves that $\tilde{F}(x) = \tilde{G}(x)$. This completes the proof. \square

Since the proof of equidistribution (4.1) uses the obstinate kernel method based on the formal power series, it is natural to ask for a bijective proof.

5 Euler numbers

5.1 Entringer–Eulerian statistics on $\mathbf{I}_n(000)$

As introduced by Simion and Sundaram [16], a permutation $\pi \in \mathfrak{S}_n$ is called a *Simsun permutation* if it has no double descents, even after removing $n, n-1, \dots, k$ for any k . Let RS_n be the set of all Simsun permutations in \mathfrak{S}_n . Using the statistic “last”, we refine a result [5, Corollary 2] by Corteel et al.

Theorem 5.1. *Let $\text{asc}(\pi) := n - 1 - \text{des}(\pi)$ for each $\pi \in \mathfrak{S}_n$. Then,*

$$\sum_{\pi \in RS_n} t^{\text{asc}(\pi)} u^{\text{last}(\pi)} = \sum_{e \in \mathbf{I}_n(000)} t^{\text{dist}(e)} u^{\text{last}(e)+1}.$$

Proof. By combining the simple bijection in [5, Theorem 7] from $\mathbf{I}_n(000)$ to 0-1-2-increasing trees with $n+1$ vertices and a special ordering of the increasing tree representation of permutations due to Maria Monks (see [15, Page 198]). \square

It also follows from the simple bijection in [5, Theorem 7] and a result of Poupard [13, Proposition 1] that the statistic “last + 1” is *Entriangler*. Can the generating function for this Entriangler–Eulerian pair be calculated?

5.2 Double Eulerian distribution on $\mathbf{I}_n(000)$

As an application of Foata’s V-code and S-code, we can prove the following double Eulerian equidistribution.

Theorem 5.2. *Let $\text{iasc}(\pi) := \text{asc}(\pi^{-1})$ for each $\pi \in \mathfrak{S}_n$. Then,*

$$\sum_{\pi \in RS_n} s^{\text{iasc}(\pi)} t^{\text{asc}(\pi)} = \sum_{e \in \mathbf{I}_n(000)} s^{\text{asc}(e)} t^{\text{dist}(e)}.$$

The details for the proof of a set-valued extension of Theorem 5.2 will be reported in a full version of this abstract.

References

- [1] J.-L. Baril and V. Vajnovszki. “A permutation code preserving a double Eulerian bivariate”. 2016. arXiv:[1606.07913](#).
- [2] M. Bousquet-Mélou. “Four classes of pattern-avoiding permutations under one roof: generating trees with two labels”. *Electron. J. Combin.* **9** (2003), Art. #R19. [URL](#).
- [3] S. Burrill, J. Courtiel, E. Fusy, S. Melczer, and M. Mishna. “Tableau sequences, open diagrams, and Baxter families”. *European J. Combin.* **58** (2016), pp. 144–165. [DOI](#).
- [4] F. R. K Chung, R. L Graham, V. E. Hoggatt, and M. Kleiman. “The number of Baxter permutations”. *J. Combin. Theory Ser. A* **24** (1978), pp. 382–394. [DOI](#).
- [5] S. Corteel, M. Martinez, C. D. Savage, and M. Weselcouch. “Patterns in Inversion Sequences I”. *Discrete Math. Theor. Comput. Sci.* **18** (2016), Art. #2. [URL](#).
- [6] D. Dumont. “Interprétations combinatoires des nombres de Genocchi”. *Duke Math. J.* **41** (1974), pp. 305–318. [DOI](#).
- [7] D. Foata. “Distributions eulériennes et mahoniennes sur le groupe des permutations”. *Higher Combinatorics (Berlin, West Germany, September 1977)*. NATO Advanced Study Institutes Series, Vol. 31. Springer, 1977, pp. 27–49. [DOI](#).
- [8] D. Kim and Z. Lin. “A sextuple equidistribution arising in Pattern Avoidance”. 2016. arXiv:[1612.02964](#).
- [9] S. Kitaev. *Patterns in Permutations and Words*. Springer, 2011.
- [10] D. Kremer. “Permutations with forbidden subsequences and a generalized Schröder number”. *Discrete Math.* **218** (2000), pp. 121–130. [DOI](#).
- [11] T. Mansour and M. Shattuck. “Pattern avoidance in inversion sequences”. *Pure Math. Appl.* **25** (2015), pp. 157–176. [DOI](#).
- [12] M. A. Martinez and C. D. Savage. “Patterns in Inversion Sequences II: Inverse Sequences Avoiding Triples of Relations”. 2016. arXiv:[1609.08106](#).
- [13] C. Poupard. “De nouvelles significations énumératives de nombres d’Entringer”. *Discrete Math.* **38** (1982), pp. 265–271. [DOI](#).
- [14] C. D. Savage and M. Visontai. “The s -Eulerian polynomials have only real roots”. *Trans. Amer. Math. Soc.* **367** (2014), pp. 1441–1466. [DOI](#).
- [15] R. Stanley. *Enumerative Combinatorics, Vol. 2*. 2nd ed. Cambridge Studies in Advanced Mathematics, Vol. 49. Cambridge University Press, 2012. [URL](#).
- [16] S. Sundaram. “The homology of partitions with an even number of blocks”. *J. Algebraic Combin.* **4** (1995), pp. 69–92. [DOI](#).