

A KAZHDAN GROUP WITH AN INFINITE OUTER AUTOMORPHISM GROUP

Traian Preda

Abstract. D. Kazhdan has introduced in 1967 the Property (T) for local compact groups (see [3]). In this article we prove that for $n \geq 3$ and $m \in \mathbb{N}$ the group $SL_n(\mathbf{K}) \times \mathcal{M}_{n,m}(\mathbf{K})$ is a Kazhdan group having the outer automorphism group infinite.

Definition 1. ([1]) Let (π, \mathcal{H}) be a unitary representation of a topological group G .
(i) For a subset Q of G and real number $\varepsilon > 0$, a vector $\xi \in \mathcal{H}$ is (Q, ε) -invariant if :

$$\sup_{x \in Q} \|\pi(x)\xi - \xi\| < \varepsilon \|\xi\|.$$

(ii) The representation (π, \mathcal{H}) almost has invariant vectors if it has (Q, ε) - invariant vectors for every compact subset Q of G and every $\varepsilon > 0$. If this holds, we write $1_G \prec \pi$.

(iii) The representation (π, \mathcal{H}) has non - zero invariant vectors if there exists $\xi \neq 0$ in \mathcal{H} such that $\pi(x)\xi = \xi$ for all $g \in G$. If this holds, we write $1_G \subset \pi$.

Definition 2. ([3]) Let G be a topological group.

G has Kazhdan's Property (T), or is a Kazhdan group, if there exists a compact subset Q of G and $\varepsilon > 0$ such that, whenever a unitary representation π of G has a (Q, ε) - invariant vector, then π has a non-zero invariant vector.

Proposition 3. ([1]) Let G be a topological group. The following statements are equivalent:

- (i) G has Kazhdan's Property (T);
- (ii) whenever a unitary representation (π, \mathcal{H}) of G weakly contains 1_G , it contains 1_G (in symbols: $1_G \prec \pi$ implies $1_G \subset \pi$).

Definition 4. Let \mathbf{K} be a field. An absolute value on \mathbf{K} is a real - valued function $x \rightarrow |x|$ such that, for all x and y in \mathbf{K} :

- (i) $|x| \geq 0$ and $|x| = 0 \Leftrightarrow x = 0$

2010 Mathematics Subject Classification: 22D10; 22D45.

Keywords: Representations of topological groups; Kazhdan Property (T); Mautner's lemma.

<http://www.utgjiu.ro/math/sma>

$$(ii) |xy| = |x||y|$$

$$(iii) |x + y| \leq |x| + |y|.$$

An absolute value defines a topology on \mathbf{K} given by the metric

$$d(x, y) = |x - y|.$$

Definition 5. A field \mathbf{K} is a local field if \mathbf{K} can be equipped with an absolute value for which \mathbf{K} is locally compact and not discrete.

Example 6. $\mathbf{K} = \mathbb{R}$ and $\mathbf{K} = \mathbb{C}$ with the usual absolute value are local fields.

Example 7. ([1] and [2]) Groups with Property (T):

a) Compact groups, $SL_n(\mathbb{Z})$ for $n \geq 3$.

b) $SL_n(\mathbf{K})$ for $n \geq 3$ and \mathbf{K} a local field.

Lemma 8. (Mautner's lemma)([1])

Let G be a topological group, and let (π, \mathcal{H}) be a unitary representation of G . Let $x \in G$ and assume that there exists a net $(y_i)_i$ in G such that $\lim_i y_i x y_i^{-1} = e$. If ξ is a vector in \mathcal{H} which is fixed by y_i for all i , then ξ is fixed by x .

Theorem 9. Let \mathbf{K} be a local field. The group $SL_n(\mathbf{K})$ acts on $\mathcal{M}_{n,m}(\mathbf{K})$ by left multiplication $(g, A) \rightarrow gA$, $g \in SL_n(\mathbf{K})$ and $A \in \mathcal{M}_{n,m}(\mathbf{K})$.

Then the semi - direct product $SL_n(\mathbf{K}) \ltimes \mathcal{M}_{n,m}(\mathbf{K})$ has Property (T) for $(\forall)n \geq 3$ and $(\forall)m \in \mathbb{N}$.

Proof. Let (π, \mathcal{H}) be a unitary representation of $G = SL_n(\mathbf{K}) \ltimes \mathcal{M}_{n,m}(\mathbf{K})$ almost having invariant vectors. Since $SL_n(\mathbf{K})$ has Property (T), there exists a non - zero vector $\xi \in \mathcal{H}$ which is $SL_n(\mathbf{K})$ - invariant.

Since \mathbf{K} is non - discret, there exists a net $(\lambda_i)_i$ in \mathbf{K} with $\lambda_i \neq 0$ and such that $\lim_i \lambda_i = 0$.

Let $\Delta_{pq}(x) \in \mathcal{M}_{n,m}(\mathbf{K})$ the matrix with x as (p,q) - entry and 0 elsewhere and $(A_i)_{\alpha\beta} \in SL_n(\mathbf{K})$ the matrix:

$$(A_i)_{\alpha,\beta} = \begin{cases} \lambda_i & \text{if } \alpha = \beta \text{ and } \alpha = p \\ \lambda_i^{-1} & \text{if } \alpha = \beta \text{ and } \alpha = (p+1) \bmod (n+1) + [p/n] \\ 1 & \text{if } \alpha = \beta \text{ and } \alpha \notin \{p, (p+1) \bmod (n+1) + [p/n]\} \\ 0 & \text{if } \alpha \neq \beta \end{cases} \quad (0.1)$$

$\Rightarrow A_i \Delta_{pq}(x) = \delta_{pq}(\lambda_i x)$, where $\delta_{pq}(\lambda_i x) \in \mathcal{M}_{n,m}(\mathbf{K})$ is the matrix with $\lambda_i x$ as (p, q) - entry and 0 elsewhere.

Then $\lim_i A_i \Delta_{pq}(x) = 0_{n,m}$.

Since in G we have

$$(A_i, 0_{n,m})(I_n, \Delta_{pq}(x))(A_i, 0_{n,m})^{-1} = (I_n, A_i \Delta_{pq}(x))$$

and since $\xi \in \mathcal{H}$ is $(A_i, 0_{n,m})$ - invariant \Rightarrow

\Rightarrow from Mautner's Lemma that ξ is $\Delta_{pq}(x)$ - invariant.

Since $\Delta_{pq}(x)$ generates the group $\mathcal{M}_{n,m}(\mathbf{K}) \Rightarrow \xi$ is G - invariant and G has Property (T). \square

Corollary 10. *The groups $SL_n(\mathbf{K}) \times \mathbf{K}^n$ and $SL_n(\mathbb{R}) \times \mathcal{M}_n(\mathbb{R})$ has Property (T), $(\forall)n \geq 3$.*

Proposition 11. *For $\delta \in SL_n(\mathbb{Z})$, let $S_\delta : \Gamma \rightarrow \Gamma$, $S_\delta((\alpha, A)) = (\alpha, A\delta)$, $(\forall)(\alpha, A) \in \Gamma$. Then:*

a) $S_\delta \in \text{Aut}(\Gamma)$.

b) $\Phi : SL_n(\mathbb{Z}) \rightarrow \text{Aut}(\Gamma)$, $\Phi(\delta) = S_\delta$ is a group homomorphism.

c) $S_\delta \in \text{Int}(\Gamma)$ if and only if $\delta \in \{\pm I\}$. In particular, the outer automorphism of Γ is infinit.

Proof. a) $S_\delta((\alpha_1, A_1) \cdot (\alpha_2, A_2)) = S_\delta((\alpha_1, A_1)) \cdot S_\delta((\alpha_2, A_2)) \Leftrightarrow$

$$\Leftrightarrow S_\delta((\alpha_1\alpha_2, A_1 + \alpha_1 A_2)) = (\alpha_1, A_1\delta) \cdot (\alpha_2, A_2\delta) \Leftrightarrow$$

$$\Leftrightarrow (\alpha_1\alpha_2, (A_1 + \alpha_1 A_2)\delta) = (\alpha_1\alpha_2, A_1\delta + \alpha_1 A_2\delta)$$

Analogous $S_{\delta^{-1}}$ is morfism and $S_\delta \cdot S_{\delta^{-1}} = S_{\delta^{-1}} \cdot S_\delta = I_\Gamma$.

$$\text{b) } \Phi(\delta_1 \cdot \delta_2) = \Phi(\delta_1) \cdot \Phi(\delta_2) \Leftrightarrow S_{\delta_1 \cdot \delta_2} = S_{\delta_1} \cdot S_{\delta_2}.$$

c) Assume that $S_\delta \in \text{Int}(\Gamma) \Rightarrow (\exists)(\alpha_0, A_0) \in \Gamma$ such that

$$S_\delta((\alpha, A)) = (\alpha_0, A_0)(\alpha, A)(\alpha_0, A_0)^{-1}, (\forall)(\alpha, A) \in \Gamma.$$

$$\Rightarrow (\alpha, A\delta) = (\alpha_0\alpha\alpha_0^{-1}, A_0 + \alpha_0 A - \alpha_0\alpha\alpha_0^{-1}A_0) \Rightarrow$$

$$\Rightarrow \text{i) } \alpha = \alpha_0\alpha\alpha_0^{-1}, (\forall)\alpha \in SL_n(\mathbb{Z}) \Rightarrow \alpha \in \{\pm I_n\}$$

$$\Rightarrow \text{ii) } A\delta = A_0 \pm A - \alpha A_0, (\forall)\alpha \in SL_n(\mathbb{Z}), (\forall)A \in \mathcal{M}_n(\mathbb{Z}) \Rightarrow A_0 = 0_n \text{ and } \delta = \pm I_n.$$

$$\Rightarrow \text{Out}(\Gamma) = \text{Aut}(\Gamma) / \text{Int}(\Gamma) \text{ is infinite.}$$

\square

References

- [1] B. Bekka, P. de la Harpe, A. Valette, *Kazhdan's Property (T)*, Monography, Cambridge University Press, 2008. [MR2415834](#).
- [2] P. de la Harpe, A. Valette, *La propriété (T) de Kazhdan pour les groupes localement compacts*, Astérisque **175**, Soc. Math. France, 1989. [MR1023471\(90m:22001\)](#). [Zbl 0759.22001](#).
- [3] D. Kazhdan, *Connection of the dual space of a group with the structure of its closed subgroups*, Funct. Anal. Appl. **1** (1967), 63 - 657. [MR209390](#). [Zbl 0168.27602](#).

Surveys in Mathematics and its Applications **7** (2012), 27 – 30

<http://www.utgjiu.ro/math/sma>

Traian Preda
University of Bucharest,
Str. Academiei nr.14, București,
Romania.
e-mail: traianpr@yahoo.com

Surveys in Mathematics and its Applications **7** (2012), 27 – 30
<http://www.utgjiu.ro/math/sma>