Theory and Applications of Categories, Vol. 4, No. 4, 1998, pp. 73–81.

SIMPLICIAL AND CATEGORICAL DIAGRAMS, AND THEIR EQUIVARIANT APPLICATIONS

RUDOLF FRITSCH AND MAREK GOLASIŃSKI

Transmitted by Michael Barr

ABSTRACT. We show that the homotopy category of simplicial diagrams $\mathbb{I}$-$\mathbb{SS}$ indexed by a small category $\mathbb{I}$ is equivalent to a homotopy category of $\mathbb{SS} \downarrow N\mathbb{I}$ simplicial sets over the nerve $N\mathbb{I}$. Then their equivalences, by means of the nerve functor $N : \mathbb{C}at \to \mathbb{SS}$ from the category $\mathbb{C}at$ of small categories, with respective homotopy categories associated to $\mathbb{C}at$ are established. Consequently, an equivariant simplicial version of the Whitehead Theorem is derived.

In his remarkable paper [14], Thomason shows the equivalence of the homotopy categories of $\mathbb{C}at$, the category of small categories, and $\mathbb{SS}$, the category of simplicial sets, by means of the nerve functor $N : \mathbb{C}at \to \mathbb{SS}$ and one of its homotopy inverses (see [7, 9] for details). By [11], the homotopy structure on $\mathbb{C}at$ induces, for every small category $\mathbb{I}$, homotopy structures on the category $\mathbb{C}at \downarrow \mathbb{I}$ of small categories over $\mathbb{I}$ and the category $\mathbb{I}$-$\mathbb{C}at$ of contravariant functors from $\mathbb{I}$ to $\mathbb{C}at$. From [10], it follows that there is a pair of adjoint functors $D : \mathbb{C}at \downarrow \mathbb{I} \to \mathbb{I}$-$\mathbb{C}at$ and $\mathbb{I}\int - : \mathbb{I}$-$\mathbb{C}at \to \mathbb{C}at \downarrow \mathbb{I}$ which establishes an equivalence of respective homotopy categories. Similarly, by [5] the homotopy category of simplicial sets on which a fixed simplicial group G acts is equivalent to the homotopy category of simplicial sets over the classifying complex $\overline{W}G$. From this it follows the well-known fact that the homotopy category of topological spaces on which a fixed discrete group G acts is equivalent to the homotopy category of spaces over the classifying space $K(G, 1)$.

We were influenced by these papers to search for a link between the comma category $\mathbb{SS} \downarrow N\mathbb{I}$ and the category $\mathbb{I}$-$\mathbb{SS}$ of contravariant functors from $\mathbb{I}$ to $\mathbb{SS}$. In Section 1 we define, by means of [3, p.327], a pair of adjoint functors $A : \mathbb{SS} \downarrow N\mathbb{I} \to \mathbb{I}$-$\mathbb{SS}$ and $B : \mathbb{I}$-$\mathbb{SS} \to \mathbb{SS} \downarrow N\mathbb{I}$, and examine in Proposition 1.4 their homotopy properties. Let $\overline{N} : \mathbb{C}at \downarrow \mathbb{I} \to \mathbb{SS} \downarrow N\mathbb{I}$ and $\widetilde{N} : \mathbb{I}$-$\mathbb{C}at \to \mathbb{I}$-$\mathbb{SS}$ be the associated functors to the nerve one $N : \mathbb{C}at \to \mathbb{SS}$. Then by Theorem 1.5, from the diagram of functors

$$\begin{array}{ccc} \mathbb{C}at \downarrow \mathbb{I} & \underset{\mathbb{I}\int -}{\overset{D}{\rightleftarrows}} & \mathbb{I}\text{-}\mathbb{C}at \\ \Big\downarrow{\scriptstyle \overline{N}} & & \Big\downarrow{\scriptstyle \widetilde{N}} \\ \mathbb{SS} \downarrow N\mathbb{I} & \underset{B}{\overset{A}{\rightleftarrows}} & \mathbb{I}\text{-}\mathbb{SS} \end{array}$$

Received by the editors 1998 April 24.

Published on 1998 July 1.

1991 Mathematics Subject Classification: Primary 55P15, 55U10; secondary 18G30, 55P91.

Key words and phrases: comma category, Grothendieck construction, homotopy colimit, pull-back, simplicial set, small category.

© Rudolf Fritsch and Marek Golasiński 1998. Permission to copy for private use granted.

we derive equivalences of respective homotopy categories to generalize the result presented in [8].

Let G be a discrete group, $\mathcal{O}(G)$ the associated category of canonical orbits (see e.g., [1, 2]) and G-$\mathbb{SS}$ the category of G-simplicial sets. Then, there is a canonical functor $\Phi : G\text{-}\mathbb{SS} \to \mathcal{O}(G)\text{-}\mathbb{SS}$ which establishes (cf. [4, 6]) an equivalence of respective homotopy categories. In Section 2, we apply our previous results to deduce an equivalence of the homotopy categories of G-$\mathbb{SS}$ and $\mathbb{SS} \downarrow N\mathcal{O}(G)$. Furthermore, a G-simplicial version of the Whitehead Theorem is derived (cf. [2]).

The second author acknowledges the hospitality and financial support of the Mathematisches Institut, der Universität München during the time this work was done.

1. Main results

Let $\mathbb{I}$ be a small category. As mentioned in the introduction, in [10] a pair of adjoint functors is given

$$\mathbb{C}at \downarrow \mathbb{I} \underset{\mathbb{I}\int\text{-}}{\overset{D}{\rightleftarrows}} \mathbb{I}\text{-}\mathbb{C}at$$

such that the adjunction maps $\eta : \mathrm{id} \to (\mathbb{I}\int\text{-})D$ and $\xi : D(\mathbb{I}\int\text{-}) \to \mathrm{id}$ are weak equivalences. The functor $\mathbb{I}\int\text{-}$ is defined by the Grothendieck construction (see e.g., [13]). If $F : \mathbb{I} \to \mathbb{C}at$ is a contravariant functor then objects in $\mathbb{I}\int F$ are pairs (i, X) with i an object in $\mathbb{I}$ and X an object in $F(i)$, and with morphisms $(\alpha, x) : (i_1, X_1) \to (i_2, X_2)$ consisting of a morphism $\alpha : i_1 \to i_2$ in $\mathbb{I}$ and a morphism $x : X_1 \to F(\alpha)(X_2)$. Composition is defined by $(\alpha, x)(\alpha', x') = (\alpha\alpha', F(\alpha')(x)x')$. For $\alpha : \mathbb{C} \to \mathbb{I}$ (an object in $\mathbb{C}at \downarrow \mathbb{I}$) and an object i in $\mathbb{I}$, $D\alpha(i) = i \downarrow \alpha$ is the comma category and the functor D is defined in an obvious way on morphisms in $\mathbb{I}$.

By [11], the simplicial closed model category structure in the sense of Quillen on the category $\mathbb{C}at$ considered in [14] determines such a structure on the categories $\mathbb{C}at \downarrow \mathbb{I}$ and $\mathbb{I}$-$\mathbb{C}at$, respectively. More precisely, from [11] one deduces immediately

1.1. Proposition. *The categories $\mathbb{C}at \downarrow \mathbb{I}$ and $\mathbb{I}$-$\mathbb{C}at$ with fibrations, cofibrations and weak equivalences of small categories as defined below, are simplicial closed model categories in the sense of Quillen* (*see* [11]) :

1) *the model category structure on $\mathbb{C}at \downarrow \mathbb{I}$ is the one induced by the usual one on the category of small categories $\mathbb{C}at$* (*see* [14]). *In particular, the weak equivalences are the weak equivalences over $\mathbb{I}$*;

2) *a map in $\mathbb{I}$-$\mathbb{C}at$ is a fibration if it is an object-wise fibration* (*resp. weak equivalence*) *in $\mathbb{C}at$* (*see* [14]) *and is a cofibration if it has the left lifting property with respect to maps which are simultaneously fibrations and weak equivalences.*

Let now [1] denote the small category associated with the ordered set $\{0 < 1\}$. Then, any natural transformation $\phi : F_0 \to F_1$ of functors $F_0, F_1 : \mathbb{B} \to \mathbb{C}$ determines a functor $\Theta : \mathbb{B} \times [1] \to \mathbb{C}$. Therefore, for an object $\alpha : \mathbb{B} \to \mathbb{I}$ in $\mathbb{C}at \downarrow \mathbb{I}$ define the functor

$\alpha \times [1] : \mathbb{B} \times [1] \to \mathbb{I}$ such that $\alpha \times [1] = \alpha\pi$, where $\pi : \mathbb{B} \times [1] \to \mathbb{B}$ is the projection functor. Furthermore, let $j_0, j_1 : \alpha \to \alpha \times [1]$ denote the canonical inclusions and let $\alpha : \mathbb{B} \to \mathbb{I}$ and $\beta : \mathbb{C} \to \mathbb{I}$ be objects in $\mathbb{C}at \downarrow \mathbb{I}$. We say that maps $F_0, F_1 : \alpha \to \beta$ are *homotopic* if there is a map $\Theta : \alpha \times [1] \to \beta$ such that $\Theta j_0 = F_0$ and $\Theta j_1 = F_1$. Similarly, for an object $G : \mathbb{I} \to \mathbb{C}at$ in the category $\mathbb{I}$-$\mathbb{C}at$ let $G \times [1] : \mathbb{I} \to \mathbb{C}at$ denote the contravariant functor which is defined by $(G \times [1])(i) = G(i) \times [1]$, for any object i in $\mathbb{I}$ and in the obvious way on morphisms in $\mathbb{I}$. For functors $G, H : \mathbb{I} \to \mathbb{C}at$, two maps $\psi_0, \psi_1 : G \to H$ are *homotopic* if there is a map $\Psi : G \times [1] \to H$ in the category $\mathbb{I}$-$\mathbb{C}at$ such that $\Psi j_0 = \psi_0$ and $\Psi j_1 = \psi_1$, where $j_0, j_1 : G \to G \times [1]$ are the canonical inclusions. Then, we may state the following result.

1.2. PROPOSITION. 1) *The functors* $D : \mathbb{C}at \downarrow \mathbb{I} \longrightarrow \mathbb{I}$-$\mathbb{C}at$ *and* $\mathbb{I}\int - : \mathbb{I}$-$\mathbb{C}at \longrightarrow \mathbb{C}at \downarrow \mathbb{I}$ *preserve weak equivalences.*

2) *If* $\alpha : \mathbb{B} \to \mathbb{I}$ *and* $\beta : \mathbb{C} \to \mathbb{I}$ *are objects in* $\mathbb{C}at \downarrow \mathbb{I}$ *and maps* F_0, $F_1 : \alpha \to \beta$ *are homotopic then the induced functors* DF_0, $DF_1 : \mathbb{I} \to \mathbb{C}at$ *are also homotopic. Conversely, for* $G, H : \mathbb{I} \to \mathbb{C}at$ *and homotopic maps* ψ_0, $\psi_1 : G \to H$ *the induced maps* $\mathbb{I}\int \psi_0 : \mathbb{I}\int G \to \mathbb{I}$ *and* $\mathbb{I}\int \psi_1 : \mathbb{I}\int H \to \mathbb{I}$ *are also homotopic.*

PROOF. 1) Let $F_0, F_1 : \mathbb{I} \to \mathbb{C}at$ be contravariant functors with a natural transformation $F_0 \to F_1$ and such that the functors $F_0(i) \to F_1(i)$ are weak equivalences, for any object i in the category $\mathbb{I}$ and consider the induced functor $\mathbb{I}\int F_0 \to \mathbb{I}\int F_1$. Put N for the nerve functor from the category $\mathbb{C}at$ to the category of simplicial sets $\mathbb{SS}$ and hocolim for the homotopy colimit functor (see [1]) on the category of diagrams of simplicial sets. Then, there is a commutative diagram

$$\begin{array}{ccc} \operatorname{hocolim} NF_0 & \longrightarrow & N(\mathbb{I}\int F_0) \\ \downarrow & & \downarrow \\ \operatorname{hocolim} NF_1 & \longrightarrow & N(\mathbb{I}\int F_1) \end{array}$$

with the horizontal maps as homotopy equivalences in the light of [13]. But by [3, p.335], the map $\operatorname{hocolim} NF_0 \to \operatorname{hocolim} NF_1$ is a weak equivalence. Thus, the map $N(\mathbb{I}\int F_0) \to N(\mathbb{I}\int F_1)$ is a weak equivalence as well.

Let now

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow{F} & \mathbb{D} \\ & {\scriptstyle\alpha}\searrow \quad \swarrow{\scriptstyle\beta} & \\ & \mathbb{I} & \end{array}$$

be a map in the category $\mathbb{C}at \downarrow \mathbb{I}$ with F as a weak equivalence. Then by [12], the comma category $d \downarrow F$ is contractible, for any object d in $\mathbb{D}$. For any object i in $\mathbb{I}$, consider the induced functor

$$i \downarrow \alpha \xrightarrow{i\downarrow F} i \downarrow \beta.$$

Then, for any object $\bar{d} = (d, i \to \beta(d))$ in $i \downarrow \beta$ we have $\bar{d} \downarrow (i \downarrow F) = d \downarrow F$. Hence, the comma category $\bar{d} \downarrow (i \downarrow F)$ is contractible, for any object $\bar{d}$ in the category $i \downarrow \beta$ and again by [12] we get that the functor $i \downarrow F$ is a weak homotopy equivalence.

2) The proof is straightforward. ∎

Denote by $\mathbb{SS} \downarrow N\mathbb{I}$ the category $\mathbb{SS}$ over the nerve $N\mathbb{I}$ of the small category $\mathbb{I}$ and by $\mathbb{I}$-$\mathbb{SS}$ the category of contravariant functors from a small category $\mathbb{I}$ to the category $\mathbb{SS}$ of simplicial sets. Then, the proposition below immediately follows from [11].

1.3. Proposition. *The categories $\mathbb{I}$-$\mathbb{SS}$ and $\mathbb{SS} \downarrow N\mathbb{I}$ with fibrations, cofibrations and weak equivalences as defined below, are simplicial closed model categories in the sense of Quillen (see* [11]*).*

1) *The model category structure on the category $\mathbb{SS} \downarrow N\mathbb{I}$ is the one induced by the usual one on the category of simplicial sets (see* [11]*). In particular, the weak equivalences are the weak homotopy equivalences over $N\mathbb{I}$;*

2) *a map in $\mathbb{I}$-$\mathbb{SS}$ is a fibration (resp. weak equivalence) if it is an object-wise fibration (resp. weak equivalence) in $\mathbb{SS}$ (see* [11]*) and is a cofibration if it has the left lifting property with respect to maps which are simultaneously fibrations and weak equivalences.*

Similarly as in the categories $\mathbb{C}at \downarrow \mathbb{I}$ and $\mathbb{I}$-$\mathbb{C}at$, we define a homotopy notion in the categories $\mathbb{SS} \downarrow N\mathbb{I}$ and $\mathbb{I}$-$\mathbb{SS}$. Then, we have the following

1.4. Proposition. *There is a pair of adjoint functors $A : \mathbb{SS} \downarrow N\mathbb{I} \to \mathbb{I}$-$\mathbb{SS}$ (the left adjoint) and $B : \mathbb{I}$-$\mathbb{SS} \to \mathbb{SS} \downarrow N\mathbb{I}$ (the right adjoint) such that in the terminology of the previous proposition:*

1) *both functors send weak equivalences into weak equivalences and homotopic maps into homotopic ones;*

2) *for every object Θ in $\mathbb{SS} \downarrow N\mathbb{I}$ and every object F in $\mathbb{I}$-$\mathbb{SS}$, the adjunction maps $\Theta \to BA(\Theta)$ and $AB(F) \to F$ are weak equivalences.*

Proof. Given an object $\Theta : \mathbb{X} \to N\mathbb{I}$ in the category $\mathbb{SS} \downarrow N\mathbb{I}$, one defines the contravariant functor $A\Theta : \mathbb{I} \to \mathbb{SS}$ as follows. For an object i in $\mathbb{I}$, let $i \downarrow \mathbb{I}$ be the category over i and $i \downarrow \mathbb{I} \to \mathbb{I}$ the natural projection functor. Then, the simplicial set $(A\Theta)(i)$ is given by the pull-back

$$\begin{array}{ccc} (A\Theta)(i) & \longrightarrow & N(i \downarrow \mathbb{I}) \\ \downarrow & & \downarrow \\ \mathbb{X} & \xrightarrow{\Theta} & N\mathbb{I}. \end{array}$$

Moreover, for a map $i \to i'$ in $\mathbb{I}$, we get the induced simplicial map $(A\Theta)(i') \to (A\Theta)(i)$. Given an object F in the category $\mathbb{I}$-$\mathbb{SS}$, one defines BF as follows. Let $\operatorname{hocolim} F$ be the homotopy colimit of F and define BF as the simplicial map $\operatorname{hocolim} F \to N\mathbb{I}$ given by the natural projection.

The adjunction of a map $f : A\Theta \to F$ in the category $\mathbb{I}$-$\mathbb{SS}$ is determined as follows. Associate, with the induced map $\bar{f} : A\Theta \to F \times N(\text{-} \downarrow \mathbb{I})$ by the maps f and $A\Theta \to N(- \downarrow \mathbb{I})$ given by the pull-back above, the simplicial map $\tilde{f} = \operatorname{colim} \bar{f} : \operatorname{colim} A\Theta \to \operatorname{colim} F \times$

$N(\text{-}\downarrow \mathbb{I})$ over the nerve $N\mathbb{I}$. But, $\operatorname{colim} F \times N(\text{-}\downarrow \mathbb{I}) = \operatorname{hocolim} F$ by [3, Chap. XII] and $\operatorname{colim} A\Theta = \coprod_i (A\Theta)(i)/\sim$, where i runs over all objects in $\mathbb{I}$ and $A\Theta(\alpha)x \sim x$ for $\alpha : i' \to i$ and $x \in A\Theta(i)$. An n-simplex in $\mathbb{X}$ determines an simplex $\Theta(x) = (i_0 \to \cdots \to i_n)$ in the nerve $N\mathbb{I}$. Then, $\widetilde{\Theta(x)} = (i_0 = i_0 \to \cdots \to i_n)$ is a simplex in $N(i_0 \downarrow \mathbb{I})$ and we get an injection $\eta : \mathbb{X} \to \operatorname{colim} A\Theta$ such that $\eta(x) = (x, \widetilde{\Theta(x)})$. On the other hand, for any $(x, \sigma) \in (A\Theta)(i)$ we have $(x, \sigma) \sim (x, \widetilde{\Theta(x)})$. Thus, the map $\eta : \mathbb{X} \to \operatorname{colim} A\Theta$ is an isomorphism. Finally, we have a map $\tilde{f} : \mathbb{X} \to \operatorname{hocolim} F$ such that the diagram



commutes.

Given a map $g : \Theta \to BF$ in the category $\mathbb{SS} \downarrow N\mathbb{I}$ one can also get an adjunction $\tilde{g} : A\Theta \to F$. For an object i in $\mathbb{I}$, we define a simplicial map $\tilde{g}(i) : A\Theta(i) \to F(i)$. By [1, p.338], the simplicial set $\operatorname{hocolim} F$ is isomorphic to the diagonal of the double simplicial set $\coprod_* F$ which in dimension n consists of the union $\coprod_n F = \coprod_{i_0 \to \cdots \to i_n} F(i_n)$. Thus, for an n-simplex x in $\mathbb{X}$ we have $g(x) = y$ with $y \in F(i_n)$ and $\Theta(x) = (i_0 \overset{\alpha_1}{\to} \cdots \overset{\alpha_n}{\to} i_n)$. Hence, for $(x, (i \overset{\alpha}{\to} i_0 \overset{\alpha_1}{\to} \cdots \overset{\alpha_n}{\to} i_n))$ in $(A\Theta)(i)$ we may define $\tilde{g}(i)(x, (i \overset{\alpha}{\to} i_0 \overset{\alpha_0}{\to} \cdots \overset{\alpha_n}{\to} i_n)) = F(\alpha_n \cdots \alpha_0 \alpha)(y)$.

1) Let now $\Theta : \mathbb{X} \to N\mathbb{I}$ and $\Theta' : \mathbb{X}' \to N\mathbb{I}$ be two objects in the category $\mathbb{SS} \downarrow \mathbb{I}$ and $f : \Theta \to \Theta'$ a map such that the associated simplicial map $\mathbb{X} \to \mathbb{X}'$ is a weak homotopy equivalence. Then, for an object i in $\mathbb{I}$ the induced maps $A\Theta(i) \to \mathbb{X}$ and $A\Theta'(I) \to \mathbb{X}'$ have the same homotopy fibre (up to homotopy). Hence, from the long homotopy exact sequences determined by $A\Theta(i) \to \mathbb{X}$ and $A\Theta'(i) \to \mathbb{X}'$, it follows that the map $Af(i) : A\Theta(i) \to A\Theta'(i)$ is a weak homotopy equivalence. Thus, the induced map $Af : A\Theta \to A\Theta'$ in the category $\mathbb{I}$-$\mathbb{SS}$ is a weak homotopy equivalence.

On the other hand, if $f : F \to F'$ is a map in the category $\mathbb{I}$-$\mathbb{SS}$ such that the simplicial maps $f(i) : F(i) \to F'(i)$ are weak homotopy equivalences for any object i in $\mathbb{I}$ then by [3, p.335], one gets that the induced map $\operatorname{hocolim} f : \operatorname{hocolim} F \to \operatorname{hocolim} F'$ is a weak homotopy equivalence. Thus, the map $Bf : BF \to BF'$ is a weak homotopy equivalence. The proof of the preservation of homotopic maps by functors A and B is straightforward.

2) Let $|\text{–}|$ be the geometric realization functor. For an object $\Theta : \mathbb{X} \to N\mathbb{I}$ in the category $\mathbb{SS} \downarrow N\mathbb{I}$, the adjunction map $\tau_\Theta : \Theta \to BA\Theta$ is defined as follows. For n-simplex x in $\mathbb{X}$ let $\Theta(x) = (i_0 \to \cdots \to i_n)$ be an n-simplex in the nerve $N\mathbb{I}$ and $\widetilde{\Theta(x)} = (i_0 = i_0 \to \cdots \to i_n)$ an associated n-simplex in $N(i_0 \downarrow \mathbb{I})$. Then, $(x, \widetilde{\Theta(x)})$ is an n-simplex in $A\Theta$ and $\tau_\Theta(x) = (x, \widetilde{\Theta(x)})$. Moreover, there is a natural projection $\pi_\Theta : \operatorname{hocolim} A\Theta \to \mathbb{X}$ such that $\pi_\Theta \tau_\Theta = \mathrm{id}_\Theta$ and it is a routine to check that the map $|\tau_\Theta||\pi_\Theta|$ is homotopic to the identity map $\mathrm{id}_{|\operatorname{hocolim} A\Theta|}$.

For an object F in the category $\mathbb{I}$-$\mathbb{SS}$, we define the adjunction map $\theta_F : ABF \to F$. Let Θ : hocolim $F \to N\mathbb{I}$ be the natural projection and consider the simplicial map $\theta_F(i) : A\Theta(i) \to F(i)$, for an object i in $\mathbb{I}$ defined as follows. If $(x, (i \overset{\alpha}{\to} i_0 \overset{\alpha_1}{\to} \cdots \overset{\alpha_n}{\to} i_n))$ is an n-simplex in $A\Theta(i)$ then $(x, (i_0 \overset{\alpha_1}{\to} \cdots \overset{\alpha_n}{\to} i_n))$ is an n-simplex in hocolim F with x in $F(i_n)$. Then, $\theta_F(i)(x, (i \overset{\alpha}{\to} i_0 \overset{\alpha_1}{\to} \cdots \overset{\alpha_n}{\to} i_n)) = F(\alpha_n \cdots \alpha_0 \alpha)(x)$. Moreover, there is an injection $\sigma(i) : F(i) \to A\Theta(i)$ given by $\sigma(i)(x) = (x, \widetilde{\mathrm{id}_i})$, for an n-simplex x in $F(i)$ with $\widetilde{\mathrm{id}_i}$ the degenerated n-simplex determined by an object i in $\mathbb{I}$. Of course $\theta_F(i)\sigma(i) = \mathrm{id}_{F(i)}$ and it is also easy to check that the map $|\sigma(i)||\theta_F(i)|$ is homotopic to the identity map $\mathrm{id}_{|A\Theta(i)|}$ for any object i in $\mathbb{I}$, and the proof is completed. ∎

The nerve functor $N : \mathbb{C}at \to \mathbb{SS}$ determines functors $\overline{N} : \mathbb{C}at \downarrow \mathbb{I} \to \mathbb{SS} \downarrow N\mathbb{I}$ and $\widetilde{N} : \mathbb{I}$-$\mathbb{C}at \to \mathbb{I}$-$\mathbb{SS}$, respectively. Then, the diagram

$$\begin{array}{ccc} \mathbb{C}at \downarrow \mathbb{I} & \xrightarrow{D} & \mathbb{I}\text{-}\mathbb{C}at \\ \downarrow{\scriptstyle \overline{N}} & & \downarrow{\scriptstyle \widetilde{N}} \\ \mathbb{SS} \downarrow N\mathbb{I} & \xrightarrow{A} & \mathbb{I}\text{-}\mathbb{SS} \end{array}$$

commutes and by [13], the diagram

$$\begin{array}{ccc} \mathbb{C}at \downarrow \mathbb{I} & \xleftarrow{\mathbb{I}\int -} & \mathbb{I}\text{-}\mathbb{C}at \\ \downarrow{\scriptstyle \overline{N}} & & \downarrow{\scriptstyle \widetilde{N}} \\ \mathbb{SS} \downarrow N\mathbb{I} & \xleftarrow[B]{} & \mathbb{I}\text{-}\mathbb{SS} \end{array}$$

commutes up to homotopy. Moreover, by [7, 9] there are homotopy inverses $\Gamma : \mathbb{SS} \to \mathbb{C}at$, for the nerve functor $N : \mathbb{C}at \to \mathbb{SS}$ such that for any small category $\mathbb{I}$ there is a weak equivalence $\Gamma N\mathbb{I} \to \mathbb{I}$. Thus, we get induced functors $\overline{\Gamma} : \mathbb{SS} \downarrow N\mathbb{I} \to \mathbb{C}at \downarrow \mathbb{I}$ and $\widetilde{\Gamma} : \mathbb{I}$-$\mathbb{SS} \to \mathbb{I}$-$\mathbb{C}at$, respectively. Consequently, we may generalize the result presented in [8] and summarize this section with the following

1.5. THEOREM. *For any small category $\mathbb{I}$, the diagram of above functors*

$$\begin{array}{ccc} \mathbb{C}at \downarrow \mathbb{I} & \underset{\mathbb{I}\int -}{\overset{D}{\rightleftarrows}} & \mathbb{I}\text{-}\mathbb{C}at \\ \overline{\Gamma} \uparrow \downarrow \overline{N} & & \widetilde{\Gamma} \uparrow \downarrow \widetilde{N} \\ \mathbb{SS} \downarrow N\mathbb{I} & \underset{B}{\overset{A}{\rightleftarrows}} & \mathbb{I}\text{-}\mathbb{SS} \end{array}$$

determines equivalences of respective homotopy categories.

In particular, if $\mathbb{I}$ is the category associated with a discrete group G then the nerve $N\mathbb{I}$ is the classifying complex $\overline{W}G$ and $\mathbb{I}$-$\mathbb{C}at$ (resp. $\mathbb{I}$-$\mathbb{SS}$) is the category of right G-small categories (resp. right G-simplicial sets). Thus from the above, we may also deduce an equivalence of respective homotopy categories (cf. [5]).

2. Equivariant applications

Let G be a discrete group and $\mathcal{O}(G)$ the associated category of canonical orbits; its objects are the left cosets G/H as H ranges over all subgroups of G and morphisms are G-maps $G/H \to G/K$ with respect to left translation. If G-$\mathbb{SS}$ denotes the category of G-simplicial sets then there is a canonical functor $\Phi : G\text{-}\mathbb{SS} \to \mathcal{O}(G)\text{-}\mathbb{SS}$ defined on objects by $\Phi(\mathbb{X})(G/H) = \mathbb{X}^H$ and in the obvious way on morphisms in G-$\mathbb{SS}$, where $\mathbb{X}^H$ is the H-fixed point simplicial subset of $\mathbb{X}$. Then (cf. [1]), the category G-$\mathbb{SS}$ is a closed model category; a map $f : \mathbb{X} \to \mathbb{Y}$ is a *fibration* (resp. *weak equivalence*) if the induced map $\Phi(f) : \Phi(\mathbb{X}) \to \Phi(\mathbb{Y})$ is a fibration (resp. weak equivalence) in $\mathcal{O}(G)$-$\mathbb{SS}$. *Cofibrations* in G-$\mathbb{SS}$ are determined by means of the left lifting property with respect to maps which are simultaneously fibrations and weak equivalences. From [4, 6], one may derive that the functor $\Phi : G\text{-}\mathbb{SS} \to \mathcal{O}(G)\text{-}\mathbb{SS}$ establishes an equivalence of respective homotopy categories. Therefore, Proposition 1.4 yields

2.1. Corollary. *If G is a discrete group then the homotopy categories of G-$\mathbb{SS}$ and $\mathbb{SS} \downarrow N\mathcal{O}(G)$ are equivalent.*

Recall that a G-simplicial set $\mathbb{X}$ is called *fibrant* (resp. *cofibrant*) if the unique map $\mathbb{X} \to *$ (resp. $\emptyset \to \mathbb{X}$) is a fibration (resp. cofibration) in the category G-$\mathbb{SS}$, where $\emptyset$ is the empty simplicial set and $*$ a single point simplicial set with the trivial action of G.

2.2. Lemma. *Every G-simplicial set $\mathbb{X}$ is cofibrant in the category G-$\mathbb{SS}$.*

Proof. Let $p : \mathbb{E} \to \mathbb{B}$ be a fibration and a weak equivalence in the category G-$\mathbb{SS}$, and $f : \mathbb{X} \to \mathbb{B}$ a G-map. The n-skeleton $\mathbb{X}^{(n)}$, for $n \geq 0$ is a G-simplicial subset of $\mathbb{X}$ and $\mathbb{X} = \mathrm{colim}_n\, \mathbb{X}^{(n)}$. Let $f^{(n)}$ denote the restriction of f to $\mathbb{X}^{(n)}$; we proceed by induction to show an existence of a G-map $\tilde{f}^{(n)} : \mathbb{X}^{(n)} \to \mathbb{E}$ with $p\tilde{f}^{(n)} = f^{(n)}$, for $n \geq 0$.

Of course, there is a G-map $\tilde{f}^{(0)} \to \mathbb{E}$ such $p\tilde{f}^{(0)} = f^{(0)}$. Suppose there is a G-map $\tilde{f}^{(n-1)} : \mathbb{X}^{(n-1)} \to \mathbb{E}$ with $p\tilde{f}^{(n-1)} = f^{(n-1)}$ and put $\Delta[n]$ for the standard simplicial n-simplex, and $\dot{\Delta}[n]$ for its boundary. Then, there is a pushout diagram

$$\begin{array}{ccc} \coprod_x G/H_x \times \dot{\Delta}[n] & \longrightarrow & \mathbb{X}^{(n-1)} \\ \downarrow & & \downarrow \\ \coprod_x G/H_x \times \Delta[n] & \longrightarrow & \mathbb{X}^{(n)}, \end{array}$$

where x runs over all non-degenerate n-simplexes in $\mathbb{X}$ and H_x is the isotropy subgroup of an n-simplex x. By means of the right lifting property of the map $p : \mathbb{E} \to \mathbb{B}$ (see e.g., [11, p.2.1]) and the map $\tilde{f}^{(n-1)} : \mathbb{X}^{(n-1)} \to \mathbb{E}$, we get a G-map $\tilde{f}^{(n)} : \mathbb{X}^{(n)} \to \mathbb{E}$ such that $p\tilde{f}^{(n)} = f^{(n)}$ and its restriction to $\mathbb{X}^{(n-1)}$ is equal to $\tilde{f}^{(n-1)}$. Therefore, the sequence of G-map $\tilde{f}^{(n)} \to \mathbb{E}$, for $n \geq 0$ determines a G-map $\tilde{f} : \mathbb{X} \to \mathbb{E}$ with $p\tilde{f} = f$ and the proof is completed. ∎

Let now $f : \mathbb{X} \to \mathbb{Y}$ be a weak equivalence in the category G-$\mathbb{SS}$. Then, from [11, p.3.15] it follows that the map $[f, \mathbb{E}]_G : [\mathbb{Y}, \mathbb{E}]_G \to [\mathbb{X}, \mathbb{E}]_G$ of G-homotopy classes is bijective, for any fibrant G-simplicial set $\mathbb{E}$. We are now in a position to state the following G-simplicial version of the Whitehead Theorem (cf.[2]).

2.3. COROLLARY. *Let $f : \mathbb{X} \to \mathbb{Y}$ be a G-map of fibrant G-simplicial sets. Then the following conditions are equivalent:*

1) *the G-map $f : \mathbb{X} \to \mathbb{Y}$ is a weak equivalence;*
2) *the G-map $f : \mathbb{X} \to \mathbb{Y}$ is a G-homotopy equivalence;*
3) *the map in the category $\mathbb{SS} \downarrow N\mathcal{O}(G)$, induced by the commutative diagram*



is a homotopy equivalence.

PROOF. Observe that by Proposition 1.4, we get 2) $\Rightarrow$ 3) and 3) $\Rightarrow$ 1). We show now the implication 1) $\Rightarrow$ 2). By Lemma 2.2 and [11, p.3.15], the induced maps $[f, \mathbb{X}]_G : [\mathbb{Y}, \mathbb{X}]_G \to [\mathbb{X}, \mathbb{X}]_G$ and $[f, \mathbb{Y}]_G : [\mathbb{Y}, \mathbb{Y}]_G \to [\mathbb{X}, \mathbb{Y}]_G$ of G-homotopy classes are bijective. Therefore, the G-map $f : \mathbb{X} \to \mathbb{Y}$ is a G-homotopy equivalence, as claimed. ∎

References

[1] M. Arkowitz, M. Golasiński, *Co-H-structures on equivariant Moore spaces*, Fund. Math. 146 (1994), 59-67.

[2] G.E., Bredon, *Equivariant cohomology theories*, Lecture Notes in Math. 34, Springer-Verlag (1967).

[3] A.K. Bousfield, D.M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Math. 304, Springer-Verlag, Berlin (1972).

[4] J.-M., Cordier, T. Porter, *Categorical aspect of equivariant homotopy*, Applied Categorical Structures 4 (1996), 195-212.

[5] E. Dror, W.G. Dwyer, D.M. Kan, *Equivariant maps which are self homotopy equivalence*, Proc. Amer. Math. Soc. 80 (1980), 670-672.

[6] A.. Elmendorf, *Systems of fixed point sets*, Trans. Amer. Math. Soc. 277 (1983), 275-284.

[7] R. Fritsch, D.M. Latch, *Homotopy inverses for nerve*, Math. Z. 177 (1981), 147-179.

[8] D.M. Latch, *a Fibred homotopy equivalence and homology theories for the category of small categories*, J. Pure Appl. Algebra 15 (1976), 247-269.

[9] _____ , R.W. Thomason, W.S. Wilson, *Simplicial sets from categories*, Math. Z. 164 (1979), 195-214.

[10] M. Nunes, *Cohomological results in monoid and category theory via classifying spaces*, J. Pure Appl. Algebra 101 (1995), 213-244.

[11] D.G. Quillen, *Homotopical algebra*, Lecture Notes in Math. 43, Springer-Verlag (1967).

[12] _____ , *Higher Algebraic K-theory, I*, Lecture Notes in Math. 341, Springer-Verlag, Berlin (1973), 85-147.

[13] R.W. Thomason, *Homotopy colimits in the category of small categories*, Math. Proc. Phil. Soc. 85 (1976), 91-109.

[14] _____ , $\mathbb{C}at$ *as a closed model category*, Cahiers Topologie Géom. Différentielle 21 (1980), 305-324.

Mathematisches Institut der Universität München
Theresienstasse 39, D-80333 München, Germany
and
Faculty of Mathematics and Informatics, Nicholas Copernicus University
87-100 Toruń, Chopina 12/18, Poland
Email: fritsch@rz.mathematik.uni-muenchen.de and marek@mat.uni.torun.pl

This article may be accessed via WWW at http://www.tac.mta.ca/tac/ or by anonymous ftp at ftp://ftp.tac.mta.ca/pub/tac/html/volumes/1998/n4/n4.{dvi,ps}

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

The method of distribution of the journal is via the Internet tools `WWW/ftp`. The journal is archived electronically and in printed paper format.

Subscription information. Individual subscribers receive (by e-mail) abstracts of articles as they are published. Full text of published articles is available in .dvi and Postscript format. Details will be e-mailed to new subscribers and are available by `WWW/ftp`. To subscribe, send e-mail to `tac@mta.ca` including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, `rrosebrugh@mta.ca`.

Information for authors. The typesetting language of the journal is TEX, and LATEX is the preferred flavour. TEX source of articles for publication should be submitted by e-mail directly to an appropriate Editor. They are listed below. Please obtain detailed information on submission format and style files from the journal's WWW server at `URL http://www.tac.mta.ca/tac/` or by anonymous ftp from `ftp.tac.mta.ca` in the directory `pub/tac/info`. You may also write to `tac@mta.ca` to receive details by e-mail.

Editorial board.
John Baez, University of California, Riverside: `baez@math.ucr.edu`
Michael Barr, McGill University: `barr@math.mcgill.ca`
Lawrence Breen, Université de Paris 13: `breen@math.univ-paris13.fr`
Ronald Brown, University of North Wales: `r.brown@bangor.ac.uk`
Jean-Luc Brylinski, Pennsylvania State University: `jlb@math.psu.edu`
Aurelio Carboni, Università della Calabria: `carboni@unical.it`
P. T. Johnstone, University of Cambridge: `ptj@pmms.cam.ac.uk`
G. Max Kelly, University of Sydney: `kelly_m@maths.usyd.edu.au`
Anders Kock, University of Aarhus: `kock@mi.aau.dk`
F. William Lawvere, State University of New York at Buffalo: `wlawvere@acsu.buffalo.edu`
Jean-Louis Loday, Université de Strasbourg: `loday@math.u-strasbg.fr`
Ieke Moerdijk, University of Utrecht: `moerdijk@math.ruu.nl`
Susan Niefield, Union College: `niefiels@union.edu`
Robert Paré, Dalhousie University: `pare@mscs.dal.ca`
Andrew Pitts, University of Cambridge: `ap@cl.cam.ac.uk`
Robert Rosebrugh, Mount Allison University: `rrosebrugh@mta.ca`
Jiri Rosicky, Masaryk University: `rosicky@math.muni.cz`
James Stasheff, University of North Carolina: `jds@charlie.math.unc.edu`
Ross Street, Macquarie University: `street@macadam.mpce.mq.edu.au`
Walter Tholen, York University: `tholen@mathstat.yorku.ca`
Myles Tierney, Rutgers University: `tierney@math.rutgers.edu`
Robert F. C. Walters, University of Sydney: `walters_b@maths.usyd.edu.au`
R. J. Wood, Dalhousie University: `rjwood@mscs.dal.ca`