

TENSOR-TRIANGULATED CATEGORIES AND DUALITIES

BAPTISTE CALMÈS AND JENS HORNBOSTEL

ABSTRACT. In a triangulated closed symmetric monoidal category, there are natural dualities induced by the internal Hom. Given a monoidal exact functor f^* between two such categories and adjoint couples (f^*, f_*) , $(f_*, f^!)$, we establish the commutative diagrams necessary for f^* and f_* to respect certain dualities, for a projection formula to hold between them (as duality preserving exact functors) and for classical base change and composition formulas to hold when such duality preserving functors are composed. This framework allows us to define push-forwards for Witt groups, for example.

Contents

1	Adjunctions and consequences	141
2	Dualities	156
3	Consequences of the closed monoidal structure	159
4	Functors between closed monoidal categories	163
5	Relations between functors	173
6	Various reformulations	187
A	Signs in the category of complexes	195

Introduction

Several cohomology theories on schemes use the concept of a duality in their definition. In fact, these cohomology theories are usually defined at a categorical level; a group is associated to a category with some additional structure, including a duality which is then a contravariant endofunctor of order two (up to a natural isomorphism) on the category. The main example that we have in mind is the Witt group, defined for a triangulated category with duality, the (coherent or locally free) Witt groups of a scheme then being defined by using one of the various bounded derived categories of a scheme endowed with a duality coming from the derived functor RHom . Of course, most of the structural properties of these cohomology theories should be proved at a categorical level. For example, Paul Balmer has proved localization for Witt groups directly at the level of triangulated categories by using localization properties of such categories [B00]. In the rest of the article, we only discuss the example of Witt groups, but everything works

Received by the editors 2008-06-03 and, in revised form, 2009-04-09.

Transmitted by Martin Hyland. Published on 2009-04-23.

2000 Mathematics Subject Classification: 18D10.

Key words and phrases: closed monoidal category, commutative diagram, duality, Witt group.

© Baptiste Calmès and Jens Hornbostel, 2009. Permission to copy for private use granted.

exactly the same way for Grothendieck-Witt groups. The reader should also have in mind that similar considerations should apply to hermitian K-theory and Chow-Witt theory, etc.

We are interested in the functoriality of Witt groups along morphisms of schemes. At the categorical level, this means that we are given functors between categories (such as the derived functors of f^* or f_*) and that we would like to use them to induce morphisms between Witt groups. Let us elaborate on this theme. The basic ideas are:

- a functor between categories respecting the duality induces a morphism of groups and
- two such morphisms can be compared if there is a morphism of functors between two such functors, again respecting the duality.

By “respecting the duality”, we actually mean the following. Let (\mathcal{C}, D, ϖ) be a category with duality, which means that $D : \mathcal{C} \rightarrow \mathcal{C}$ is a contravariant functor, that $\varpi : Id_{\mathcal{C}} \rightarrow D^2$ is a morphism of functors¹ and that for any object A , we have

$$D(\varpi_A) \circ \varpi_{DA} = Id_{DA}. \tag{1}$$

A duality preserving functor from $(\mathcal{C}_1, D_1, \varpi_1)$ to $(\mathcal{C}_2, D_2, \varpi_2)$ is a pair $\{F, \phi\}$ where $F : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ is a functor and $\phi : FD \rightarrow DF$ is a morphism of functors, such that the diagram of morphism of functors

$$\begin{array}{ccc} F & \xrightarrow{F\varpi_1} & FD_1D_1 \\ \varpi_2F \downarrow & & \downarrow \phi_{D_1} \\ D_2D_2F & \xrightarrow{D_2\phi} & D_2FD_1 \end{array} \tag{2}$$

commutes. In practice, the functor F is usually given. For example, in the context of schemes, it can be the derived functor of the pull-back or of the push-forward along a morphism of schemes. It then remains to find an interesting ϕ and to check the commutativity of diagram (2), which can be very intricate.

A morphism of duality preserving functors $\{F, \phi\} \rightarrow \{G, \psi\}$ is a morphism of functors $\rho : F \rightarrow G$ such that the diagram

$$\begin{array}{ccc} FD_1 & \xrightarrow{\rho_{D_1}} & GD_1 \\ \phi \downarrow & & \downarrow \psi \\ D_2F & \xleftarrow{D_2\rho} & D_2G \end{array} \tag{3}$$

commutes. This situation with two functors arises when we want to compare what two different functors (respecting the dualities) yield as Witt group morphisms. For example,

¹By default, all dualities considered in this article are *weak*, which means that we do not require ϖ to be an isomorphism, because we are only interested in the commutativity of diagrams.

we might want to compare the composition of two pull-backs and the pull-back of the composition, in the case of morphisms of schemes. In this kind of situation, usually $\{F, \phi\}$ and $\{G, \psi\}$ are already given, ρ has to be found and the commutative diagram (3) has to be proved.

It turns out that in the context of closed symmetric monoidal categories, this type of diagram commutes for certain dualities, functors and morphisms of functors arising in a natural way. The main object of this article is to prove it. The main results of the paper are in sections 4 and 5 and the reader might want to have a quick look at them before reading the other sections.

Our original motivation for dealing with the questions of this paper was to define push-forward morphisms for Witt groups with respect to proper maps of schemes and study their properties, which in turn should be useful both for general theorems and concrete computations of Witt groups of schemes. The article [CH08] uses the results of this article to establish these push-forwards. We refer the reader to loc. cit. for details. As Witt groups are the main application that we have in mind, we have stated in corollaries of the main theorems what they imply for Witt groups. The proofs of these corollaries are trivial, and just rely on the classical propositions 2.2.3 and 2.2.4. This should be seen as some help for the reader familiar with cohomology theories and geometric intuition. The reader not interested in Witt groups or other Witt-like cohomology theories might just skip the corollaries. In all of them, the triangulated categories are assumed to be $\mathbb{Z}[1/2]$ -linear (to be able to define triangulated Witt groups). Also note that our formalism applies to other settings such as Witt groups in topology.

We now give a more precise description of the functors and morphisms of functors we consider.

DUALITIES. Assume we are given a closed symmetric monoidal category \mathcal{C} where the tensor product is denoted by \otimes and its “right adjoint”, the internal Hom, is denoted by $[-, -]$. Then, as recalled in Section 3, fixing an object K and setting $D_K = [-, K]$, a natural morphism of functors $\varpi_K : Id \rightarrow D_K D_K$ can be defined using the symmetry and the adjunction of the tensor product and the internal Hom and the formula (1) is satisfied. This is well known and recalled here just for the sake of completeness.

PULL-BACK. Assume now that we are given a monoidal functor denoted by f^* (by analogy with the algebro-geometric case) from \mathcal{C}_1 to \mathcal{C}_2 , then, for any object K , as explained above, we can consider the categories with duality $(\mathcal{C}_1, D_K, \varpi_K)$ and $(\mathcal{C}_2, D_{f^*K}, \varpi_{f^*K})$. There is a natural morphism of functors $\beta_K : f^* D_K \rightarrow D_{f^*K} f^*$ such that diagram (2) is commutative (see Proposition 4.1.1 and Theorem 4.1.2). In other words, $\{f^*, \beta_K\}$ is a duality preserving functor from $(\mathcal{C}_1, D_K, \varpi_K)$ to $(\mathcal{C}_2, D_{f^*K}, \varpi_{f^*K})$.

PUSH-FORWARD. Assume furthermore that that we are given a right adjoint f_* , then there is a natural morphism of projection $\pi : f_*(-) \otimes * \rightarrow f_*(- \otimes f^*(*))$ (see Proposition 4.2.5). When f_* also has a right adjoint $f^!$, we can consider the categories with duality $(\mathcal{C}_1, D_K, \varpi_K)$ and $(\mathcal{C}_2, D_{f^!K}, \varpi_{f^!K})$. There is a natural isomorphism $\zeta_K : f_* D_{f^!K} \rightarrow D_K f_*$

such that diagram (2) is commutative. In other words, $\{f_*, \zeta_K\}$ is a duality preserving functor from $(\mathcal{C}_2, D_{f^!K}, \varpi_{f^!K})$ to $(\mathcal{C}_1, D_K, \varpi_K)$.

PRODUCT. Theorems involving products of dualities are stated. Given a pair of categories with duality $(\mathcal{C}_1, D_1, \varpi_1)$ and $(\mathcal{C}_2, D_2, \varpi_2)$, there is an obvious structure of a category with duality $(\mathcal{C}_1 \times \mathcal{C}_2, D_1 \times D_2, \varpi_1 \times \varpi_2)$. If \mathcal{C} is closed monoidal symmetric, there is a natural morphism of (bi)functors $\tau_{K,M} : D_K \otimes D_M \rightarrow D(- \otimes -)_{K \otimes M}$ (see Definition 4.4.1). Proposition 4.4.6 recalls that $\{- \otimes -, \tau_{K,M}\}$ is a duality preserving functor from $(\mathcal{C} \times \mathcal{C}, D_K \times D_M, \varpi_K \times \varpi_M)$ to $(\mathcal{C}, D_{K \otimes M}, \varpi_{K \otimes M})$. This gives a product on Witt groups.

We now explain relations between the functors $\{f^*, \beta\}$, $\{f_*, \zeta\}$ and $\{\otimes, \tau\}$ in different contexts. Before going any further, let us remark that a morphism $\iota : K \rightarrow M$ induces a morphism of functors $\tilde{\iota} : D_K \rightarrow D_M$ which is easily shown to yield a duality preserving functor $I_\iota = \{Id, \tilde{\iota}\}$ from $(\mathcal{C}, D_K, \varpi_K)$ to $(\mathcal{C}, D_M, \varpi_M)$.

COMPOSITION. There is a natural way to compose duality preserving functors given by $\{F, f\} \circ \{G, g\} := \{FG, gF \circ fG\}$. Suppose that we are given a pseudo contravariant functor $(-)^*$ from a category \mathcal{B} to a category where the objects are closed symmetric monoidal categories and the morphisms are monoidal functors. As usual, ‘‘pseudo’’ means that we have almost a functor, except that we only have a natural isomorphism $a_{g,f} : f^*g^* \simeq (gf)^*$ instead of an equality. Then an obvious question arises: can we compare $\{(gf)^*, \beta_K\}$ with $\{f^*, \beta_{g^*K}\} \circ \{g^*, \beta_K\}$ for composable morphisms f and g in \mathcal{B} . The answer is that the natural isomorphism a is a morphism of duality preserving functors from $I_{a_{g,f,K}} \circ \{f^*, \beta_{g^*K}\} \circ \{g^*, \beta_K\}$ to $\{(gf)^*, \beta_K\}$, i.e. the diagram (3) commutes (see 5.1.3) with $\rho = a$. The correction by $I_{a_{g,f,K}}$ is necessary for otherwise, strictly speaking, the target categories of the two duality preserving functors are not equipped with the same duality.

A similar composition question arises when some $f \in \mathcal{B}$ are such that there are adjunctions (f^*, f_*) and $(f_*, f^!)$. On the subcategory \mathcal{B}' of such f , there is a way of defining natural (with respect to the adjunctions) pseudo functor structures $b : (gf)_* \simeq g_*f_*$ and $c : f^!g^! \simeq (gf)^!$, and we can compare $\{(gf)_*, \zeta_K\}$ and $\{g_*, \zeta_{g^!K}\} \circ \{f_*, \zeta_{f^!K}\}$ using the natural morphism of functors b up to a small correction of the duality using c as above (see Theorem 5.1.9).

BASE CHANGE. A ‘‘base change’’ question arises when we are given a commutative diagram

$$\begin{array}{ccc} V & \xrightarrow{\bar{g}} & Y \\ \bar{f} \downarrow & & \downarrow f \\ X & \xrightarrow{g} & Z \end{array}$$

in \mathcal{B} such that g and \bar{g} are in \mathcal{B}' . Then, there is a natural morphism $\varepsilon : f^*g_* \rightarrow \bar{g}_*\bar{f}^*$ (see Section 5.2). When ε is an isomorphism, there is a natural morphism $\gamma : \bar{f}^*g^! \rightarrow \bar{g}^!f^*$. Starting from an object $K \in \mathcal{C}_Z$, the morphism of functors γ is used to define a duality preserving functor I_{γ_K} from $(\mathcal{C}_V, D_{\bar{f}^*g^!K}, \varpi_{\bar{f}^*g^!K})$ to $(\mathcal{C}_V, D_{\bar{g}^!f^*K}, \varpi_{\bar{g}^!f^*K})$. How can we

compare $\{f^*, \beta_K\} \circ \{g_*, \zeta_K\}$ and $\{\bar{g}_*, \zeta_{f^*K}\} \circ I_{\gamma_K} \circ \{\bar{f}^*, \beta_{g^!K}\}$? Theorem 5.2.1 proves that ε defines a duality preserving functor from the first to the second.

PULL-BACK AND PRODUCT. Given a monoidal functor f^* the isomorphism making f^* a monoidal functor and called $\alpha : f^*(-) \otimes f^*(-) \rightarrow f^*(- \otimes -)$ defines a duality preserving functor $I_{\alpha_{K,M}}$ from $(\mathcal{C}, D_{f^*K \otimes f^*M}, \varpi_{f^*K \otimes f^*M})$ to $(\mathcal{C}, D_{f^*(K \otimes M)}, \varpi_{f^*(K \otimes M)})$. Proposition 5.4.1 shows that the morphism of functors α then defines a duality preserving morphism of functors from $I_{\alpha_{K,M}} \circ \{- \otimes -, \tau_{f^*K, f^*M}\} \circ \{f^* \times f^*, \beta_K \times \beta_M\}$ to $\{f^*, \beta_{K \otimes M}\} \circ \{- \otimes -, \tau_{K,M}\}$. It essentially means that the pull-back is a ring morphism on Witt groups.

PROJECTION FORMULA. Given a monoidal functor f^* with adjunctions (f^*, f_*) , $(f_*, f^!)$ such that the projection morphism π is an isomorphism, there is a natural morphism of functors $\theta : f^!(-) \otimes f^*(-) \rightarrow f^!(- \otimes -)$. This defines a duality preserving functor $I_{\theta_{K,M}}$ from $(\mathcal{C}_2, D_{f^!K \otimes f^*M}, \varpi_{f^!K \otimes f^*M})$ to $(\mathcal{C}_2, D_{f^!(K \otimes M)}, \varpi_{f^!(K \otimes M)})$. Theorem 5.5.1 proves that π is a duality preserving morphism of functors from $\{- \otimes -, \tau_{K,M}\} \circ \{f_* \times Id, \zeta_K \times id\}$ to $\{f_*, \zeta_{K \otimes M}\} \circ I_{\theta_{K,M}} \circ \{- \otimes -, \tau_{f^!K, f^*M}\} \circ \{Id \times f^*, id \times \beta_M\}$.

SUSPENSIONS AND TRIANGULATED STRUCTURES. As it is required for the example of Witt groups, all the results are proved as well in the setting of *triangulated* symmetric monoidal categories, functors and morphisms of functors respecting the triangulated structure. This means essentially two things. First, the categories are endowed with an autoequivalence T (called suspension in the article) and the tensor products defining monoidal structures respect this suspension, i.e. such a tensor product comes equipped with a pair of isomorphism of functors such that a certain diagram anticommutes (see Definition 1.4.12). As proved in the article, it follows that all the other functors (or bifunctors) mentioned above can be endowed by construction (see Point 1 of Propositions 1.5.3 and 1.5.8) with morphisms of functors to commute with the suspensions in a suitable way involving commutative diagrams (similar to (2) with T instead of D). All the morphisms of functors involved then also respect the suspension, i.e. satisfy commutative diagrams similar to (3). This is important, because in the main applications, checking this kind of thing by hand amounts to checking signs involved in the definitions of complexes or maps of complexes, and it has been the source of errors in the literature. Here, we avoid such potential errors by construction (there are no signs and no complexes). Second, using these morphisms of functors to deal with the suspensions, the functors should respect the collection of distinguished triangles. This is easy and has nothing to do with commutative diagrams. It is again obtained by construction (see Point 2 of Propositions 1.5.3 and 1.5.8), except for the first variable of the internal Hom, for which it has to be assumed. The reader who is not interested in triangulated categories, but rather only in monoidal structures can just forget about all the parts of the statements involving T . What remains is the monoidal structure.

These functors and morphisms between them are discussed in the first five sections of this article. Section 6 is devoted to reformulations of the main results when the base category \mathcal{B} is enriched in order that the objects also specify the duality: they are pairs (X, K) where $K \in X^*$ is used to form the duality D_K . The morphisms can then be

chosen in different ways (see Sections 6.1, 6.2 and 6.3). These sections are directed towards geometric applications. In particular, the object ω_f introduced at the beginning of 6.1 has a clear geometric interpretation in the case of morphisms of schemes: it is the relative canonical sheaf. Finally, we have included an appendix to recall closed symmetric monoidal structures on categories involving chain complexes and to relate the sign choices involved with the ones made by various authors working with Witt groups.

REMARKS ON THE PROOFS. Nearly every construction of a morphism in the article is based on variations on a single lemma on adjunctions, namely Lemma 1.2.6. Its refinements are Theorem 1.2.8, Lemma 1.3.5 and Theorem 1.5.10. Similarly, to establish commutativities of natural transformations arising from by Lemma 1.2.6 the main tool is Lemma 1.2.7. The whole Section 1 is devoted to these formal results about adjunctions.

In some applications, certain natural morphisms considered in this article have to be isomorphisms. For example, to define Witt groups, one requires that the morphism of functors ϖ_K is an isomorphism. This is not important to prove the commutativity of the diagrams, therefore it is not part of our assumptions. Nevertheless, in some cases, it might be useful to know that if a particular morphism is an isomorphism, then another one is also automatically an isomorphism. When possible, we have included such statements, for example in Propositions 4.3.1 and 4.3.3. The attribute “strong” in Definitions 2.1.1, 2.2.1 and 2.2.2 is part of this philosophy. By contrast, two isomorphism assumptions are important to the abstract setting regardless of applications. This is the case of the assumption on π , since the essential morphism θ is *defined* using the inverse of π , as well as the assumption on ε whose inverse is involved in the definition of γ . These isomorphism assumptions are listed at the beginning of Section 4 and are recalled in every theorem where they are used.

Considering possible future applications (e.g. motivic homotopy categories or the stable homotopy category), we have presented some aspects in as much generality as possible. Conversely, we do not attempt to provide a complete list of articles where parts of the general framework we study has already been considered. See however [FHM03] for a recent reference written by homotopy theorists which contains many further references.

Finally, let us mention a question a category theorist might ask when reading such a paper: are there coherence theorems, in the spirit of [KML71], that would prove systematically the needed commutative diagrams? As far as the authors know, there are no such coherence theorems available for the moment. Although it is certainly an interesting question, it is unclear (to the authors) how to formulate coherence statements. The interested reader might consider this article as a source of inspiration for future coherence theorems.

1. Adjunctions and consequences

1.1. NOTATIONS AND CONVENTIONS. The opposite category of a category \mathcal{C} is denoted by \mathcal{C}^o .

When F and G are functors with same source and target, we denote a morphism of functors between them as $t : F \rightarrow G$. When $s : G \rightarrow H$ is another one, their composition is denoted by $s \circ t$. When $F_1, F_2 : \mathcal{C} \rightarrow \mathcal{D}$, $G_1, G_2 : \mathcal{D} \rightarrow \mathcal{E}$, $f : F_1 \rightarrow F_2$ and $g : G_1 \rightarrow G_2$, let gf denote the morphism of functors defined by $(fg)_A = G_2(f_A) \circ g_{F_1(A)} = g_{F_2(A)} \circ G_1(f_A)$ on any object A . When $F_1 = F_2 = F$ and $f = id_F$ (resp. $G_1 = G_2 = G$ and $g = id_G$), we usually use the notation gF (resp. Gf). With this convention, $gf = G_2f \circ gF_1 = gF_2 \circ G_1f$. When a commutative diagram is obtained by this equality or other properties immediate from the definition of a morphism of functors, we just put an \boxed{mf} label on it and avoid further justification. To save space, it may happen that when labeling maps in diagrams, we drop the functors from the notation, and just keep the important part, that is the morphism of functors (thus FgH might be reduced to g). Many of the commutative diagrams in the article will be labeled by a symbol in a box (letters or numbers, such as in \boxed{H} or $\boxed{3}$). When they are used in another commutative diagram, eventually after applying a functor to them, we just label them with the same symbol, so that the reader recognizes them, but without further comment.

1.2. USEFUL PROPERTIES OF ADJUNCTIONS. This section is devoted to easy facts and theorems about adjunctions, that are repeatedly used throughout this article. All these facts are obvious, and we only prove the ones that are not completely classical. A good reference for the background on categories and adjunctions as discussed here is [ML98].

1.2.1. DEFINITION. *An adjoint couple (L, R) is the data consisting of two functors $L : \mathcal{C} \rightarrow \mathcal{D}$ and $R : \mathcal{D} \rightarrow \mathcal{C}$ and equivalently:*

- *a bijection $\text{Hom}(LA, B) \simeq \text{Hom}(A, RB)$, functorial in $A \in \mathcal{C}$ and $B \in \mathcal{D}$, or*
- *two morphism of functors $\eta : Id_{\mathcal{C}} \rightarrow RL$ and $\epsilon : LR \rightarrow Id_{\mathcal{D}}$, called respectively unit and counit, such that the resulting compositions $R \xrightarrow{\eta^R} RLR \xrightarrow{R\epsilon} R$ and $L \xrightarrow{L\eta} LRL \xrightarrow{\epsilon L} L$ are identities.*

In the couple, L is called the left adjoint and R the right adjoint. When we want to specify the unit and counit of the couple and the categories involved, we say (L, R, η, ϵ) is an adjoint couple from \mathcal{C} to \mathcal{D} .

When the commutativity of a diagram follows by one of the above compositions giving the identity, we label it \boxed{adj} .

1.2.2. REMARK. Adjunctions between functors that are contravariant can be considered in two different ways, by taking the opposite category of the source of L or R . This does *not* lead to the same notion, essentially because if (L, R) is an adjoint couple, then (R°, L°) is an adjoint couple (instead of (L°, R°)). For this reason, we only use covariant functors in adjoint couples.

1.2.3. LEMMA. Let (L, R, η, ϵ) and $(L', R', \eta', \epsilon')$ be two adjoint couples between the same categories \mathcal{C} and \mathcal{D} , and let $l : L \rightarrow L'$ (resp. $r : R \rightarrow R'$) be an isomorphism. Then, there is a unique isomorphism $r : R \rightarrow R'$ (resp. $l : L \rightarrow L'$) such that $\eta' = rl \circ \eta$ and $\epsilon' = \epsilon \circ l^{-1}r^{-1}$. In particular, a right (resp. left) adjoint is unique up to unique isomorphism.

PROOF. The morphism r is given by the composition $R'\epsilon \circ R'l^{-1}R \circ \eta'R$ and its inverse by the composition $R\epsilon' \circ RlR' \circ \eta R'$. ■

1.2.4. LEMMA. An equivalence of categories is an adjoint couple (F, G, a, b) for which the unit and counit are isomorphisms. In particular, (G, F, b, a) is also an adjoint couple.

1.2.5. LEMMA. Let (L, R, η, ϵ) (resp. $(L', R', \eta', \epsilon')$) be an adjoint couple from \mathcal{C} to \mathcal{D} (resp. from \mathcal{D} to \mathcal{E}). Then $(L'L, RR', R\eta'L \circ \eta, \epsilon' \circ L'\epsilon R')$ is an adjoint couple from \mathcal{C} to \mathcal{E} .

We now turn to a series of less standard results, nevertheless very easy.

1.2.6. LEMMA. (mates) Let H, H', J_1, K_1, J_2 and K_2 be functors with sources and targets as on the following diagram.

$$\begin{array}{ccc} \mathcal{C}_1 & \xrightarrow{H} & \mathcal{C}_2 \\ J_1 \updownarrow K_1 & & J_2 \updownarrow K_2 \\ \mathcal{C}'_1 & \xrightarrow{H'} & \mathcal{C}'_2 \end{array}$$

Consider adjoint couples $(J_i, K_i, \eta_i, \epsilon_i)$ for $i = 1, 2$. Assume $a : J_2H' \rightarrow HJ_1$ (resp. $b : H'K_1 \rightarrow K_2H$) is a morphism of functors. Then there exists a unique morphism of functors $b : H'K_1 \rightarrow K_2H$ (resp. $a : J_2H' \rightarrow HJ_1$) such that the diagrams

$$\begin{array}{ccc} J_2H'K_1 & \xrightarrow{J_2b} & J_2K_2H \\ aK_1 \downarrow & \boxed{H} & \downarrow \epsilon_2H \\ HJ_1K_1 & \xrightarrow{H\epsilon_1} & H \end{array} \quad \text{and} \quad \begin{array}{ccc} H' & \xrightarrow{H'\eta_1} & H'K_1J_1 \\ \eta_2H' \downarrow & \boxed{H'} & \downarrow bJ_1 \\ K_2J_2H' & \xrightarrow{K_2a} & K_2HJ_1 \end{array}$$

are commutative. Furthermore, given two morphisms of functors a and b , the commutativity of one diagram is equivalent to the commutativity of the other one. In this situation, we say that a and b are mates with respect to the rest of the data.


PROOF. We only prove that the existence of a implies the uniqueness and existence of b , the proof of the other case is similar. Assume that b exists and makes the diagrams commutative. The commutative diagram

$$\begin{array}{ccccc} H'K_1 & \xrightarrow{H'\eta_1K_1} & H'K_1J_1K_1 & \xrightarrow{H'K_1\epsilon_1} & H'K_1 \\ \eta_2H'K_1 \downarrow & & \downarrow & \boxed{mf} & \downarrow b \\ K_2J_2H'K_1 & \xrightarrow{K_2aK_1} & K_2HJ_1K_1 & \xrightarrow{K_2H\epsilon_1} & K_2H \end{array}$$


in which the upper horizontal composition is the identity of $H'K_1$ (by adjunction) shows that b has to be given by the composition

$$H'K_1 \xrightarrow{\eta_2 H'K_1} K_2 J_2 H'K_1 \xrightarrow{K_2 a K_1} K_2 H J_1 K_1 \xrightarrow{K_2 H \epsilon_1} K_2 H .$$

This proves uniqueness. Now let b be given by the above composition. The commutative diagram




proves $\square H$ and the commutative diagram




proves $\square H'$. The fact that the commutativity of one of the diagrams implies commutativity to the other is left to the reader. ■

1.2.7. LEMMA. *Let us consider a cube of functors and morphisms of functors*



that is commutative in the following sense: The morphism between the two outer compositions of functors from p to q given by the composition of the three morphisms of functors of the front is equal to the composition of the three morphism of functors of the back. Assume that the vertical maps have right adjoints. Then, by Lemma 1.2.6 applied to the vertical squares, we obtain the following cube (the top and bottom squares have not changed).



This cube is commutative (in the sense just defined, using r and s instead of p and q).

PROOF. This is straightforward, using the commutative diagrams of Lemma 1.2.6, and left to the reader. ■

We now use Lemma 1.2.6 to prove a theorem, which doesn't contain a lot more than the lemma, but is stated in a convenient way for future reference in the applications we are interested in.

1.2.8. THEOREM. Let $L, R, L', R', F_1, G_1, F_2, G_2$ be functors whose sources and targets are specified by the diagram

$$\begin{array}{ccc} \mathcal{C}_1 & \begin{array}{c} \xrightarrow{L} \\ \xleftarrow{R} \end{array} & \mathcal{C}_2 \\ F_1 \uparrow \downarrow G_1 & & F_2 \uparrow \downarrow G_2 \\ \mathcal{C}'_1 & \begin{array}{c} \xrightarrow{L'} \\ \xleftarrow{R'} \end{array} & \mathcal{C}'_2 \end{array}$$

We will study morphisms of functors $f_L, f'_L, g_L, g'_L, f_R, f'_R, g_R$ and g'_R whose sources and targets will be as follows:

$$\begin{array}{ccc} LF_1 & \begin{array}{c} \xrightarrow{f_L} \\ \xleftarrow{f'_L} \end{array} & F_2L' \\ F_1R' & \begin{array}{c} \xrightarrow{f'_R} \\ \xleftarrow{f_R} \end{array} & RF_2 \end{array} \quad \begin{array}{ccc} L'G_1 & \begin{array}{c} \xrightarrow{g'_L} \\ \xleftarrow{g_L} \end{array} & G_2L \\ G_1R & \begin{array}{c} \xrightarrow{g_R} \\ \xleftarrow{g'_R} \end{array} & R'G_2 \end{array}$$

Let us consider the following diagrams, in which the maps and their directions will be the obvious ones induced by the eight maps above and the adjunctions accordingly to the different cases discussed below.

$$\begin{array}{ccc} F_2L'G_1 & \text{---} & LF_1G_1 \\ \downarrow & \boxed{L} & \downarrow \\ F_2G_2L & \text{---} & L \\ \\ F_1R'G_2 & \text{---} & RF_2G_2 \\ \downarrow & \boxed{R} & \downarrow \\ F_1G_1R & \text{---} & R \\ \\ G_1 & \text{---} & G_1RL \\ \downarrow & \boxed{G_1} & \downarrow \\ R'L'G_1 & \text{---} & R'G_2L \\ \\ F_1 & \text{---} & F_1R'L' \\ \downarrow & \boxed{F_1} & \downarrow \\ RLF_1 & \text{---} & RF_2L' \end{array} \quad \begin{array}{ccc} L' & \text{---} & G_2F_2L' \\ \downarrow & \boxed{L'} & \downarrow \\ L'G_1F_1 & \text{---} & G_2LF_1 \\ \\ R' & \text{---} & G_1F_1R' \\ \downarrow & \boxed{R'} & \downarrow \\ R'G_2F_2 & \text{---} & G_1RF_2 \\ \\ L'G_1R & \text{---} & G_2LR \\ \downarrow & \boxed{G_2} & \downarrow \\ L'R'G_2 & \text{---} & G_2 \\ \\ LF_1R' & \text{---} & F_2L'R' \\ \downarrow & \boxed{F_2} & \downarrow \\ LRF_2 & \text{---} & F_2 \end{array}$$

Then

1. Let (G_i, F_i) , $i = 1, 2$ be adjoint couples. Let g_L (resp. f_L) be given, then there is a unique f_L (resp. g_L) such that \boxed{L} and $\boxed{L'}$ are commutative. Let g_R (resp. f_R) be given, then there is a unique f_R (resp. g_R) such that \boxed{R} and $\boxed{R'}$ are commutative.
- 1'. Let (F_i, G_i) , $i = 1, 2$ be adjoint couples. Let g'_L (resp. f'_L) be given, then there is a unique f'_L (resp. g'_L) such that \boxed{L} and $\boxed{L'}$ are commutative. Let g'_R (resp. f'_R) be given, then there is a unique f'_R (resp. g'_R) such that \boxed{R} and $\boxed{R'}$ are commutative.
2. Let (L, R) and (L', R') be adjoint couples. Let f_L (resp. f'_R) be given, then there is a unique f'_R (resp. f_L) such that $\boxed{F_1}$ and $\boxed{F_2}$ are commutative. Let g'_L (resp. g_R) be given, then there is a unique g_R (resp. g'_L) such that $\boxed{G_1}$ and $\boxed{G_2}$ are commutative.
3. Assuming (G_i, F_i) , $i = 1, 2$, (L, R) and (L', R') are adjoint couples, and $g_L, g'_L = g_L^{-1}$ are given (resp. f_R and $f'_R = f_R^{-1}$). By 1 and 2, we obtain f_L and g_R (resp. g_R and f_L). We then may construct f_R and f'_R (resp. g_L and g'_L) which are inverse to each other.
- 3'. Assuming (F_i, G_i) , $i = 1, 2$, (L, R) and (L', R') are adjoint couples, f_L and $f'_L = f_L^{-1}$ are given (resp. g_R and $g'_R = g_R^{-1}$). By 1' and 2, we obtain g'_L and f'_R (resp. f'_R and g'_L). We then may construct g'_R and g_R (resp. f'_L and f_L) which are inverse to each other.

PROOF. Points 1, 1' and 2 are obvious translations of the previous lemma. We only prove Point 3, since 3' is dual to it. Let (L, R, η, ϵ) , $(L', R', \eta', \epsilon')$ and $(G_i, F_i, \eta_i, \epsilon_i)$, $i = 1, 2$, be the adjoint couples. Using 1 and 2, we first obtain f_L and g_R , as well as the commutative diagrams \boxed{L} , $\boxed{L'}$ (both involving $g_L = (g'_L)^{-1}$), $\boxed{G_1}$ and $\boxed{G_2}$ (both involving $g'_L = (g_L)^{-1}$). The morphisms of functors f'_R and f_R are respectively defined by the compositions

$$F_1 R' \xrightarrow{\eta_{F_1 R'}} R L F_1 R' \xrightarrow{R f_L R'} R F_2 L' R' \xrightarrow{R F_2 \epsilon'} R F_2$$

and

$$R F_2 \xrightarrow{\eta_{R F_2}} F_1 G_1 R F_2 \xrightarrow{F_1 g_R F_2} F_1 R' G_2 F_2 \xrightarrow{F_1 R' \epsilon_2} F_1 R'.$$

We compute $f_R \circ f'_R$ as the upper right composition of the following commutative diagram

$$\begin{array}{ccccccc}
 F_1 R' & \xrightarrow{\eta} & R L F_1 R' & \xrightarrow{f_L} & R F_2 L' R' & \xrightarrow{\epsilon'} & R F_2 \\
 \eta_1 \downarrow & \boxed{mf} & \downarrow & \boxed{mf} & \downarrow & \boxed{mf} & \downarrow \eta_1 \\
 F_1 G_1 F_1 R' & \longrightarrow & F_1 G_1 R L F_1 R' & \longrightarrow & F_1 G_1 R F_2 L' R' & \longrightarrow & F_1 G_1 R F_2 \\
 & \boxed{G_1} & \downarrow & \boxed{mf} & \downarrow & \boxed{mf} & \downarrow g_R \\
 & & F_1 R' G_2 L F_1 R' & \longrightarrow & F_1 R' G_2 F_2 L' R' & \longrightarrow & F_1 R' G_2 F_2 \\
 & \searrow \eta' & \downarrow (g'_L)^{-1} = g_L & \boxed{L'} & \downarrow & \boxed{mf} & \downarrow \epsilon_2 \\
 & & F_1 R' L' G_1 F_1 R' & \xrightarrow{\epsilon_1} & F_1 R' L' R' & \xrightarrow{\epsilon'} & F_1 R'
 \end{array}$$

The lower left composition in the above diagram is the identity because it appears as the upper right composition of the commutative diagram

$$\begin{array}{ccccc}
 F_1 R' & \xrightarrow{\eta_1} & F_1 G_1 F_1 R' & \xrightarrow{\eta'} & F_1 R' L' G_1 F_1 R' . \\
 & \searrow & \boxed{adj} \downarrow & \boxed{mf} & \downarrow \epsilon_1 \\
 & & F_1 R' & \xrightarrow{\quad} & F_1 R' L' R' \\
 & & & \searrow & \downarrow \epsilon' \\
 & & & & F_1 R'
 \end{array}$$

The composition $f'_R \circ f_R = id$ is proved in a similar way, involving the diagrams \boxed{L} and $\boxed{G_2}$. ■

The reader has certainly noticed that there is a statement 2' which we didn't spell out because we don't need it.

1.3. BIFUNCTORS AND ADJUNCTIONS. We have to deal with couples of bifunctors that give adjoint couples of usual functors when one of the entries in the bifunctors is fixed. We need to explain how these adjunctions are functorial in this entry. The standard example for that is the classical adjunction between tensor product and internal Hom.

1.3.1. DEFINITION. *Let $\mathcal{X}, \mathcal{C}, \mathcal{C}'$ be three categories, and let $L : \mathcal{X} \times \mathcal{C}' \rightarrow \mathcal{C}$ and $R : \mathcal{X} \times \mathcal{C} \rightarrow \mathcal{C}'$ be bifunctors. We say that (L, R) form an adjoint couple of bifunctors (abbreviated as ACB) from \mathcal{C}' to \mathcal{C} with parameter in \mathcal{X} if we are given adjoint couples $(L(X, -), R(X, -), \eta_X, \epsilon_X)$ for every X and if furthermore η and ϵ are generalized transformations in the sense of [EK66], i.e. for any morphism $f : A \rightarrow B$ in \mathcal{X} , the diagrams*

$$\begin{array}{ccc}
 L(A, R(B, C)) \xrightarrow{L(f, id)} L(B, R(B, C)) & & C' \xrightarrow{\eta_A} R(A, L(A, C')) \\
 \downarrow L(id, R(f, id)) \quad \boxed{gen} \quad \downarrow \epsilon_B & & \eta_B \downarrow \quad \boxed{gen} \quad \downarrow R(id, L(f, id)) \\
 L(A, R(A, C)) \xrightarrow{\epsilon_A} C & & R(B, L(B, C')) \xrightarrow{R(f, id)} R(A, L(B, C'))
 \end{array}$$

commute. We sometimes use the notation $(L(*, -), R(*, -))$, where the $*$ is the entry in \mathcal{X} and write

$$\mathcal{C} \begin{array}{c} \xrightarrow{L} \\ \mathcal{X} \\ \xleftarrow{R} \end{array} \mathcal{C}'$$

in diagrams.

1.3.2. EXAMPLE. Let $\mathcal{C} = \mathcal{C}' = \mathcal{X}$ be the category of modules over a commutative ring. The tensor product (with the variables switched) and the internal Hom form an ACB, with the usual unit and counit.

1.3.3. LEMMA. *Let $F : \mathcal{X}' \rightarrow \mathcal{X}$ be a functor, and (L, R) be an ACB with parameter in \mathcal{X} . Then $(L(F(*), -), R(F^o(*), -))$ is again an ACB in the obvious way.*

1.3.4. LEMMA. Let (L, R) be an ACB from \mathcal{C}' to \mathcal{C} with parameter in \mathcal{X} and let (F, G) be an adjoint couple from \mathcal{D} to \mathcal{C}' , then $(L(*, F(-)), GR(*, -))$ is an ACB with unit and counit defined in the obvious and natural way.

PROOF. Left to the reader. \blacksquare

We now give a version of Lemma 1.2.6 for ACBs.

1.3.5. LEMMA. For $i = 1$ or 2 , let $(J_i, K_i, \eta_i, \epsilon_i)$ be an ACB from \mathcal{C}'_i to \mathcal{C}_i with parameter in \mathcal{X} , let $H : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ and $H' : \mathcal{C}'_1 \rightarrow \mathcal{C}'_2$ be functors. Let $a : J_2(*, H'(-)) \rightarrow HJ_1(*, -)$ (resp. $b : H'K_1(*, -) \rightarrow K_2(*, H(-))$) be a morphism of bifunctors. Then there exists a unique morphism of bifunctors

$$b : H'K_1(*, -) \rightarrow K_2(*, H(-)) \quad (\text{resp. } a : J_2(*, H'(-)) \rightarrow HJ_1(*, -))$$

such that the diagrams

$$\begin{array}{ccc} J_2(X, H'K_1(X, -)) & \xrightarrow{J_2 b} & J_2(X, K_2(X, H(-))) \\ \downarrow a_{K_1} & \boxed{H} & \downarrow \eta_2 H \\ HJ_1(X, K_1(X, -)) & \xrightarrow{H\epsilon_1} & H \end{array}$$

and

$$\begin{array}{ccc} H' & \xrightarrow{H'\eta_1} & H'K_1(X, J_1(X, -)) \\ \downarrow \eta_2 H' & \boxed{H'} & \downarrow b_{J_1} \\ K_2(X, J_2(X, H'(-))) & \xrightarrow{K_2 a} & K_2(X, HJ_1(X, -)) \end{array}$$

are commutative for every morphism $X \in \mathcal{X}$.

PROOF. We only consider the case when a is given. For every $X \in \mathcal{X}$, we apply Lemma 1.2.6 to obtain $b : H'K_1(X, -) \rightarrow K_2(X, H(-))$. It remains to prove that this is functorial in X . This is an easy exercise on commutative diagrams which we leave to the reader (it involves that both η_1 and ϵ_2 are generalized transformations as in Definition 1.3.1). \blacksquare

1.3.6. LEMMA. Let (L, R, η, ϵ) and $(L', R', \eta', \epsilon')$ be ACBs, and let $l : L \rightarrow L'$ (resp. $r : R \rightarrow R'$) be an isomorphism of bifunctors. There is a unique isomorphism of bifunctors $r : R \rightarrow R'$ (resp. $l : L \rightarrow L'$) such that $\eta'_X = r_X R(X, l)\eta_X$ and $\epsilon'_X = \epsilon_X L(X, r_X^{-1})l_X^{-1}$ for every $X \in \mathcal{X}$. In other words, a right (resp. left) bifunctor adjoint is unique up to unique isomorphism.

PROOF. We only give the proof in case l is given, the case with r given is similar. Apply Lemma 1.3.5 with H and H' being identity functors. Starting with l^{-1} , we get r , and starting with l we get a morphism $r' : R' \rightarrow R$. The morphisms r and r' are inverse to each other as in the usual proof of the uniqueness of adjunction (which applies to every parameter $X \in \mathcal{X}$). \blacksquare

1.4. SUSPENDED AND TRIANGULATED CATEGORIES. We recall here what we need about triangulated categories. The reason why we make a distinction between suspended categories and triangulated ones is because all the commutative diagrams that we are interested in are just related to the suspension, and not to the exactness of the functors involved. So when we need to prove the commutativity of those diagrams, we forget about the exactness of our functors, and just think of them as suspended functors, in the sense described below.

1.4.1. DEFINITION. *A suspended category is an additive category \mathcal{C} together with an adjoint couple (T, T^{-1}) from \mathcal{C} to \mathcal{C} which is an equivalence of category (the unit and counit are isomorphisms).*

1.4.2. REMARK. We assume furthermore in all what follows that TT^{-1} and $T^{-1}T$ are the identity of \mathcal{C} and that the unit and counit are also the identity. This assumption is not true in some suspended (triangulated) categories arising in stable homotopy theory. Nevertheless, it simplifies the exposition which is already sufficiently technical. When working in an example where this assumption does not hold, it is of course possible to make the modifications to get this even more general case.

Between suspended categories $(\mathcal{C}, T_{\mathcal{C}})$ and $(\mathcal{D}, T_{\mathcal{D}})$, we use suspended functors:

1.4.3. DEFINITION. *A suspended functor (F, f) from \mathcal{C} to \mathcal{D} is a functor F together with an isomorphism of functors $f : FT_{\mathcal{C}} \rightarrow T_{\mathcal{D}}F$. We sometimes forget about f in the notation.*

Without the assumption in Remark 1.4.2, we would need yet another isomorphism $f' : FT^{-1} \rightarrow T^{-1}F$ and compatibility diagrams analogous to the ones in Lemma 1.2.6. Then, we would have to carry those compatibilities in our constructions. Again, this would not be a problem, just making things even more tedious.

Suspended functors can be composed in an obvious way, and (T, id_{T^2}) and (T^{-1}, id_{Id}) are suspended endofunctors of \mathcal{C} that we call T and T^{-1} for short.

1.4.4. DEFINITION. *To a suspended functor F , one can associate “shifted” ones, composing F by T or T^{-1} several times on either sides. The isomorphisms $T^i FT^j \simeq T^k FT^l$ with $i + j = k + l$ constructed using $f, f^{-1}, T^{-1}T = Id$ and $TT^{-1} = Id$ all coincide, so whenever we use one, we label it “ f ” without further mention.*

1.4.5. DEFINITION. *The opposite suspended category \mathcal{C}^o of a suspended category \mathcal{C} is given the suspension $(T_{\mathcal{C}}^{-1})^o$.*

With this convention, we can deal with contravariant suspended functors in two different ways (depending where we put the “op”), and this yields essentially the same thing, using the definition of shifted suspended functors.

1.4.6. DEFINITION. A morphism of suspended functors $h : (F, f) \rightarrow (G, g)$ is a morphism of functors $h : F \rightarrow G$ such that the diagram

$$\begin{array}{ccc} FT & \xrightarrow{f} & TF \\ hT \downarrow & \boxed{\text{sus}} & \downarrow Th \\ GT & \xrightarrow{g} & TG \end{array}$$

is commutative.

1.4.7. LEMMA. The composition of two morphisms of suspended functors yields a morphism of suspended functors.

PROOF. Straightforward. ■

A triangulated category is a suspended category with the choice of some exact triangles, satisfying some axioms. This can be found in text books as [W94] (see also the nice introduction in [B00, Section 1]). We include the enriched octahedron axiom in the list of required axioms as it is suitable to deal with Witt groups, as explained in loc. cit.

1.4.8. DEFINITION. (see for example [GN03, 1.1]) Let $(F, f) : \mathcal{C} \rightarrow \mathcal{D}$ be a covariant (resp. contravariant) suspended functor. We say that (F, f) is δ -exact ($\delta = \pm 1$) if for any exact triangle

$$A \xrightarrow{u} B \xrightarrow{v} C \xrightarrow{w} TA$$

the triangle

$$FA \xrightarrow{Fu} FB \xrightarrow{Fv} FC \xrightarrow{\delta f_A \circ Fw} TFA$$

respectively

$$FC \xrightarrow{Fv} FB \xrightarrow{Fu} FA \xrightarrow{\delta f_C \circ FT^{-1}w} TFC$$

is exact.

1.4.9. REMARK. With this definition, T and T^{-1} are (-1) -exact functors, because of the second axiom of triangulated categories, and the composition of exact functors multiplies their signs. Thus, if F is δ -exact, then $T^i FT^j$ is $(-1)^{i+j} \delta$ -exact.

To define morphisms between exact functors F and G , the signs δ_F and δ_G of the functors have to be taken into account, so that the morphism of functors induces a morphism between the triangles obtained by applying F or G to a triangle and making the sign modifications.

1.4.10. DEFINITION. We say that $h : F \rightarrow G$ is a morphism of exact functors if the diagram $\boxed{\text{sus}}$ in Definition 1.4.6 is $\delta_F \delta_G$ commutative.

On the other hand, we have the following lemma.

1.4.11. LEMMA. Let $h : F \rightarrow G$ be an isomorphism of suspended functors such that $\boxed{\text{sus}}$ is ν -commutative. Assume F is δ -exact. Then G is $\delta \nu$ -exact.

PROOF. For any triangle

$$A \xrightarrow{u} B \xrightarrow{v} C \xrightarrow{w} TA$$

the triangle

$$GA \xrightarrow{Gu} GB \xrightarrow{Gv} FC \xrightarrow{\delta g_A \circ Gw} TGA$$

is easily shown to be isomorphic to

$$FA \xrightarrow{Fu} FB \xrightarrow{Fv} FC \xrightarrow{\nu \delta f_A \circ Fw} TFA$$

■

We also need to deal with bifunctors from two suspended categories to another one. These are just suspended functors in each variable, with a compatibility condition. Examples are the internal Hom or the tensor product in triangulated categories.

1.4.12. DEFINITION. Let \mathcal{C}_1 , \mathcal{C}_2 and \mathcal{D} be suspended categories. A suspended bifunctor from $\mathcal{C}_1 \times \mathcal{C}_2$ to \mathcal{D} is a triple (B, b_1, b_2) where $B : \mathcal{C}_1 \times \mathcal{C}_2 \rightarrow \mathcal{D}$ is a functor and two morphisms of functors $b_1 : B(T(-), *) \rightarrow TB(-, *)$ and $b_2 : B(-, T(*)) \rightarrow TB(-, *)$, such that the diagram

$$\begin{array}{ccc} B(TA, TC) & \xrightarrow{b_{2,TA,C}} & TB(TA, C) \\ b_{1,A,TC} \downarrow & -1 & \downarrow b_{1,A,C} \\ TB(A, TC) & \xrightarrow{b_{2,A,C}} & T^2B(A, C) \end{array}$$

anti-commutes for every A and C .

1.4.13. REMARK. As in Definition 1.4.4, we have shifted versions of b_1 (or b_2) which we will sometimes label “ b_1 ” (or “ b_2 ”) below. It is important to describe precisely in which order morphisms are applied when combining b_1 and b_2 .

1.4.14. DEFINITION. A morphism of suspended bifunctors from a suspended bifunctor (B, b_1, b_2) to a suspended bifunctor (B', b'_1, b'_2) is a morphism of functors $f : B \rightarrow B'$ such that the two diagrams

$$\begin{array}{ccccc} B(TA, C) & \xrightarrow{b_{1,A,C}} & TB(A, C) & \xleftarrow{b_{2,A,C}} & B(A, TC) \\ f_{TA,C} \downarrow & & \downarrow Tf_{A,C} & & \downarrow f_{A,TC} \\ B'(TA, C) & \xrightarrow{b'_{1,A,C}} & TB'(A, C) & \xleftarrow{b'_{2,A,C}} & B'(A, TC) \end{array}$$

are commutative for every A and C .

By composing with a usual suspended functor to \mathcal{C}_1 or \mathcal{C}_2 or from \mathcal{D} , we get other suspended bifunctors (the verification is easy). But, if we do that several times, using different functors, the order in which the suspended functors to \mathcal{C}_1 or \mathcal{C}_2 are used *does* matter. For example, as with usual suspended functors, it is possible to define shifted versions by composing with the suspensions in each category as mentioned in Remark

1.4.13. This can be useful. Unfortunately, according to the order in which we do this (if we mix functors to \mathcal{C}_1 and to \mathcal{C}_2), we don't get the same isomorphism of functors, even though we get the same functors in the pair. One has to be careful about that.

1.5. SUSPENDED ADJUNCTIONS. As with usual functors, there is a notion of adjunction well suited for suspended functors.

1.5.1. DEFINITION. *A suspended adjoint couple (L, R) is an adjoint couple in the usual sense in which L and R are suspended functors and the unit and counit are morphisms of suspended functors.*

1.5.2. DEFINITION. *When (L, R) is an adjoint couple of suspended functors, using Lemma 1.2.5 we obtain shifted versions $(T^i L T^j, T^{-j} R T^{-i})$. Using Definition 1.4.4, we obtain isomorphisms exchanging the T 's. Applying Lemma 1.2.3 to them, we get an adjoint couple of suspended bifunctors $(T^i L T^j, T^{-i} R T^{-j})$.*

The following proposition seems to be well-known.

1.5.3. PROPOSITION. *Let (L, R) be an adjoint couple from \mathcal{C} to \mathcal{D} (of usual functors) and let (L, l) be a suspended functor. Then*

1. *there is a unique isomorphism of functors $r : RT \rightarrow TR$ that turns (R, r) into a suspended functor and (L, R) into a suspended adjoint couple.*
2. *if furthermore \mathcal{C} and \mathcal{D} are triangulated and (L, l) is δ -exact, then (R, r) is also δ -exact (with the same δ).*

(When starting with a suspended functor (R, r) , the dual statement holds.)

PROOF. Point 1 is a direct corollary of Point 3 of Theorem 1.2.8, by taking $L = L'$, $R = R'$, $F_1 = T_{\mathcal{C}}$, $G_1 = T_{\mathcal{C}}^{-1}$, $F_2 = T_{\mathcal{D}}$, $G_2 = T_{\mathcal{D}}^{-1}$, $g_L = (g'_L)^{-1} = T^{-1} l T^{-1}$. This gives $f_R = r$. The commutative diagrams $\boxed{F_1}$ and $\boxed{F_2}$ exactly tell us that the unit and counit are suspended morphisms of functors with this choice of r . To prove Point 2, we have to show that the pair (R, r) is exact. Let

$$A \xrightarrow{u} B \xrightarrow{v} C \xrightarrow{w} TA$$

be an exact triangle. We want to prove that the triangle

$$RA \xrightarrow{u} RB \xrightarrow{v} RC \xrightarrow{r_A \circ R w} TRA$$

is exact. We first complete $RA \xrightarrow{u} RB$ as an exact triangle

$$RA \xrightarrow{u} RB \xrightarrow{v'} C' \xrightarrow{w'} TRA$$

and we prove that this triangle is in fact isomorphic to the previous one. To do so, one completes the incomplete morphism of triangles

$$\begin{array}{ccccccc} LRA & \xrightarrow{LRu} & LRB & \xrightarrow{LRv} & LC' & \xrightarrow{f_{RA} \circ LRw} & TLRA \\ \downarrow & & \downarrow & & \downarrow h & & \downarrow \\ A & \xrightarrow{u} & B & \xrightarrow{v} & C & \xrightarrow{w} & TA \end{array}$$

Looking at the adjoint diagram, we see that $ad(h) : C' \rightarrow RC$ is an isomorphism by the five lemma for triangulated categories. ■

1.5.4. **THEOREM.** *Lemma 1.2.6, Lemma 1.2.7 and Theorem 1.2.8 hold when we replace every functor by a suspended functor, every adjoint couple by a suspended adjoint couple and every morphism of functor by a morphism of suspended functors.*

PROOF. The same proofs hold, since they only rely on operations and properties of functors and morphism of functors, such as composition or commutative diagrams, that exist and behave the same way in the suspended case. ■

We now adapt the notion of an adjoint couple of bifunctors to suspended categories.

1.5.5. **DEFINITION.** *Let (L, R) be an ACB from C' to C with parameter in \mathcal{X} , where C, C' and \mathcal{X} are suspended categories. Assume moreover that (L, l_1, l_2) and (R, r_1, r_2) are suspended bifunctors. We say that (L, R) is a suspended adjoint couple of bifunctors if*

1. $((L(X, -), l_2), (R(X, -), r_2))$ is a suspended adjoint couple of functors for every parameter X ,
2. the following diagrams commute:

$$\begin{array}{ccc}
 TC & \xrightarrow{\eta_{X,TC}} & R(X, L(X, TC)) \xrightarrow{\text{“}r_1\text{”}} TR(TX, L(X, TC)) \\
 \downarrow T\eta_{TX,C} & & \downarrow TR(TX, l_2) \\
 TR(TX, L(TX, C)) & \xrightarrow{TR(X, l_1)} & TR(TX, TL(X, C)) \\
 \\
 TL(X, R(TX, C)) & \xrightarrow{l_1^{-1}} & L(TX, R(TX, C)) \\
 \downarrow l_2^{-1} & & \downarrow \epsilon_{TX,C} \\
 L(X, TR(TX, C)) & \xrightarrow{\text{“}r_1\text{”}^{-1}} & L(X, R(X, C)) \xrightarrow{\epsilon_{X,C}} C
 \end{array}$$

1.5.6. **REMARK.** Note that (2) ensures the compatibility of the suspension functor on the parameter with the other ones.

1.5.7. **LEMMA.** *Let $(F, f) : \mathcal{X}' \rightarrow \mathcal{X}$ be a suspended functor, and (L, R) a suspended ACB with parameter in \mathcal{X} . Then $(L(F(*), -), R(F^o(*), -))$ is again a suspended ACB in the obvious way.*

PROOF. Left to the reader. ■

The following proposition is an analogue of Proposition 1.5.3 for suspended bifunctors.

1.5.8. PROPOSITION. *Let (L, R) be an ACB such that (L, l_1, l_2) (resp. (R, r_1, r_2)) is a suspended bifunctor. Then*

1. *there exist unique r_1, r_2 (resp. l_1, l_2) such that (R, r_1, r_2) (resp. (L, l_1, l_2)) is a suspended bifunctor and (L, R) is a suspended ACB,*
2. *if $(L(X, -), l_2)$ is δ -exact for some object X , then so is $(R(X, -), r_2)$.*

PROOF. The existence and uniqueness of r_2 follows (for every parameter X) from Point 1 of Proposition 1.5.3. That r_1 is also natural in the first variable follows from a large diagram of commutative squares of type \boxed{mf} and \boxed{gen} . This yields in particular morphisms of suspended functors η_X and ϵ_X for every X . For the existence and uniqueness of r_1 , we use Lemma 1.3.5 applied to $H' = T$, $H = Id$, $J_1 = L(*, -)$, $K_1 = R(*, -)$, $J_2 = L(T^{-1}(*), -)$, $K_2 = R(T^{-1}(*), -)$ and $a = "l_1" "l_2"$ (the order is important). We obtain two diagrams which are easily seen to be equivalent to the ones required for (L, R) to be a suspended ACB (Point 2 of Definition 1.5.5). The anticommutativity required by Definition 1.4.12 for r_2 and r_1 may be proved using the anticommutativity of l_1 and l_2 and the fact that η_X and $\epsilon_{T^{-1}X}$ are morphisms of suspended functors. Point 2 is proved in the same way as Point 2 of Proposition 1.5.3. \blacksquare

We need a version of Lemma 1.3.5 for suspended bifunctors.

1.5.9. LEMMA. *Lemma 1.3.5 holds when all the functors, bifunctors and adjunctions become suspended ones.*

PROOF. The proof of Lemma 1.3.5 works since it only involves compositions and commutative diagrams that exist in the suspended case. \blacksquare

Finally, there is also a version of Theorem 1.2.8 for (suspended) bifunctors, which is our main tool for the applications. For this reason, we state it in full detail.

1.5.10. THEOREM. *Let $L, R, L', R', F_1, G_1, F_2, G_2$ be (suspended) functors whose sources and targets are specified by the diagram (recall the notation of Definition 1.3.1),*

$$\begin{array}{ccc}
 \mathcal{C}_1 & \begin{array}{c} \xrightarrow{L} \\ \xleftarrow{R} \end{array} & \mathcal{C}_2 \\
 F_1 \uparrow \mathcal{X} \downarrow G_1 & & F_2 \uparrow \mathcal{X} \downarrow G_2 \\
 \mathcal{C}'_1 & \begin{array}{c} \xrightarrow{L'} \\ \xleftarrow{R'} \end{array} & \mathcal{C}'_2
 \end{array}$$

and let $f_L, f'_L, g_L, g'_L, f_R, f'_R, g_R$ and g'_R be morphisms of bifunctors (resp. suspended bifunctors) whose sources and targets will be as follows:

$$\begin{array}{cc}
 LF_1(*, -) \begin{array}{c} \xleftarrow{f_L} \\ \xrightarrow{f'_L} \end{array} F_2(*, L'(-)) & L'G_1(*, -) \begin{array}{c} \xleftarrow{g'_L} \\ \xrightarrow{g_L} \end{array} G_2(*, L(-)) \\
 F_1(*, R'(-)) \begin{array}{c} \xleftarrow{f'_R} \\ \xrightarrow{f_R} \end{array} RF_2(*, -) & G_1(*, R(-)) \begin{array}{c} \xleftarrow{g_R} \\ \xrightarrow{g'_R} \end{array} R'G_2(*, -)
 \end{array}$$

Let us consider the following diagrams, in which the maps and their directions will be the only obvious ones in all the cases discussed below.

$$\begin{array}{ccc}
 F_2(X, L'G_1(X, -)) & \xrightarrow{\quad} & LF_1(X, G_1(X, -)) & & G_1(C, -) & \xrightarrow{\quad} & G_1(C, RL(-)) \\
 \downarrow & & \boxed{L} & & \downarrow & & \boxed{G_1} & & \downarrow \\
 F_2(X, G_2(X, L(-))) & \xleftarrow{\quad} & L & & R'L'G_1(C, -) & \xrightarrow{\quad} & R'G_2(C, L(-)) \\
 \\
 F_1(X, R'G_2(X, -)) & \xrightarrow{\quad} & RF_2(X, G_2(X, -)) & & F_1(C, -) & \xrightarrow{\quad} & F_1(C, R'L'(-)) \\
 \downarrow & & \boxed{R} & & \downarrow & & \boxed{F_1} & & \downarrow \\
 F_1(X, G_1(X, R(-))) & \xleftarrow{\quad} & R & & RLF_1(C, -) & \xrightarrow{\quad} & RF_2(C, L'(-)) \\
 \\
 L' & \xleftarrow{\quad} & G_2(X, F_2(X, L'(-))) & & L'G_1(C, R(-)) & \xrightarrow{\quad} & G_2(C, LR(-)) \\
 \uparrow & & \boxed{L'} & & \downarrow & & \boxed{G_2} & & \downarrow \\
 L'G_1(X, F_1(X, -)) & \xrightarrow{\quad} & G_2(X, LF_1(X, -)) & & L'R'G_2(C, -) & \xrightarrow{\quad} & G_2(C, -) \\
 \\
 R' & \xleftarrow{\quad} & G_1(X, F_1(X, R'(-))) & & LF_1(C, R'(-)) & \xrightarrow{\quad} & F_2(C, L'R'(-)) \\
 \uparrow & & \boxed{R'} & & \downarrow & & \boxed{F_2} & & \downarrow \\
 R'G_2(X, F_2(X, -)) & \xrightarrow{\quad} & G_1(X, RF_2(X, -)) & & LRF_2(C, -) & \xrightarrow{\quad} & F_2(C, -)
 \end{array}$$

Then

1. Let (G_i, F_i) , $i = 1, 2$ be ACBs (resp. suspended ACBs). Let g_L (resp. f_L) be given, then there is a unique (suspended) f_L (resp. g_L) such that \boxed{L} and $\boxed{L'}$ are commutative for any X . Let g_R (resp. f_R) be given, then there is a unique (suspended) f_R (resp. g_R) such that \boxed{R} and $\boxed{R'}$ are commutative for any X .
2. Let (L, R) and (L', R') be adjoint couples (resp. suspended adjoint couples). Let f_L (resp. f'_R) be given, then there is a unique (suspended) f'_R (resp. f_L) such that $\boxed{F_1}$ and $\boxed{F_2}$ are commutative. Let g'_L (resp. g_R) be given, then there is a unique (suspended) g_R (resp. g'_L) such that $\boxed{G_1}$ and $\boxed{G_2}$ are commutative.
3. Assuming (G_i, F_i) , $i = 1, 2$, are ACBs (resp. suspended ACBs), (L, R) and (L', R') are (suspended) adjoint couples, and g_L and $g'_L = g_L^{-1}$ are given (resp. f_R and $f'_R = f_R^{-1}$). By 1 and 2, we obtain f_L and g_R (resp. g_R and f_L). We then may construct f_R and f'_R (resp. g_L and g'_L) which are inverse to each other.

PROOF. The proof is the same as for Theorem 1.2.8, but using Lemma 1.3.5 (or 1.5.9) instead of Lemma 1.2.6 for Points 1 and 2. \blacksquare

1.5.11. **REMARK.** We didn't state the analogues of Points 1' and 2' of Theorem 1.2.8 in this context because we don't need them.

2. Dualities

We now introduce our main subject of interest: duality. As before, we state everything for the suspended (or triangulated) case and the usual case in a uniform way.

2.1. CATEGORIES WITH DUALITY.

2.1.1. **DEFINITION.** *A category with duality is a triple (\mathcal{C}, D, ϖ) where \mathcal{C} is a category with an adjoint couple $(D, D^\circ, \varpi, \varpi^\circ)$ from \mathcal{C} to \mathcal{C}° . When ϖ is an isomorphism, D is an equivalence of categories and we say that the duality is strong. A suspended (resp. triangulated) category with duality is defined in the same way, but (D, d) is a suspended (resp. δ -exact) functor on a suspended (resp. triangulated) category \mathcal{C} , and the adjunction is suspended.*

2.1.2. **REMARK.** Observe that the standard condition $(D^\circ \varpi^\circ) \circ (\varpi^\circ D) = id_{D^\circ}$ is satisfied by the definition of an adjunction. The only difference between our definition and Balmer's definition [B00, Def. 2.2] is that we don't require the isomorphism $d : DT \rightarrow T^{-1}D$ to be an equality in the suspended or triangulated case. Assuming that duality and suspension strictly commute is as bad as assuming that the internal Hom and the suspension strictly commute, or (by adjunction) that the tensor product and the suspension strictly commute. This is definitely a too strong condition when checking strict commutativity of diagrams in some derived category. Dropping all signs in this setting when defining these isomorphisms just by saying "take the canonical ones" may even lead to contradictions as the results of the Appendix show. When $d = id$, we say that the duality is *strict*.

2.1.3. **DEFINITION.** *Let (\mathcal{C}, D, ϖ) be a triangulated category with duality for which (D, d) is δ -exact. By Definition 1.5.2 we get $T(D, D^\circ) = (TD, TD^\circ, \varpi', (\varpi')^\circ)$, the shifted adjoint couple. We define the suspension of (\mathcal{C}, D, ϖ) as $T(\mathcal{C}, D, \varpi) = (\mathcal{C}, TD, -\delta\varpi')$. Note that (TD, Td) is $(-\delta)$ -exact because T is (-1) -exact.*

2.1.4. **REMARK.** This is the definition of [B00, Definition 2.8] adapted to cover the non strict case, and the next one generalizes [B00, Definition 2.13] to the non strict case.

2.1.5. **DEFINITION.** *For any triangulated category with strong duality (\mathcal{C}, D, ϖ) , we define its i -th Witt group W^i by $W^i(\mathcal{C}, D, \varpi) := W(T^i(\mathcal{C}, D, \varpi))$ (extending [B00, 2.4 and Definitions 2.12 and 2.13] in the obvious way). If D and ϖ are understood, we sometimes also write $W^i(\mathcal{C})$ or $W^i(\mathcal{C}, D)$ for short.*

2.1.6. **REMARK.** In concrete terms, this means that the condition of loc. cit. for an element in $W^1(\mathcal{C})$ represented by some ϕ to be symmetric is that $(TdT^{-1}) \circ \phi = (D^\circ T\phi) \circ \varpi$ whereas in the strict case the (TdT^{-1}) may be omitted.

2.1.7. **REMARK.** To define a Witt group, the duality has to be strong (that is ϖ has to be an isomorphism), but this is not necessary to prove all sorts of commutative diagrams, so there is no reason to assume it here in general.

2.2. DUALITY PRESERVING FUNCTORS AND MORPHISMS.

2.2.1. **DEFINITION.** A duality preserving functor from a (suspended, triangulated) category with duality $(\mathcal{C}_1, D_1, \varpi_1)$ to another one $(\mathcal{C}_2, D_2, \varpi_2)$ is a pair $\{F, f\}$ where F is a (suspended, δ -exact) functor from \mathcal{C}_1 to \mathcal{C}_2 , and $f : FD_1^o \rightarrow D_2^o F^o$ is a morphism (in \mathcal{C}_2) of (suspended, δ -exact) functors where f and f^o are mates as in Lemma 1.2.6 (resp. Theorem 1.5.4) when setting $J_1 = D_1, K_1 = D_1^o, J_2 = D_2, K_2 = D_2^o, H = F^o, H' = F, a = f^o$ and $b = f$. In other words, the diagram \boxed{H} (equivalent to $\boxed{H'}$) of Lemma 1.2.6 must commute:

$$\begin{array}{ccc} F & \xrightarrow{F\varpi_1} & FD_1^o D_1 \\ \varpi_2 F \downarrow & \boxed{P} & \downarrow f D_1 \\ D_2^o D_2 F & \xrightarrow{D_2^o f^o} & D_2^o F^o D_1 \end{array}$$

We sometimes denote the functor simply by $\{F\}$ if f is understood. When f is an isomorphism, we say the functor is strong duality preserving.

When the dualities are both strict and strong and the functor is strongly duality preserving, then this coincides with the usual definition (see for example [G02, Definition 2.6] where Point 2 corresponds to the fact that f is a morphism of suspended functors in our definition and is only used in the suspended case). Duality preserving functors are composed in the obvious way by setting $\{F', f'\}\{F, f\} = \{F'F, f'F^o \circ F'f\}$.

2.2.2. **DEFINITION.** A morphism of (suspended, exact) duality preserving functors from $\{F, f\}$ to $\{F', f'\}$ (with same source and target) is a morphism of the underlying (suspended, exact) functors $\rho : F \rightarrow F'$ such that the diagram

$$\begin{array}{ccc} FD_1^o & \xrightarrow{f} & D_2^o F^o \\ \rho D_1^o \downarrow & \boxed{M} & \downarrow D_2^o \rho \\ F'D_1^o & \xrightarrow{f'} & D_2^o (F')^o \end{array}$$

commutes. We say that such a morphism is strong if the underlying morphism of functors is an isomorphism.

Composing two morphisms between duality preserving functors obviously gives another one, and composition preserves being strong.

The proofs of following two propositions are straightforward (see also [G02, Theorem 2.7] for a proof of the first one in the strict case).

2.2.3. **PROPOSITION.** A 1-exact strong duality preserving functor $\{F, f\}$ between triangulated categories with strong dualities induces a morphism on Witt groups by sending an element represented by a form $\psi : A \rightarrow D_1(A)$ to the class of the form $f_A \circ F(\psi)$.

2.2.4. PROPOSITION. *A duality preserving isomorphism between 1-exact strong duality preserving functors ensures that they induce the same morphism on Witt groups.*

The following definition and lemma will be used when deriving applications of the main theorems : often, the strongness assumptions on the functors and morphisms necessary to define Witt groups, morphisms of Witt groups and to compare them only hold on subcategories that are not stable by the tensor product.

2.2.5. DEFINITION.

1. *Let (\mathcal{C}, D, ϖ) be a triangulated category with duality and let \mathcal{A} be a full triangulated subcategory of \mathcal{C} preserved by D . Then, $(\mathcal{A}, D|_{\mathcal{A}}, \varpi|_{\mathcal{A}})$ is trivially a triangulated category with duality, and we say that the duality of \mathcal{C} restricts to \mathcal{A} .*
2. *Let $\{F, f\} : (\mathcal{C}_1, D_1, \varpi_1) \rightarrow (\mathcal{C}_2, D_2, \varpi_2)$ be a duality preserving functor between triangulated categories with dualities as in Definition 2.2.1. Assume that there are full triangulated subcategories $\mathcal{A}_i \subset \mathcal{C}_i$, $i = 1, 2$ such that D_i restricts to \mathcal{A}_i and such that $F|_{\mathcal{A}_1}$ factors through \mathcal{A}_2 . Then we say that the duality preserving pair $\{F, f\}$ restricts to the subcategories \mathcal{A}_1 and \mathcal{A}_2 .*
3. *Let ρ be a morphism between two such restricting functors $\{F, f\}$ and $\{F', f'\}$, as in Definition 2.2.2, then the restriction of ρ automatically defines a morphism of exact duality preserving functors between the restricted functors.*

2.2.6. LEMMA. *Let $\{F, f\} : (\mathcal{C}_1, D_1, \varpi_1) \rightarrow (\mathcal{C}_2, D_2, \varpi_2)$ be a duality preserving functor between triangulated categories with duality that restricts to the subcategories \mathcal{A}_1 and \mathcal{A}_2 .*

1. *Assume that the restricted dualities on \mathcal{A}_1 and \mathcal{A}_2 are strong and that $f|_{F(\mathcal{A}_1^\circ)}$ is an isomorphism. Then the restriction of the duality preserving pair $\{F, f\}$ to \mathcal{A}_1 and \mathcal{A}_2 is a strongly duality preserving functor between triangulated categories with strong duality and therefore induces a morphism*

$$W(\mathcal{A}_1, D_1|_{\mathcal{A}_1}, \varpi_1|_{\mathcal{A}_1}) \rightarrow W(\mathcal{A}_2, D_2|_{\mathcal{A}_2}, \varpi_2|_{\mathcal{A}_2})$$

on Witt groups.

2. *If a morphism ρ between two such duality preserving functors is strong when restricted to \mathcal{A}_1 , then they induce the same morphisms on Witt groups.*

PROOF. One has to check that certain diagrams in \mathcal{A}_2 are commutative. This follows as they are already commutative in \mathcal{C}_2 by assumption. Now Propositions 2.2.3 and 2.2.4 prove the claims. ■

3. Consequences of the closed monoidal structure

We now recall a few notions on tensor products and internal Hom functors (denoted by $[-, *]$) and prove very basic facts related to the suspension. A category satisfying the axioms of this section deserves to be called a “(suspended or triangulated) closed symmetric monoidal category”. Then we prove that the dualities defined using the internal Hom on such a category are naturally equipped with the necessary data to define (triangulated) categories with dualities.

3.1. TENSOR PRODUCT AND INTERNAL HOM. Let (\mathcal{C}, \otimes) be a symmetric monoidal category (see [ML98, Chapter VII]) with an internal Hom $[-, -]$ adjoint to the tensor product. More precisely, we assume that $(- \otimes *, [* , -])$ is an ACB. Let $s = s^{-1}$ denote the symmetry isomorphism and call this datum a “closed symmetric monoidal category”.

When talking about a “suspended closed symmetric monoidal category”, we assume that we have a suspended bifunctor $(- \otimes *, tp_1, tp_2)$ (see Definition 1.4.12) such that the diagram

$$\begin{array}{ccc} (TA \otimes B) \otimes C & \longrightarrow & TA \otimes (B \otimes C) \\ \begin{array}{c} tp_1 \otimes id \downarrow \\ T(A \otimes B) \otimes C \\ tp_1 \downarrow \end{array} & \boxed{\text{assoc}} & \downarrow tp_1 \\ T((A \otimes B) \otimes C) & \longrightarrow & T(A \otimes (B \otimes C)) \end{array}$$

commutes, as well as the two similar ones in which the suspension starts on one of the other variables. We also assume that the diagram

$$\begin{array}{ccc} T(-) \otimes * & \xrightarrow{s} & * \otimes T(-) \\ \begin{array}{c} tp_1 \downarrow \\ T(- \otimes *) \\ \xrightarrow{Ts} \end{array} & \boxed{s} & \downarrow tp_2 \\ T(- \otimes *) & \xrightarrow{Ts} & T(* \otimes -) \end{array}$$

commutes. By Proposition 1.5.8, we get morphisms

$$th_1 : [T^{-1}(*), -] \rightarrow T[* , -] \qquad th_2 : [* , T(-)] \rightarrow T[* , -]$$

that make $([* , -], th_1, th_2)$ a suspended bifunctor and $(- \otimes *, [* , -])$ a suspended ACB (Definition 1.5.5). Using s , we obtain a new suspended ACB $(* \otimes -, [* , -])$ from the previous one.

If \mathcal{C} is triangulated, we furthermore assume that \otimes is exact in both variables (by symmetry it suffices to check this for one of them). By Proposition 1.5.8, $[* , -]$ is suspended in both variables and automatically exact in the second variable. We assume furthermore that it is exact in the first variable, and say that we have a “triangulated closed symmetric monoidal category”.

The morphisms

$$ev_{A,K}^l : [A, K] \otimes A \rightarrow K \qquad coev_{A,K}^l : K \rightarrow [A, K \otimes A]$$

respectively

$$\text{ev}_{A,K}^r : A \otimes [A, K] \rightarrow K \quad \text{coev}_{A,K}^r : K \rightarrow [A, A \otimes K]$$

induced by the counit and the unit of the (suspended) ACB $(- \otimes *, [* , -])$ (or resp. $(* \otimes -, [* , -])$) are called the left (resp. right) evaluation and coevaluation.

3.1.1. LEMMA. *The following diagrams are commutative.*

$$\begin{array}{ccc} (T[A, K]) \otimes A \xrightarrow{tp_1} T([A, K] \otimes A) & & A \otimes (T[A, K]) \xrightarrow{tp_2} T(A \otimes [A, K]) \\ th_2 \otimes id \uparrow \quad \boxed{1} & & id \otimes th_2 \uparrow \quad \boxed{2} \\ [A, TK] \otimes A \xrightarrow{\text{ev}_{A,TK}^l} TK & & A \otimes [A, TK] \xrightarrow{\text{ev}_{A,TK}^r} TK \end{array}$$

$$\begin{array}{ccc} TK \xrightarrow{\text{coev}_{A,TK}^l} [A, TK \otimes A] & & TK \xrightarrow{\text{coev}_{A,TK}^r} [A, A \otimes TK] \\ T\text{coev}_{A,K}^l \downarrow \quad \boxed{3} & & T\text{coev}_{A,K}^r \downarrow \quad \boxed{4} \\ T[A, K \otimes A] \xrightarrow{th_2^{-1}} [A, T(A \otimes K)] & & T[A, A \otimes K] \xrightarrow{th_2^{-1}} [A, T(K \otimes A)] \end{array}$$

$$\begin{array}{ccc} (T^{-1}[A, K]) \otimes TA \xrightarrow{tp_2} T(T^{-1}[A, K] \otimes A) \xrightarrow{tp_1^{-1}} [A, K] \otimes A \\ T^{-1}th_{1,TA,K} \otimes id \downarrow \quad \boxed{5} & & \downarrow \text{ev}_{A,K}^l \\ [TA, K] \otimes TA \xrightarrow{\text{ev}_{TA,K}^l} K \end{array}$$

$$\begin{array}{ccc} (TA \otimes T^{-1}[A, K]) \xrightarrow{tp_1} T(A \otimes T^{-1}[A, K]) \xrightarrow{tp_2^{-1}} A \otimes [A, K] \\ id \otimes T^{-1}th_{1,TA,K} \downarrow \quad \boxed{6} & & \downarrow \text{ev}_{A,K}^r \\ TA \otimes [TA, K] \xrightarrow{\text{ev}_{TA,K}^r} K \end{array}$$

$$\begin{array}{ccc} K \xrightarrow{\text{coev}_{TA,K}^l} [TA, K \otimes TA] \\ \text{coev}_{A,K}^l \downarrow \quad T^{-1}th_2^{-1} \quad \boxed{7} & & \downarrow T^{-1}th_{1,TA,K}^{-1} \\ [A, K \otimes A] \xrightarrow{T^{-1}th_2^{-1}} T^{-1}[A, T(K \otimes A)] \xrightarrow{tp_2^{-1}} T^{-1}[A, K \otimes TA] \end{array}$$

$$\begin{array}{ccc} K \xrightarrow{\text{coev}_{TA,K}^r} [TA, TA \otimes K] \\ \text{coev}_{A,K}^r \downarrow \quad T^{-1}th_2^{-1} \quad \boxed{8} & & \downarrow T^{-1}th_{1,TA,TA \otimes K}^{-1} \\ [A, A \otimes K] \xrightarrow{T^{-1}th_2^{-1}} T^{-1}[A, T(A \otimes K)] \xrightarrow{tp_1^{-1}} T^{-1}[A, TA \otimes K] \end{array}$$

PROOF. This is a straightforward consequence of Point 1 of Definition 1.5.5 for the first four diagrams and of Point 2 of Definition 1.5.5 for the other four. \blacksquare

3.2. **BIDUAL ISOMORPHISM.** We still assume that (\mathcal{C}, \otimes) is a monoidal category with an internal Hom as in the previous section. We now show that the functor $D_K = [-, K]$ naturally defines a duality on the category \mathcal{C} , and that in the suspended case that the dualities D_{TK} and TD_K are naturally isomorphic.

To form the adjoint couple $(D_K, D_K^o, \varpi_K, \varpi_K^o)$, we define the bidual morphism of functors $\varpi_K : Id \rightarrow D_K^o D_K$ as the image of the right evaluation by the adjunction $(- \otimes *, [* , -])$ isomorphism

$$\mathrm{Hom}(A \otimes [A, K], K) \xrightarrow{\sim} \mathrm{Hom}(A, [[A, K], K]).$$

It is functorial in A and defines a morphism of functors from Id to $D_K^o D_K$. Note that its definition uses the adjunction $(- \otimes *, [* , -])$ and the right evaluation, which is not the counit of this adjunction but of the one obtained from it by using s ; so the fact that the monoidal category is symmetric is essential, here. One cannot proceed with only one of these adjunctions. In the suspended case, D_K becomes a suspended functor via $T^{-1}th_{1,-,K}^{-1}T : D_K T \rightarrow T^{-1}D_K$.

3.2.1. **PROPOSITION.** *In the suspended (or triangulated) case, ϖ_K is a morphism of suspended (or exact) functors.*

PROOF. First note that in the exact case, whatever the sign of D_K is, $D_K^o D_K$ is 1-exact, so there is no sign involved in the diagram

$$\begin{array}{ccc} TA & \xrightarrow{\varpi_K T} & [[TA, K], K] \\ T\varpi_K \downarrow & & \downarrow T^{-1}th_{1,TA,K} \\ T[[A, K], K] & \xrightarrow{th_{1,[A,K],K}^{-1}} & [T^{-1}[A, K], K] \end{array}$$

that we have to check (see Definitions 1.4.10 and 1.4.6). It is obtained by (suspended) adjunction from Diagram [6](#) in Lemma 3.1.1 ■

3.2.2. **DEFINITION.** *We say that K is a dualizing object when ϖ_K is an isomorphism of (suspended) functors.*

3.2.3. **PROPOSITION.** *The functors $[-, *] : \mathcal{C} \times \mathcal{C}^o \rightarrow \mathcal{C}^o$ and $[-, *]^o : \mathcal{C}^o \times \mathcal{C} \rightarrow \mathcal{C}$ form a (suspended) ACB with unit ϖ and counit ϖ^o .*

PROOF. We first have to prove that $(D_K, D_K^o, \varpi_K, \varpi_K^o)$ is an adjoint couple in the usual sense. We already know that ϖ_K is a suspended morphism. Consider the following diagram, in which all vertical maps are isomorphisms. Let $f^\sharp : \mathrm{Hom}(F', G) \rightarrow \mathrm{Hom}(F, G)$ and $f_\sharp : \mathrm{Hom}(G, F) \rightarrow \mathrm{Hom}(G, F')$ be the maps induced by $f : F \rightarrow F'$. The unlabeled

morphisms are just adjunction bijections, and we set $\varpi_{A,K} := (\varpi_K)_A$.

$$\begin{array}{ccc}
\mathrm{Hom}([A, K], [A, K]) & \xleftarrow{(\varpi_{A,K}, Id_K)^\#} & \mathrm{Hom}([A, K], [[[A, K], K], K]) \\
\uparrow & & \uparrow \\
\mathrm{Hom}([A, K] \otimes A, K) & \xleftarrow{(Id_{[A,K]} \otimes \varpi_{A,K})^\#} & \mathrm{Hom}([A, K] \otimes [[A, K], K], K) \\
\uparrow s_{A,[A,K]}^\# & & \uparrow s_{[[A,K],K],[A,K]}^\# \\
\mathrm{Hom}(A \otimes [A, K], K) & \xleftarrow{(\varpi_{A,K} \otimes Id_{[A,K]})^\#} & \mathrm{Hom}([A, K], K \otimes [A, K], K) \\
\downarrow & & \downarrow \\
\mathrm{Hom}(A, [[A, K], K]) & \xleftarrow{(\varpi_{A,K})^\#} & \mathrm{Hom}([A, K], K, [[A, K], K])
\end{array}$$

The diagram commutes by functoriality of s and the adjunction bijections. Now $Id_{[[A,K],K]}$ in the lower right set is sent to $\varpi_{[A,K],K}$ in the upper right set, which is in turn sent to $[\varpi_{A,K}, K] \circ \varpi_{[A,K],K}$ in the upper left set. But $Id_{[[A,K],K]}$ is also sent to $\varpi_{A,K}$ in the lower left set, which is sent to $Id_{[A,K]}$ in the upper left set by definition of $\varpi_{A,K}$. This proves the two required formulas (see Definition 1.2.1) for the composition of the unit and the counit in the adjoint couple (which are identical in this case). We leave to the reader the easy fact that $\varpi_{A,K}$ is a generalized transformation (in K) (just using that the unit of the adjunction $(- \otimes *, [* , -])$ is one). The adjoint couple is then a suspended adjoint couple by Proposition 3.2.1. This proves Point 1 of Definition 1.5.5. Point 2 is proved using diagrams [6] and [7]. ■

3.2.4. COROLLARY. *The triple $(\mathcal{C}, D_K, \varpi_K)$ is a (suspended) category with duality (see Definition 2.1.1). When \mathcal{C} is triangulated closed, D_K is exact, and $(\mathcal{C}, D_K, \varpi_K)$ is a triangulated category with duality which we often denote by \mathcal{C}_K for short. When K is a dualizing object, the duality is strong.*

3.2.5. PROPOSITION. *The functors D_K and D_{TK} are exact. The isomorphism $th_2 : [-, T(*)] \rightarrow T[-, *]$ defines a suspended duality preserving functor $\{Id_{\mathcal{C}}, th_{2,-,K}\}$ from \mathcal{C}_{TK} to $T(\mathcal{C}, D_K, \varpi_K)$. This functor is an isomorphism of triangulated categories with duality and therefore induces an isomorphism on Witt groups.*

PROOF. We know by 1.4.9 that TD_K is (-1) -exact. Diagram [sus] should therefore be anti-commutative (see Definition 1.4.10). It follows from the fact that $([-, *], th_1, th_2)$ is a suspended bifunctor. A square obtained by adjunction from [2] in Lemma 3.1.1 then implies that th_2 defines a duality preserving functor (see Definition 2.2.1). ■

We conclude this section by a trivial lemma for future reference.

3.2.6. LEMMA. *Let $\iota : K \rightarrow M$ be a morphism. Then $I_\iota = \{Id, \tilde{\iota}\}$, where $\tilde{\iota} : D_K \rightarrow D_M$ is induced by ι , is a duality preserving functor. This respects composition: if $\kappa : M \rightarrow N$ is another morphism, then $I_{\kappa\iota} = I_\kappa I_\iota$. Let ι be an isomorphism, then if K is dualizing and D_K δ -exact, the same is true for M and D_M , and I_ι induces an isomorphism on Witt groups denoted by I_ι^W .*

4. Functors between closed monoidal categories

Assume from now on that all categories \mathcal{C} (maybe with an index) are symmetric monoidal and equipped with an internal Hom, satisfying the set-up of the previous section. We say that a functor $f^* : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ is a symmetric monoidal (suspended, exact in both variables) functor when it comes equipped with an isomorphism of (suspended) bifunctors $\alpha : f^*(-) \otimes f^*(*) \rightarrow f^*(- \otimes *)$ and, when a unit $\mathbf{1}$ for the tensor product is considered, an isomorphism $f^*(\mathbf{1}) \simeq \mathbf{1}$ making the standard diagrams commutative (see [ML98, section XI.2] for the details where such functors are called *strongly monoidal*).

We will consider the following assumptions (used in the definition of some morphisms of functors):

(**A_f**) The functor $f^* : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ is symmetric monoidal (suspended, exact in both variables).

(**B_f**) We have a functor $f_* : \mathcal{C}_2 \rightarrow \mathcal{C}_1$ that fits into an adjoint couple $(f^*, f_*, \eta_*, \epsilon_*)$.

(**C_f**) We have a functor $f^! : \mathcal{C}_1 \rightarrow \mathcal{C}_2$ that fits into an adjoint couple $(f_*, f^!, \eta_*^!, \epsilon_*^!)$.

(**D_f**) The morphism $\pi : f_*(-) \otimes * \rightarrow f_*(- \otimes f^*(*))$ from Proposition 4.2.5 is an isomorphism (the “projection formula” isomorphism).

(**E_{f,g}**) The morphism of functors ε of Section 5.2 is an isomorphism.

In the following, we will define several natural transformations and establish commutative diagrams involving them. Since there are so many of them, for the convenience of the reader, we include a Table 1, which displays (from the left to the right): the name of the natural transformation, its source and target functor, the necessary assumptions to define the natural transformation and where it is defined.

4.1. THE MONOIDAL FUNCTOR f^* . In this section, we obtain duality preserving functors and morphisms related to a monoidal functor f^* .

4.1.1. PROPOSITION. *Under Assumption (**A_f**), there is a unique morphism*

$$\beta : f^*[*, -] \rightarrow [f^*(*), f^*(-)]$$

of (suspended) bifunctors such that the diagrams

$$\begin{array}{ccccc} [f^*X, f^*(- \otimes X)] & \xleftarrow{\beta} & f^*[X, - \otimes X] & & f^*(-) \longleftarrow [f^*X, f^*(-)] \otimes f^*X \\ \alpha \uparrow & & \uparrow & & \uparrow \beta \\ [f^*X, f^*(-) \otimes f^*X] & \xleftarrow{\quad} & f^*(-) & & f^*([X, -] \otimes X) \xleftarrow{\alpha} f^*[X, -] \otimes f^*X \end{array}$$

commute for every $X \in \mathcal{C}_1$. For any A and B , β is given by the composition

$$f^*[A, B] \xrightarrow{\text{coev}^l} [f^*A, f^*[A, B] \otimes f^*A] \xrightarrow{\alpha} [f^*A, f^*([A, B] \otimes A)] \xrightarrow{\text{ev}^l} [f^*A, f^*B].$$

PROOF. Apply Lemma 1.3.5 to $H = H' = f^*$, $J_1 = - \otimes *$, $K_1 = [*, -]$, $J_2 = (- \otimes f^*(*))$, $K_2 = [f^*(*), -]$ and $a = \alpha$ (recall that in an ACB, the variable denoted $*$ is the parameter, as explained in Definition 1.3.1). Then define $\beta = b$. ■

α	$f^*(-) \otimes f^*(*) \rightarrow f^*(- \otimes *)$	(\mathbf{A}_f)	Section 4
β	$f^*[-, *] \rightarrow [f^*(-), f^*(*)]$	(\mathbf{A}_f)	Prop. 4.1.1
λ	$f_*(-) \otimes f_*(*) \rightarrow f_*(- \otimes *)$	$(\mathbf{A}_f) (\mathbf{B}_f)$	Prop. 4.2.1
μ	$f_*[-, *] \rightarrow [f_*(-), f_*(*)]$	$(\mathbf{A}_f) (\mathbf{B}_f)$	Prop. 4.2.2
π	$f_*(-) \otimes * \rightarrow f_*(- \otimes f^*(*))$	$(\mathbf{A}_f) (\mathbf{B}_f)$	Prop. 4.2.5
κ	$[-, f_*(*)] \rightarrow f_*[f^*(-), *]$	$(\mathbf{A}_f) (\mathbf{B}_f)$	Prop. 4.2.5
ζ	$f_*[* , f^!(-)] \rightarrow [f_*(*) , -]$	$(\mathbf{A}_f) (\mathbf{B}_f) (\mathbf{C}_f)$	Thm. 4.2.9
θ	$f^!(-) \otimes f^*(*) \rightarrow f^!(- \otimes *)$	$(\mathbf{A}_f) (\mathbf{B}_f) (\mathbf{C}_f) (\mathbf{D}_f)$	Prop. 4.3.1
ν	$f^![* , -] \rightarrow [f^*(*) , f^!(*)]$	$(\mathbf{A}_f) (\mathbf{B}_f) (\mathbf{C}_f) (\mathbf{D}_f)$	Prop. 4.3.1
$\tau_{K,M}$	$D_K \otimes D_M \rightarrow D_{K \otimes M}$		Def. 4.4.1
ξ	$\bar{g}^* f^* \rightarrow \bar{f}^* g^*$	$(\mathbf{A}_f) (\mathbf{A}_{\bar{f}}) (\mathbf{A}_g) (\mathbf{A}_{\bar{g}})$ $(\mathbf{B}_g) (\mathbf{B}_{\bar{g}}) (\mathbf{C}_g) (\mathbf{C}_{\bar{g}})$	Section 5.2
ε	$f^* g_* \rightarrow \bar{g}_* \bar{f}^*$	$(\mathbf{A}_f) (\mathbf{A}_{\bar{f}}) (\mathbf{A}_g) (\mathbf{A}_{\bar{g}})$ $(\mathbf{B}_g) (\mathbf{B}_{\bar{g}}) (\mathbf{C}_g) (\mathbf{C}_{\bar{g}})$	Section 5.2
γ	$\bar{f}^* \bar{g}^! \rightarrow g^! f^*$	$(\mathbf{A}_f) (\mathbf{A}_{\bar{f}}) (\mathbf{A}_g) (\mathbf{A}_{\bar{g}}) (\mathbf{B}_g)$ $(\mathbf{B}_{\bar{g}}) (\mathbf{C}_g) (\mathbf{C}_{\bar{g}}) (\mathbf{E}_{f,g})$	Section 5.2

Table 1: Morphisms of functors

4.1.2. THEOREM. (existence of the pull-back) Under Assumption (\mathbf{A}_f) , the morphism

$$\beta_K : f^* D_K^\circ \rightarrow D_{f^*K}^\circ (f^*)^\circ$$

defines a duality preserving functor $\{f^*, \beta_K\}$ of (suspended, triangulated) categories with duality from $(\mathcal{C}_1)_K$ to $(\mathcal{C}_2)_{f^*K}$.

4.1.3. COROLLARY. (Pull-back for Witt groups) When the dualities and the duality preserving functor are strong (i.e. ϖ_K, ϖ_{f^*K} and β_K are isomorphisms), $\{f^*, \beta_K\}$ induces a morphism of Witt groups $f_W^* : W^*(\mathcal{C}_1, K) \rightarrow W^*(\mathcal{C}_2, f^*K)$ by Proposition 2.2.3.

PROOF PROOF OF THEOREM 4.1.2. We need to show that the diagram

$$\begin{array}{ccc} f^* & \xrightarrow{f^* \varpi_K} & f^* D_K^\circ D_K \\ \varpi_K f^* \downarrow & & \downarrow \beta_K D_K \\ D_{f^*K}^\circ D_{f^*K} f^* & \xrightarrow{D_{f^*K}^\circ \beta_K^\circ} & D_{f^*K}^\circ (f^*)^\circ D_K \end{array}$$

4.2.2. PROPOSITION. Assume (\mathbf{A}_f) and (\mathbf{B}_f) . Then, there is a unique morphism

$$\mu : f_*[* , -] \rightarrow [f_*(*) , f_*(-)]$$

of (suspended) bifunctors such that the diagrams

$$\begin{array}{ccc} [f_*X, f_*(- \otimes X)] & \xleftarrow{\mu} & f_*[X, - \otimes X] & & f_*(-) & \xleftarrow{\quad} & [f_*X, f_*(-)] \otimes f_*X \\ \lambda \uparrow & & \uparrow & & \uparrow & & \uparrow \mu \otimes id \\ [f_*X, f_*(-) \otimes f_*X] & \xleftarrow{\quad} & f_*(-) & & f_*([X, -] \otimes X) & \xleftarrow{\lambda} & f_*[X, -] \otimes f_*X \end{array}$$

[13] [14]

commute for every $X \in \mathcal{C}_1$. For any A and B , μ is given by the composition

$$f_*[A, B] \xrightarrow{\text{coev}^l} [f_*A, f_*[A, B] \otimes f_*A] \xrightarrow{\lambda} [f_*A, f_*([A, B] \otimes A)] \xrightarrow{\text{ev}^l} [f_*A, f_*B].$$

PROOF. Apply Lemma 1.3.5 (resp. Lemma 1.5.9) to the functors $J_1 = (- \otimes *)$, $K_1 = [* , -]$, $J_2 = (- \otimes f_*(*))$, $K_2 = [f_*(*) , -]$, $H = H' = f_*$ and $a = \lambda$. ■

4.2.3. LEMMA. The diagram

$$\begin{array}{ccc} f_*A \otimes f_*B & \xrightarrow{\lambda} & f_*(A \otimes B) \\ s(f_* \otimes f_*) \downarrow & & \downarrow f_*s \\ f_*B \otimes f_*A & \xrightarrow{\lambda} & f_*(B \otimes A) \end{array}$$

[15]

is commutative.

PROOF. Apply Lemma 1.2.7 to the cube

$$\begin{array}{ccccc} \mathcal{C}_2 \times \mathcal{C}_2 & \xrightarrow{\otimes} & \mathcal{C}_2 & & \\ \uparrow f^* \times f^* & \searrow x & \nearrow c & \searrow Id & \\ \mathcal{C}_2 \times \mathcal{C}_2 & \xrightarrow{\otimes} & \mathcal{C}_2 & & \\ \uparrow id & \searrow f^* \times f^* & \nearrow \alpha^{-1} & \searrow f^* & \\ \mathcal{C}_1 \times \mathcal{C}_1 & \xrightarrow{\otimes} & \mathcal{C}_1 & & \\ \uparrow x & \searrow c & \nearrow Id & \searrow f^* & \\ \mathcal{C}_1 \times \mathcal{C}_1 & \xrightarrow{\otimes} & \mathcal{C}_1 & & \end{array}$$

where x is the functor exchanging the components. Note that the morphism of functors $f_*(- \otimes *) \rightarrow f_*(-) \otimes f_*(*)$ obtained on the front and back squares indeed coincides with λ by construction. ■

4.2.4. PROPOSITION. In the suspended case, under Assumption (\mathbf{B}_f) there is a unique way of turning f_* into a suspended functor such that (f^*, f_*) is a suspended adjoint couple. If further Assumption (\mathbf{C}_f) holds, then there is a unique way of turning $f^!$ into a suspended functor such that $(f_*, f^!)$ is a suspended adjoint couple.

PROOF. Both results follow directly from Proposition 1.5.3. ■

4.2.5. PROPOSITION. Under assumptions (\mathbf{A}_f) and (\mathbf{B}_f) , there is a unique morphism

$$\pi : f_*(-) \otimes * \rightarrow f_*(- \otimes f^*(*))$$

and a unique isomorphism

$$\kappa : [* , f_*(-)] \rightarrow f_*[f^*(*) , -]$$

of (suspended) bifunctors such that the diagrams

$$\begin{array}{ccc}
 [X, f_*(- \otimes f^* X)] & \xleftarrow{\kappa^{-1}} & f_*[f^* X, - \otimes f^* X] & & f^* & \xleftarrow{\quad} & [X, f_*(-)] \otimes X \\
 \uparrow \pi & & \uparrow & & \uparrow & & \uparrow \kappa^{-1} \\
 [X, f_*(-) \otimes X] & \xleftarrow{\quad} & f_* & & f_*([f^* X, -] \otimes f^* X) & \xleftarrow{\pi} & f_*[f^* X, -] \otimes X \\
 \\
 - \otimes X & \xrightarrow{\quad} & f_* f^*(-) \otimes X & & f^*(f_*(-) \otimes X) & \xrightarrow{\alpha^{-1}} & f^* f^*(-) \otimes f^* X \\
 \downarrow & & \downarrow \pi & & \downarrow \pi & & \downarrow \\
 f_* f^*(- \otimes X) & \xrightarrow{\alpha^{-1}} & f_*(f^*(-) \otimes f^* X) & & f^* f^*(- \otimes f^* X) & \xrightarrow{\quad} & - \otimes f^* X \\
 \\
 [X, -] & \xrightarrow{\quad} & [X, f_* f^*(-)] & & f^*[X, f_*(-)] & \xrightarrow{\beta} & [f^* X, f^* f^*(-)] \\
 \downarrow & & \downarrow \kappa & & \downarrow \kappa & & \downarrow \\
 f_* f^*[X, -] & \xrightarrow{\beta} & f_*[f^* X, f^*(-)] & & f^* f^*[f^* X, -] & \xrightarrow{\quad} & [f^* X, -]
 \end{array}$$

commute for any $X \in \mathcal{C}_1$. For any A and B , π is given by the composition

$$f_* A \otimes B \xrightarrow{\eta_*^*} f_* f^*(f_* A \otimes B) \xrightarrow{\alpha} f_*(f^* f_* A \otimes f^* B) \xrightarrow{\epsilon_*^*} f_*(A \otimes f^* B)$$

κ by

$$[A, f_* B] \xrightarrow{\eta_*^*} f_* f^*[A, f_* B] \xrightarrow{\beta} f_*[f^* A, f^* f_* B] \xrightarrow{\epsilon_*^*} f_*[f^* A, B]$$

and κ^{-1} by

$$f_*[f^* A, B] \xrightarrow{\text{coev}^l} [A, f_*[f^* A, B] \otimes A] \xrightarrow{\pi} [A, f_*([f^* A, B] \otimes f^* A)] \xrightarrow{\text{ev}^l} [A, f_* B].$$

PROOF. Apply Point 3 of Theorem 1.5.10 with $L = L' = f^*$, $R = R' = f_*$, $G_1 = - \otimes *$, $G_2 = - \otimes f^*(*)$, $F_1 = [* , -]$, $F_2 = [f^*(*) , -]$, $g_L = (g'_L)^{-1} = \alpha$. Then define $\pi = g_R$ and $\kappa = f'_R$. \blacksquare

4.2.6. LEMMA. *The composition*

$$f_*(-) \otimes (-) \xrightarrow{id \otimes \eta_*^*} f_*(-) \otimes f_*f^*(-) \xrightarrow{\lambda} f_*(- \otimes f^*(-))$$

coincides with π .

PROOF. This follows from the commutative diagram (for any A and B in \mathcal{C}_2)

$$\begin{array}{ccccc}
 f_*A \otimes B & \longrightarrow & f_*A \otimes f_*f^*B & & \\
 \downarrow & \boxed{mf} & \downarrow & & \searrow \lambda \\
 f_*f^*(f_*A \otimes B) & \longrightarrow & f_*f^*(f_*A \otimes f_*f^*B) & & \\
 \alpha^{-1} \downarrow & \boxed{mf} & \downarrow \alpha^{-1} & & \\
 f_*(f^*f_*A \otimes f^*B) & \longrightarrow & f_*(f^*f_*A \otimes f^*f_*f^*B) & & \\
 \downarrow & \boxed{mf} & \downarrow & & \\
 f_*(A \otimes f^*B) & \longrightarrow & f_*(A \otimes f^*f_*f^*B) & \longrightarrow & f_*(A \otimes f^*B) \\
 & & \boxed{adj} & & \\
 & & Id & &
 \end{array}$$

A large curved arrow labeled π goes from the top-left node to the bottom-right node. A large curved arrow labeled λ goes from the top-right node to the bottom-right node.

in which the curved maps are indeed λ and π by construction. ■

4.2.7. LEMMA. *The composition*

$$f_*(-) \otimes f_*(-) \xrightarrow{\pi} f_*(- \otimes f^*f_*(-)) \xrightarrow{f_*(id \otimes \epsilon_*^*)} f_*(- \otimes -)$$

coincides with λ .

PROOF. This follows directly from the construction of π and diagram [11](#). ■

4.2.8. LEMMA. *The composition*

$$f_*[f^*(-), *] \xrightarrow{\mu} [f_*f^*(-), f_*(*)] \xrightarrow{\eta_*^*} [-, f_*(*)]$$

coincides with κ .

PROOF. This follows from the commutative diagram (for any A in \mathcal{C}_1 and B in \mathcal{C}_2)

$$\begin{array}{ccc}
 f_*[f^*A, B] & \xrightarrow{coev^l} & [A, f_*[f^*A, B] \otimes A] \\
 \text{coev}^l \downarrow & \boxed{gen} & \downarrow \eta_*^* \\
 [f_*f^*A, f_*[f^*A, B] \otimes f_*f^*A] & \xrightarrow{\eta_*^*} & [A, f_*[f^*A, B] \otimes f_*f^*A] \\
 \lambda \downarrow & \boxed{mf} & \downarrow \lambda \\
 [f_*f^*A, f_*([f^*A, B] \otimes f^*A)] & \longrightarrow & [A, f_*([f^*A, B] \otimes f^*A)] \\
 \text{ev}^l \downarrow & \boxed{mf} & \downarrow \text{ev}^l \\
 [f_*f^*A, f_*B] & \xrightarrow{\eta_*^*} & [A, f_*B]
 \end{array}$$

A large curved arrow labeled μ goes from the top-left node to the bottom-left node. A large curved arrow labeled π goes from the top-right node to the bottom-right node.

in which the left curved arrow is μ by construction, the right one is π by Lemma 4.2.6 and the composition from the top left corner to the bottom right one along the upper right corner is then κ by construction. ■

4.2.9. **THEOREM.** (*existence of the push-forward*) Under assumptions (\mathbf{A}_f) , (\mathbf{B}_f) and (\mathbf{C}_f) , let ζ be the (suspended, exact) bifunctor defined by the composition

$$f_*[-, f^!(-)] \xrightarrow{\mu} [f_*(-), f_*f^!(-)] \xrightarrow{\epsilon_*^!} [f_*(-), -].$$

Then $\{f_*, \zeta_K\}$ is a (suspended, exact) duality preserving functor from $(\mathcal{C}_2, D_{f^!K}, \varpi_{f^!K})$ to $(\mathcal{C}_1, D_K, \varpi_K)$.

PROOF. We need to prove that the diagram $\square P$ in Definition 2.2.1 is commutative. We consider the commutative diagram

$$\begin{array}{ccc}
 f_*A & \xrightarrow{\text{coev}^l} & f_*[[A, f^!K], A \otimes [A, f^!K]] \xrightarrow{\text{ev}^r} f_*[[A, f^!K], f^!K] \\
 \text{coev}^l \downarrow & \boxed{13} & \mu \downarrow \\
 [f_*[A, f^!K], f_*A \otimes f_*[A, f^!K]] & \xrightarrow{\quad} & [f_*[A, f^!K], f_*(A \otimes [A, f^!K])] \\
 \text{id} \otimes \mu \downarrow & \boxed{14'} & \text{ev}^r \downarrow \\
 [f_*[A, f^!K], f_*A \otimes [f_*A, f_*f^!K]] & \xrightarrow{\text{ev}^r} & [f_*[A, f^!K], f_*f^!K] \quad \boxed{mf} \\
 \text{id} \otimes \epsilon_*^! \downarrow & \boxed{mf} & \epsilon_*^! \downarrow \\
 [f_*[A, f^!K], f_*A \otimes [f_*A, K]] & \xrightarrow{\text{ev}^r} & [f_*[A, f^!K], K]
 \end{array}$$

μ (curved arrow from top right to bottom right)

where $\boxed{14'}$ is obtained from $\boxed{14}$ by using s . The top row is $f_*\varpi_{f^!K}$ and the composition from the top right corner to the bottom one is ζ , both by definition. The result follows if we prove that the composition from the top left corner to the bottom right one, going through the bottom left corner, is $\varpi_K f_*$ followed by $D_K^o \zeta^o$. This follows from the commutative diagram

$$\begin{array}{ccc}
 f_*A & \xrightarrow{\text{coev}^l} & [f_*[A, f^!K], f_*A \otimes f_*[A, f^!K]] \\
 \text{coev}^l \downarrow & \boxed{gen} & \downarrow \text{id} \otimes \zeta \\
 [[f_*A, K], f_*A \otimes [f_*A, K]] & \xrightarrow{\zeta} & [f_*[A, f^!K], f_*A \otimes [f_*A, K]] \\
 \text{ev}^r \downarrow & \boxed{mf} & \downarrow \text{ev}^r \\
 [[f_*A, K], K] & \xrightarrow{\zeta} & [f_*[A, f^!K], K]
 \end{array}$$

in which the left vertical composition is $\varpi_K f_*$ and the composition along the upper right corner is the composition along the lower left corner of the previous diagram. \blacksquare

4.3. **WHEN THE PROJECTION MORPHISM IS INVERTIBLE.** In this section, we prove that the projection morphism is an isomorphism if and only if the morphism ζ is. In that case, we obtain a new morphism θ that will be used in Section 5.5 to state a projection formula.

4.3.1. **PROPOSITION.** Under assumptions (\mathbf{A}_f) , (\mathbf{B}_f) and (\mathbf{C}_f) , there is a unique morphism

$$\nu' : [f^*(*), f^!(-)] \rightarrow f^![*, -]$$

of (suspended) bifunctors such that the diagrams

$$\begin{array}{ccccc}
 [f^*X, -] & \longrightarrow & [f^*X, f^!f_*(-)] & f_*[f^*X, f^!(-)] & \xrightarrow{\kappa^{-1}} & [X, f_*f^!(-)] \\
 \downarrow & & \downarrow \nu' & \nu' \downarrow & & \downarrow \\
 f^!f_*[f^*X, -] & \xrightarrow{\kappa^{-1}} & f^![X, f_*(-)] & f_*f^![X, -] & \longrightarrow & [X, -]
 \end{array}$$

[22]
[23]

commute for any $X \in \mathcal{C}_1$. The morphism of bifunctors ν' is invertible if and only if π is invertible (Assumption **(D_f)**). In that case, let ν denote the inverse of ν' , and there is a unique morphism of (suspended) bifunctors

$$\theta : f^!(-) \otimes f^*(*) \rightarrow f^!(- \otimes *)$$

such that the diagrams

$$\begin{array}{ccc}
 [f^*X, f^!(- \otimes X)] & \xleftarrow{\nu} & f^![X, - \otimes X] \\
 \theta \uparrow & & \uparrow \\
 [f^*X, f^!(-) \otimes f^*X] & \xleftarrow{\quad} & f^!
 \end{array}
 \quad
 \begin{array}{ccc}
 f^! & \xleftarrow{\quad} & [f^*X, f^!(-)] \otimes f^*X \\
 \uparrow & & \uparrow \nu \\
 f^!([X, -] \otimes X) & \xleftarrow{\theta} & f^![X, -] \otimes f^*X
 \end{array}$$

$$\begin{array}{ccc}
 - \otimes f^*X & \longrightarrow & f^!f_*(-) \otimes f^*X & f_*(f^!(-) \otimes f^*X) & \xrightarrow{\pi^{-1}} & f_*f^!(-) \otimes X \\
 \downarrow & & \downarrow \theta & \theta \downarrow & & \downarrow \\
 f^!f_*(- \otimes f^*X) & \xrightarrow{\pi^{-1}} & f^!(f_*(-) \otimes X) & f_*f^!(- \otimes X) & \longrightarrow & - \otimes X
 \end{array}$$

[24]
[25]

[26]
[27]

commute for any $X \in \mathcal{C}_1$. For any A and B , ν' is given by the composition

$$[f^*A, f^!B] \xrightarrow{\eta_*^!} f^!f_*[f^*A, f^!B] \xrightarrow{\kappa^{-1}} f^![A, f_*f^!B] \xrightarrow{\epsilon_*^!} f^![A, B]$$

and θ by

$$f^!A \otimes f^*B \xrightarrow{\eta_*^!} f^!f_*(f^!A \otimes f^*B) \xrightarrow{\pi^{-1}} f^!(f_*f^!A \otimes B) \xrightarrow{\epsilon_*^!} f^!(A \otimes B)$$

or equivalently by

$$f^!A \otimes f^*B \xrightarrow{\text{coev}^!} f^![B, A \otimes B] \otimes f^*B \xrightarrow{\nu} [f^*B, f^!(A \otimes B)] \otimes f^*B \xrightarrow{\text{ev}^!} f^!(A \otimes B).$$

PROOF. Apply Point 2 of Theorem 1.5.10 with $L = L' = f_*$, $R = R' = f^!$, $G_1 = - \otimes f^*(*)$, $G_2 = - \otimes *$, $F_1 = [f^*(*), -]$, $F_2 = [* , -]$ and $f_L = \kappa^{-1}$ to obtain $\nu' = f'_R$ and diagrams [22] and [23]. Then Point 3 of the same Theorem tells us that $\nu' = f'_R$ is invertible if and only if $\pi = g_L$ is (Note that π and κ^{-1} indeed correspond to each other through Point 1 of the Theorem by construction). In that case, we can use Point 2 to obtain $\theta = g_R$ from $\pi^{-1} = g'_L$ and diagrams [24], [25], [26] and [27]. ■

In particular, under Assumption (\mathbf{D}_f) , this gives for each K an isomorphism of functors $\nu_K : f^! D_K^o \rightarrow D_{f^! K}^o (f^*)^o$ (in \mathcal{C}_2).

4.3.2. LEMMA. *The composition*

$$f_*[-, f^!(-)] \xrightarrow{\epsilon_*^*} f_*[f^* f_*(-), f^!(-)] \xrightarrow{\nu'} f_* f^![f_*(-), -] \xrightarrow{\epsilon_*^!} [f_*(-), -]$$

coincides with ζ .

PROOF. This follows from the commutative diagram (for every A in \mathcal{C}_2 and K in \mathcal{C}_1)

$$\begin{array}{ccccc} f_*[A, f^!K] & \longrightarrow & f_*[f^* f_* A, f^!K] & \xrightarrow{\nu'} & f_* f^![f_* A, K] \\ \mu \downarrow & \boxed{mf} & \downarrow \mu & & \downarrow \epsilon_*^! \\ [f_* A, f_* f^!K] & \longrightarrow & [f_* f^* f_* A, f_* f^!K] & \boxed{23} & \\ & \searrow \boxed{adj} & \downarrow \eta_*^* & & \\ & & [f_* A, f_* f^!K] & \xrightarrow{\epsilon_*^!} & [f_* A, K] \end{array}$$

identifying the vertical composition in the middle with κ^{-1} by Lemma 4.2.8 to recognize $\boxed{23}$. \blacksquare

4.3.3. PROPOSITION. *Under assumptions (\mathbf{A}_f) , (\mathbf{B}_f) , (\mathbf{C}_f) , ζ is an isomorphism if and only if (\mathbf{D}_f) holds.*

4.3.4. COROLLARY. *(Push-forward for Witt groups) When the dualities $D_{f^! K}$ and D_K are strong and Assumption (\mathbf{D}_f) is satisfied, then $\{f_*, \zeta_K\}$ is a strong duality preserving functor and thus induces a morphism of Witt groups*

$$f_*^W : W^*(\mathcal{C}_2, f^! K) \rightarrow W^*(\mathcal{C}_1, K).$$

PROOF. This follows from Definition 2.1.5, Theorem 4.2.9, Proposition 4.3.3 and Proposition 2.2.3. \blacksquare

PROOF OF PROPOSITION 4.3.3. Let us first recall that $([-, *], [-, *]^o, \varpi, \varpi^o)$ is a (suspended) ACB from \mathcal{C}_1 to \mathcal{C}_1^o with parameter in \mathcal{C}_1^o by Proposition 3.2.3, and thus $([-, (f^!)^o(*)], [-, f^!(*)]^o, \varpi, \varpi^o)$ is a (suspended) ACB from \mathcal{C}_2 to \mathcal{C}_2^o with parameter in \mathcal{C}_1^o by Lemma 1.5.7. We then apply Theorem 1.5.10 with the square

$$\begin{array}{ccc} \mathcal{C}_1 & \xrightleftharpoons{f^*} & \mathcal{C}_2 \\ \uparrow [-, *]^o \Big| \mathcal{C}_1^o & \begin{array}{c} \downarrow [-, *] \\ \xrightarrow{f_*} \\ \downarrow (f^!)^o \end{array} & \uparrow [-, (f^!)^o(*)] \Big| \mathcal{C}_1^o \\ \mathcal{C}_1^o & \xrightleftharpoons{f_*^o} & \mathcal{C}_2^o \end{array}$$

starting with $g'_L = (\nu')^o$. By Point 2 of loc. cit., we obtain a unique g_R such that diagrams $\boxed{G1}$ and $\boxed{G2}$ commute. This g_R coincides with ζ^o by Lemma 4.3.2. By Point 1 of loc. cit., we then obtain a unique f_R such that diagrams \boxed{R} and $\boxed{R'}$ commute. But f_R has to

coincide with ζ by uniqueness, since those commutative diagrams were already proven in Theorem 4.2.9 with ζ and ζ^o . By Point 3 of loc. cit., f_R is an isomorphism if and only if $g'_L = (\nu')^o$ is, which is an isomorphism if and only if (\mathbf{D}_f) holds by Proposition 4.3.1. We have thus proved that ζ is an isomorphism if and only if (\mathbf{D}_f) holds. ■

4.4. PRODUCTS. We now define one more classical morphism related to products for Witt groups. It is easily checked that since $(-\otimes*, [*,-])$ forms an ACB from \mathcal{C} to \mathcal{C} with parameter in \mathcal{C} , $((-1 \otimes *_1) \times (-2 \otimes *_2), [*_1, -1] \times [*_2, -2])$ forms an ACB from $\mathcal{C} \times \mathcal{C}$ to $\mathcal{C} \times \mathcal{C}$ with parameter in $\mathcal{C} \times \mathcal{C}$, and $(-\otimes(*_1 \otimes *_2), [*_1 \otimes *_2, -])$ forms an ACB from \mathcal{C} to \mathcal{C} with parameter in $\mathcal{C} \times \mathcal{C}$. We can thus consider the situation of Lemma 1.3.5 applied to the square

$$\begin{array}{ccc} \mathcal{C} \times \mathcal{C} & \xrightarrow{\quad \otimes \quad} & \mathcal{C} \\ \begin{array}{c} \uparrow \scriptstyle{(-1 \otimes *_1) \times (-2 \otimes *_2)} \\ \mathcal{C} \times \mathcal{C} \\ \downarrow \scriptstyle{[*_1, -1] \times [*_2, -2]} \end{array} & & \begin{array}{c} \uparrow \scriptstyle{-\otimes(*_1 \otimes *_2)} \\ \mathcal{C} \\ \downarrow \scriptstyle{[*_1 \otimes *_2, -]} \end{array} \end{array}$$

and to the morphism of bifunctors $\varrho : (-1 \otimes -2) \otimes (*_1 \otimes *_2) \rightarrow (-1 \otimes *_1) \otimes (-2 \otimes *_2)$. We thus obtain its mate, a morphism of bifunctors

$$\tau : [*_1, -1] \otimes [*_2, -2] \rightarrow [*_1 \otimes *_2, -1 \otimes -2]$$

which, for any A, B, K, M is given by the composition

$$\begin{array}{ccc} [A, K] \otimes [B, M] & \xrightarrow{\text{coev}^t} & [A \otimes B, ([A, K] \otimes [B, M]) \otimes (A \otimes B)] \\ & & \downarrow e \\ [A \otimes B, K \otimes M] & \xleftarrow{\text{ev}^t \otimes \text{ev}^t} & [A \otimes B, ([A, K] \otimes A) \otimes ([B, M] \otimes B)] \end{array}$$

and obtain the following commutative diagrams

$$\begin{array}{ccc} ([X_1, K] \otimes [X_2, M]) \otimes (X_1 \otimes X_2) & \xrightarrow{\tau} & [X_1 \otimes X_2, K \otimes M] \otimes (X_1 \otimes X_2) \\ \downarrow e & \boxed{28} & \downarrow \text{ev}^t \\ ([X_1, K] \otimes X_1) \otimes ([X_2, M] \otimes X_2) & \xrightarrow{\text{ev}^t \otimes \text{ev}^t} & K \otimes M \end{array}$$

$$\begin{array}{ccc} K \otimes M & \xrightarrow{\text{coev}^t \otimes \text{coev}^t} & [X_1, K \otimes X_1] \otimes [X_2, M \otimes X_2] \\ \text{coev}^t \downarrow & \boxed{29} & \downarrow \tau \\ [X_1 \otimes X_2, (K \otimes M) \otimes (X_1 \otimes X_2)] & \xrightarrow{e} & [X_1 \otimes X_2, (K \otimes X_1) \otimes (M \otimes X_2)] \end{array}$$

as well.

4.4.1. DEFINITION. Consider objects $K, M \in \mathcal{C}$. We write $\tau_{K,M}$ for the morphism of functors $D_K(-1) \otimes D_M(-2) \rightarrow D_{K \otimes M}(-1 \otimes -2)$ defined above where $-1 = K$ and $-2 = M$.

The proofs of the next two results are left to the reader. They are not difficult although they require large commutative diagrams.

4.4.2. PROPOSITION. *The morphism $\tau_{K,M}$ is a morphism of suspended bifunctors.*

The proof of the following lemma uses the unit of the tensor product.

4.4.3. LEMMA. *If K is dualizing, M is dualizing and $\tau_{K,M}$ is an isomorphism, then the object $K \otimes M$ is dualizing. If $\beta_{f,K}$, $\beta_{f,M}$ and $\tau_{K,M}$ are isomorphisms, then $\beta_{f,K \otimes M}$ is an isomorphism.*

In [GN03], Gille and Nenashev define two natural products for Witt groups. These products coincide up to a sign. We just choose one of them (the left product, for example), and refer to it as *the* product, but everything works fine with the other one too. Let us recall the basic properties of the product, rephrasing [GN03] in our terminology.

4.4.4. THEOREM. ([GN03, Definition 1.11 and Theorem 2.9]) *Let \mathcal{C}_1 , \mathcal{C}_2 and \mathcal{C}_3 be triangulated categories with dualities D_1 , D_2 and D_3 . Let $(B, b_1, b_2) : \mathcal{C}_1 \times \mathcal{C}_2 \rightarrow \mathcal{C}_3$ be a suspended bifunctor (see Definition 1.4.12) and $d : B(D_1^o \times D_2^o) \rightarrow D_3^o B^o$ be an isomorphism of suspended bifunctors (see Definition 1.4.14) that makes $\{B, d\}$ a duality preserving functor (see Definition 2.2.1, here $\mathcal{C}_1 \times \mathcal{C}_2$ is endowed with the duality $D_1 \times D_2$). Then $\{B, d\}$ induces a product*

$$W(\mathcal{C}_1) \times W(\mathcal{C}_2) \rightarrow W(\mathcal{C}_3).$$

The following proposition is not stated in [GN03], but easily follows from the construction of the product.

4.4.5. PROPOSITION. *Let $\sigma : B \rightarrow B'$ be an isomorphism of suspended bifunctors that is duality preserving. Then B and B' induce the same product on Witt groups.*

Let us now apply this to our context.

4.4.6. PROPOSITION. (product) *The morphism of functors $\tau_{K,M}$ turns the functor \otimes into a duality preserving functor $\{-1 \otimes -2, \tau_{K,M}\}$ from $(\mathcal{C} \times \mathcal{C}, D_K \times D_M, \varpi_K \times \varpi_M)$ to $(\mathcal{C}, D_{K \otimes M}, \varpi_{K \otimes M})$.*

4.4.7. COROLLARY. (product for Witt groups) *By Theorem 4.4.4, when the dualities and the functor are strong (i.e. ϖ_K , ϖ_M and $\tau_{K,M}$ are isomorphisms, and thus $\varpi_{K \otimes M}$ as well by Lemma 4.4.3), $\{-1 \otimes -2, \tau_{K,M}\}$ induces a product*

$$W(\mathcal{C}, D_K) \times W(\mathcal{C}, D_M) \longrightarrow W(\mathcal{C}, D_{K \otimes M})$$

on Witt groups.

5. Relations between functors

In this section, we prove the main results on composition, base change, and the projection formula.

5.1. COMPOSITION. This section studies the behavior of pull-backs and push-forwards with respect to composition. Let \mathcal{K} be a category whose objects are (suspended, triangulated) closed categories as in Section 3.1, and whose morphisms are (suspended, exact) functors. Let \mathcal{B} be another category, and let $(-)^*$ be a contravariant pseudo functor from \mathcal{B} to \mathcal{K} , i.e. a functor, except that instead of having equalities $f^*g^* = (gf)^*$ when f and g are composable, we only have isomorphisms of (suspended) functors $a_{g,f} : f^*g^* \rightarrow (gf)^*$. We also require that $(-)^*$ sends the identity of an object to the identity. Let X^* be denoted by \mathcal{C}_X . When, moreover, the diagram

$$\begin{array}{ccc} f^*g^*h^* & \xrightarrow{a_{g,f}} & (gf)^*h^* \\ a_{h,g} \downarrow & \boxed{30} & \downarrow a_{h,gf} \\ f^*(hg)^* & \xrightarrow{a_{hg,f}} & (hgf)^* \end{array}$$

is commutative, we say that the pseudo functor is associative.

5.1.1. REMARK. An example for this setting is to take for \mathcal{B} the category of schemes (or regular schemes) and $\mathcal{C}_X = D^b(\text{Vect}(X))$ (or $\mathcal{C}_X = D^b(O_X - \text{Mod})$).

We require that $(-)^*$ is a monoidal associative pseudo functor, which means that each f^* is symmetric monoidal (Assumption (\mathbf{A}_f)) and that the diagram

$$\begin{array}{ccc} g^*f^* \otimes g^*f^* & \xrightarrow{\alpha} & g^*(f^* \otimes f^*) \xrightarrow{\alpha} g^*f^*(- \otimes -) \\ a \otimes a \downarrow & \boxed{31} & \downarrow a \\ (fg)^* \otimes (fg)^* & \xrightarrow{\alpha} & (fg)^*(- \otimes -) \end{array}$$

commutes for any two composable f and g .

5.1.2. NOTATION. Let X be an object in \mathcal{B} and K an object in \mathcal{C}_X . Let $\mathcal{C}_{X,K}$ denote the (suspended, triangulated) category with duality $(\mathcal{C}_X, D_K, \varpi_K)$ obtained by Corollary 3.2.4. When K is dualizing, let $W^i(X, K)$ denote the i -th shifted Witt group of $\mathcal{C}_{X,K}$.

We have already seen in Theorem 4.1.2 that under Assumption (\mathbf{A}_f) , the couple $\{f^*, \beta_K\}$ is a duality preserving functor between (suspended, triangulated) categories with dualities. Recall the I -notation of Lemma 3.2.6.

5.1.3. THEOREM. (composition of pull-backs) For any two composable f and g in \mathcal{B} , the isomorphism of functors $a_{g,f} : f^*g^* \rightarrow (gf)^*$ is a morphism of duality preserving functors from $I_{a_{g,f,K}}\{f^*, \beta_{f,g^*K}\}\{g^*, \beta_{g,K}\}$ to $\{(gf)^*, \beta_{gf,K}\}$.

5.1.4. COROLLARY. (composition of pull-backs for Witt groups) By Proposition 2.2.4, for composable morphisms f and g in \mathcal{B} , the pull-back on Witt groups defined in Corollary 4.1.3 satisfies $(gf)^*_W = I_{a_{g,f,K}}^W f^*_W g^*_W$.

PROOF OF THEOREM 5.1.3. The claim amounts to checking that the diagram

$$\begin{array}{ccccc} f^*g^*[-, K] & \xrightarrow{\beta} & f^*[g^*(-), g^*K] & \xrightarrow{\beta} & [f^*g^*(-), f^*g^*K] \\ \downarrow a & & & & \downarrow a \\ (gf)^*[-, K] & \xrightarrow{\beta} & [(gf)^*(-), (gf)^*K] & \xrightarrow{\alpha} & [f^*g^*(-), (gf)^*K] \end{array}$$

is commutative. Since by construction of β (Proposition 4.1.1), α and β are mates, this commutativity follows from Lemma 1.2.7 applied to the cube

$$\begin{array}{ccc} \begin{array}{ccc} \mathcal{C}_Z & \xrightarrow{(gf)^*} & \mathcal{C}_X \\ \downarrow g^* & \searrow a & \downarrow Id \\ \mathcal{C}_Y & \xrightarrow{f^*} & \mathcal{C}_X \\ \downarrow \alpha & \nearrow \alpha & \downarrow \alpha \\ \mathcal{C}_Z & & \mathcal{C}_Y \\ \downarrow g^* & & \downarrow f^* \\ \mathcal{C}_Y & \xrightarrow{f^*} & \mathcal{C}_X \end{array} & & \begin{array}{ccc} \mathcal{C}_Z & \xrightarrow{(gf)^*} & \mathcal{C}_X \\ \downarrow -\otimes K & \searrow -\otimes (gf)^*K & \downarrow Id \\ \mathcal{C}_Z & \xrightarrow{(gf)^*} & \mathcal{C}_X \\ \downarrow g^* & \searrow a & \downarrow Id \\ \mathcal{C}_Y & \xrightarrow{f^*} & \mathcal{C}_X \\ \downarrow -\otimes f^*g^*K & \nearrow id \otimes a & \downarrow -\otimes f^*g^*K \end{array} \end{array}$$

which is commutative as Diagram 31 is. ■

A covariant pseudo functor $(-)_*$ is defined in the same way as a contravariant one, except that we are given isomorphisms in the other direction: $b_{g,f} : (gf)_* \rightarrow g_*f_*$.

5.1.5. DEFINITION. Let $(-)^*$ (resp. $(-)_*$) be a (suspended, exact) contravariant (resp. covariant) pseudo functor from \mathcal{B} to \mathcal{K} . We say that $((-)^*, (-)_*)$ is an adjoint couple of pseudo functors if (f^*, f_*) is a (suspended, exact) adjoint couple for every f (in particular $(-)^*$ and $(-)_*$ coincide on objects), and the diagrams

$$\begin{array}{ccc} Id & \xrightarrow{\quad} & (gf)_*(gf)^* \\ \downarrow & \boxed{32} & \downarrow b \\ g_*f_*f^*g^* & \xrightarrow{a} & g_*f_*(gf)^* \end{array} \quad \text{and} \quad \begin{array}{ccc} f^*g^*(gf)_* & \xrightarrow{b} & f^*g^*g_*f_* \\ \downarrow a & \boxed{33} & \downarrow \\ (gf)^*(gf)_* & \xrightarrow{\quad} & Id \end{array}$$

commute for any composable f and g .

Spelling out the symmetric notion when the left adjoint pseudofunctor is covariant and the right one contravariant is left to the reader.

As usual, the right (or left) adjoint is unique up to unique isomorphism.

5.1.6. LEMMA. Let $(-)^*$ be a contravariant pseudo-functor. Assume that for any f^* , we are given a right (suspended, exact) adjoint f_* which is the identity when f is the identity. Then there is a unique collection of isomorphisms $b_{g,f} : (gf)_* \rightarrow g_*f_*$ such that $((-)^*, (-)_*)$ forms an adjoint couple of pseudo functors.

PROOF. Apply Theorem 1.2.8 (or 1.5.4 in the suspended case) to $(L, R) = (f^*, f_*)$, $(F_1, G_1) = (g^*, g_*)$, $(F_2, G_2) = (Id, Id)$ and $(L', R') = ((gf)^*, (gf)_*)$. This gives the required isomorphism $(gf)_* \rightarrow g_*f_*$ and the diagrams 32 and 33. ■

5.1.7. LEMMA. *The right (resp. left) adjoint of an associative pseudo functor is associative.*

PROOF. Left to the reader. ■

5.1.8. LEMMA. *Let $((-)^*, (-)_*)$ be an adjoint pair of pseudo-functors (left contravariant and right covariant). If $(-)^*$ satisfies the commutativity of [31], then the diagram*

$$\begin{array}{ccc} (gf)_*A \otimes (gf)_*B & \xrightarrow{\lambda} & (gf)_*(A \otimes B) \\ \downarrow b \otimes b & \boxed{34} & \downarrow b \\ g_*f_*A \otimes g_*f_*B & \xrightarrow{\lambda} g_*(f_*A \otimes f_*B) \xrightarrow{\lambda} & g_*f_*(A \otimes B) \end{array}$$

is commutative for every A and B .

PROOF. The commutative diagram

$$\begin{array}{ccccc} g^*(A \otimes B) & \xrightarrow{\alpha^{-1}} & g^*A \otimes g^*B & \xrightarrow{\eta_*^* \otimes \eta_*^*} & f_*f^*g^*A \otimes f_*f^*g^*B \\ \eta_*^* \downarrow & \boxed{mf} & \downarrow & \boxed{11} & \downarrow \lambda \\ f_*f^*g^*(A \otimes B) & \longrightarrow & f_*f^*(g^*A \otimes g^*B) & \longrightarrow & f_*(f^*g^*A \otimes f^*g^*B) \\ f_*a \downarrow & & \boxed{31} & & \boxed{mf} \\ f_*(gf)^*(A \otimes B) & \xrightarrow{\alpha^{-1}} & f_*((gf)^*A \otimes (gf)^*B) & \xleftarrow{\lambda} & f_*(gf)^*A \otimes f_*(gf)^*B \end{array}$$

$f_*a \otimes f_*a$ (curved arrow from $f_*(gf)^*A \otimes f_*(gf)^*B$ to $f_*(gf)^*(A \otimes B)$)

shows that the cube

is commutative. We apply Lemma 1.2.7 to it and obtain a new commutative cube which is Diagram [34]. Note that the morphism of functors on the left face of our first cube indeed gives b in the resulting cube, by construction of b . ■

Let us now consider the subcategory \mathcal{B}' of \mathcal{B} with the same objects, but whose morphisms are only those $f : X \rightarrow Y$ satisfying assumptions (\mathbf{A}_f) , (\mathbf{B}_f) and (\mathbf{C}_f) . We then choose successive right adjoints f_* and $f^!$ for each morphism f (they are unique up to unique isomorphism, and we choose $(Id_X)_* = Id_X^! = Id_{\mathcal{C}_X}$ for simplicity), and by Lemma 5.1.6, using $(-)^*$, we turn them into (suspended, exact) pseudo functors

$$(-)_* : \mathcal{B}' \rightarrow \mathcal{K} \quad (-)^! : \mathcal{B}' \rightarrow \mathcal{K}$$

with structure morphisms $b_{g,f}$ and $c_{g,f}$. They are associative by Lemma 5.1.7.

5.1.9. **THEOREM.** For morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ in \mathcal{B}' and K an object of \mathcal{C}_Z , the isomorphism of functors $b_{g,f} : (gf)_* \rightarrow g_*f_*$ defined above is a morphism of duality preserving functors from $\{(gf)_*, \zeta_K\} I_{c_{g,f,K}}$ to $\{g_*, \zeta_K\}\{f_*, \zeta_{g^!K}\}$.

5.1.10. **COROLLARY.** (composition of push-forwards) By Proposition 2.2.4, for composable morphisms f and g in \mathcal{B}' , the push-forward on Witt groups defined in Corollary 4.3.4 satisfies $(gf)_*^W I_{c_{g,f,K}}^W = g_*^W f_*^W$.

PROOF OF THEOREM 5.1.9. We have to prove that the diagram \boxed{M} of Definition 2.2.2 which is here

$$\begin{array}{ccccc}
 (gf)_*[A, f^!g^!K] & \xrightarrow{c} & (gf)_*[A, (gf)^!K] & \xrightarrow{\zeta} & [(gf)_*A, K] \\
 \downarrow b & & \boxed{35} & & \uparrow b^\circ \\
 g_*f_*[A, f^!g^!K] & \xrightarrow{\zeta} & g_*[f_*A, g^!K] & \xrightarrow{\zeta} & [g_*f_*A, K]
 \end{array}$$

is commutative (for any A and B). The commutative diagram $\boxed{34}$ may be written as the commutative cube

$$\begin{array}{ccccc}
 \mathcal{C}_Z & \xrightarrow{Id} & \mathcal{C}_Z & & \mathcal{C}_Z \\
 \uparrow Id & & \uparrow id & & \uparrow Id \\
 \mathcal{C}_Z & \xrightarrow{Id} & \mathcal{C}_Z & & \mathcal{C}_Z \\
 \uparrow Id & & \uparrow Id & & \uparrow Id \\
 \mathcal{C}_X & \xrightarrow{Id} & \mathcal{C}_X & \xrightarrow{(gf)_*} & \mathcal{C}_Z \\
 \uparrow g_*f_*(-\otimes B) & & \uparrow (gf)_*(-\otimes B) & & \uparrow -\otimes(gf)_*B \\
 \mathcal{C}_X & \xrightarrow{Id} & \mathcal{C}_X & \xrightarrow{f_*} & \mathcal{C}_Y \\
 \uparrow Id & & \uparrow Id & & \uparrow Id \\
 \mathcal{C}_X & \xrightarrow{Id} & \mathcal{C}_X & \xrightarrow{(gf)_*} & \mathcal{C}_Z
 \end{array}$$

out of which we get a new commutative cube by Lemma 1.2.7, whose commutativity is equivalent to that of Diagram $\boxed{35}$. \blacksquare

5.2. **BASE CHANGE.** The next fundamental theorem that we will prove is a base change formula. In this section, we fix a commutative diagram in \mathcal{B} with g and \bar{g} in \mathcal{B}' .

$$\begin{array}{ccc}
 V & \xrightarrow{\bar{g}} & Y \\
 \bar{f} \downarrow & \boxed{36} & \downarrow f \\
 X & \xrightarrow{g} & Z
 \end{array}$$

Using Lemma 1.2.6 (or its suspended version 1.5.4) with $J_1 = g^*$, $K_1 = g_*$, $J_2 = \bar{g}^*$, $K_2 = \bar{g}_*$, $H = \bar{f}^*$, $H' = f^*$ and the isomorphism of (suspended) functors

$$\xi : \bar{g}^* f^* \rightarrow (f\bar{g})^* = (g\bar{f})^* \rightarrow \bar{f}^* g^*$$

for a , we obtain its mate, a morphism

$$\varepsilon : f^* g_* \rightarrow \bar{g}_* \bar{f}^*$$

and two commutative diagrams. Assuming $(\mathbf{E}_{f,g})$ (i.e. ε is an isomorphism) and applying the same lemma to $J_1 = g_*$, $K_1 = g^!$, $J_2 = \bar{g}_*$, $K_2 = \bar{g}^!$, $H = f^*$, $H' = \bar{f}^*$ and $a = \varepsilon^{-1}$, we obtain a morphism

$$\gamma : \bar{f}^* g^! \rightarrow \bar{g}^! f^*$$

and two commutative diagrams.

$$\begin{array}{ccc} \bar{g}_* \bar{f}^* g^! & \xrightarrow{\gamma} & \bar{g}_* \bar{g}^! f^* \\ \varepsilon^{-1} \downarrow & \boxed{37} & \downarrow \\ f^* g_* g^! & \longrightarrow & f^* \end{array} \qquad \begin{array}{ccc} \bar{f}^* & \longrightarrow & \bar{f}^* g^! g_* \\ \downarrow & \boxed{38} & \downarrow \gamma \\ \bar{g}^! \bar{g}_* \bar{f}^* & \xrightarrow{\varepsilon^{-1}} & \bar{g}^! f^* g_* \end{array}$$

5.2.1. THEOREM. (base change) *Let f and \bar{f} be morphisms in \mathcal{B} and g and \bar{g} be morphisms in \mathcal{B}' fitting in the commutative diagram $\boxed{36}$ such that Assumption $(\mathbf{E}_{f,g})$ is satisfied. Then, the isomorphism of functors $\varepsilon : f^* g_* \rightarrow \bar{g}_* \bar{f}^*$ from $\{f^*, \beta_K\} \{g_*, \zeta_K\}$ to $\{\bar{g}_*, \zeta_{f^* K}\} I_{\gamma_K} \{\bar{f}^*, \beta_{\bar{g}^! K}\}$ is duality preserving.*

This together with Proposition 2.2.4 immediately implies the following.

5.2.2. COROLLARY. *If in the situation of the theorem above $\gamma_K : \bar{f}^* g^! K \rightarrow \bar{g}^! f^* K$ is an isomorphism, then the pull-backs and push-forwards on Witt groups defined in corollaries 4.1.3 and 4.3.4 satisfy $f_W^* g_*^W = \bar{g}_*^W I_{\gamma_K}^W \bar{f}_W^*$.*

PROOF PROOF OF THEOREM 5.2.1. We have to prove that the diagram

$$\begin{array}{ccccc} f^* g_* [A, g^! K] & \xrightarrow{\zeta} & f^* [g_* A, K] & \xrightarrow{\beta} & [f^* g_* A, f^* K] \\ \varepsilon \downarrow & & \boxed{39} & & \uparrow \varepsilon^o \\ \bar{g}_* \bar{f}^* [A, g^! K] & \xrightarrow{\beta} & \bar{g}_* [\bar{f}^* A, \bar{f}^* g^! K] & \xrightarrow{\gamma} & \bar{g}_* [\bar{f}^* A, \bar{g}^! f^* K] & \xrightarrow{\zeta} & [\bar{g}_* \bar{f}^* A, f^* K] \end{array}$$

(corresponding to Diagram \boxed{M} of Definition 2.2.2) is commutative. We first prove the two following lemmas.

5.2.3. LEMMA. *The composition*

$$\bar{g}_* (\bar{f}^* (-) \otimes \bar{f}^* (*)) \xrightarrow{\alpha} \bar{g}_* \bar{f}^* (- \otimes *) \xrightarrow{\varepsilon^{-1}} f^* g_* (- \otimes *)$$

and the composition

$$\bar{f}^* [* , g^! (-)] \xrightarrow{\beta} [\bar{f}^* (*), \bar{f}^* g^! (-)] \xrightarrow{\gamma} [\bar{f}^* (*), \bar{g}^! f^* (-)]$$

are mates in Lemma 1.3.5 when $J_1 = g_ (- \otimes *)$, $K_1 = [* , g^! (-)]$, $J_2 = \bar{g}_* (- \otimes \bar{f}^* (*))$, $K_2 = [\bar{f}^* (*), \bar{g}^! (-)]$, $H = f^*$ and $H' = \bar{f}^*$.*

PROOF. By uniqueness, this follows from the proof of one of the commutative diagrams in Lemma 1.3.5. We choose \boxed{H} which is as follows.

$$\begin{array}{ccccc}
 \bar{g}_*(\bar{f}^*[X, g^!A] \otimes \bar{f}^*X) & \xrightarrow{\beta} & \bar{g}_*([\bar{f}^*X, \bar{f}^*g^!A] \otimes \bar{f}^*X) & \xrightarrow{\gamma} & \bar{g}_*([\bar{f}^*X, \bar{g}^!f^*A] \otimes \bar{f}^*X) \\
 \alpha \downarrow & \boxed{10} & \downarrow & \boxed{mf} & \downarrow \\
 \bar{g}_*\bar{f}^*([X, g^!A] \otimes X) & \xrightarrow{\quad} & \bar{g}_*\bar{f}^*g^!A & \xrightarrow{\gamma} & \bar{g}_*g^!f^*A \\
 \varepsilon^{-1} \downarrow & \boxed{mf} & \varepsilon^{-1} \downarrow & \boxed{37} & \downarrow \\
 f^*g_*([X, g^!A] \otimes X) & \xrightarrow{\quad} & f^*g_*g^!A & \xrightarrow{\quad} & f^*A
 \end{array}$$

■

5.2.4. LEMMA. *The composition*

$$\bar{g}_*(-) \otimes f^*g_*(*) \xrightarrow{id \otimes \varepsilon} \bar{g}_*(-) \otimes \bar{g}_*\bar{f}^*(*) \xrightarrow{\lambda} \bar{g}_*(- \otimes \bar{f}^*(*))$$

and the composition

$$\bar{g}_*[\bar{f}^*(*), \bar{g}^!(-)] \xrightarrow{\zeta} [\bar{g}_*\bar{f}^*(*), -] \xrightarrow{\varepsilon^o} [f^*g_*(*), -]$$

are mates in Lemma 1.3.5 when using the functors $J_1 = \bar{g}_*(- \otimes \bar{f}^*(*))$, $K_1 = [\bar{f}^*(*), \bar{g}^!(-)]$, $J_2 = (- \otimes f^*g_*(*))$, $K_2 = [f^*g_*(*), -]$, $H = Id$ and $H' = \bar{g}_*$.

PROOF. Similarly to the proof of the previous lemma, this follows from the commutative diagram


$$\begin{array}{ccccc}
 \bar{g}_*[\bar{f}^*X, \bar{g}^!A] \otimes f^*g_*X & \xrightarrow{\zeta \otimes id} & [\bar{g}_*\bar{f}^*X, A] \otimes f^*g_*X & \xrightarrow{\varepsilon^o} & [f^*g_*X, A] \otimes f^*g_*X \\
 id \otimes \varepsilon \downarrow & \boxed{mf} & \downarrow & \boxed{gen} & \downarrow \\
 \bar{g}_*[\bar{f}^*X, \bar{g}^!A] \otimes \bar{g}_*\bar{f}^*X & \xrightarrow{\zeta \otimes id} & [\bar{g}_*\bar{f}^*X, A] \otimes \bar{g}_*\bar{f}^*X & & \\
 \lambda \downarrow & & \boxed{14''} & \searrow & \\
 \bar{g}_*([\bar{f}^*X, \bar{g}^!A] \otimes \bar{f}^*X) & \xrightarrow{\quad} & \bar{g}_*\bar{g}^!A & \xrightarrow{\quad} & A
 \end{array}$$

where $\boxed{14''}$ is easily obtained from $\boxed{14'}$ by looking at the definition of ζ in Theorem 4.2.9. ■


The commutative diagram $\boxed{31}$ easily implies (by definition of ξ) that the diagram

$$\begin{array}{ccccc}
 \bar{g}^*(f^*A \otimes f^*B) & \xrightarrow{\alpha^{-1}} & \bar{g}^*f^*A \otimes \bar{g}^*f^*B & \xrightarrow{\xi \otimes \xi} & \bar{f}^*g^*A \otimes \bar{f}^*g^*B \\
 \alpha \downarrow & & & & \downarrow \alpha \\
 \bar{g}^*f^*(A \otimes B) & \xrightarrow{\xi} & \bar{f}^*g^*(A \otimes B) & \xrightarrow{\alpha^{-1}} & \bar{f}^*(g^*A \otimes g^*B)
 \end{array}$$


is commutative for any A and B . It can be written as the commutative cube



out of which Lemma 1.2.7 gives the commutative cube



The commutativity of the last cube implies the commutativity of the following cube for any A .



where x is the first composition in Lemma 5.2.3 and y is the first composition in Lemma 5.2.4. Applying Lemma 1.2.7 to this new cube gives a cube in which the mates of x and y may be described using lemmas 5.2.3 and 5.2.4, and whose commutativity is the one of Diagram [39]. ■

Let us now establish two more commutative diagrams that will be useful in applications. In geometric situations they allow us to check whether θ is an isomorphism by restricting to open subsets.

5.2.5. PROPOSITION. *The diagram*

$$\begin{array}{ccccc}
 f^*g_*(-) \otimes f^*(*) & \xrightarrow{\varepsilon} & \bar{g}_*\bar{f}^*(-) \otimes f^*(*) & \xrightarrow{\pi} & \bar{g}_*(\bar{f}^*(-) \otimes \bar{g}^*f^*(*)) \\
 \downarrow \alpha & & \boxed{40} & & \downarrow \xi \\
 & & & & \bar{g}_*(\bar{f}^*(-) \otimes \bar{f}^*g^*(*)) \\
 & & & & \downarrow \alpha \\
 f^*(g_*(-) \otimes *) & \xrightarrow{\pi} & f^*g_*(- \otimes g^*(*)) & \xrightarrow{\varepsilon} & \bar{g}_*\bar{f}^*(- \otimes g^*(*))
 \end{array}$$

is commutative. If furthermore ε is an isomorphism (Assumption $(\mathbf{E}_{f,g})$), γ is defined and the diagram

$$\begin{array}{ccccc}
 \bar{f}^*g^!(-) \otimes \bar{f}^*g^*(*) & \xrightarrow{\gamma \otimes \xi^{-1}} & \bar{g}^!f^*(-) \otimes \bar{g}^*f^*(*) & \xrightarrow{\theta} & \bar{g}^!(f^*(-) \otimes f^*(*)) \\
 \alpha \downarrow & & \boxed{41} & & \downarrow \alpha \\
 \bar{f}^*(g^!(-) \otimes g^*(*)) & \xrightarrow{\theta} & \bar{f}^*g^!(- \otimes *) & \xrightarrow{\gamma} & \bar{g}^!f^*(- \otimes *)
 \end{array}$$

is commutative.

PROOF. To get the first diagram, apply Lemma 1.2.7 to the cube

which is commutative by [31] (using $f\bar{g} = g\bar{f}$). This gives a cube whose commutativity is the one of [40]. Exchanging the top and bottom faces of this new cube (thus reversing the vertical arrows), inverting the front (ε), back ($\varepsilon \times Id$) and sides (π) morphisms of functors and applying again Lemma 1.2.7 to this new cube shows that [41] is commutative too. ■

5.3. ASSOCIATIVITY OF PRODUCTS. We now establish a few commutative diagrams implied by the compatibility of f^* with the associativity of the tensor product. For simplicity, we hide in diagrams all associativity morphisms and bracketing concerning the tensor product (as if the tensor product was strictly associative).

Since f^* is monoidal, the diagram (involving α)

$$\begin{array}{ccc} f^*A \otimes f^*B \otimes f^*C & \longrightarrow & f^*(A \otimes B) \otimes f^*C \\ \downarrow & \boxed{42} & \downarrow \\ f^*A \otimes f^*(B \otimes C) & \longrightarrow & f^*(A \otimes B \otimes C) \end{array}$$

is commutative.

5.3.1. LEMMA. *Under assumptions (\mathbf{A}_f) and (\mathbf{B}_f) , the diagram (involving λ)*

$$\begin{array}{ccc} f_*A \otimes f_*B \otimes f_*C & \longrightarrow & f_*(A \otimes B) \otimes f_*C \\ \downarrow & \boxed{43} & \downarrow \\ f_*A \otimes f_*(B \otimes C) & \longrightarrow & f_*(A \otimes B \otimes C) \end{array}$$

is commutative

PROOF. Left to the reader. ■

5.3.2. PROPOSITION. *Under assumptions (\mathbf{A}_f) and (\mathbf{B}_f) , the following diagrams are commutative.*

$$\begin{array}{ccc} f_*A \otimes B \otimes C & \xrightarrow{\pi} & f_*(A \otimes f^*B) \otimes C \\ \pi \downarrow & \boxed{44} & \downarrow \pi \\ f_*(A \otimes f^*(B \otimes C)) & \xrightarrow{\alpha^{-1}} & f_*(A \otimes f^*B \otimes f^*C) \\ \\ f_*A \otimes f_*B \otimes C & \xrightarrow{\lambda \otimes id} & f_*(A \otimes B) \otimes C \\ id \otimes \pi \downarrow & \boxed{45} & \downarrow \pi \\ f_*(A \otimes f_*(B \otimes f^*C)) & \xrightarrow{\lambda} & f_*(A \otimes B \otimes f^*C) \end{array}$$

PROOF. Using Lemma 4.2.6 to decompose π , Diagram 44 is

$$\begin{array}{ccccc} f_*A \otimes B \otimes C & \xrightarrow{\eta^*} & f_*A \otimes f_*f^*B \otimes C & \xrightarrow{\lambda} & f_*(A \otimes f^*B) \otimes C \\ \eta^* \downarrow & & \eta^* \downarrow & \boxed{mf} & \downarrow \eta^* \\ \boxed{11} & & f_*A \otimes f_*f^*B \otimes f_*f^*C & \xrightarrow{\lambda} & f_*(A \otimes f^*B) \otimes f_*f^*C \\ & & \lambda \downarrow & \boxed{43} & \downarrow \lambda \\ f_*A \otimes f_*f^*(B \otimes C) & \xrightarrow{\alpha^{-1}} & f_*A \otimes f_*(f^*B \otimes f^*C) & \xrightarrow{\lambda} & f_*(A \otimes f^*B \otimes f^*C) \\ & \searrow \lambda & \boxed{mf} & \nearrow \alpha^{-1} & \\ & & f_*(A \otimes f^*(B \otimes C)) & & \end{array}$$

and Diagram 45 is

$$\begin{array}{ccccc} f_*A \otimes f_*B \otimes C & \xrightarrow{\eta^*} & f_*A \otimes f_*B \otimes f_*f^*C & \xrightarrow{\lambda} & f_*A \otimes f_*(B \otimes f^*C) \\ \lambda \downarrow & & \downarrow \lambda & \boxed{43} & \downarrow \lambda \\ f_*(A \otimes B) \otimes C & \xrightarrow{\eta^*} & f_*(A \otimes B) \otimes f_*f^*C & \xrightarrow{\lambda} & f_*(A \otimes B \otimes f^*C) \end{array}$$

■

5.4. THE MONOIDAL FUNCTOR f^* AND PRODUCTS. We have the following diagram of duality preserving functors

$$\begin{array}{ccc} (\mathcal{C}_1)_K \times (\mathcal{C}_1)_M & \xrightarrow{\{-\otimes^*, \tau_{K,M}\}} & (\mathcal{C}_1)_{K \otimes M} \\ \{f^* \times f^*, \beta_K \times \beta_M\} \downarrow & \{ -\otimes^*, \tau_{f^*K, f^*M} \} & \downarrow \{f^*, \beta_{K \otimes M}\} \\ (\mathcal{C}_2)_{f^*K} \times (\mathcal{C}_2)_{f^*M} & \xrightarrow{\quad} (\mathcal{C}_2)_{f^*K \otimes f^*M} \xrightarrow{I_{\alpha_{K,M}}} & (\mathcal{C}_2)_{f^*(K \otimes M)} \end{array}$$

where $I_\alpha : (\mathcal{C}_2)_{f^*K \otimes f^*M} \rightarrow (\mathcal{C}_2)_{f^*(K \otimes M)}$ is the duality preserving functor induced by α by Lemma 3.2.6).

5.4.1. PROPOSITION. (the pull-back respects the product) Under Assumption (\mathbf{A}_f) , the isomorphism of suspended bifunctors $\alpha : f^*(-) \otimes f^*(*) \rightarrow f^*(- \otimes *)$ is an isomorphism of duality preserving functors between the two duality preserving functors defined by the compositions above.

5.4.2. COROLLARY. The pull-back and the product defined in Corollary 4.1.3 and Proposition 4.4.6 satisfy

$$f_W^*(x.y) = I_{\alpha_{K,M}}^W(f_W^*(x).f_W^*(y))$$

for all $x \in W((\mathcal{C}_1)_K)$ and $y \in W((\mathcal{C}_1)_M)$ by Proposition 2.2.4 provided all necessary assumptions about being strong are satisfied.

PROOF PROOF OF PROPOSITION 5.4.1. We start with a lemma.

5.4.3. LEMMA. The composition

$$f^*(-) \otimes (f^*A \otimes f^*B) \xrightarrow{id \otimes \alpha} f^*(-) \otimes f^*(A \otimes B) \xrightarrow{\alpha} f^*(- \otimes (A \otimes B))$$

and the composition

$$f^*[A \otimes B, -] \xrightarrow{\beta} [f^*(A \otimes B), f^*(-)] \xrightarrow{\alpha^\circ} [f^*A \otimes f^*B, -]$$

are mates in Lemma 1.2.6 when $J_1 = - \otimes (A \otimes B)$, $K_1 = [A \otimes B, -]$, $J_2 = - \otimes (f^*A \otimes f^*B)$, $K_2 = [f^*A \otimes f^*B, -]$, $H = f^*$ and $H' = f^*$.

PROOF. By uniqueness, it suffices to establish the commutativity of diagram \boxed{H} which is

$$\begin{array}{ccc} f^*[A \otimes B, -] \otimes (f^*A \otimes f^*B) & \xrightarrow{\beta \otimes id} & [f^*(A \otimes B), f^*(-)] \otimes (f^*A \otimes f^*B) \\ \downarrow id \otimes \alpha & \boxed{mf} & \downarrow id \otimes \alpha \\ f^*[A \otimes B, -] \otimes f^*(A \otimes B) & \longrightarrow & [f^*(A \otimes B), f^*(-)] \otimes f^*(A \otimes B) \\ \downarrow \alpha & \boxed{10} & \downarrow \alpha \\ f^*([A \otimes B, -] \otimes (A \otimes B)) & \xrightarrow{coev^l} & [f^*A \otimes f^*B, f^*(-)] \otimes (f^*A \otimes f^*B) \\ \downarrow coev^l & & \downarrow coev^l \\ f^*(-) & \xleftarrow{coev^l} & f^*(-) \end{array} \quad \begin{array}{c} \curvearrowright \\ \alpha^\circ \\ \curvearrowleft \end{array}$$

■

Now, according to Definition 2.2.2, we need to prove that the following diagram is commutative for any A, B, K and M .

$$\begin{array}{ccc}
f^*[A, K] \otimes f^*[B, M] \xrightarrow{\beta \otimes \beta} [f^*A, f^*K] \otimes [f^*B, f^*M] \xrightarrow{\tau} [f^*A \otimes f^*B, f^*K \otimes f^*M] & & \\
\downarrow \alpha & \boxed{46} & \downarrow \alpha \\
f^*([A, K] \otimes [B, M]) \xrightarrow{\tau} f^*[A \otimes B, K \otimes M] \xrightarrow{\beta} [f^*(A \otimes B), f^*(K \otimes M)] & & [f^*A \otimes f^*B, f^*(K \otimes M)] \\
& & \uparrow \alpha \circ
\end{array}$$

We apply Lemma 1.2.7 to the cube

$$\begin{array}{ccc}
\begin{array}{ccc}
\mathcal{C}_1 \times \mathcal{C}_1 & \xrightarrow{-\otimes -} & \mathcal{C}_1 \\
\downarrow \alpha \times \alpha & \searrow f^* \times f^* & \downarrow f^* \\
\mathcal{C}_2 \times \mathcal{C}_2 & \xrightarrow{-\otimes -} & \mathcal{C}_2 \\
\downarrow \alpha \times \alpha & \searrow f^* \times f^* & \downarrow f^* \\
\mathcal{C}_1 \times \mathcal{C}_1 & \xrightarrow{-\otimes -} & \mathcal{C}_1 \\
\downarrow \alpha \times \alpha & \searrow f^* \times f^* & \downarrow f^* \\
\mathcal{C}_2 \times \mathcal{C}_2 & \xrightarrow{-\otimes -} & \mathcal{C}_2
\end{array} & &
\begin{array}{ccc}
\mathcal{C}_1 \times \mathcal{C}_1 & \xrightarrow{-\otimes -} & \mathcal{C}_1 \\
\downarrow \alpha \times \alpha & \searrow f^* \times f^* & \downarrow f^* \\
\mathcal{C}_2 \times \mathcal{C}_2 & \xrightarrow{-\otimes -} & \mathcal{C}_2 \\
\downarrow \alpha \times \alpha & \searrow f^* \times f^* & \downarrow f^* \\
\mathcal{C}_1 \times \mathcal{C}_1 & \xrightarrow{-\otimes -} & \mathcal{C}_1 \\
\downarrow \alpha \times \alpha & \searrow f^* \times f^* & \downarrow f^* \\
\mathcal{C}_2 \times \mathcal{C}_2 & \xrightarrow{-\otimes -} & \mathcal{C}_2
\end{array}
\end{array}$$

which is easily shown to be commutative using the compatibility of f^* with the symmetry and the associativity of the tensor product (diagrams [15] and [42]). This yields a new cube whose commutativity is the one of [46] using Lemma 5.4.3 to recognize the right face. ■

5.5. PROJECTION FORMULA. The last theorem we want to prove is a projection formula.

5.5.1. THEOREM. (projection formula) Let \mathcal{C}_1 and \mathcal{C}_2 be closed categories and let f^* be a monoidal functor (Assumption (\mathbf{A}_f)) from \mathcal{C}_1 to \mathcal{C}_2 satisfying Assumptions (\mathbf{B}_f) , (\mathbf{C}_f) and (\mathbf{D}_f) . Let K and M be objects of \mathcal{C}_1 . Then, π is a morphism of duality preserving functors from

$$\{Id \otimes Id, \tau_{K,M}\} \{f_* \times Id, \zeta_K \times id\}$$

to

$$\{f_*, \zeta_{K \otimes M}\} I_{\theta_{K,M}} \{Id \otimes Id, \tau_{f^!K, f^*M}\} \{Id \times f^*, id \times \beta_M\}.$$

5.5.2. COROLLARY. (projection formula for Witt groups) By Proposition 2.2.4, under strongness assumptions necessary for their existence and when furthermore the morphism $\theta_{K,M} : f^!K \otimes f^*M \rightarrow f^!(K \otimes M)$ is an isomorphism, the pull-back, push-forward and product on Witt groups defined in corollaries 4.1.3, 4.3.4 and Proposition 4.4.6 satisfy the equality

$$f_*^W(I_{\theta_{K,M}}^W(x \cdot f_W^*(y))) = f_*^W(x) \cdot y$$

in $W((\mathcal{C}_1)_{K \otimes M})$ for all $x \in W((\mathcal{C}_2)_{f^!K})$ and $y \in W((\mathcal{C}_1)_M)$.

PROOF OF THEOREM 5.5.1. According to Definition 2.2.2, we have to prove that the following diagram is commutative for any A, B, K and M .

$$\begin{array}{ccccc}
 f_*[A, f^!K] \otimes [B, M] & \xrightarrow{\zeta} & [f_*A, K] \otimes [B, M] & \xrightarrow{\tau} & [f_*A \otimes B, K \otimes M] \\
 \pi \downarrow & & \boxed{47} & & \uparrow \pi^o \\
 f_*([A, f^!K] \otimes f^*[B, M]) & & & & [f_*(A \otimes f^*B), K \otimes M] \\
 \beta \downarrow & & & & \uparrow \zeta \\
 f_*([A, f^!K] \otimes [f^*B, f^*M]) & \xrightarrow{\tau} & f_*[A \otimes f^*B, f^!K \otimes f^*M] & \xrightarrow{\theta} & f_*[A \otimes f^*B, f^!(K \otimes M)]
 \end{array}$$

Decomposing ζ as in its definition (see Theorem 4.2.9) and using a few \boxed{mf} diagrams and Diagram $\boxed{27}$, it is easy to reduce this to the commutativity of the following diagram (with $N = f^!K$).

$$\begin{array}{ccccc}
 f_*[A, N] \otimes [B, M] & \xrightarrow{\mu} & [f_*A, f_*N] \otimes [B, M] & \xrightarrow{\tau} & [f_*A \otimes B, f_*N \otimes M] \\
 \pi \downarrow & & \boxed{48} & & \uparrow \pi^o \\
 f_*([A, N] \otimes f^*[B, M]) & & & & [f_*(A \otimes f^*B), f_*N \otimes M] \\
 \beta \downarrow & & & & \uparrow \pi^{-1} \\
 f_*([A, N] \otimes [f^*B, f^*M]) & \xrightarrow{\tau} & f_*[A \otimes f^*B, N \otimes f^*M] & \xrightarrow{\mu} & [f_*(A \otimes f^*B), f_*(N \otimes f^*M)]
 \end{array}$$

which requires the following preliminary lemmas.

5.5.3. LEMMA. *The composition*

$$\begin{array}{ccc}
 f_*(-_1 \otimes -_2) \otimes f_*(A \otimes f^*B) & \xrightarrow{\lambda} & f_*((-_1 \otimes -_2) \otimes (A \otimes f^*B)) \\
 & & \downarrow e \\
 & & f_*((-_1 \otimes A) \otimes (-_2 \otimes f^*B))
 \end{array}$$

and the composition

$$f_*([A, -_1] \otimes [f^*B, -_2]) \xrightarrow{\tau} f_*[A \otimes f^*B, -_1 \otimes -_2] \xrightarrow{\mu} [f_*(A \otimes f^*B), f_*(-_1 \otimes -_2)]$$

are mates in Lemma 1.2.6 when $J_1 = (-_1 \otimes A) \times (-_2 \otimes f^*B)$, $K_1 = [A, -_1] \times [f^*B, -_2]$, $J_2 = - \otimes f_*(A \otimes f^*B)$, $K_2 = [f_*(A \otimes f^*B), -]$, and $H = H' = f_*(-_1 \otimes -_2)$.

PROOF. By uniqueness, it suffices to prove Diagram \boxed{H} which is here

$$\begin{array}{ccc}
 f_*([A, -_1] \otimes [f^*B, -_2]) \otimes f_*(A \otimes f^*B) & \xrightarrow{\tau} & f_*[A \otimes f^*B, -_1 \otimes -_2] \otimes f_*(A \otimes f^*B) \\
 \lambda \downarrow & \boxed{mf} & \lambda \downarrow \\
 f_*([A, -_1] \otimes [f^*B, -_2]) \otimes (A \otimes f^*B) & \xrightarrow{\tau} & f_*([A \otimes f^*B, -_1 \otimes -_2] \otimes (A \otimes f^*B)) \xrightarrow{\mu} \\
 e \downarrow & \boxed{28} & \uparrow \text{ev}^l \quad \boxed{14} \\
 f_*([A, -_1] \otimes A) \otimes ([f^*B, -_2] \otimes f^*B) & & [f_*(A \otimes f^*B), f_*(-_1 \otimes -_2)] \otimes f_*(A \otimes f^*B) \\
 \text{ev}^l \downarrow & & \swarrow \text{ev}^l \\
 f_*(-_1 \otimes -_2) & &
 \end{array}$$

■

5.5.4. LEMMA. *The composition*

$$(-_1 \otimes -_2) \otimes f_*(A \otimes f^*B) \xrightarrow{\pi^{-1}} (-_1 \otimes -_2) \otimes (f_*A \otimes B) \xrightarrow{e} (-_1 \otimes f_*A) \otimes (-_2 \otimes B)$$

and the composition

$$[f_*A, -_1] \otimes [B, -_2] \xrightarrow{\tau} [f_*A \otimes B, -_1 \otimes -_2] \xrightarrow{(\pi^{-1})^\circ} [f_*(A \otimes f^*B), -_1 \otimes -_2]$$

are mates in Lemma 1.2.6 when $J_1 = (-_1 \otimes f_*A) \times (-_2 \otimes B)$, $K_1 = [f_*A, -_1] \times [B, -_2]$, $J_2 = - \otimes f_*(A \otimes f^*B)$, $K_2 = [f_*(A \otimes f^*B), -]$, and $H = H' = \otimes$.

PROOF. By uniqueness, it suffices to prove that diagram \boxed{H} commutes, which is

$$\begin{array}{ccc}
 ([f_*A, -_1] \otimes [B, -_2]) \otimes f_*(A \otimes f^*B) & \xrightarrow{\tau} & [f_*A \otimes B, -_1 \otimes -_2] \otimes f_*(A \otimes f^*B) \\
 \pi^{-1} \downarrow & \boxed{mf} & \pi^{-1} \downarrow \\
 ([f_*A, -_1] \otimes [B, -_2]) \otimes (f_*A \otimes B) & \xrightarrow{\tau} & [f_*A \otimes B, -_1 \otimes -_2] \otimes (f_*A \otimes B) & (\pi^{-1})^\circ \\
 e \downarrow & \boxed{28} & \swarrow \text{ev}^l & \boxed{gen} \\
 ([f_*A, -_1] \otimes f_*A) \otimes ([B, -_2] \otimes B) & & [f_*(A \otimes f^*B), -_1 \otimes -_2] \otimes f_*(A \otimes f^*B) \\
 \text{ev}^l \downarrow & & \swarrow \text{ev}^l \\
 (-_1 \otimes -_2) & & (-_1 \otimes -_2)
 \end{array}$$


■

The cube

The diagram shows a cube with nodes $C_1 \times C_1$, $C_1 \times C_2$, $C_2 \times C_2$, C_1 , C_2 , and C_2 . The front face (left) has nodes $C_1 \times C_1$, $C_1 \times C_2$, $C_2 \times C_2$, and C_1 . The back face (right) has nodes $C_1 \times C_1$, $C_1 \times C_2$, $C_2 \times C_2$, and C_1 . The top face has nodes $C_2 \times C_2$, C_2 , and C_2 . Morphisms include \otimes , $f^* \times f^*$, f^* , e , α^{-1} , $(-\otimes M) \times (-\otimes B)$, $(-\otimes f^*M) \times (-\otimes f^*B)$, $(-\otimes f^*M \otimes f^*B)$, $(-\otimes M \otimes B)$, $(id \otimes \alpha^{-1}) \circ \alpha^{-1}$, and f^* .

is commutative by a classical exercise on monoidal functors, symmetry and associativity. Applying Lemma 1.2.7 to it, we get a new cube, in which the morphism of functors on the right face is $(id \otimes \alpha^{-1}) \circ \pi$, simply because π is the mate of α^{-1} (by Proposition 4.2.5) and $(id \otimes \alpha^{-1})$ is just a change in the parameter. The commutativity of the cube thus

obtained can be rewritten as the commutativity of the cube



where x and y are the first compositions in lemmas 5.5.3 and 5.5.4. Applying one more time Lemma 1.2.7 to this new cube yields another one whose commutativity is the one of [48]. ■

6. Various reformulations

This section is devoted to reformulations of the main theorems in view of applications. The corollaries on Witt groups show that these reformulations are useful. The reformulations are obtained by changing the duality at the source or at the target using functors I_ι (see Lemma 3.2.6) with ι adapted to the situation. For this reason, deriving those reformulations from the original theorems is very easy and most of the times follows from simple commutative diagrams involving functors of the form I_ι . We therefore leave all proofs of Sections 6.1, 6.2 and 6.3 as exercises for the reader.

6.1. REFORMULATIONS USING $f^!$ OF UNIT OBJECTS. In this section, we use the unit objects in the monoidal categories to formulate the main theorems in a different way. In the application to Witt groups and algebraic geometry, this will relate $f^!$ to f^* using canonical sheaves. Let \mathcal{B}'' denote the subcategory of \mathcal{B}' in which the morphisms f are such that (\mathbf{D}_f) is satisfied (π is an isomorphism).

For any morphism $f : X \rightarrow Y$ in \mathcal{B}' , we define

$$\omega'_f = f^!(\mathbf{1}_Y).$$

To every morphism f , we associate a number d_f such that $d_{gf} = d_f + d_g$ for composable morphisms f and g . For example, for every object X we may choose a number d_X and set $d_f = d_X - d_Y$. We define

$$\omega_f = T^{-d_f} f^!(\mathbf{1}_Y).$$

6.1.1. REMARK. In applications to schemes, d_f can be the relative dimension of a morphism f and d_X can be the relative dimension of a smooth X over a base scheme. In that case, ω'_f is isomorphic to a shifted line bundle and ω_f to a line bundle (the canonical

sheaf). This explains why we introduce d_f and ω_f . On the other hand, it is always possible to set $d_f = 0$ for any morphism f , in which case $\omega_f = \omega'_f$, thus statements in terms of ω_f also apply to ω'_f .

For any two composable morphisms f and g in \mathcal{B}'' , let us denote by $i'_{g,f}$ the composition

$$\omega'_f \otimes f^*(\omega'_g) = f^!(\mathbf{1}_Y) \otimes f^*(\omega'_g) \xrightarrow{\theta} f^!(\omega'_g) = f^!g^!(\mathbf{1}_Z) \xrightarrow{\zeta} (gf)^!(\mathbf{1}_Z) = \omega'_{gf}$$

and

$$i_{g,f} : \omega_f \otimes f^*(\omega_g) \longrightarrow \omega_{gf}$$

the composition obtained by the same chain of morphisms and then desuspending.

6.1.2. LEMMA. For any composable morphisms $X \xrightarrow{f} Y \xrightarrow{g} Z \xrightarrow{h} V$ in \mathcal{B}'' , the diagram of isomorphisms

$$\begin{array}{ccc} \omega_f \otimes f^*(\omega_{hg}) & \xleftarrow{id \otimes f^* i_{h,g}} & \omega_f \otimes f^*(\omega_g \otimes g^*(\omega_h)) \\ \omega_{hgf} \xleftarrow{i_{hg,f}} & & \downarrow (id \otimes a) \circ (id \otimes \alpha^{-1}) \\ \omega_{gf} \otimes (gf)^*(\omega_h) & \xleftarrow{i_{g,f} \otimes id} & \omega_f \otimes f^*(\omega_g) \otimes (gf)^*(\omega_h) \end{array}$$

is commutative. In other words, $i_{g,f}$ (as well as $i'_{g,f}$) satisfies a cocycle condition.

We can then reformulate the main theorems 4.2.9, 5.1.9, 5.2.1 and 5.5.1 and their corollaries for Witt groups as follows.

6.1.3. DEFINITION. (push-forward) Let $f : X \rightarrow Y$ be a morphism in \mathcal{B}'' and K be an object in \mathcal{C}_Y . We define a duality preserving functor $\mathcal{C}_{X, \omega'_f \otimes f^* K} \rightarrow \mathcal{C}_{Y, K}$ by the composition $\{f_*, \zeta_K\} I_{\theta_{1,K}}$. We denote it simply by $\{f_*\}$ in the rest of the section (K is always understood).

6.1.4. DEFINITION. (push-forward for Witt groups) When the dualities are strong and $\theta_{1,K}$ is an isomorphism, $\{f_*\}$ induces a push-forward

$$f_*^W : W^{i+d_f}(X, \omega_f \otimes f^* K) \rightarrow W^i(Y, K)$$

on Witt groups (recall Proposition 3.2.5 to switch from suspending ω'_f to the suspended duality used in Definition 2.1.5 of shifted Witt groups).

6.1.5. THEOREM. (composition of push-forwards) Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be morphisms in \mathcal{B}'' and K be an object of \mathcal{C}_Z . The morphism $b_{g,f} : (gf)_* \rightarrow g_* f_*$ (see Section 5.1) is duality preserving between the compositions of duality preserving functors in the diagram

$$\begin{array}{ccc} \mathcal{C}_{X, \omega'_f \otimes f^*(\omega'_g \otimes g^* K)} & \xrightarrow{I_\iota} & \mathcal{C}_{X, \omega'_{gf} \otimes (gf)^* K} \\ \{f_*\} \downarrow & \swarrow b_{g,f} & \downarrow \{(gf)_*\} \\ \mathcal{C}_{Y, \omega'_g \otimes g^* K} & \xrightarrow{\{g_*\}} & \mathcal{C}_{Z, K} \end{array}$$

where ι is the composition

$$\omega'_f \otimes f^*(\omega'_g \otimes g^* K) \xrightarrow{Id \otimes \alpha^{-1}} \omega'_f \otimes f^* \omega'_g \otimes f^* g^* K \xrightarrow{i'_{g,f} \otimes a_{g,f}} \omega'_{gf} \otimes (gf)^* K.$$

6.1.6. **COROLLARY.** (*composition of push-forwards for Witt groups*) For f and g as in the theorem, the push-forwards on Witt groups of Definition 6.1.4 satisfy

$$g_*^W f_*^W = (gf)_*^W I_\iota^W.$$

6.1.7. **THEOREM.** (*base change*) In the situation of Theorem 5.2.1, with furthermore g and \bar{g} in \mathcal{B}'' , the morphism of functors $\varepsilon : g_* f^* \rightarrow \bar{f}^* \bar{g}_*$ is duality preserving between the two compositions of duality preserving functors in the diagram

$$\begin{array}{ccccc} & & \mathcal{C}_{V, \bar{f}^*(\omega'_g \otimes g^* K)} & \xrightarrow{I_\iota} & \mathcal{C}_{V, \omega'_g \otimes \bar{g}^* f^* K} & & \\ & \nearrow \{\bar{f}^*\} & & \uparrow \varepsilon & & \searrow \{\bar{g}_*\} & \\ \mathcal{C}_{X, \omega'_g \otimes g^* K} & \xrightarrow{\{g_*\}} & \mathcal{C}_{Z, K} & \xrightarrow{\{f_*\}} & \mathcal{C}_{Y, f^* K} & & \end{array}$$

where ι is defined as the composition

$$\begin{array}{ccc} \bar{f}^*(\omega'_g \otimes g^* K) & \xrightarrow{\alpha^{-1}} & \bar{f}^* \omega'_g \otimes \bar{f}^* g^* K \\ & \text{Id} \otimes \xi \downarrow & \\ \bar{f}^* \omega'_g \otimes \bar{g}^* f^* K & \xrightarrow{\gamma_1 \otimes \text{Id}} & \bar{g}^! f^* \mathbf{1} \otimes \bar{g}^* f^* K \simeq \omega'_g \otimes \bar{g}^* f^* K. \end{array}$$

6.1.8. **COROLLARY.** (*base change for Witt groups*) In the situation of the theorem, when furthermore $\gamma_1 : \bar{f}^* g^! \mathbf{1} \rightarrow \bar{g}^! f^* \mathbf{1}$ is an isomorphism, the push-forwards and pull-backs for Witt groups of Definition 6.1.4 and Corollary 4.1.3 satisfy

$$f_W^* g_*^W = \bar{g}_*^W I_\iota^W \bar{f}_W^*.$$

6.1.9. **THEOREM.** (*projection formula*) In the situation of Theorem 5.5.1, π is a morphism of duality preserving functors between the two compositions in the diagram

$$\begin{array}{ccccc} \mathcal{C}_{\omega'_f \otimes f^* K} \times \mathcal{C}_{f^* M} & \xrightarrow{\{\otimes\}} & \mathcal{C}_{\omega'_f \otimes f^* K \otimes f^* M} & \xrightarrow{I_{\text{Id} \otimes \alpha_{K, M}}} & \mathcal{C}_{\omega'_f \otimes f^*(K \otimes M)} \\ \{Id \times f^*\} \uparrow & & \pi \uparrow \uparrow & & \downarrow \{f_*\} \\ \mathcal{C}_{\omega'_f \otimes f^* K} \times \mathcal{C}_M & \xrightarrow{\{f_* \times Id\}} & \mathcal{C}_K \times \mathcal{C}_M & \xrightarrow{\{\otimes\}} & \mathcal{C}_{K \otimes M} \end{array}$$

6.1.10. **COROLLARY.** (*projection formula for Witt groups*) In the situation of the theorem, the push-forward, pull-back and product on Witt groups of Definition 6.1.4, Corollary 4.1.3 and Proposition 4.4.6 satisfy

$$f_*^W I_{\text{Id} \otimes \alpha_{K, M}}^W (x \cdot f_W^*(y)) = f_*^W (x) \cdot y$$

in $W^{i+j}(Y, K \otimes M)$ for any $x \in W^{i+d_f}(X, \omega_f \otimes f^* K)$ and $y \in W^j(Y, M)$.

6.2. **REFORMULATIONS USING MORPHISMS TO CORRECT THE DUALITIES.** We now reformulate the main results using a convenient categorical setting which allows us to display the previous results in an even nicer way. This will be useful in applications.

We define new categories.

6.2.1. **DEFINITION.** Let \mathcal{B}^* be the category whose objects are pairs (X, K) with $X \in \mathcal{B}$, $K \in \mathcal{C}_X$, and whose morphisms from (X, K) to (Y, L) are pairs (f, ϕ) where $f : X \rightarrow Y$ is a morphism in \mathcal{B} and $\phi : f^*L \rightarrow K$ is a morphism in \mathcal{C}_X . The composition is defined by $(g, \psi)(f, \phi) = (gf, \phi \circ f^*(\psi) \circ (a_{g,f})^{-1})$. The identity morphism on (X, K) is (Id_X, Id_K) and the composition is clearly associative.

6.2.2. **DEFINITION.** Let $\mathcal{B}^!$ be the category whose objects are pairs (X, K) with $X \in \mathcal{B}^!$, $K \in \mathcal{C}_X$, and whose morphisms from (X, K) to (Y, M) are pairs (f, ϕ) where $f : X \rightarrow Y$ is a morphism in $\mathcal{B}^!$ and $\phi : K \rightarrow f^!M$ is a morphism in \mathcal{C}_X . The composition is defined by $(g, \psi)(f, \phi) = (gf, c_{g,f} \circ f^!(\psi) \circ \phi)$. The identity morphism on (X, K) is (Id_X, Id_K) and the composition is clearly associative.

For applications to Witt groups, we need the objects defining the dualities to be dualizing and the duality preserving functors to be strong. We thus define two more categories.

6.2.3. **DEFINITION.** Let \mathcal{B}_W^* denote the subcategory of \mathcal{B}^* in which the objects (X, K) are such that K is dualizing (ϖ_K is an isomorphism) and the morphisms (f, ϕ) from (X, K) to (Y, L) are such that ϕ and $\beta_{f,L}$ are isomorphisms.

6.2.4. **DEFINITION.** Let $\mathcal{B}_W^!$ denote the subcategory of $\mathcal{B}^!$ in which the objects (X, K) are such that K is dualizing (ϖ_K is an isomorphism) and the morphisms (f, ϕ) are such that ϕ is an isomorphism.

Using the categories $\mathcal{B}^!$ and \mathcal{B}^* and their subcategories, the main theorems 4.1.2, 4.2.9, 5.1.3, 5.1.9, 5.2.1, 5.5.1 and their corollaries on Witt groups can be rephrased as follows.

6.2.5. **DEFINITION.** (pull-back) For any morphism $(f, \phi) : (X, K) \rightarrow (Y, L)$ in \mathcal{B}^* , we define a duality preserving functor $|f, \phi|^* : \mathcal{C}_{Y,L} \rightarrow \mathcal{C}_{X,K}$ by the composition

$$\mathcal{C}_{Y,L} \xrightarrow{\{f^*, \beta_L\}} \mathcal{C}_{X, f^*L} \xrightarrow{I_\phi} \mathcal{C}_{X,K}.$$

6.2.6. **DEFINITION.** (pull-back on Witt groups) For any morphism (f, ϕ) from (X, K) to (Y, L) in \mathcal{B}_W^* , the duality preserving functor $|f, \phi|$ induces a morphism

$$|f, \phi|_W^* : W^i(Y, L) \rightarrow W^i(X, K)$$

on Witt groups.

6.2.7. **THEOREM.** (composition of pull-backs) Let (f, ϕ) and (g, ψ) be composable morphisms in \mathcal{B}^* . The morphism $a_{g,f} : f^*g^* \rightarrow (gf)^*$ is a morphism of duality preserving functors as in the following diagram.

$$\begin{array}{ccc} & \mathcal{C}_{Y,L} & \\ \begin{array}{c} \nearrow |g, \psi|^* \\ \searrow |f, \phi|^* \end{array} & \Downarrow a_{g,f} & \\ \mathcal{C}_{Z,M} & \xrightarrow{|(g, \psi)(f, \phi)|^*} & \mathcal{C}_{X,K} \end{array}$$

6.2.8. **COROLLARY.** (*composition of pull-backs for Witt groups*) Let (f, ϕ) and (g, ψ) be composable morphisms in \mathcal{B}_W^* . Then the pull-backs for Witt groups of Definition 6.2.6 satisfy $|(g, \psi)(f, \phi)|_W^* = |f, \phi|_W^* |g, \psi|_W^*$. In other words, W^* is a contravariant functor from \mathcal{B}_W^* to graded abelian groups.

6.2.9. **DEFINITION.** (*push-forward*) For any morphism $(f, \phi) : (X, K) \rightarrow (Y, L)$ in $\mathcal{B}^!$, we define a duality preserving functor $|f, \phi|_* : \mathcal{C}_{X, K} \rightarrow \mathcal{C}_{Y, L}$ by the composition

$$\mathcal{C}_{X, K} \xrightarrow{I_\phi} \mathcal{C}_{X, f^!L} \xrightarrow{\{f_*, \zeta_L\}} \mathcal{C}_{Y, L}.$$

6.2.10. **DEFINITION.** (*push-forward for Witt groups*) For any morphism (f, ϕ) from (X, K) to (Y, L) in $\mathcal{B}_W^!$, the duality preserving functor $|f, \phi|_*$ induces a morphism on Witt groups $|f, \phi|_*^W : W^i(X, K) \rightarrow W^i(X, L)$.

6.2.11. **THEOREM.** (*composition of push-forwards*) Let (f, ϕ) and (g, ψ) be composable morphisms in $\mathcal{B}^!$. The morphism $b_{g, f} : (gf)_* \rightarrow g_* f_*$ is a morphism of duality preserving functors as in the following diagram.

$$\begin{array}{ccc} & \mathcal{C}_{Y, L} & \\ |f, \phi|_* \nearrow & \uparrow b_{g, f} & \searrow |g, \psi|_* \\ \mathcal{C}_{X, K} & \xrightarrow{|(g, \psi)(f, \phi)|_*} & \mathcal{C}_{Z, M} \end{array}$$

6.2.12. **COROLLARY.** (*composition of push-forwards for Witt groups*) Let (f, ϕ) and (g, ψ) be composable morphisms in $\mathcal{B}_W^!$. Then the push-forwards for Witt groups of Definition 6.2.10 satisfy $|(g, \psi)(f, \phi)|_*^W = |g, \psi|_*^W |f, \phi|_*^W$. In other words, W^* is a covariant functor from $\mathcal{B}_W^!$ to graded abelian groups.

6.2.13. **THEOREM.** (*base change*) Let (f, ϕ) and $(\bar{f}, \bar{\phi})$ be morphisms in \mathcal{B}^* , (g, ψ) , and $(\bar{g}, \bar{\psi})$ be morphisms in $\mathcal{B}^!$ fitting in the diagram

$$\begin{array}{ccc} (V, N) & \xrightarrow{(\bar{g}, \bar{\psi})} & (Y, L) \\ (\bar{f}, \bar{\phi}) \downarrow & & \downarrow (f, \phi) \\ (X, M) & \xrightarrow{(g, \psi)} & (Z, K) \end{array}$$

such that $f\bar{g} = g\bar{f} \in \mathcal{B}$, such that Assumption $(\mathbf{E}_{f, g})$ is satisfied and such that the diagram

$$\begin{array}{ccccc} & \bar{\phi} & \bar{f}^* M & \xrightarrow{\bar{f}^*(\psi)} & \bar{f}^* g^! K \\ N & \swarrow & & & \downarrow \gamma_K \\ & \bar{\psi} & \bar{g}^! L & \xleftarrow{\bar{g}^!(\phi)} & \bar{g}^! f^* K \end{array}$$

is commutative. Then the morphism $\varepsilon : f^*g_* \rightarrow \bar{g}_*f^*$ is duality preserving between the compositions of duality preserving functors in the diagram.

$$\begin{array}{ccc} \mathcal{C}_{V,N} & \xrightarrow{|\bar{g},\bar{\psi}|_*} & \mathcal{C}_{Y,L} \\ |\bar{f},\bar{\phi}|^* \uparrow & \swarrow \varepsilon & \uparrow |f,\phi|^* \\ \mathcal{C}_{X,M} & \xrightarrow{|g,\psi|_*} & \mathcal{C}_{Z,K} \end{array}$$

6.2.14. **COROLLARY.** (base change for Witt groups) In the situation of the theorem, assuming furthermore that (f, ϕ) and $(\bar{f}, \bar{\phi})$ are in \mathcal{B}_W^* , (g, ψ) and $(\bar{g}, \bar{\psi})$ are in $\mathcal{B}_W^!$ and $\gamma_K : \bar{f}^*g^!K \rightarrow \bar{g}^!f^*K$ is an isomorphism, the pull-backs and push-forwards on Witt groups of definitions 6.2.6 and 6.2.10 satisfy $|\bar{g}, \bar{\psi}|_*^W |\bar{f}, \bar{\phi}|^* = |f, \phi|_*^W |g, \psi|^*{}^W$.

6.2.15. **THEOREM.** (projection formula) Let $f : X \rightarrow Y$ be a morphism in \mathcal{B}'' . Let $K, M \in \mathcal{C}_X$ and $L, N \in \mathcal{C}_Y$ be objects,

$$\phi : f^*L \rightarrow K, \quad \psi : M \rightarrow f^!N \quad \text{and} \quad \chi : M \otimes K \rightarrow f^!(N \otimes L)$$

be morphisms such that

$$\begin{array}{ccc} M \otimes K & \xrightarrow{\chi} & f^!(N \otimes L) \\ Id \otimes \phi \uparrow & & \uparrow \theta_{N,L} \\ M \otimes f^*L & \xrightarrow{\psi \otimes Id} & f^!N \otimes f^*L \end{array}$$

is commutative.

Then, the morphism π is duality preserving between the compositions in the diagram

$$\begin{array}{ccccc} & & \mathcal{C}_{X,M} \times \mathcal{C}_{X,K} & \xrightarrow{\{\otimes\}} & \mathcal{C}_{X,M \otimes K} \\ Id \times |f,\phi|^* \nearrow & & \uparrow \pi & & \searrow |f,\chi|_* \\ \mathcal{C}_{X,M} \times \mathcal{C}_{Y,L} & \xrightarrow{|f,\psi|_* \times Id} & \mathcal{C}_{Y,N} \times \mathcal{C}_{Y,L} & \xrightarrow{\{\otimes\}} & \mathcal{C}_{Y,N \otimes L} \end{array}$$

6.2.16. **COROLLARY.** (projection formula for Witt groups) In the situation of the theorem, assuming furthermore that (f, ϕ) is in \mathcal{B}_W^* and (f, ψ) and (f, χ) are in $\mathcal{B}_W^!$, the push-forward, pull-back and product for Witt groups of Definition 6.2.10, Definition 6.2.6 and Proposition 4.4.6 satisfy $|f, \chi|_*^W (x \cdot |f, \phi|_*^W (y)) = |f, \psi|_*^W (x) \cdot y$ in $W^{i+j}(Y, N \otimes L)$ for any $x \in W^i(X, M)$ and $y \in W^j(Y, L)$.

6.3. **REFORMULATIONS USING A FINAL OBJECT.** When the category \mathcal{B} has a final object denoted by Pt that is also a final object for the subcategory \mathcal{B}'' , there is a reformulation of the main results using absolute (that is relative to the final object) canonical objects rather than relative ones as in section 6.1. For any object $X \in \mathcal{B}'$, let p_X denote the unique morphism $X \rightarrow \text{Pt}$. We define $d_X = d_{p_X}$, $\omega'_X = \omega'_{p_X}$ and $\omega_X = \omega_{p_X}$.

6.3.1. **DEFINITION.** Let \mathcal{B}_* be the category whose objects are pairs (X, K) with $X \in \mathcal{B}''$, $K \in \mathcal{C}_X$, and whose morphisms from (X, K) to (Y, L) are pairs (f, ϕ) where $f : X \rightarrow Y$ is a morphism in \mathcal{B}'' and $\phi : K \rightarrow f^*L$ is a morphism in \mathcal{C}_X . The composition is defined by $(g, \psi)(f, \phi) = (gf, b_{g,f})f^*(\psi)\phi$. The identity of (X, K) is (Id_X, Id_K) and the composition is clearly associative.

6.3.2. DEFINITION. Let \mathcal{B}_*^W be the subcategory of \mathcal{B}_* in which the objects (X, K) are such that $\omega_X \otimes K$ is dualizing and the morphisms $(f, \phi) : (X, K) \rightarrow (Y, L)$ are such that ι defined as the composition

$$\omega'_X \otimes K \xrightarrow{(c_{f,p_Y})^{-1} \otimes \phi} f^! \omega'_Y \otimes f^* L \xrightarrow{\theta_{\omega'_Y, L}} f^!(\omega'_Y \otimes L).$$

is an isomorphism.

6.3.3. DEFINITION. (push-forward) For any morphism $(f, \phi) : (X, K) \rightarrow (Y, L)$ in \mathcal{B}_* , we define a duality preserving functor $|f, \phi|_* : \mathcal{C}_{X, \omega'_X \otimes K} \rightarrow \mathcal{C}_{Y, \omega'_Y \otimes L}$ by the composition

$$\mathcal{C}_{X, \omega'_X \otimes K} \xrightarrow{I_L} \mathcal{C}_{X, f^!(\omega'_Y \otimes L)} \xrightarrow{\{f_*, \zeta_{\omega'_Y \otimes L}\}} \mathcal{C}_{Y, L}.$$

where ι is defined as in Definition 6.3.2 (but not necessarily an isomorphism).

6.3.4. DEFINITION. (push-forward for Witt groups) Let (f, ϕ) be a morphism in \mathcal{B}_*^W . Then the duality preserving functor $|f, \phi|_*$ of Definition 6.3.3 induces a morphism

$$|f, \phi|_*^W : W^{i+d_X}(X, \omega_X \otimes K) \rightarrow W^{i+d_Y}(Y, \omega_Y \otimes L)$$

on Witt groups.

6.3.5. THEOREM. (composition of push-forwards) Let (f, ϕ) and (g, ψ) be composable morphisms in \mathcal{B}_* . Then the morphism $b_{g,f} : (gf)_* \rightarrow g_* f_*$ is a morphism of duality preserving functors as in the following diagram.

$$\begin{array}{ccc} & \mathcal{C}_{Y, \omega'_Y \otimes L} & \\ & \uparrow b_{g,f} & \\ \mathcal{C}_{X, \omega'_X \otimes K} & \xrightarrow{|f, \phi|_*} & \mathcal{C}_{Z, \omega'_Z \otimes M} \\ & \downarrow |(g, \psi)(f, \phi)|_* & \end{array}$$

6.3.6. COROLLARY. (composition of push-forwards for Witt groups) Let (f, ϕ) and (g, ψ) be composable morphisms in \mathcal{B}_*^W . Then the push-forwards of Definition 6.3.4 satisfy $|(g, \psi)(f, \phi)|_*^W = |g, \psi|_*^W |f, \phi|_*^W$. In other words, W^* is a covariant functor from \mathcal{B}_*^W to graded abelian groups.

To state a base change theorem, we need the following. Let (f, ϕ) and $(\bar{f}, \bar{\phi})$ be morphisms in \mathcal{B}^* and (g, ψ) and $(\bar{g}, \bar{\psi})$ be morphisms in \mathcal{B}_* with sources and targets as on the diagram

$$\begin{array}{ccccc} (V, \omega'_V \otimes N) & & (V, N) \xrightarrow{(\bar{g}, \bar{\psi})} & (Y, L) & (Y, \omega'_Y \otimes L) \\ (\bar{f}, \bar{\phi}) \downarrow & & & & \downarrow (f, \phi) \\ (X, \omega'_X \otimes M) & & (X, M) \xrightarrow{(g, \psi)} & (Z, K) & (Z, \omega'_Z \otimes K) \end{array}$$

such that $f\bar{g} = g\bar{f} \in \mathcal{B}$ and such that Assumption $(\mathbf{E}_{f,g})$ is satisfied. The morphism ϕ induces a morphism “ ϕ ” defined by the composition

$$\begin{array}{ccccc} \bar{f}^*(\omega'_X \otimes g^*K) & \xrightarrow{\alpha^{-1}} & \bar{f}^*\omega'_X \otimes \bar{f}^*g^*K & \xrightarrow{\bar{f}^*(c_{p_Z,g})^{-1} \otimes Id} & \bar{f}^*g^!\omega'_Z \otimes \bar{f}^*g^*K \\ \downarrow \text{“}\phi\text{”} & & & & \downarrow \gamma \otimes \xi^{-1} \\ \bar{g}^!(\omega'_Y \otimes L) & \xleftarrow{\bar{g}^!(\phi)} & \bar{g}^!f^*(\omega'_Z \otimes K) & \xleftarrow{\alpha \circ \theta} & \bar{g}^!f^*\omega'_Z \otimes \bar{g}^*f^*K \end{array}$$

The morphism $\bar{\psi}$ induces a morphism “ $\bar{\psi}$ ” defined by the composition

$$\omega'_Y \otimes N \xrightarrow{(c_{p_Y,\bar{g}})^{-1} \otimes \bar{\psi}} \bar{g}^!\omega'_Y \otimes \bar{g}^*L \xrightarrow{\theta} \bar{g}^!(\omega'_Y \otimes L)$$

6.3.7. THEOREM. (*base change*) *Let (f, ϕ) , $(\bar{f}, \bar{\phi})$, (g, ψ) and $(\bar{g}, \bar{\psi})$ be as above. We assume that the diagram*

$$\begin{array}{ccc} \omega'_Y \otimes N & \xrightarrow{\text{“}\bar{\psi}\text{”}} & \bar{g}^!(\omega'_Y \otimes L) \\ \bar{\phi} \uparrow & & \uparrow \text{“}\phi\text{”} \\ \bar{f}^*(\omega'_X \otimes M) & \xrightarrow{\psi} & \bar{f}^*(\omega_X \otimes g^*K) \end{array}$$

*is commutative. Then, $\varepsilon : f^*g_* \rightarrow \bar{g}_*\bar{f}^*$ is a morphism of duality preserving functors as in the following diagram.*

$$\begin{array}{ccc} \mathcal{C}_{V, \omega'_Y \otimes N} & \xrightarrow{|\bar{g}, \bar{\psi}|_*} & \mathcal{C}_{Y, \omega'_Y \otimes L} \\ |\bar{f}, \bar{\phi}|^* \uparrow & \swarrow \varepsilon & \uparrow |f, \phi|^* \\ \mathcal{C}_{X, \omega'_X \otimes M} & \xrightarrow{|\bar{g}, \bar{\psi}|_*} & \mathcal{C}_{Z, \omega'_Z \otimes K} \end{array}$$

6.3.8. COROLLARY. (*base change for Witt groups*) *In the situation of the theorem, assuming furthermore that (f, ϕ) and $(\bar{f}, \bar{\phi})$ are in \mathcal{B}_W^* and (g, ψ) and $(\bar{g}, \bar{\psi})$ are in \mathcal{B}_*^W and $\gamma_{\omega_Z} : \bar{f}^*g^!\omega_Z \rightarrow \bar{g}^!f^*\omega_Z$ is an isomorphism. Then the pull-backs and push-forwards on Witt groups of definitions 6.2.6 and 6.3.4 satisfy*

$$|\bar{g}, \bar{\psi}|_*^W |\bar{f}, \bar{\phi}|_W^* = |f, \phi|_W^* |g, \psi|_*^W.$$

6.3.9. THEOREM. (*projection formula*) *Let $f : X \rightarrow Y$ be a morphism in \mathcal{B}'' . Let $K, M \in \mathcal{C}_X$ and $L, N \in \mathcal{C}_Y$ be objects,*

$$\phi : f^*L \rightarrow K, \quad \psi : M \rightarrow f^*N \quad \text{and} \quad \chi : M \otimes K \rightarrow f^*(N \otimes L)$$

be morphisms such that

$$\begin{array}{ccc} M \otimes K & \xrightarrow{\chi} & f^*(N \otimes L) \\ \psi \otimes Id \downarrow & & \downarrow \alpha^{-1} \\ f^*N \otimes K & \xrightarrow{Id \otimes \phi} & f^*N \otimes f^*L \end{array}$$

is commutative. Then, the morphism π is duality preserving between the compositions in the diagram

$$\begin{array}{ccccc}
 & & \mathcal{C}_{X,\omega'_X \otimes M} \times \mathcal{C}_{X,K} & \xrightarrow{\{\otimes\}} & \mathcal{C}_{X,\omega'_X \otimes M \otimes K} & & \\
 & \nearrow^{Id \times |f, \phi|^*} & & \uparrow \pi & \searrow^{|f, \chi|_*} & & \\
 \mathcal{C}_{X,\omega'_X \otimes M} \times \mathcal{C}_{Y,L} & \xrightarrow{|f, \psi|_* \times Id} & \mathcal{C}_{Y,N} \times \mathcal{C}_{Y,L} & \xrightarrow{\{\otimes\}} & \mathcal{C}_{Y,N \otimes L} & &
 \end{array}$$

6.3.10. COROLLARY. (*projection formula for Witt groups*) In the situation of the theorem, assuming furthermore that (f, ϕ) is in \mathcal{B}_W^* and (f, ψ) and (f, χ) are in \mathcal{B}_*^W . Then the pull-back, push-forwards and product for Witt groups of definitions 6.2.6, 6.3.4 and Proposition 4.4.6 satisfy $|f, \chi|_*^W(x \cdot |f, \phi|_W^*(y)) = |f, \psi|_*^W(x) \cdot y$.

A. Signs in the category of complexes

Let \mathcal{E} be an exact category \mathcal{E} admitting infinite countable direct sums and products, with a tensor product \bullet adjoint to an internal Hom (denoted by h) in the sense of Definition 1.3.1. In this section, we explain how certain signs have to be chosen in order to induce a suspended closed symmetric monoidal structure on the category of chain complexes of \mathcal{E} (resp. the homotopy category, the derived category). We use homological complexes, as it is the usual convention in the articles about Witt groups, so the differential of a complex is

$$d_i^A : A_i \rightarrow A_{i-1}.$$

The suspension functor T is

$$(TA)_n = A_{n-1}$$

and the tensor product and the internal Hom are given by

$$(A \otimes B)_n = \bigoplus_{i+j=n} A_i \bullet B_j$$

and

$$[A, B]_n = \prod_{j-i=n} h(A_i, B_j).$$

In table 2, we give a possible choice of signs for the translation functor, tensor product, the associativity morphism (denoted by *asso*), the symmetry morphism and the adjunction morphism (denoted by *ath*), and what it induces on the internal Hom using Proposition 1.5.8. In table 3 further below, we state the compatibility that these signs must satisfy to ensure that all axioms of suspended closed symmetric monoidal categories considered hold.

Balmer [B00], [B01], Gille and Nenashev [G02] [GN03] always consider strict dualities, that is $\epsilon^{th1} = 1$. The signs chosen in [B01, 2.6] imply that $\epsilon_{i,0}^{1h} = 1$. The choices made by [GN03, Example 1.4] are $\epsilon_{i,j}^{1\otimes} = 1$ and $\epsilon_{i,j}^{2\otimes} = (-1)^i$. In [G02, p. 111] the signs $\epsilon_{i,j}^{1h} = 1$ and

Definition of	Sign	Choice	Locus
TA	ϵ_i^T	-1	$d_{i+1}^{TA} = \epsilon_i^T d_i A$
$A \otimes B$	$\epsilon_{i,j}^{1\otimes}$ $\epsilon_{i,j}^{2\otimes}$	1 $(-1)^i$	$\epsilon_{i,j}^{1\otimes} d_i^A \bullet id_{B_j}$ $\epsilon_{i,j}^{2\otimes} id_{A_i} \bullet d_j^B$
$tp_{1,A,B}$	$\epsilon_{i,j}^{tp1}$	1	$\epsilon_{i,j}^{tp1} id_{A_i \bullet B_j}$
$tp_{2,A,B}$	$\epsilon_{i,j}^{tp2}$	$(-1)^i$	$\epsilon_{i,j}^{tp2} id_{A_i \bullet B_j}$
$asso_{A,B,C}$	$\epsilon_{i,j,k}^{asso}$	1	$\epsilon_{i,j,k}^{asso}((A_i \bullet B_j) \bullet C_k \rightarrow A_i \bullet (B_j \bullet C_k))$
$s_{A,B}$	$\epsilon_{i,j}^s$	$(-1)^{ij}$	$\epsilon_{i,j}^s(A_i \bullet B_j \rightarrow B_j \bullet A_i)$
$ath_{A,B,C}$	$\epsilon_{i,j}^{ath}$	$(-1)^{i(i-1)/2}$	$\epsilon_{i,j}^{ath}(\text{Hom}(A_i \bullet B_j, C_{i+j})$ $\rightarrow \text{Hom}(A_i, h(B_j, C_{i+j})))$
$[A, B]$	$\epsilon_{i,j}^{1h}$ $\epsilon_{i,j}^{2h}$	1 $(-1)^{i+j+1}$	$\epsilon_{i,j}^{1h}(d_{i+1}^A)^\#$ $\epsilon_{i,j}^{2h}(d_j^B)^\#$
$th_{1,A,B}$	$\epsilon_{i,j}^{th1}$	1	$\epsilon_{i,j}^{th1} id_{h(A_i, B_j)}$
$th_{2,A,B}$	$\epsilon_{i,j}^{th2}$	$(-1)^{i+j}$	$\epsilon_{i,j}^{th2} id_{h(A_i, B_j)}$

Table 2: Sign definitions

$\epsilon_{i,j}^{2h} = (-1)^{i+j+1}$ are chosen. Finally, the sign chosen for ϖ in [G02, p. 112] corresponds via our definition of ϖ (see Section 3.2) to the equality $\epsilon_{j-i,i}^{ath} \epsilon_{i,j-i}^{ath} \epsilon_{j-i,i}^s = (-1)^{j(j-1)/2}$. It is possible to choose the signs in a way compatible with all these choices and our formalism. It is given in the third column of Table 2. More precisely, we have the following theorem.

A.0.11. THEOREM. *Let $a, b \in \{+1, -1\}$. Then*

$$\begin{array}{ll}
\epsilon_{i,j}^{1\otimes} = 1 & \epsilon_{i,j}^{tp1} = a \\
\epsilon_{i,j}^{2\otimes} = (-1)^i & \epsilon_{i,j}^{tp2} = a(-1)^i \\
\epsilon_{i,j}^{1h} = 1 & \epsilon_{i,j}^{th1} = 1 \\
\epsilon_{i,j}^{2h} = (-1)^{i+j+1} & \epsilon_{i,j}^{th2} = a(-1)^{i+j} \\
\epsilon_{i,j}^{ath} = b(-1)^{i(i-1)/2} & \epsilon_{i,j}^s = (-1)^{ij} \\
\epsilon_i^T = -1 &
\end{array}$$

satisfies all equalities of Table 3 as well as $\epsilon_{j-i,i}^{ath} \epsilon_{i,j-i}^{ath} \epsilon_{j-i,i}^s = (-1)^{j(j-1)/2}$. Therefore, for any exact category \mathcal{E} the category of chain complexes $Ch(\mathcal{E})$ and its bounded variant $Ch_b(\mathcal{E})$ may be equipped with the entire structure of suspended symmetric monoidal category discussed in Section 3.1. Moreover, all signs may be chosen in a compatible way with all the above sign choices of Balmer, Gille and Nenashev.

PROOF. Straightforward. ■

These structures trivially pass to the homotopy category. If the exact category \mathcal{E} one considers has enough injective and projective objects, one obtains a left derived functor of the tensor product and a right derived functor of the internal Hom which are exact in both variables.

References

- [B00] P. Balmer, *Triangular Witt Groups Part 1: The 12-Term Localization Exact Sequence*, K-Theory **4** (2000), no. 19, 311–363.
- [B01] P. Balmer, *Triangular Witt groups. II. From usual to derived*, Math. Z. **236** (2001), no. 2, 351–382.
- [CH08] B. Calmès and J. Hornbostel, *Push-forwards for Witt groups of schemes*, [arXiv:0806.0571](https://arxiv.org/abs/0806.0571) (2008), to appear in Comment. Math. Helv.
- [EK66] S. Eilenberg and G. M. Kelly, *A generalization of the functional calculus*, J. Algebra **3** (1966), 366–375.
- [FHM03] H. Fausk, P. Hu, and J. P. May, *Isomorphisms between left and right adjoints*, Theory Appl. Categ. **11** (2003), No. 4, 107–131 (electronic).
- [G02] S. Gille, *On Witt groups with support*, Math. Annalen **322** (2002), 103–137.
- [GN03] S. Gille and A. Nenashev, *Pairings in triangular Witt theory*, J. Algebra **261** (2003), no. 2, 292–309.
- [KML71] G. M. Kelly and S. Mac Lane, *Coherence in closed categories*, Journal of Pure and Applied Algebra **1** (1971), no. 1, 97–140.
- [ML98] S. Mac Lane, *Categories for the working mathematician*, second ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998.
- [W94] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.

*Department of Pure Mathematics and Mathematical Statistics
University of Cambridge
Cambridge CB3 0WB, UK*

*Hausdorff Center for Mathematics
Universität Bonn
53115 Bonn, Germany*

Email: b.calmes@dpms.cam.ac.uk
jens.hornbostel@hcm.uni-bonn.de

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/22/6/22-06.{dvi,ps,pdf}>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION. Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS. The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\epsilon$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

$\text{T}_{\text{E}}\text{X}$ TECHNICAL EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT $\text{T}_{\text{E}}\text{X}$ EDITOR. Gavin Seal, McGill University: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown \(at\) btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Aurelio Carboni, Università dell' Insubria: aurelio.carboni@uninsubria.it

Valeria de Paiva, Cuill Inc.: valeria@cuill.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Martin Hyland, University of Cambridge: M.Hyland@dpms.cam.ac.uk

P. T. Johnstone, University of Cambridge: ptj@dpms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, University of Western Sydney: s.lack@uws.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@acsu.buffalo.edu

Jean-Louis Loday, Université de Strasbourg: loday@math.u-strasbg.fr

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Brooke Shipley, University of Illinois at Chicago: bshipley@math.uic.edu

James Stasheff, University of North Carolina: jds@math.unc.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca

	compatibility	reason
1	$\epsilon_{i,j}^{1\otimes} \epsilon_{i,j-1}^{1\otimes} \epsilon_{i,j}^{2\otimes} \epsilon_{i-1,j}^{2\otimes} = -1$	$A \otimes B$ is a complex
2	$\epsilon_{i,j}^{\otimes 1} \epsilon_{i,j+k}^{\otimes 1} \epsilon_{i+j,k}^{\otimes 1} \epsilon_{i,j,k}^{asso} \epsilon_{i-1,j,k}^{asso}$	$asso$ is a morphism
3	$\epsilon_{i,j}^{\otimes 2} \epsilon_{j,k}^{\otimes 1} \epsilon_{i+j,k}^{\otimes 1} \epsilon_{i,j+k}^{\otimes 2} \epsilon_{i,j,k}^{asso} \epsilon_{i,j-1,k}^{asso}$	
4	$\epsilon_{i+j,k}^{\otimes 2} \epsilon_{j,k}^{\otimes 2} \epsilon_{i,j+k}^{\otimes 2} \epsilon_{i,j,k}^{asso} \epsilon_{i,j,k-1}^{asso}$	
5	$\epsilon_{i,j,k}^{asso} \epsilon_{i,j+k,l}^{asso} \epsilon_{j,k,l}^{asso} \epsilon_{i,j,k+l}^{asso} \epsilon_{i+j,k,l}^{asso} = 1$	the pentagon of [ML98, p. 252] commutes
6	$\epsilon_{i,j}^{1\otimes} \epsilon_{j,i}^{2\otimes} \epsilon_{i,j}^s \epsilon_{i-1,j}^s = 1$	$s_{A,B}$ is a morphism
7	$\epsilon_{j,i}^{1\otimes} \epsilon_{i,j}^{2\otimes} \epsilon_{i,j}^s \epsilon_{i,j-1}^s = 1$	
8	$\epsilon_{i,j}^s \epsilon_{j,i}^s = 1$	s is self-inverse
9	$\epsilon_{j,k}^s \epsilon_{i,k}^s \epsilon_{i+j,k}^s \epsilon_{i,j,k}^{asso} \epsilon_{k,i,j}^{asso} \epsilon_{i,k,j}^{asso} = 1$	the hexagons of [ML98, p. 253] commutes
10	$\epsilon_i^T \epsilon_{i+j}^T \epsilon_{i,j}^{1\otimes} \epsilon_{i+1,j}^{1\otimes} \epsilon_{i,j}^{tp1} \epsilon_{i-1,j}^{tp1} = 1$	$tp_{1,A,B}$ is a morphism
11	$\epsilon_{i+j}^T \epsilon_{i,j}^{2\otimes} \epsilon_{i+1,j}^{2\otimes} \epsilon_{i,j}^{tp1} \epsilon_{i,j-1}^{tp1} = 1$	
12	$\epsilon_j^T \epsilon_{i+j}^T \epsilon_{i,j}^{2\otimes} \epsilon_{i,j+1}^{2\otimes} \epsilon_{i,j}^{tp2} \epsilon_{i,j-1}^{tp2} = 1$	$tp_{2,A,B}$ is a morphism
13	$\epsilon_{i+j}^T \epsilon_{i,j}^{1\otimes} \epsilon_{i,j+1}^{1\otimes} \epsilon_{i,j}^{tp2} \epsilon_{i-1,j}^{tp2} = 1$	
14	$\epsilon_{i,j}^{tp1} \epsilon_{i,j+1}^{tp1} \epsilon_{i,j}^{tp2} \epsilon_{i+1,j}^{tp2} = -1$	the square in Definition 1.4.12 anti-commutes
15	$\epsilon_{i,j}^{tp1} \epsilon_{i+j,k}^{tp1} \epsilon_{i,j+k}^{tp1} \epsilon_{i,j,k}^{asso} \epsilon_{i+1,j,k}^{asso} = 1$	\boxed{assoc} et al. commute
16	$\epsilon_{i,j}^{tp2} \epsilon_{i+j,k}^{tp1} \epsilon_{i,j+k}^{tp2} \epsilon_{j,k}^{tp1} \epsilon_{i,j,k}^{asso} \epsilon_{i,j+1,k}^{asso} = 1$	
17	$\epsilon_{i+j,k}^{tp2} \epsilon_{i,j+k}^{tp2} \epsilon_{j,k}^{tp2} \epsilon_{i,j,k}^{asso} \epsilon_{i,j,k+1}^{asso} = 1$	
18	$\epsilon_{i,j}^{tp1} \epsilon_{j,i}^{tp2} \epsilon_{i,j}^s \epsilon_{i+1,j}^s = 1$	the square \boxed{s} commutes
19	$\epsilon_{i,j}^{1h} \epsilon_{i,j-1}^{1h} \epsilon_{i,j}^{2h} \epsilon_{i+1,j}^{2h} = -1$	$[A, B]$ is a complex
20	$\epsilon_{i,j}^{1\otimes} \epsilon_{i,j}^{2\otimes} \epsilon_{j-1,i+j-1}^{1h} \epsilon_{i,j-1}^{ath} \epsilon_{i-1,j}^{ath} = -1$	ath is well defined
21	$\epsilon_{i,j}^{1\otimes} \epsilon_{j,i+j}^{2h} \epsilon_{i-1,j}^{ath} \epsilon_{i,j}^{ath} = 1$	

Table 3: Sign definitions