Theory and Applications of Categories, Vol. 6, No. 6, pp. 77–82.

COMPARING COEQUALIZER AND EXACT COMPLETIONS

Dedicated to Joachim Lambek
on the occasion of his 75th birthday.

M. C. PEDICCHIO AND J. ROSICKÝ

ABSTRACT. We characterize when the coequalizer and the exact completion of a category $\mathscr{C}$ with finite sums and weak finite limits coincide.

Introduction

Our aim is to compare two well known completions: the coequalizer completion $\mathscr{C}_{\mathrm{coeq}}$ of a small category $\mathscr{C}$ with finite sums (see [P]) and the exact completion of a small category $\mathscr{C}$ with weak finite limits (see [CV]). For a category $\mathscr{C}$ with finite sums and weak finite limits, $\mathscr{C}_{\mathrm{ex}}$ is always a full subcategory of $\mathscr{C}_{\mathrm{coeq}}$. We characterize when the two completions are equivalent - it turns out that this corresponds to a finiteness condition expressed in terms of reflexive and symmetric graphs in $\mathscr{C}$.

1. Two completions

For a small category $\mathscr{C}$ with finite sums, the *coequalizer completion* of $\mathscr{C}$ is a category $\mathscr{C}_{\mathrm{coeq}}$ with finite colimits together with a finite sums preserving functor $G_{\mathscr{C}} : \mathscr{C} \to \mathscr{C}_{\mathrm{coeq}}$ such that, for any finite sums preserving functor $F : \mathscr{C} \to \mathscr{X}$ into a finitely cocomplete category, there is a unique finite colimits preserving functor $\overline{F} : \mathscr{C}_{\mathrm{coeq}} \to \mathscr{X}$ with $\overline{F} \cdot G_{\mathscr{C}} = F$. This construction has been described by Pitts (cf. [BC]).

For a small category $\mathscr{C}$ with weak limits, the *exact completion* $E_{\mathscr{C}} : \mathscr{C} \to \mathscr{C}_{\mathrm{ex}}$ can be characterized by a universal property as well (see [CV]). In a special case when $\mathscr{C}$ has finite limits, $E_{\mathscr{C}}$ is a finite limits preserving functor into an exact category $\mathscr{C}_{\mathrm{ex}}$ such that, for any finite limits preserving functor $F : \mathscr{C} \to \mathscr{X}$ into an exact category, there is a unique functor $F' : \mathscr{C}_{\mathrm{ex}} \to \mathscr{X}$ which preserves finite limits and regular epimorphisms such that $F' \cdot E_{\mathscr{C}} = F$. Following [HT], $\mathscr{C}_{\mathrm{ex}}$ can be described as a full subcategory of $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$ and $E_{\mathscr{C}}$ as the codomain restriction of the Yoneda embedding $Y : \mathscr{C} \to \mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$. To explain it, we recall that a functor $H : \mathscr{C}^{\mathrm{op}} \to \mathbf{Set}$ is *weakly representable* if it admits

Second author partially supported by the Grant Agency of the Czech Republic under the grant No. 201/99/0310 and by the Italian CNR.

Received by the editors 1999 January 12 and, in revised form, 1999 August 27.

Published on 1999 November 30.

1991 Mathematics Subject Classification: 18A99.

Key words and phrases: exact completion, coequalizer completion, variety.

© M. C. Pedicchio and J. Rosický 1999. Permission to copy for private use granted.

a regular epimorphism $\gamma : YC \to H$ from a representable functor. Then $\mathscr{C}_{\mathrm{ex}}$ consists of those weakly representable functors H admitting $\gamma : YC \to H$ whose kernel pair

$$K \underset{\beta}{\overset{\alpha}{\rightrightarrows}} YC \xrightarrow{\ \gamma\ } H \tag{1}$$

has K weakly representable.

We will start by showing that $\mathscr{C}_{\mathrm{coeq}}$ can be presented as a full subcategory of $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$ too, with $G_{\mathscr{C}}$ being the codomain restriction of Y. The full subcategory of $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$ consisting of all finite products preserving functors will be denoted by $FP(\mathscr{C}^{\mathrm{op}})$. It is well known that $FP(\mathscr{C}^{\mathrm{op}})$ is a variety (see [AR] 3.17).

1.1. Lemma. *Let $\mathscr{C}$ be a category with finite sums. Then $\mathscr{C}_{\mathrm{coeq}}$ is equivalent to the full subcategory of $FP(\mathscr{C}^{\mathrm{op}})$ consisting of finitely presentable objects in $FP(\mathscr{C}^{\mathrm{op}})$.*

Proof. By the universal property of $\mathscr{C}_{\mathrm{coeq}}$, we get

$$FP(\mathscr{C}^{\mathrm{op}}) \approx \mathbf{Lex}((\mathscr{C}_{\mathrm{coeq}})^{\mathrm{op}})$$

where, on the right, there is the full subcategory of $\mathbf{Set}^{(\mathscr{C}_{\mathrm{coeq}})^{\mathrm{op}}}$ consisting of all finite limits preserving functors. The result thus follows from [AR] 1.46. ∎

1.2. Proposition. *Let $\mathscr{C}$ be a category with finite sums and weak finite limits. Then $\mathscr{C}_{\mathrm{ex}}$ is equivalent to a full subcategory of $\mathscr{C}_{\mathrm{coeq}}$.*

Proof. Let $H \in \mathscr{C}_{\mathrm{ex}}$ and consider the corresponding diagram (1). There is a regular epimorphism $\delta : YD \to K$ (because K is weakly representable) and we obtain a coequalizer

$$YD \underset{\overline{\beta}}{\overset{\overline{\alpha}}{\rightrightarrows}} YC \xrightarrow{\ \gamma\ } H \tag{2}$$

where $\overline{\alpha} = \alpha\delta$ and $\overline{\beta} = \beta\delta$. Since YD is a regular projective in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$, the graph $(\overline{\alpha}, \overline{\beta})$ is reflexive (it means the existence of $\varphi : YC \to YD$ with $\overline{\alpha}\varphi = \overline{\beta}\varphi = \mathrm{id}_{Y(C)}$). Following [PW], (2) is a coequalizer in $FP(\mathscr{C}^{\mathrm{op}})$. Therefore, H is finitely presentable in $FP(\mathscr{C}^{\mathrm{op}})$ (cf. [AR] 1.3). Hence, using Lemma 1.1, H belongs to $\mathscr{C}_{\mathrm{coeq}}$. ∎

When $\mathscr{C}$ has finite sums, objects of $\mathscr{C}$ are precisely finitely generated free algebras in the variety $FP(\mathscr{C}^{\mathrm{op}})$. The condition of having weak finite limits too, is a very restrictive one. We give another formulation of it.

1.3. Proposition. *Let $\mathscr{C}$ have finite sums. Then $\mathscr{C}$ has weak finite limits iff finite limits of objects of $\mathscr{C}$ in $FP(\mathscr{C}^{\mathrm{op}})$ are finitely generated.*

Proof. Let $D : \mathscr{D} \to \mathscr{C}$ be a finite diagram and $(\delta_d : A \to YD_d)_{d \in \mathscr{D}}$ its limit in $FP(\mathscr{C}^{\mathrm{op}})$.

Assume that A is finitely generated. Then there is a regular epimorphism $\pi : YC \to A$ where $C \in \mathscr{C}$. Consider a cone $(f_d : X \to D_d)_{d \in \mathscr{D}}$ in $\mathscr{C}$. There is a unique $\varphi : YX \to A$ with $\delta_d\varphi = Yf_d$ for all $d \in \mathscr{D}$. Since π is a regular epimorphism (and YX regular

projective) φ factorizes through π and therefore $(\delta_d\pi : YC \to YD_d)_{d\in\mathscr{D}}$ is a weak limit of YD in $Y(\mathscr{C})$. Hence D has a weak limit in $\mathscr{C}$.

Conversely, assume that D has a weak limit $(\gamma_d : C \to D_d)_{d\in\mathscr{D}}$ in $\mathscr{C}$. There is a unique $\pi : YC \to A$ with $\delta_d\pi = Y\gamma_d$ for all $d \in \mathscr{D}$. Consider $\varphi : YX \to A$, $X \in \mathscr{C}$. There exists $\varphi : X \to C$ such that $Y(\gamma_d\psi) = \delta_d\varphi$ for all $d \in \mathscr{D}$. Hence $\varphi = \pi\psi$, which implies that π is a regular epimorphism (because $YX, X \in \mathscr{C}$ are finitely generated free algebras in the variety $FP(\mathscr{C}^{\mathrm{op}})$). Hence A is finitely generated. ∎

2. When do they coincide?

2.1. Construction. Let $\mathscr{C}$ be a category with weak finite limits. Let $r_0, r_1 : C_1 \to C_0$ be a reflexive and symmetric graph in $\mathscr{C}$. It means that there are morphisms $d : C_0 \to C_1$ and $s : C_1 \to C_1$ with $r_0d = r_1d = \mathrm{id}_{C_0}$ and $r_1s = r_0$, $r_0s = r_1$. We form a weak pullback

$$\begin{array}{ccccc} & & C_2 & & \\ & \overline{r}_0\swarrow & & \searrow\overline{r}_1 & \\ C_1 & & & & C_1 \\ & r_1\searrow & & \swarrow r_0 & \\ & & C_0 & & \end{array} \tag{3}$$

By taking $r_i^2 = r_i\bar{r}_i$, $i = 0, 1$, we get the graph

$$C_2 \underset{r_1^2}{\overset{r_0^2}{\rightrightarrows}} C_1$$

This graph is reflexive: $d^2 : C_0 \to C_2$ is given by $\bar{r}_0d^2 = \bar{r}_1d^2 = d$. It is also symmetric: $s^2 : C_2 \to C_2$ is given by $\bar{r}_0s^2 = s\bar{r}_1$ and $\bar{r}_1s^2 = s\bar{r}_0$. By iterating this procedure, we get reflexive and symmetric graphs

$$C_n \underset{r_1^n}{\overset{r_0^n}{\rightrightarrows}} C_1$$

for $n = 1, 2, \ldots$.

2.2. Definition. *Let $\mathscr{C}$ have weak finite limits. We say that a reflexive and symmetric graph $r_0, r_1 : C_1 \to C_0$ has a bounded transitive hull if there is $n \in \mathbb{N}$ such that, for any $m > n$, there exists $f_m : C_m \to C_n$ with $r_0^n f_m = r_0^m$ and $r_1^n f_m = r_1^m$.*

It is easy to check that the definition does not depend on the choice of weak finite limits. Evidently, if $\mathscr{C}$ has finite limits, Definition 2.2 means that the pseudoequivalence generated by the graph $r_0, r_1 : C_1 \to C_0$ is equal to $r_0^n, r_1^n : C_n \to C_0$.

2.3. Theorem. *Let $\mathscr{C}$ have finite sums and weak finite limits. Then $\mathscr{C}_{\mathrm{ex}}$ is equivalent to $\mathscr{C}_{\mathrm{coeq}}$ iff any reflexive and symmetric graph in $\mathscr{C}$ has a bounded transitive hull.*

PROOF. I. Let $r_0, r_1 : C_1 \to C_0$ be a reflexive and symmetric graph in $\mathscr{C}$. Consider the coequalizer

$$YC_1 \underset{Yr_1}{\overset{Yr_0}{\rightrightarrows}} YC_0 \xrightarrow{\ \gamma\ } H$$

in $\mathbf{Set}^{\mathscr{C}^{op}}$. Put $H_1 = YC_1$, $H_0 = YC_0$ and $\gamma_i = Yr_i$ for $i = 0, 1$. Let γ_0^n, $\gamma_1^n : H_n \to H_0$ be iterations of the graph (γ_0, γ_1) constructed as before, by using pullbacks in $\mathbf{Set}^{\mathscr{C}^{op}}$. There are morphisms $\xi_n : YC_n \to H_n$ such that $\xi_1 = \mathrm{id}_{H_0}$ and $\bar{\gamma}_i^n \xi_{n+1} = \xi_n Y(\bar{r}_i^n)$ for $i = 1, 2$ and $n = 0, 1, \ldots$:



By induction, we will prove that ξ_n are regular epimorphisms in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$. Assume that ξ_n is a regular epimorphism and consider the pullback



in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$. There are morphisms $\varphi : YC_{n+1} \to G$ and $\psi : G \to H_{n+1}$ such that $\psi\varphi = \xi_{n+1}$, $\varrho_i\varphi = Y\bar{r}_i^n$ and $\bar{\gamma}_i^n\psi = \xi_n\varrho_i$ for $i = 0, 1$. Since ξ_n is a regular epimorphism in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$, ψ is a regular epimorphism in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$. Furthermore, it follows from the proof of Proposition 1.3 that φ is a regular epimorphism in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$ too. Hence ξ_{n+1} is a regular epimorphism in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$.

Let I be the relation in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$ determined by the graph (γ_0, γ_1). It means that I is

given by the regular epi-monopair factorization



in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$. Let $\iota_0^n, \iota_1^n : I^n \to H_0$ be the composition of n copies of I. Then I^n is the relation generated by the graph $\gamma_0^n, \gamma_1^n : H_n \to H_0$. Since $\xi_n : YC_n \to H_n$ is a regular epimorphism, I^n is also the relation generated by the graph $Yr_0^n, Yr_1^n : YC_n \to YC_0 = H_0$.

II. Now, assume that $\mathscr{C}$ has bounded transitive hulls of reflexive and symmetric graphs. We are going to prove that $\mathscr{C}_{\mathrm{ex}} \approx \mathscr{C}_{\mathrm{coeq}}$. Let $H : \mathscr{C}^{\mathrm{op}} \to \mathbf{Set}$ belong to $\mathscr{C}_{\mathrm{coeq}}$. Following Proposition 1.2, it suffices to prove that H belongs to $\mathscr{C}_{\mathrm{ex}}$. Since $FP(\mathscr{C}^{\mathrm{op}})$ is a variety and, following Lemma 1.1, H is finitely presentable, H is presented by a coequalizer

$$H_1 \underset{\gamma_1}{\overset{\gamma_0}{\rightrightarrows}} H_0 \xrightarrow{\ \gamma\ } H \tag{4}$$

of a reflexive and symmetric graph in $FP(\mathscr{C}^{\mathrm{op}})$ where H_1 and H_0 are free algebras in $FP(\mathscr{C}^{\mathrm{op}})$ over finitely many generators (cf. [AR], Remark 3.13). Since finitely presentable free algebras in $FP(\mathscr{C}^{\mathrm{op}})$ are precisely finite sums of representable functors and $G_{\mathscr{C}} : \mathscr{C} \to \mathscr{C}_{\mathrm{coeq}}$ preserves finite sums, the functors H_0 and H_1 are representable, $H_i = YC_i$, $i = 0, 1$. Hence we get a reflexive and symmetric graph $r_0, r_1 : C_1 \to C_0$ in $\mathscr{C}$ such that $\gamma_i = Yr_i$, $i = 0, 1$. Since (4) is a coequalizer in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$ too (by [PW]), it suffices to show that the kernel pair

$$K \underset{\beta}{\overset{\alpha}{\rightrightarrows}} YC_0 \xrightarrow{\ \gamma\ } H$$

has K weakly representable.

Since the graph (r_0, r_1) has a bounded transitive hull, there is n such that, for any $m > n$, there exists a graph morphism $Yf_m : YC_m \to YC_n$ from the graph (Yr_0^m, Yr_1^m) to the graph (Yr_0^n, Yr_1^n). Following I., they induce morphisms $I^m \to I^n$ of the corresponding relations. Hence I^n is an equivalence relation and, consequently, it yields a kernel pair of γ

$$I^n \underset{\iota_1^n}{\overset{\iota_0^n}{\rightrightarrows}} H_0 \xrightarrow{\ \gamma\ } H$$

Hence $K \cong I^n$ and since I^n is a quotient of YC_n, K is weakly representable.

III. Conversely, let $\mathscr{C}_{\mathrm{ex}} \approx \mathscr{C}_{\mathrm{coeq}}$ and consider a reflexive and symmetric graph $r_0, r_1 : C_1 \to C_0$ in $\mathscr{C}$. Take a coequalizer

$$YC_1 \underset{Yr_1}{\overset{Yr_0}{\rightrightarrows}} YC_0 \xrightarrow{\ \gamma\ } H$$

in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$. Following [PW], it is a coequalizer in $FP(\mathscr{C}^{\mathrm{op}})$ as well and, using Lemma 1.1, we get that $H \in \mathscr{C}_{\mathrm{coeq}}$. Hence $H \in \mathscr{C}_{\mathrm{ex}}$ and therefore the kernel pair

$$K \underset{\beta}{\overset{\alpha}{\rightrightarrows}} YC_0 \xrightarrow{\ \gamma\ } H$$

has K weakly representable. Hence K is finitely generated in $\mathbf{Set}^{\mathscr{C}^{\mathrm{op}}}$ (see [AR] 1.69). Since K is a union of the chain of compositions I^n, $n = 0, 1, \ldots$, there is n such that $K \cong I^n$. Hence $I^m \cong I^n$ for all $m \geq n$. Following I., I^m is the relation generated by the graph (Yr_0^m, Yr_1^m). Since YC_m are regular projectives, there are graph morphisms $YC_m \to YC_n$ for all $m > n$. Hence, there are graph morphisms $f_m : C_m \to C_n$ for all $m > n$. We have proved that $\mathscr{C}$ has bounded transitive hulls of reflexive and symmetric graphs. ∎

2.4. Example. 1) Let $\mathscr{V}$ be a variety in which finitely generated algebras are closed under finite products and subalgebras (like sets, vector spaces or abelian groups). Let $\mathscr{C}$ be the full subcategory of $\mathscr{V}$ consisting of finitely generated free algebras. Then $\mathscr{C}_{\mathrm{ex}} \approx \mathscr{C}_{\mathrm{coeq}}$.

At first, following [AR] 3.16, $\mathscr{V} \cong FP(\mathscr{C}^{\mathrm{op}})$ and $Y : \mathscr{C} \to FP(\mathscr{C}^{\mathrm{op}})$ corresponds to the inclusion $\mathscr{C} \subseteq \mathscr{V}$. Consider a reflexive and symmetric graph $r_0, r_1 : C_1 \to C_0$ in $\mathscr{C}$. The equivalence relation $K \subseteq C_0 \times C_0$ determined by it is finitely generated (as a subalgebra of $C_0 \times C_0$). Following III. of the proof of 2.3, the graph (r_0, r_1) has a bounded transitive hull.

Remark that $\mathscr{C}$ has weak finite limits (following Proposition 1.3).

2) On the other hand, it is easy to find examples of a small category $\mathscr{C}$ such that $\mathscr{C}_{\mathrm{ex}} \approx \mathscr{C}_{\mathrm{coeq}}$ does not hold. It suffices to consider the category $\mathscr{C}$ of countable sets (and the infinite path as a reflexive and symmetric graph in it) and to use Theorem 2.3.

References

[AR] J. Adámek and J. Rosický, *Locally presentable and accessible categories,* Cambridge University Press 1994

[BC] M. Bunge and A. Carboni, The symmetric topos, *Jour. Pure Appl. Algebra* 105 (1995), 233-249

[CV] A. Carboni and E. M. Vitale, Regular and exact completions, *Jour. Pure Appl. Algebra* 125 (1998), 79-117

[HT] H. Hu and W. Tholen, A note on free regular and exact completions and their infinitary generalizations, *Theory and Applications of Categories,* 2 (1996), 113-132

[PW] M. C. Pedicchio and R. Wood, A simple characterization theorem of theories of varieties, to appear

[P] A. Pitts The lex reflection of a category with finite products, unpublished notes 1996

University of Trieste
Ple Europa 1
34100 Trieste, Italy

Masaryk University
*Janáčkovo nám. 2*a
66295 Brno, Czech Republic

Email: pedicchi@univ.trieste.it and rosicky@math.muni.cz

This article may be accessed via WWW at http://www.tac.mta.ca/tac/ or by anonymous ftp at ftp://ftp.tac.mta.ca/pub/tac/html/volumes/6/n6/n6.{dvi,ps}

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

The method of distribution of the journal is via the Internet tools `WWW/ftp`. The journal is archived electronically and in printed paper format.

Subscription information. Individual subscribers receive (by e-mail) abstracts of articles as they are published. Full text of published articles is available in .dvi and Postscript format. Details will be e-mailed to new subscribers and are available by `WWW/ftp`. To subscribe, send e-mail to `tac@mta.ca` including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, `rrosebrugh@mta.ca`.

Information for authors. The typesetting language of the journal is TEX, and LATEX is the preferred flavour. TEX source of articles for publication should be submitted by e-mail directly to an appropriate Editor. They are listed below. Please obtain detailed information on submission format and style files from the journal's WWW server at `URL http://www.tac.mta.ca/tac/` or by anonymous ftp from `ftp.tac.mta.ca` in the directory `pub/tac/info`. You may also write to `tac@mta.ca` to receive details by e-mail.

Editorial board.
John Baez, University of California, Riverside: `baez@math.ucr.edu`
Michael Barr, McGill University: `barr@barrs.org`
Lawrence Breen, Université de Paris 13: `breen@math.univ-paris13.fr`
Ronald Brown, University of North Wales: `r.brown@bangor.ac.uk`
Jean-Luc Brylinski, Pennsylvania State University: `jlb@math.psu.edu`
Aurelio Carboni, Università dell Insubria: `carboni@fis.unico.it`
P. T. Johnstone, University of Cambridge: `ptj@pmms.cam.ac.uk`
G. Max Kelly, University of Sydney: `maxk@maths.usyd.edu.au`
Anders Kock, University of Aarhus: `kock@imf.au.dk`
F. William Lawvere, State University of New York at Buffalo: `wlawvere@acsu.buffalo.edu`
Jean-Louis Loday, Université de Strasbourg: `loday@math.u-strasbg.fr`
Ieke Moerdijk, University of Utrecht: `moerdijk@math.ruu.nl`
Susan Niefield, Union College: `niefiels@union.edu`
Robert Paré, Dalhousie University: `pare@mscs.dal.ca`
Andrew Pitts, University of Cambridge: `ap@cl.cam.ac.uk`
Robert Rosebrugh, Mount Allison University: `rrosebrugh@mta.ca`
Jiri Rosicky, Masaryk University: `rosicky@math.muni.cz`
James Stasheff, University of North Carolina: `jds@charlie.math.unc.edu`
Ross Street, Macquarie University: `street@math.mq.edu.au`
Walter Tholen, York University: `tholen@mathstat.yorku.ca`
Myles Tierney, Rutgers University: `tierney@math.rutgers.edu`
Robert F. C. Walters, University of Sydney: `walters_b@maths.usyd.edu.au`
R. J. Wood, Dalhousie University: `rjwood@mscs.dal.ca`