



The solutions of a class of operator equations based on different inequality

Xiaofang Yan, Chuanxi Zhu*, Zhaoqi Wu

Department of Mathematics, Nanchang University, Nanchang, 330031, P. R. China.

Communicated by C. Park

Abstract

In this paper, by using random fixed point index theory, some new boundary conditions based on strictly convex or strictly concave functions are established and some new theorems for the solutions of a class of random semi-closed 1-set-contractive operator equations $A(\omega, x) = \mu x$ are obtained, which extend and generalize some corresponding results of Wang [S. Wang, Fixed Point Theory Appl., **2011** (2011), 7 pages]. Finally, an application to a class of random nonlinear integral equations is given to illustrate the usability of the obtained results. ©2016 All rights reserved.

Keywords: Real Banach space, random semi-closed 1-set-contractive operator, random topological degree.
2010 MSC: 47H10, 60H25.

1. Introduction and preliminaries

Random nonlinear analysis is an important mathematical branch, which is mainly concerned with the study of random nonlinear operators and their properties and is widely applied in studying various classes of random differential equations and random integral equations ([1, 6]). In [2], Li introduced the random fixed point index theory of random semi-closed 1-set-contractive operators. Since then, the problem of random nonlinear operators are extensively studied by using such theory (see Refs. [3, 4, 5, 7, 8, 9, 10, 11]). In this paper, by using random fixed point index theory, some new boundary conditions based on strictly convex or strictly concave functions are established, and some new theorems for the solutions of a class of random semi-closed 1-set-contractive operator equations $A(\omega, x) = \mu x$ are obtained in real separable Banach spaces, which extend and generalize some corresponding results in previous literatures [7].

*Corresponding author

Email addresses: xiaoxiaoyan_green@163.com (Xiaofang Yan), chuanxizhu@126.com (Chuanxi Zhu), wuzhaoqi_conquer@163.com (Zhaoqi Wu)

Received 2015-04-02

In this paper, let (Ω, U, γ) be a complete probability measure space, $\gamma(\Omega) = 1$. Let E be a separable real Banach space, (E, B) be a measurable space, where B denotes the σ -algebra generated by all the subsets of E , and D be a bounded open subset in E , ∂D the boundary of D in E .

Definition 1.1 ([2]). A random continuous and bounded operator $A : \Omega \times \overline{D} \rightarrow E$ is said to be a E -valued random semi-closed 1-set-contractive operator, if for almost all $\omega \in \Omega$, $A(\omega, \cdot)$ is a semi-closed 1-set-contractive operator in \overline{D} , (if $I - A$ is a closed operator and for every $\omega \in \Omega, \alpha(A(\omega, D)) \leq \alpha(D)$, see [5]), and for every $x \in \overline{D}$, $A(\cdot, x) : \Omega \rightarrow E$ is an E -valued random operator.

Let D be a nonempty subset of R . If $\varphi : D \rightarrow R$ is a real function such that

$$\varphi[tx + (1 - t)y] < t\varphi(x) + (1 - t)\varphi(y)$$

for all $x, y \in D, x \neq y, t \in (0, 1)$, then φ is called a strictly convex function on D . If $\varphi : D \rightarrow R$ is a real function such that

$$\varphi[tx + (1 - t)y] > t\varphi(x) + (1 - t)\varphi(y)$$

for all $x, y \in D, x \neq y, t \in (0, 1)$, then φ is called a strictly concave function on D .

Lemma 1.2 ([2]). Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X , and $\theta \in D$. Suppose that $A : \Omega \times \overline{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator, such that

$$A(\omega, x) \neq \alpha x$$

for all $(\omega, x) \in \Omega \times \partial D$, and $\alpha > \mu$, where $\mu \geq 1$. Then random operator equations $A(\omega, x) = \mu x$ has a random solution in D .

2. Main results

Theorem 2.1. Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X and $\theta \in D$. Suppose that $A : \Omega \times \overline{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator and $\theta \notin A(\omega, \partial D), \forall \omega \in \Omega$. Moreover, if there exist a strictly convex function $\varphi : R^+ \rightarrow R^+$ with $\varphi(0) = 0$ and a real function $\phi : R^+ \rightarrow R^+$ with $\phi(t) \geq 1$ for all $t > 1$, such that

$$\lambda_1 \varphi(\|A(\omega, x) - \mu x\|) \geq \lambda_2 \varphi(\|A(\omega, x)\|) \phi(\|A(\omega, x) + \mu x\| \cdot \|\mu x\|^{-1}) - \lambda_1 \varphi(\|\mu x\|) \tag{2.1}$$

for all $(\omega, x) \in \Omega \times \partial D$, where $0 < \lambda_1 \leq \lambda_2$. Then random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D .

Proof. By Lemma 1.2, we only need to prove that $A(\omega, x) \neq \alpha x, \forall (\omega, x) \in \Omega \times \partial D, \alpha > \mu$, where $\mu \geq 1$. Suppose this is not true. Then there exists $(\omega_0, x_0) \in \Omega \times \partial D$ and $\alpha_0 > \mu \geq 1$, such that $A(\omega_0, x_0) = \alpha_0 x_0$, i.e., $x_0 = \alpha_0^{-1} A(\omega_0, x_0)$. From (2.1), we have

$$\begin{aligned} \lambda_1 \varphi(\|A(\omega_0, x_0) - \alpha_0^{-1} \mu A(\omega_0, x_0)\|) &\geq \lambda_2 \varphi(\|A(\omega_0, x_0)\|) \phi(\|A(\omega_0, x_0) + \alpha_0^{-1} \mu A(\omega_0, x_0)\| \\ &\quad \cdot \|\alpha_0^{-1} \mu A(\omega_0, x_0)\|^{-1}) - \lambda_1 \varphi(\|\alpha_0^{-1} \mu A(\omega_0, x_0)\|), \end{aligned}$$

which implies that

$$\begin{aligned} &\lambda_1 \varphi((1 - \alpha_0^{-1} \mu) \|A(\omega_0, x_0)\|) + \lambda_1 \varphi(\|\alpha_0^{-1} \mu A(\omega_0, x_0)\|) \\ &\geq \lambda_2 \varphi(\|A(\omega_0, x_0)\|) \cdot \phi[(1 + \alpha_0^{-1} \mu) \cdot \|A(\omega_0, x_0)\| \cdot \alpha_0 \mu^{-1} \|A(\omega_0, x_0)\|^{-1}] \\ &= \lambda_2 \varphi(\|A(\omega_0, x_0)\|) \cdot \phi(\alpha_0 \mu^{-1} + 1). \end{aligned} \tag{2.2}$$

By the strict convexity of φ and $\varphi(0) = 0, A(\omega_0, x_0) \neq \theta$, we obtain

$$\begin{aligned} & \lambda_1\varphi[(1 - \alpha_0^{-1}\mu)\|A(\omega_0, x_0)\| + \alpha_0^{-1}\mu\|\theta\|] + \lambda_1\varphi(\alpha_0^{-1}\mu\|A(\omega_0, x_0)\|) + (1 - \alpha_0^{-1}\mu)\|\theta\|) \\ & < (1 - \alpha_0^{-1}\mu)\lambda_1\varphi(\|A(\omega_0, x_0)\|) + \alpha_0^{-1}\mu\lambda_1\varphi(\|\theta\|) + \alpha_0^{-1}\mu\lambda_1\varphi(\|A(\omega_0, x_0)\|) + (1 - \alpha_0^{-1}\mu)\lambda_1\varphi(\|\theta\|) \quad (2.3) \\ & = \lambda_1\varphi(\|A(\omega_0, x_0)\|). \end{aligned}$$

It is easy to see from (2.2)and(2.3),

$$\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi[(\alpha_0\mu^{-1} + 1)] < \lambda_1\varphi(\|A(\omega_0, x_0)\|).$$

However, since $\alpha_0 > \mu \geq 1$, we have $1 + \alpha_0\mu^{-1} > 1$, and thus $\phi(\alpha_0\mu^{-1} + 1) \geq 1$. Noting that $0 < \lambda_1 \leq \lambda_2$, we have $\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi[(\alpha_0^{-1}\mu + 1)] \geq \lambda_1\varphi(\|A(\omega_0, x_0)\|)$, which is a contradiction. It follows from Lemma 1.2 that random operator equations $A(\omega, x) = \mu x(\mu \geq 1)$ has a random solution in D . \square

Remark 2.2. If there exist a convex function $\varphi : R^+ \rightarrow R^+$, with $\varphi(0) = 0$ and a real function $\phi : R^+ \rightarrow R^+$, with $\phi(t) > 1$, for all $t > 1$ satisfying (2.1), the conclusion of Theorem2.1 also hold.

Corollary 2.3. *Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X and $\theta \in D$. Suppose that $A : \Omega \times \bar{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator and $\theta \notin A(\omega, \partial D), \forall \omega \in \Omega$. Moreover, if there exist $\alpha \in (-\infty, 0) \cup (1, +\infty)$ and $\beta \geq 0$, such that*

$$\lambda_1\|A(\omega, x) - \mu x\|^\alpha \|\mu x\|^\beta \geq \lambda_2\|A(\omega, x)\|^\alpha \|A(\omega, x) + \mu x\|^\beta - \lambda_1\|\mu x\|^{\alpha+\beta}$$

for all $(\omega, x) \in \Omega \times \partial D$, where $0 < \lambda_1 \leq \lambda_2$. Then random operator equations $A(\omega, x) = \mu x(\mu \geq 1)$ has a random solution in D .

Proof. Let $\varphi(t) = t^\alpha, \phi(t) = t^\beta$. Then $\varphi(t)$ is a strictly convex function with $\varphi(0) = 0$ and $\phi(t) \geq 1$ for all $t > 1$. Therefore, by Theorem 2.1, the conclusions follows immediately. \square

Remark 2.4. Letting $\beta = 0, \alpha = 2, \lambda_1 = \lambda_2 = 1$ and $\mu = 1$ in Corollary2.3, $A(\omega_0, \cdot) = A$ and A is completely continuous operator. Thus Corollary 2.3 becomes the famous *Altman's* theorem. Thus, Corollary 2.3 generalizes *Altman's* theorem.

Theorem 2.5. *Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X , and $\theta \in D$. Suppose that $A : \Omega \times \bar{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator and $\theta \notin A(\omega, \partial D), \forall \omega \in \Omega$. Moreover, if there exist a strictly concave function $\varphi : R^+ \rightarrow R^+$ with $\varphi(0) = 0$ and a real function $\phi : R^+ \rightarrow R^+$ with $\phi(t) \leq 1$ for all $t > 1$, such that*

$$\lambda_1\varphi(\|A(\omega, x) - \mu x\|) \leq \lambda_2\varphi(\|A(\omega, x)\|)\phi(\|A(\omega, x) + \mu x\| \cdot \|\mu x\|^{-1}) - \lambda_1\varphi(\|\mu x\|) \quad (2.4)$$

for all $(\omega, x) \in \Omega \times \partial D$, where $\lambda_1 \geq \lambda_2 > 0$. Then random operator equations $A(\omega, x) = \mu x(\mu \geq 1)$ has a random solution in D .

Proof. By Lemma 1.2, we only need to prove that $A(\omega, x) \neq \alpha x, \forall (\omega, x) \in \Omega \times \partial D, \alpha > \mu$, where $\mu \geq 1$. Suppose this is not true. Then there exist $(\omega_0, x_0) \in \Omega \times \partial D$ and $\alpha_0 > \mu \geq 1$, such that $A(\omega_0, x_0) = \alpha_0 x_0$, i.e., $x_0 = \alpha_0^{-1}A(\omega_0, x_0)$. From (2.4), we have

$$\begin{aligned} \lambda_1\varphi(\|A(\omega_0, x_0) - \alpha_0^{-1}\mu A(\omega_0, x_0)\|) & \leq \lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi(\|A(\omega_0, x_0) + \alpha_0^{-1}\mu A(\omega_0, x_0)\| \\ & \cdot \|\alpha_0^{-1}\mu A(\omega_0, x_0)\|^{-1}) - \lambda_1\varphi(\|\alpha_0^{-1}\mu A(\omega_0, x_0)\|), \end{aligned}$$

which implies that

$$\begin{aligned} & \lambda_1\varphi[(1 - \alpha_0^{-1}\mu)\|A(\omega_0, x_0)\|] + \lambda_1\varphi[\|\alpha_0^{-1}\mu A(\omega_0, x_0)\|] \\ & \leq \lambda_2\varphi(\|A(\omega_0, x_0)\|) \cdot \phi[(1 + \alpha_0^{-1}\mu) \cdot \|A(\omega_0, x_0)\| \cdot \alpha_0\mu^{-1}\|A(\omega_0, x_0)\|^{-1}] \quad (2.5) \\ & = \lambda_2\varphi(\|A(\omega_0, x_0)\|) \cdot \phi(\alpha_0\mu^{-1} + 1). \end{aligned}$$

By the strict concavity of φ and $\varphi(0) = 0$, we obtain

$$\begin{aligned} &\lambda_1\varphi[(1 - \alpha_0^{-1}\mu)\|A(\omega_0, x_0)\| + \alpha_0^{-1}\mu\|\theta\|] + \lambda_1\varphi[\alpha_0^{-1}\mu\|A(\omega_0, x_0)\|] + (1 - \alpha_0^{-1}\mu)\|\theta\| \\ &>(1 - \alpha_0^{-1}\mu)\lambda_1\varphi(\|A(\omega_0, x_0)\|) + \alpha_0^{-1}\mu\lambda_1\varphi(\|\theta\|) + \alpha_0^{-1}\mu\lambda_1\varphi(\|A(\omega_0, x_0)\|) + (1 - \alpha_0^{-1}\mu)\lambda_1\varphi(\|\theta\|) \quad (2.6) \\ &=\lambda_1\varphi(\|A(\omega_0, x_0)\|). \end{aligned}$$

It is easy to see from (2.5) and (2.6) that

$$\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi[(\alpha_0\mu^{-1} + 1)] > \lambda_1\varphi(\|A(\omega_0, x_0)\|).$$

However, since $\alpha_0 > \mu \geq 1$, we have $\alpha_0\mu^{-1} + 1 > 1$ and thus $\phi(\alpha_0\mu^{-1} + 1) \leq 1$. Noting that $\lambda_1 \geq \lambda_2 > 0$, we have $\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi[(\alpha_0^{-1}\mu + 1)] \leq \lambda_1\varphi(\|A(\omega_0, x_0)\|)$, which is a contradiction. It follows from Lemma 1.2 that random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D . \square

Remark 2.6. If there exist a concave function $\varphi : R^+ \rightarrow R^+$ with $\varphi(0) = 0$ and a real function $\phi : R^+ \rightarrow R^+$, with $\phi(t) < 1$ for all $t > 1$ satisfying (2.4), the conclusion of Theorem 2.5 also hold.

Corollary 2.7. *Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X and $\theta \in D$. Suppose that $A : \Omega \times \overline{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator and $\theta \notin A(\omega, \partial D)$, for all $\omega \in \Omega$, where $\lambda_1 \geq \lambda_2 > 0$. Moreover, if there exist $\alpha \in (0, 1)$ and $\beta \leq 0$, such that*

$$\lambda_1\|A(\omega, x) - \mu x\|^\alpha \|\mu x\|^\beta \leq \lambda_2\|A(\omega, x)\|^\alpha \|A(\omega, x) + \mu x\|^\beta - \lambda_1\|\mu x\|^{\alpha+\beta}$$

for all $(\omega, x) \in \Omega \times \partial D$, $\lambda_1 \geq \lambda_2 > 0$. Then random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D .

Proof. Let $\varphi(t) = t^\alpha$, $\phi(t) = t^\beta$. Then $\varphi(t)$ is a strictly convex function with $\phi(t) \geq 1$ for all $t > 1$ and $\varphi(0) = 0$. Therefore, by Theorem 2.5, the conclusion follows immediately. \square

Theorem 2.8. *Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X and $\theta \in D$. Suppose that $A : \Omega \times \overline{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator and $\theta \notin A(\omega, \partial D)$, $\forall \omega \in \Omega$. Moreover, if there exist a strictly convex function $\varphi : R^+ \rightarrow R^+$ with $\varphi(0) = 0$ and a real function $\phi : R^+ \rightarrow R^+$ with $\phi(t) \geq 1$ for all $t > 1$, such that*

$$\lambda_1\varphi(\|A(\omega, x) - \mu x\|) \geq \lambda_2\varphi(\|A(\omega, x)\|)\phi(\|A(\omega, x)\| \cdot \|\mu x\|^{-1}) - \lambda_1\varphi(\|\mu x\|) \quad (2.7)$$

for all $(\omega, x) \in \Omega \times \partial D$, where $\lambda_2 \geq \lambda_1 > 0$. Then random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D .

Proof. By Lemma 1.2, we only need to prove that $A(\omega, x) \neq \alpha x$, $\forall (\omega, x) \in \Omega \times \partial D$, $\alpha > \mu \geq 1$. Suppose this is not true. Then there exists $(\omega_0, x_0) \in \Omega \times \partial D$ and $\alpha_0 > \mu \geq 1$, such that $A(\omega_0, x_0) = \alpha_0 x_0$, i.e., $x_0 = \alpha_0^{-1}A(\omega_0, x_0)$. From (2.7), we have

$$\begin{aligned} \lambda_1\varphi(\|A(\omega_0, x_0) - \alpha_0^{-1}\mu A(\omega_0, x_0)\|) &\geq \lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi(\|A(\omega_0, x_0)\| \cdot \|\alpha_0^{-1}\mu A(\omega_0, x_0)\|^{-1}) \\ &\quad - \lambda_1\varphi(\|\alpha_0^{-1}\mu A(\omega_0, x_0)\|), \end{aligned}$$

which implies that

$$\lambda_1\varphi[(1 - \alpha_0^{-1}\mu)\|A(\omega_0, x_0)\|] + \lambda_1\varphi[\|\alpha_0^{-1}\mu A(\omega_0, x_0)\|] \geq \lambda_2\varphi(\|A(\omega_0, x_0)\|) \cdot \phi(\mu^{-1}\alpha_0). \quad (2.8)$$

By the strict convexity of φ and $\varphi(0) = 0$, we obtain

$$\begin{aligned} &\lambda_1\varphi[(1 - \alpha_0^{-1}\mu)\|A(\omega_0, x_0)\| + \alpha_0^{-1}\mu\|\theta\|] + \lambda_1\varphi[\alpha_0^{-1}\mu\|A(\omega_0, x_0)\|] + (1 - \alpha_0^{-1}\mu)\|\theta\| \\ &< \lambda_1(1 - \alpha_0^{-1}\mu)\varphi(\|A(\omega_0, x_0)\|) + \lambda_1\alpha_0^{-1}\mu\varphi(\|\theta\|) + \lambda_1\alpha_0^{-1}\mu\varphi(\|A(\omega_0, x_0)\|) + \lambda_1(1 - \alpha_0^{-1}\mu)\varphi(\|\theta\|) \quad (2.9) \\ &= \lambda_1\varphi(\|A(\omega_0, x_0)\|). \end{aligned}$$

It is easy to see from (2.8) and (2.9) that

$$\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi(\alpha_0\mu^{-1}) < \lambda_1\varphi(\|A(\omega_0, x_0)\|).$$

However, since $\alpha_0 > \mu \geq 1$, we have $\alpha_0\mu^{-1} \geq 1$, and thus $\phi(\alpha_0\mu^{-1}) > 1$. Noting that $\lambda_2 \geq \lambda_1 > 0$, we have $\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi(\mu^{-1}\alpha_0) \geq \lambda_1\varphi(\|A(\omega_0, x_0)\|)$, which is a contradiction. Therefore, it follows from Lemma 1.2 that random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D . \square

Remark 2.9. If there exist a convex function $\varphi : R^+ \rightarrow R^+$ with $\varphi(0) = 0$ and a real function $\phi : R^+ \rightarrow R^+$ with $\phi(t) > 1$ for all $t > 1$ satisfying (2.7), the conclusion of Theorem 2.8 also hold.

Corollary 2.10. *Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X and $\theta \in D$. Suppose that $A : \Omega \times \overline{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator and $\theta \notin A(\omega, \partial D), \forall \omega \in \Omega$. Moreover, if there exist $\alpha \in (-\infty, 0] \cup [1, +\infty)$ and $\beta > 0$, such that*

$$\lambda_1\|A(\omega, x) - \mu x\|^\alpha \|\mu x\|^\beta \geq \lambda_2\|A(\omega, x)\|^{\alpha+\beta} - \lambda_1\|\mu x\|^{\alpha+\beta}$$

for all $(\omega, x) \in \Omega \times \partial D, \lambda_2 \geq \lambda_1 > 0$. Then random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D .

Proof. Let $\varphi(t) = t^\alpha, \phi(t) = t^\beta$. Then $\varphi(t)$ is a convex function with $\varphi(0) = 0$ and $\phi(t) > 1$ for all $t > 1$. Therefore, by theorem 2.8, the conclusion follows immediately. \square

Remark 2.11. Letting $\mu = 1$ and $\lambda_1 = \lambda_2 = 1$ in Corollary 2.10, $A(\omega, \cdot) = A$, we can obtain Theorem 2.17 of [7].

Theorem 2.12. *Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X , and $\theta \in D$. Suppose that $A : \Omega \times \overline{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator, and $\theta \notin A(\omega, \partial D), \forall \omega \in \Omega$. Moreover, if there exist a strictly concave function $\varphi : R^+ \rightarrow R^+$ with $\varphi(0) = 0$ and a real function $\phi : R^+ \rightarrow R^+$ with $\phi(t) \leq 1$ for all $t > 1$, such that*

$$\lambda_1\varphi(\|A(\omega, x) - \mu x\|) \leq \lambda_2\varphi(\|A(\omega, x)\|)\phi(\|A(\omega, x)\| \cdot \|\mu x\|^{-1}) - \lambda_1\varphi(\|\mu x\|) \tag{2.10}$$

for all $(\omega, x) \in \Omega \times \partial D$, where $0 < \lambda_2 \leq \lambda_1$. Then random operator equations $A(\omega, x) = \mu x$ has a random solution in D .

Proof. By Lemma 1.2, we only need to prove that $A(\omega, x) \neq \alpha x, \forall (\omega, x) \in \Omega \times \partial D, \alpha > \mu \geq 1$. Suppose this is not true. Then there exist $(\omega_0, x_0) \in \Omega \times \partial D$ and $\alpha_0 > \mu \geq 1$ such that $A(\omega_0, x_0) = \alpha_0 x_0$, i.e., $x_0 = \alpha_0^{-1}A(\omega_0, x_0)$. From (2.10), we have

$$\begin{aligned} \lambda_1\varphi(\|A(\omega_0, x_0) - \alpha_0^{-1}\mu A(\omega_0, x_0)\|) &\leq \lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi(\|A(\omega_0, x_0)\| \cdot \|\alpha_0^{-1}\mu A(\omega_0, x_0)\|^{-1}) \\ &\quad - \lambda_1\varphi(\|\alpha_0^{-1}\mu A(\omega_0, x_0)\|), \end{aligned}$$

which implies that

$$\lambda_1\varphi[(1 - \alpha_0^{-1}\mu)\|A(\omega_0, x_0)\|] + \lambda_1\varphi[\|\alpha_0^{-1}\mu A(\omega_0, x_0)\|] \leq \lambda_2\varphi(\|A(\omega_0, x_0)\|) \cdot \phi(\mu^{-1}\alpha_0). \tag{2.11}$$

By the strict concavity of φ and $\varphi(0) = 0$, we obtain

$$\begin{aligned} &\lambda_1\varphi[(1 - \alpha_0^{-1}\mu)\|A(\omega_0, x_0)\| + \alpha_0^{-1}\mu\|\theta\|] + \lambda_1\varphi[\alpha_0^{-1}\mu\|A(\omega_0, x_0)\|] + (1 - \alpha_0^{-1}\mu)\|\theta\| \\ &> \lambda_1(1 - \alpha_0^{-1}\mu)\varphi(\|A(\omega_0, x_0)\|) + \lambda_1\alpha_0^{-1}\mu\varphi(\|\theta\|) + \lambda_1\alpha_0^{-1}\mu\varphi(\|A(\omega_0, x_0)\|) + \lambda_1(1 - \alpha_0^{-1}\mu)\varphi(\|\theta\|) \\ &= \lambda_1\varphi(\|A(\omega_0, x_0)\|). \end{aligned} \tag{2.12}$$

It is easy to see from (2.11) and (2.12) that

$$\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi(\alpha_0\mu^{-1}) > \lambda_1\varphi(\|A(\omega_0, x_0)\|).$$

However, since $\alpha_0 > \mu \geq 1$, we have $\alpha_0\mu^{-1} > 1$, and thus $\phi(\alpha_0\mu^{-1}) \leq 1$. Noting that $\lambda_1 \geq \lambda_2 > 0$, we have

$$\lambda_2\varphi(\|A(\omega_0, x_0)\|)\phi(\mu^{-1}\alpha_0) \leq \lambda_1\varphi(\|A(\omega_0, x_0)\|),$$

which is a contradiction. Therefore, it follows from Lemma 1.2 that random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D . □

Corollary 2.13. *Let E be a separable real Banach space, X be a closed convex subset of E , D be a bounded open subset in X and $\theta \in D$. Suppose that $A : \Omega \times \overline{D} \rightarrow X$ is a random semi-closed 1-set-contractive operator and $\theta \notin A(\omega, \partial D), \forall \omega \in \Omega$. Moreover, if there exist $\alpha \in (0, 1)$ and $\beta \leq 0$, such that*

$$\lambda_1\|A(\omega, x) - \mu x\|^\alpha \|\mu x\|^\beta \leq \lambda_2\|A(\omega, x)\|^{\alpha+\beta} - \lambda_1\|\mu x\|^{\alpha+\beta}$$

for all $(\omega, x) \in \Omega \times \partial D$, where $0 < \lambda_2 \leq \lambda_1$. Then random operator equations $A(\omega, x) = \mu x (\mu \geq 1)$ has a random solution in D .

Proof. Let $\varphi(t) = t^\alpha, \phi(t) = t^\beta$. Then $\varphi(t)$ is a strictly concave function with $\varphi(0) = 0$ and $\phi(t) \leq 1$ for all $t > 1$. Therefore, by Theorem 2.12, the conclusion follows immediately. □

3. An application

In this section, we give an example to demonstrate Corollary 2.10.

Example 3.1. Let us consider the following random nonlinear integral equation

$$\mu\varphi(x) = \int_G k(\omega, x, y, \varphi(y))dy, \tag{3.1}$$

where $\mu \geq 1$, and G is a bounded open subset in \mathbb{R}^n .

Suppose that

- (i) $k(\omega, x, y, t)$ is random continuous on $\Omega \times G \times G \times \mathbb{R}^1$.
- (ii) There exist $a, b > 0$ and $b \cdot \text{mes}G < m$, where $m > 0$, such that for any $(\omega, x, y, t) \in \Omega \times G \times G \times \mathbb{R}^1$,

$$|k(\omega, x, y, t)| \leq a + b|t|.$$

Then Equation (3.1) has a random continuous solution $\varphi(\omega, x)$.

Proof. Let $C(G)$ be the Banach space of all continuous functions defined on G with norm defined by

$$\|f\|_C = \sup_{x \in G} \|f(x)\|, f \in C(G).$$

Imitating the proof of Theorem 2.1 of [4], we know that the operator

$$A(\omega, \varphi) = \int_G k(\omega, x, y, \varphi(y))dy$$

is a random compact continuous operator. Therefore, $A(\omega, \varphi) : \Omega \times C(G) \rightarrow C(G)$ is a random semi-closed 1-set-contractive operator.

Let $r = \frac{a \cdot \text{mes}G}{m - b \cdot \text{mes}G} > 0$ and $D = \{\varphi \mid \|\varphi(x)\|_C \leq r\}$ be a closed ball in $C(G)$. From (ii), we have

$$\begin{aligned} |A(\omega, \varphi)| &\leq \int_G |k(\omega, x, y, \varphi(y))| dy \leq \int_G (a + b|\varphi(y)|) dy = a \int_G dy + b \int_G |\varphi(y)| dy \\ &= a \cdot \text{mes}G + b \cdot \text{mes}G \|\varphi\|_C, \forall (\omega, \varphi) \in \Omega \times \partial D. \end{aligned}$$

Thus

$$|A(\omega, \varphi)| \leq a \cdot \text{mes}G + b \cdot \text{mes}G \|\varphi\|_C \leq a \cdot \text{mes}G + br \cdot \text{mes}G = mr, \forall (\omega, \varphi) \in \Omega \times \partial D,$$

which implies that

$$\|A(\omega, \varphi)\| \leq m\varphi, \forall (\omega, \varphi) \in \Omega \times \partial D.$$

Take $m = \frac{2\lambda_1}{\lambda_2}, \alpha = 0, \beta = 1$ in Corollary 2.10. Then all the conditions of Corollary 2.10 are satisfied. Therefore, the random nonlinear integral equation (3.1) has a random continuous solution $\varphi(\omega, x)$. □

Acknowledgements:

The authors thank the editor and the referees for their valuable comments and suggestions. The research was supported by the National Natural Science Foundation of China (11361042, 11326099, 11461045, 11071108) and the Provincial Natural Science Foundation of Jiangxi, China (20132BAB201001, 20142BAB211016, 2010GZS0147).

References

- [1] S. Itoh, *Random fixed point theorems with an application to random differential equations in Banach spaces*, J. Math. Anal. Appl., **67** (1979), 261–273.1
- [2] G. Li, *On random fixed point index and some random fixed point theorems of random 1-set-contractive operator*, Acta Math. Appl. Sin., **19** (1996), 203–212.1, 1.1, 1.2
- [3] G. Li, *The existence theorems of the random solutions for random Hammerstein equation with random kernel*, Appl. Math. Lett., **15** (2002), 121–125.1
- [4] G. Li, L. Debnath, *The existence theorems of the random solutions for random Hammerstein type nonlinear equation*, Appl. Math. Lett., **13** (2000), 111–115.1, 3
- [5] J. X. Sun, *Nonlinear Functional Analysis and Applications*, Science Press, Beijing, (2007).1, 1.1
- [6] Z. K. Wang, *An introduction to random functional analysis*, (in Chinese), Adv. Math., **5** (1962), 45–71.1
- [7] S. Wang, *The fixed point theorems of 1-set-contractive operators in Banach spaces*, Fixed Point Theory Appl., **2011** (2011), 7 pages.1, 2.11
- [8] C. Zhu, *Research on some problems for nonlinear operators*, Nonlinear Anal., **71** (2009), 4568–4571.1
- [9] C. Zhu, C. F.Chen *Calculations of random fixed point index*, J. Math. Anal. Appl., **339** (2008), 839–844.1
- [10] C. Zhu, Z. Xu, *Inequalities and solution of an operator equation*, Appl. Math. Lett., **21** (2008), 607–611.1
- [11] C. Zhu, J. D. Yin, *Calculations of a random fixed point index of a random semi-closed 1-set-contractive operator*, Math. Comput. Modelling, **51** (2010), 1135–1139.1