

Geometry & Topology Monographs

Volume 4: Invariants of knots and 3-manifolds (Kyoto 2001)

Pages 103–117

Knot invariants derived from quandles and racks

SEIICHI KAMADA

Abstract The homology and cohomology of quandles and racks are used in knot theory: given a finite quandle and a cocycle, we can construct a knot invariant. This is a quick introductory survey to the invariants of knots derived from quandles and racks.

AMS Classification 57M25, 57Q45; 55N99, 18G99

Keywords Quandles, racks, cocycle knot invariants, knot colorings.

1 Introduction

Many knot invariants can be calculated and interpreted via knot diagrams. For example, the (fundamental) quandle of a knot and the rack of a framed knot [12, 19] can be computed diagrammatically.

In 1942, Takasaki [31] introduced the notion of a *kei*, that is an algebraic object consisting of a non-empty set and a binary operation on it. A kei acts on itself by multiplication from the right side (the right translations). The notion of a kei is an abstraction of the notion of the symmetric transformations. In particular, the right translations of a kei are involutory ($(x * y) * y = x$ for any elements x, y). When we drop this condition from the kei, we obtain the notion of a *quandle*, that was introduced by Joyce [19] in 1982. He associated a quandle to a knot, called the knot quandle and proved that the knot quandle is a complete invariant of a knot (up to weak equivalence). At the same time, Matveev [23] proved a similar result independently. He called the notion a *distributive groupoid*. Kauffman studied knot quandles and knot *crystals* in [20]. In 1988, Brieskorn [1] introduced the notion of an *automorphic set* and, in 1992, Fenn and Rourke [12] introduced the notion of a *rack*. Automorphic sets and racks are the same notion, but their actions are the left translations and the right translations, respectively. They removed the idempotency condition ($x * x = x$ for any element x) from the quandle. Fenn and Rourke also generalized the notion of the knot quandle to the rack of a framed knot or a knot diagram.

Although knot quandles and racks are strong invariants, it is not easy to use them to distinguish knots by direct calculations. The situation is similar to the knot group. An easy method to use them in order to distinguish knots is to calculate the representations in a given finite quandle/rack. Fox's 3-colorings are representations of the knot quandle to the dihedral kei of order 3 and generalized Fox's colorings (n -colorings) are representations to the dihedral kei of order n [11, 16, 25].

In the 1990's and the early 2000's, the homology and cohomology theory for quandles and racks appeared [4, 6, 13, 14]. In order to distinguish two quandles or racks, we may calculate and compare their homology groups. Such methods cannot distinguish a knot and its dual knot (the mirror image with the reversed orientation), since they have isomorphic knot quandles. Carter, Jelsovsky, Kamada, Langford, and Saito (CJKLS) [4] introduced an invariant of a knot derived from a quandle cocycle. Associated with a finite quandle and a cocycle, we can construct a knot invariant. The cohomology theory of a quandle introduced in [4] is related to the homology theory of a rack introduced by Fenn, Rourke and Sanderson (FRS) in [13, 14], cf [6, 21] or section 4.

The invariant of CJKLS corresponds to the evaluation, by a fixed cocycle, of the fundamental class represented by a knot diagram in the homology group in the sense of FRS. The invariants due to CJKLS and FRS are not invariants that can be derived only from the knot quandle/rack. In fact, a trefoil knot and its mirror image have isomorphic quandles and the 2-twist spun trefoil and its orientation reversed 2-knot have isomorphic quandles. However they can be distinguished by the invariants.

This is a quick introductory survey on the invariants of knots derived from quandles and racks. The author thanks the organizers of the conference for giving him an opportunity to give a talk. The talk was introductory and concentrated on the classical knot case according to the organizer's request. However this article includes some information on the 2-knot case.

2 Keis, quandles and racks

A *kei*, X , is a non-empty set with a binary operation $(a, b) \mapsto a * b$ satisfying the following axioms:

- (K1) For any $a \in X$, $a * a = a$.
- (K2) For any $a, b \in X$, $(a * b) * b = a$.

(K3) For any $a, b, c \in X$, $(a * b) * c = (a * c) * (b * c)$.

A *quandle*, X , is a non-empty set with a binary operation $(a, b) \mapsto a * b$ satisfying the following conditions:

(Q1) For any $a \in X$, $a * a = a$.

(Q2) For any $a, b \in X$, there is a unique $c \in X$ such that $a = c * b$.

(Q3) For any $a, b, c \in X$, we have $(a * b) * c = (a * c) * (b * c)$.

The three axioms correspond respectively to the Reidemeister moves of type I, II, and III (see [12], [20], for example).

A *rack*, X , is a non-empty set with a binary operation $(a, b) \mapsto a * b$ satisfying the following conditions:

(R1) For any $a, b \in X$, there is a unique $c \in X$ such that $a = c * b$.

(R2) For any $a, b, c \in X$, we have $(a * b) * c = (a * c) * (b * c)$.

By definition, a *kei* is a quandle and a quandle is a rack:

$$\{\text{keis}\} \subset \{\text{quandles}\} \subset \{\text{racks}\}.$$

Here are some typical examples.

- Any set X with the operation $x * y = x$ for any $x, y \in X$ is a kei, which is called the *trivial kei*, the *trivial quandle* or the *trivial rack*.
- Let n be a positive integer. For elements $i, j \in \{0, 1, \dots, n - 1\}$, define $i * j \equiv 2j - i \pmod{n}$. Then $*$ defines a kei structure called the *dihedral kei*, R_n . This set can be identified with the set of reflections of a regular n -gon with conjugation as the kei operation.
- A group $X = G$ with n -fold conjugation as the quandle operation: $a * b = b^{-n} a b^n$.
- A $\Lambda (= \mathbf{Z}[T, T^{-1}])$ -module M is a quandle with $a * b = Ta + (1 - T)b$, $a, b \in M$, called an *Alexander quandle*. For example, $R_4 \cong \mathbf{Z}_2[T, T^{-1}]/(T^2 + 1)$.

See [1, 12, 19, 23] for further examples.

A subset S of a rack X is called a *generating set* if any element of X is obtained from the elements of S by applying the operation suitably. If X is a kei, we have

$$\begin{aligned} x * (y * z) &= ((x * (y * z)) * z) * z && \text{(by K2)} \\ &= ((x * z) * ((y * z) * z)) * z && \text{(by K3)} \\ &= ((x * z) * y) * z && \text{(by K2)} \end{aligned} \tag{1}$$

for any x, y, z of X . Using this, Takasaki [31] and Winker [32] (cf. [20], p. 195) proved that any element of a kei X which is expressed by elements of S can be re-expressed by using the same elements in the form as

$$(\cdots(((x_1 * x_2) * x_3) * x_4) \cdots).$$

Such an expression is denoted by $x_1 x_2 x_3 x_4 \cdots$ or $x_1^{x_2 x_3 x_4 \cdots}$ following [31] and [12]. For example,

$$\begin{aligned} (a * (b * c)) * (d * e) &= (((a * c) * b) * c) * (d * e) \\ &= (((((a * c) * b) * c) * e) * d) * e \\ &= a^{cbcede} \end{aligned} \tag{2}$$

Fenn and Rourke [12] introduced this notation for any rack X . They denote by a^b the element $a * b$ of X and assume that a^{b^c} stands for $a^{(b^c)} = a * (b * c)$ and a^{bc} stands for $(a^b)^c = (a * b) * c$. Then the axioms are stated as follows:

(R1) For any $a, b \in X$, there is a unique $c \in X$ such that $a = c^b$.

(R2) For any $a, b, c \in X$, we have

$$a^{bc} = a^{cb^c}. \tag{3}$$

The axiom R1 implies that the function $S_x : X \rightarrow X$ (acting from the right side) defined by $(u)S_x = u * x = u^x$ is a bijection. The axiom R2 implies that it is a homomorphism; $(u * v)S_x = (u)S_x * (v)S_x$ or $(u * v)^x = (u^x) * (v^x)$. We call it the *right translation* by the element x . For a word $W = x_1^{\epsilon_1} \cdots x_k^{\epsilon_k}$ consisting of elements of X and for an element $u \in X$, we denote by u^W the element $(u)S_{x_1}^{\epsilon_1} \cdots S_{x_k}^{\epsilon_k}$. In particular, $a^{b^{-1}}$ stands for the element c with $c^b = a$. Then

$$x^{yz} = x^{z^{-1}yz} \tag{4}$$

for any x, y, z of X . Using this, we see that if S is a generating set of a rack, then any element of the rack is expressed by u^W for some $u \in S$ and a word W consisting of elements of S . Refer to [12] for the details.

3 Knot quandles and colored knot diagrams

The quandle $Q(\mathcal{K})$ of a knot \mathcal{K} and the rack $R(\mathcal{K})$ of a framed knot \mathcal{K} were introduced in [19] and [12]. They can be calculated from a diagram.

Let D be a diagram in \mathbf{R}^2 of an oriented knot \mathcal{K} , and let $E = \{x_1, \dots, x_m\}$ be the set of the arcs of the diagram. We give each arc of the diagram D a specific normal direction (co-orientation) determined by use of the orientations of \mathcal{K}

and \mathbf{R}^2 . For a crossing point τ , let x_i , x_j and x_k be the three arcs around τ such that x_k is the over-arc at the crossing and x_i is one of the under-arcs away from which the normal direction of x_k points. We consider a relation

$$x_j = x_i * x_k \tag{5}$$

and call it the *crossing relation* at τ (see Fig. 1).




Figure 1

The kei $K(D)$, the quandle $Q(D)$, or the rack $R(D)$ of the diagram is defined to be the kei, the quandle or the rack that is generated by the set $E = \{x_1, \dots, x_m\}$ and the defining relations are the crossing relations of the crossings of D .

The kei $K(D)$ can be defined even if the knot \mathcal{K} is unoriented, since $x_j = x_i * x_k$ and $x_i = x_j * x_k$ are equivalent by the second axiom, K2.

By using Reidemeister moves, we see that $K(D)$ and $Q(D)$ are invariants of the knot \mathcal{K} . In fact, the quandle $Q(D)$ is isomorphic to the knot quandle $Q(\mathcal{K})$ of the knot \mathcal{K} , see [12, 19]. Similarly, $R(D)$ is preserved by Reidemeister moves of type 2 and type 3, and hence it is an invariant of the framed knot \mathcal{K} , where we assume the framing is the blackboard framing of the diagram. This is isomorphic to the rack $R(\mathcal{K})$ of the framed knot \mathcal{K} , see [12].

In general it is difficult to distinguish the isomorphism types of two given presentations of keis, quandles or racks. A convenient method is to use representations to a finite kei, quandle or rack X .

Let $\rho : K(D) \rightarrow X$ be a homomorphism. We call it a *coloring* of D by X . We color the arc x_i of the diagram D with the element $\rho(x_i)$ of X . The crossing relation $x_j = x_i * x_k$ implies $\rho(x_j) = \rho(x_i) * \rho(x_k)$ in X . This is called the *crossing relation* for the coloring or the *coloring condition*.

For example, consider the case that X is the dihedral kei $R_3 = \{0, 1, 2\}$ of order three. The crossing relation implies that the three colors $\rho(x_i)$, $\rho(x_j)$ and $\rho(x_k)$ are the same or all of them are distinct. This is the 3-coloring condition due to Fox, cf. [11, 25]. (In Exercise 6 of Chapter VI of [11], there is an additional

condition that all three colors are actually used. The 3-coloring condition together this condition is also referred to as Fox's 3-coloring condition. In this paper, we do not assume it.)




Figure 2

A typical diagram of the trefoil knot yields the presentation

$$\langle x_1, x_2, x_3 \mid x_3 = x_1 * x_2, \quad x_1 = x_2 * x_3, \quad x_2 = x_3 * x_1 \rangle \quad (6)$$

(see Fig. 2). Then the diagram has 9 Fox's 3-colorings. Three of them are trivial colorings and the other six are non-trivial colorings. Since the obvious diagram of the trivial knot has trivial colorings only, we see that the trefoil knot is not the trivial knot.

When we use the dihedral kei R_n of order n , we have Fox's n -coloring. Fox's n -coloring is studied in [25]. In general, by use of a finite kei, we can consider colorings for unoriented knot diagrams. When we use a finite quandle or rack X which is not involutory, diagrams must be oriented. A generalized n -coloring, called an (n, r) -coloring, was introduced by Silver and Williams [30]. It is interpreted as a coloring by a certain quandle X .

The knot group (the fundamental group of the knot complement) $G(\mathcal{K})$ can be calculated as the *group* $G(D)$ of a diagram D that is the group with the generating set $E = \{x_1, \dots, x_m\}$ and the Wirtinger relations $x_j = x_k^{-1} x_i x_k$ derived from the crossing points. The presentation of $G(D)$ is obtained from the presentation of $Q(D)$ by assuming $x_j = x_i * x_k$ to be $x_j = x_k^{-1} x_i x_k$. In fact, the knot group is a consequence of the knot quandle (see below or [12]).

For a quandle/rack X , assuming $x * y$ to be $y^{-1} x y$, we obtain a group. It is called the *associated group* of X and denoted by $\text{As}(X)$. More precisely $\text{As}(X)$ is the group $F(X)/N$ where $F(X)$ is the free group generated by the elements of X and N is the normal subgroup of $F(X)$ generated by the words $(x * y) y^{-1} x y$

for $x, y \in X$, see [12]. A (group) presentation of $\text{As}(X)$ is obtained from a (quandle/rack) presentation of X by reading $x * y$ as $y^{-1}xy$, cf. Lemma 4.3 of [12]. Conversely, for a group G , define a binary operation $x * y$ by $y^{-1}xy$ and obtain a quandle. Such a quandle is called the *conjugation quandle* of G and denoted by $\text{conj}(G)$.

Joyce [19] and Matveev [23] proved the following theorem, which shows that the knot quandle is truly stronger than the knot group for some composite knots.

Theorem 3.1 (Joyce [19], Matveev [23]) *Knots \mathcal{K} and \mathcal{K}' with $Q(\mathcal{K}) \cong Q(\mathcal{K}')$ are weakly equivalent; namely, $(\mathbf{R}^3, \mathcal{K})$ is homeomorphic to $(\mathbf{R}^3, \mathcal{K}')$ when we ignore the orientations of \mathbf{R}^3 and the knots.*

For an oriented knot \mathcal{K} , let $-\mathcal{K}$ denote the knot with the reversed orientation and let \mathcal{K}^* denote the mirror image. Then $Q(\mathcal{K}) \cong Q(-\mathcal{K}^*)$. If the knot \mathcal{K} is invertible (i.e., $\mathcal{K} \cong -\mathcal{K}$), or amphicheiral (i.e., $\mathcal{K} \cong \mathcal{K}^*$), then $Q(\mathcal{K})$, $Q(-\mathcal{K})$, $Q(\mathcal{K}^*)$ and $Q(-\mathcal{K}^*)$ are isomorphic to each other. Therefore any invariant which is derived only from the quandle $Q(\mathcal{K})$ cannot distinguish the trefoil and its mirror image, since the trefoil is invertible. The invariants due to CJKLS and FRS can distinguish this pair and hence they are not invariants that are derived only from the knot quandle.

4 Homologies and cohomologies of a quandle and a rack

For a rack X , let $C_n^{\text{R}}(X)$ be the free abelian group generated by n -tuples (x_1, \dots, x_n) of elements of X when n is a positive integer and put $C_n^{\text{R}}(X) = 0$. Define a homomorphism $\partial_n : C_n^{\text{R}}(X) \rightarrow C_{n-1}^{\text{R}}(X)$ by

$$\begin{aligned} \partial_n(x_1, x_2, \dots, x_n) &= \sum_{i=1}^n (-1)^i \left[(x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \right. \\ &\quad \left. - (x_1^{x_i}, x_2^{x_i}, \dots, x_{i-1}^{x_i}, x_{i+1}, \dots, x_n) \right] \end{aligned} \tag{7}$$

for $n \geq 2$ and $\partial_n = 0$ for $n \leq 1$. Then $C_*^{\text{R}}(X) = \{C_n^{\text{R}}(X), \partial_n\}$ is a chain complex. Let $C_n^{\text{D}}(X)$ be the subset of $C_n^{\text{R}}(X)$ generated by n -tuples (x_1, \dots, x_n) with $x_i = x_{i+1}$ for some $i \in \{1, \dots, n-1\}$ if $n \geq 2$; otherwise let $C_n^{\text{D}}(X) = 0$. If X is a quandle, then $\partial_n(C_n^{\text{D}}(X)) \subset C_{n-1}^{\text{D}}(X)$ and $C_*^{\text{D}}(X) = \{C_n^{\text{D}}(X), \partial_n\}$

is a sub-complex of $C_*^R(X)$. In this case, let $C_*^Q(X)$ be the quotient complex $C_*^R(X)/C_*^D(X)$. Let G be an abelian group and let W stand for R, D or Q. Then we consider the homology and cohomology groups

$$H_n^W(X; G) = H_n(C_*^W(X) \otimes G), \quad H_W^n(X; G) = H^n(\text{Hom}(C_*^W(X), G)). \quad (8)$$

Definition 4.1 The n th rack homology and rack cohomology groups [13] of a rack/quandle X with coefficient group G are $H_n^R(X; G)$ and $H_R^n(X; G)$. The n th degeneration homology and degeneration cohomology groups of a quandle X with coefficient group G are $H_n^D(X; G)$ and $H_D^n(X; G)$. The n th quandle homology and quandle cohomology groups [4] of a quandle X with coefficient group G are $H_n^Q(X; G)$ and $H_Q^n(X; G)$.

The n th cocycle group of the cochain complex $\text{Hom}(C_*^Q(X), G)$ is called the quandle n -cocycle group and denoted by $Z_n^Q(X; G)$. We will omit the coefficient group G as usual if $G = \mathbf{Z}$. Refer to [13, 14, 15, 17] for some calculations and applications of the rack homology groups, and [4, 5, 6, 7, 21, 24] for the quandle homology groups.

Theorem 4.2 (Universal Coefficient Theorem [6]) *There exist split exact sequences*

$$0 \rightarrow H_n^W(X) \otimes G \rightarrow H_n^W(X; G) \rightarrow \text{Tor}(H_{n-1}^W(X), G) \rightarrow 0 \quad (9)$$

$$0 \rightarrow \text{Ext}(H_{n-1}^W(X), G) \rightarrow H_W^n(X; G) \rightarrow \text{Hom}(H_n^W(X), G) \rightarrow 0. \quad (10)$$

By definition, there is a natural short exact sequence

$$0 \rightarrow C_*^D(X) \rightarrow C_*^R(X) \rightarrow C_*^Q(X) \rightarrow 0, \quad (11)$$

which is split (in the weak sense), namely, for each n , the short exact sequence

$$0 \rightarrow C_n^D(X) \rightarrow C_n^R(X) \rightarrow C_n^Q(X) \rightarrow 0 \quad (12)$$

is split. Therefore we have a long exact sequence

$$\dots \xrightarrow{\partial_*} H_n^D(X; G) \rightarrow H_n^R(X; G) \rightarrow H_n^Q(X; G) \xrightarrow{\partial_*} H_{n-1}^D(X; G) \rightarrow \dots \quad (13)$$

In [6], it was proved that this sequence is split into short exact sequences for small n 's and it was conjectured that so is for all n 's. This was proved by Litherland and Nelson [21].

Theorem 4.3 (Litherland and Nelson [21]) *The short exact sequence (11) is split in the strong sense, that is, there exist splitting homomorphisms compatible with the boundary maps. Thus, for each n , there is a short exact sequence*

$$0 \rightarrow H_n^D(X; G) \rightarrow H_n^R(X; G) \rightarrow H_n^Q(X; G) \rightarrow 0. \quad (14)$$

Suppose that X is a quandle with finitely many orbits, where orbits means orbits by the inner-automorphisms of X , i.e., orbits by the right action of X on itself by the right translations. (In [19] orbits by all the automorphisms of X are studied.) Let $\mathbf{Z}[\text{Orb}(X)]$ be the free abelian group generated by the orbit set, $\text{Orb}(X)$, of X . In [6, 17] it is shown that

$$H_1^D(X) = 0, \quad H_1^R(X) \cong H_1^Q(X) \cong \mathbf{Z}[\text{Orb}(X)], \quad \text{and} \quad H_2^D(X) \cong \mathbf{Z}[\text{Orb}(X)].$$

Hence

$$H_2^R(X) \cong H_2^Q(X) \oplus \mathbf{Z}[\text{Orb}(X)].$$

In [21](Theorem 2.2) it is proved that

$$H_3^R(X) \cong H_3^Q(X) \oplus H_2^Q(X) \oplus \mathbf{Z}[\text{Orb}(X) \times \text{Orb}(X)].$$

The quandle (co)homology groups of some finite quandles were calculated in [5] (Table 1). However there are errata due to the authors. They thank R. A. Litherland, S. Nelson [21] and T. Mochizuki [24] for pointing out and correcting them. Further calculations on homology/cohomology groups of finite Alexander quandles are found in their papers [21] and [24].

5 Knot invariants

Let X be a finite quandle, let G be an abelian group (written in multiplicative notation), and let $\phi \in C_Q^2(X; G)$ be a quandle 2-cocycle.

Let D be an oriented knot diagram. Recall that a homomorphism $\rho : Q(D) \rightarrow X$ is called a coloring and satisfies the crossing condition $\rho(x_j) = \rho(x_i) * \rho(x_k)$ at each crossing, where x_i, x_j and x_k are as before (see Fig. 1). For each crossing point τ , we consider an element $W_\phi(\tau, \rho)$ of G determined by

$$W_\phi(\tau, \rho) = \phi(\rho(x_i), \rho(x_k))^\epsilon, \tag{15}$$

where x_i and x_k are the arcs around τ corresponding the crossing relation $x_j = x_i * x_k$, and ϵ is the sign of the crossing point τ . Consider the element

$$\Phi_\phi(D) = \sum_\rho \prod_\tau W_\phi(\tau, \rho) \tag{16}$$

of the group ring $\mathbf{Z}G$, where ρ runs all colorings of D by X and τ runs all crossing points of D . Since ϕ is a cocycle, the value $\Phi_\phi(D)$ is preserved by Reidemeister moves. Therefore it is an invariant of the knot \mathcal{K} represented by the diagram. Thus we denote this value by $\Phi_\phi(\mathcal{K})$ and call it the *state-sum invariant* or the *quandle cocycle invariant*, [4]. If ϕ and ϕ' are cocycles

that are cohomologous then $\Phi_\phi(\mathcal{K}) = \Phi_{\phi'}(\mathcal{K})$. Thus this invariant depends on the quandle cohomology class of ϕ . When ϕ is a trivial cocycle, the element $W_\phi(\tau, \rho)$ is the trivial element of G . In this case the invariant $\Phi_\phi(\mathcal{K})$ is equal to the number of all colorings of D by X . In [8, 2], the *abelian extension* $E = E(X, G, \phi)$ is defined as $G \times X$ as a set, with the quandle operation $(g_1, x_1) * (g_2, x_2) = (g_1\phi(x_1, x_2), x_1 * x_2)$. The state-sum $\Phi_\phi(D)$ is an obstruction to extending the colorings of D by X to colorings by $E(X, G, \phi)$, see Theorems 4.1 and 4.5 of [2].

Let X be a finite quandle, let G be an abelian group (written in multiplicative notation), and let $\theta \in C_Q^3(X; G)$ be a quandle 3-cocycle.

A *shadow coloring* (or *face coloring*) of D by X is a function $\tilde{\rho} : \tilde{E} \rightarrow X$, where \tilde{E} is the set of arcs of the diagram D and the regions separated by the underlying immersed curve of the diagram, satisfying the coloring condition ($\tilde{\rho}(x_j) = \tilde{\rho}(x_i) * \tilde{\rho}(x_k)$) and the condition that if y_i and y_j are regions which are adjacent to an arc x_k and the normal direction of x_k points away from y_i toward y_j , then

$$\tilde{\rho}(y_j) = \tilde{\rho}(y_i) * \tilde{\rho}(x_k).$$

We call this condition the *face coloring condition*.

Let τ be a crossing point and let x_i and x_k be the arcs around τ appearing in the crossing relation $x_j = x_i * x_k$ at τ as before. Let y be the region which is one of the four regions around τ such that the normal directions of x_i and x_k are away from the region y . Then we give this crossing point τ an element $W_\theta(\tau, \tilde{\rho})$ of G determined by

$$W_\theta(\tau, \tilde{\rho}) = \theta(\tilde{\rho}(y), \tilde{\rho}(x_i), \tilde{\rho}(x_k))^\epsilon, \quad (17)$$

where ϵ is the sign of the crossing point. Consider the element

$$\Psi_\theta(D) = \sum_{\tilde{\rho}} \prod_{\tau} W_\theta(\tau, \tilde{\rho}) \quad (18)$$

of the group ring $\mathbf{Z}G$, where $\tilde{\rho}$ runs all shadow colorings of D by X and τ runs all crossing points of D . Since θ is a cocycle, the value $\Psi_\theta(D)$ is preserved by Reidemeister moves. It is an invariant of the knot \mathcal{K} represented by D . We denote this value by $\Psi_\theta(\mathcal{K})$. If θ and θ' are cohomologous, then $\Psi_\theta(\mathcal{K}) = \Psi_{\theta'}(\mathcal{K})$. Thus this invariant depends on the quandle 3-cohomology class of θ . When θ is a trivial cocycle, the invariant $\Psi_\theta(\mathcal{K})$ is equal to the number of the shadow colorings of D by X .

Colorings and shadow colorings are defined for oriented knotted surfaces in 4-space similarly using their diagrams in 3-space. For a knotted surface \mathcal{K}

in \mathbf{R}^4 , by modifying it slightly, we may assume that the projection $p : \mathcal{K} \rightarrow \mathbf{R}^3$ to the 3-space is a generic map. The singularity set of the projection consists of double points, triple points and branch points. By removing a small regular neighborhood of the under-curve of the double curve, we have a compact embedded surface in the 3-space. This is a diagram of the knotted surface \mathcal{K} (see [10] for details). Let D be the diagram and let $E = \{x_1, \dots, x_m\}$ be the set of sheets of the diagram. Using the orientation of \mathcal{K} and the orientation of 3-space, we give a normal direction (co-orientation) to each sheet. At a double curve of the projection $p(\mathcal{K})$, let x_i, x_j and x_k be the three sheets around the double curve such that x_k is the over-sheet and x_i and x_j are under-sheets and that the normal direction is away from x_i toward x_j . At each double curve, we consider a relation

$$x_j = x_i * x_k \tag{19}$$

and call it the *crossing relation* around the double curve (see Fig. 3). The *quandle* $Q(D)$ of the diagram is defined to be the quandle generated by $E = \{x_1, \dots, x_m\}$ and the defining relations are the crossing relations around the double curves. This quandle is isomorphic to the quandle $Q(\mathcal{K})$ of the knotted surface \mathcal{K} in the sense of [12, 19].




Figure 3

A homomorphism $\rho : Q(D) \rightarrow X$ is called a *coloring* of the diagram D by X . This is equivalent to coloring the sheets by elements of X such that the colors $\rho(x_i), \rho(x_j)$ and $\rho(x_k)$ of the three sheets x_i, x_j and x_k around a double curve satisfies $\rho(x_j) = \rho(x_i) * \rho(x_k)$.

Let X be a finite quandle, let G be an abelian group (written in multiplicative notation), and let $\theta \in C^3_Q(X; G)$ be a quandle 3-cocycle. For each triple point τ of the diagram D , we consider an element $W_\theta(\tau, \rho)$ of G determined by

$$W_\theta(\tau, \rho) = \phi(\rho(x_i), \rho(x_j), \rho(x_k))^\epsilon, \tag{20}$$

where x_i, x_j and x_k are the sheets around τ such that x_k is the upper-sheet, x_j is one of the two middle-sheets from which the normal direction of x_k is away,

and x_i is one of the four lower-sheets away from which the normal directions of x_k and x_j point, and ϵ is the sign of the triple point τ . Consider the element

$$\Phi_\theta(D) = \sum_{\rho} \prod_{\tau} W_\phi(\tau, \rho) \quad (21)$$

of the group ring $\mathbf{Z}G$, where ρ runs all colorings of D by X and τ runs all triple points of D . Since ϕ is a cocycle, the value $\Phi_\phi(D)$ is preserved by Roseman moves ([10, 26]). Therefore it is an invariant of the knotted surface \mathcal{K} represented by the diagram. We denote this value by $\Phi_\phi(\mathcal{K})$. If ϕ and ϕ' are cohomologous, then $\Phi_\phi(\mathcal{K}) = \Phi_{\phi'}(\mathcal{K})$. Thus this invariant depends on the quandle cohomology class of ϕ . When ϕ is a trivial cocycle, the invariant $\Phi_\phi(\mathcal{K})$ is equal to the number of the colorings of D by X .

6 Examples

Let X be the dihedral kei $R_3 = \{0, 1, 2\}$ of order three, let G be the cyclic group $\langle t \mid t^3 = 1 \rangle$, and let $\theta \in Z_{\mathbf{Q}}^3(R_3; \mathbf{Z}_3)$ be the quandle 3-cocycle defined by

$$\theta = \chi_{012}\chi_{021}\chi_{101}\chi_{201}\chi_{202}\chi_{102}$$

where $\chi_{abc}(x, y, z) = t$ if $(x, y, z) = (a, b, c)$; otherwise $\chi_{abc}(x, y, z) = 1$. In this situation, we consider the invariant Ψ_θ for classical knots (defined by shadow colorings) and the invariant Φ_θ for knotted surfaces (defined by colorings). Note that the values $\Psi_\theta(\mathcal{K})$ and $\Phi_\theta(\mathcal{K})$ are elements of $\mathbf{Z}\langle t \mid t^3 = 1 \rangle$.

Theorem 6.1 (Rourke and Sanderson [27]) *The invariant Ψ_θ distinguishes the trefoil $T(2, 3)$ and its mirror image $T(2, -3)$; $\Psi_\theta(T(2, 3)) \neq \Psi_\theta(T(2, -3))$.*

Theorem 6.2 (Carter et al. [9]) *A torus knot/link $T(2, n)$ is 3-colorable if and only if $n = 3k$ for some integer k . In this case, $\Psi_\theta(T(2, 3k)) = 9 + 18t^k$.*

Theorem 6.3 (Carter et al. [9]) *A torus knot/link $T(3, n)$ is 3-colorable if and only if $n = 2k$ for some integer k . In this case, if k is not a multiple of 3, then $\Psi_\theta(T(3, 2k)) = 9 + 18t^k$; otherwise $\Psi_\theta(T(3, 2k)) = 45$.*

There are four knots in the table with less than 8 crossings that are 3-colorable: 3_1 , 6_1 , 7_4 , and 7_7 . $\Psi_\theta(3_1) = \Psi_\theta(7_4) = \Psi_\theta(7_7) = 9 + 18t$ and $\Psi_\theta(6_1) = 27$, [9]. Further examples are calculated in [5, 9].

Theorem 6.4 (Carter et al. [4], Rourke and Sanderson [27]) *Let \mathcal{K} be the 2-twist spun trefoil and let $-\mathcal{K}$ be the same 2-knot with the reversed orientation. Then $\Phi_\theta(\mathcal{K}) = 3 + 6t^2$ and $\Phi_\theta(-\mathcal{K}) = 3 + 6t$. Thus the invariant Φ_θ distinguishes \mathcal{K} and $-\mathcal{K}$.*

Carter et al. [5] (Theorem 7.2, Cor. 7.3 and 7.4) and Satoh [28] (Theorem 12) gave formulae on the invariants $\Phi_\theta(\mathcal{K})$ for the twist-spun knots. The following theorem is a consequence of the formulae.

Theorem 6.5 (Carter et al. [5], Satoh[28]) *Let \mathcal{K} be the m -twist spun trefoil. Then*

$$\Phi_\theta(\mathcal{K}) = \begin{cases} 3 + 6t^2 & \text{for } m \equiv 2 \pmod{6} \\ 3 + 6t & \text{for } m \equiv 4 \pmod{6} \\ 9 & \text{for } m \equiv 0 \pmod{6} \\ 3 & \text{otherwise.} \end{cases} \quad (22)$$

Remark 6.6 (1) Here we used quandle cohomology of a finite quandle to define the invariants Φ and Ψ . We may use the rack cohomology of a quandle or a rack to define a framed knot. For the purpose of constructing a 3-manifold invariant, one should use the rack homology.

(2) To define the invariants Φ and Ψ , we used the summation over all possible colorings. When we need an invariant of a colored knot, we do not need to take the summation.

Remark 6.7 (1) Shadow coloring of a classical knot diagram is closely related to coloring a knotted surface diagram and its lower decker set. The correspondence is given in [7].

(2) Recall that a shadow coloring of a classical knot diagram is a function $\tilde{\rho} : \tilde{E} \rightarrow X$. The arcs and the regions in \tilde{E} are colored by elements of X . However, when we use an X -set Y in the sense of Fenn et al. (cf. [13, 14]), we may color the regions with elements of Y and the arcs with elements of X . Equivalently, we may introduce and use the cohomologies with twisted coefficients. So there are a lot of variations of the invariants derived from quandle/rack (co)homologies. For example, see [3].

Remark 6.8 Recently, S. Satoh and A. Shima [29] proved that if $\Phi_\theta(\mathcal{K})$ is not an integer for a knotted surface \mathcal{K} , then the minimum triple point number $t(\mathcal{K})$ of the generic projections of \mathcal{K} is greater than three, where Φ_θ is the invariant used in this section. As a corollary, they proved that $t(\mathcal{K}) = 4$ for the 2-twist spun trefoil \mathcal{K} .

References

- [1] **E Brieskorn**, *Automorphic sets and braids and singularities*, *Contemp. Math.* 78 (1988) 45–115
- [2] **J S Carter, M Elhamdadi, M, A Nikiforou, M Saito**, *Extensions of quandles and cocycle knot invariants*, *J. Knot Theory Ramifications* 12 (2003) 725–738
- [3] **J S Carter, M Elhamdadi, M Saito**, *Twisted quandle homology theory and cocycle knot invariant*, *Algebr. Geom. Topol.* 2 (2002) 95–135
- [4] **J S Carter, D Jelsovsky, S Kamada, L Langford, M Saito**, *Quandle cohomology and state-sum invariants of knotted curves and surfaces*, *Trans. Amer. Math. Soc.* 355 (2003) 3947–3989
- [5] **J S Carter, D Jelsovsky, S Kamada, M Saito**, *Computations of quandle cocycle invariants of knotted curves and surfaces*, *Adv. in Math.* 157 (2001) 36–94
- [6] **J S Carter, D Jelsovsky, S Kamada, M Saito**, *Quandle homology groups, their Betti numbers, and virtual knots*, *J. Pure Appl. Algebra* 157 (2001) 135–155
- [7] **J S Carter, S Kamada, M Saito**, *Geometric interpretations of quandle homology*, *J. Knot Theory Ramifications* 10 (2001) 345–386
- [8] **J S Carter, S Kamada, M Saito**, *Diagrammatic computations for quandles and cocycle knot invariants*, *arXiv:math.GT/0102092*
- [9] **J S Carter, S Kamada, M Saito**, *Quandles and cocycle knot invariants*, preprint
- [10] **J Carter, M Saito**, *Knotted surfaces and their diagrams*, *Mathematical Surveys and Monographs* 55, AMS, Providence, RI (1998)
- [11] **R H Crowell, R H Fox** *Introduction to Knot Theory*. Blaisdell Pub. Co. (1963)
- [12] **R Fenn, C Rourke**, *Racks and links in codimension two*, *J. Knot Theory Ramifications* 1 (1992) 343–406
- [13] **R Fenn, C Rourke, B Sanderson**, *Trunks and classifying spaces*, *Appl. Categ. Structures* 3 (1995) 321–356
- [14] **R Fenn, C Rourke, B Sanderson**, *James bundles and applications*, preprint at <http://www.maths.warwick.ac.uk/~cpr/ftp/james.ps>
- [15] **J Flower**, *Cyclic Bordism and Rack Spaces*, PhD Dissertation, Warwick (1995)
- [16] **R H Fox**, *A quick trip through knot theory*, from: “Topology of 3-Manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961)”, Prentice-Hall, Englewood Cliffs, N.J. (1962) 120–167
- [17] **M T Greene**, *Some Results in Geometric Topology and Geometry*, Ph.D. Dissertation, Warwick (1997)

- [18] **A Inoue**, *Quandle homomorphisms of knot quandles to Alexander quandles*, J. Knot Theory Ramifications 10 (2001) 813–821
- [19] **D Joyce**, *A classifying invariant of knots, the knot quandle*, J. Pure Appl. Alg. 23 (1982) 37–65
- [20] **L H Kauffman**, *Knots and Physics*, Series on Knots and Everything 1, World Scientific Publishing Co. Inc. River Edge, NJ (1991)
- [21] **R A Litherland, S Nelson**, *The Betti numbers of some finite racks*, J. Pure Appl. Algebra 178 (2003) 187–202
- [22] **P Lopes**, *Quandles at finite temperatures I*, J. Knot Theory Ramifications 12 (2003) 159–186
- [23] **S Matveev**, *Distributive groupoids in knot theory*, (Russian) Mat. Sb. (N.S.) 119(161) (1982) 78–88, 160
- [24] **T Mochizuki**, *Some calculations of cohomology groups of finite Alexander quandles*, J. Pure Appl. Algebra 179 (2003) 287–330
- [25] **J H Przytycki**, *3-coloring and other elementary invariants of knots*, from: “Knot theory”, Banach Center Publications 42, Polish Acad. Sci. Warszawa (1998) 275–295
- [26] **D Roseman**, *Reidemeister-type moves for surfaces in four dimensional space*, from: “Knot theory”, Banach Center Publications 42, Polish Acad. Sci. Warszawa (1998) 347–380
- [27] **C Rourke, B Sanderson**, *There are two 2-twist-spun trefoils*, arXiv:math.GT/0006062
- [28] **S Satoh**, *Surface diagrams of twist-spun 2-knot*, from “Knots 2000 Korea, Vol. 1 (Yongpyong)”, J. Knot Theory Ramifications 11 (2002) 413–430
- [29] **S Satoh, A Shima**, *The 2-twist-spun trefoil has the triple point number four*, Trans. Amer. Math. Soc. 356 (2004) 1007–1024
- [30] **D S Silver, S G Williams**, *Generalized n -colorings of links*, from: “Knot theory”, Banach Center Publications 42, Polish Acad. Sci. Warszawa (1998) 381–394
- [31] **M Takasaki**, *Abstraction of symmetric transformations*, (in Japanese) Tohoku Math. J. 49 (1942/43) 145–207
- [32] **S Winker**, *Quandles, knots invariants and the n -fold branched cover*, Ph.D. Thesis, University of Illinois at Chicago (1984)

Dept. of Math., Hiroshima University, Hiroshima 739-8526, JAPAN

Email: kamada@math.sci.hiroshima-u.ac.jp

URL: <http://www.math.sci.hiroshima-u.ac.jp/~kamada/index.html>

Received: 30 November 2001 Revised: 12 February 2002