

KMS States for generalized Gauge actions on Cuntz-Krieger algebras

(An application of the Ruelle-Perron-Frobenius Theorem)

Ruy Exel*

Abstract. Given a zero-one matrix A we consider certain one-parameter groups of automorphisms of the Cuntz-Krieger algebra \mathcal{O}_A , generalizing the usual gauge group, and depending on a positive continuous function H defined on the Markov space Σ_A . The main result consists of an application of Ruelle's Perron-Frobenius Theorem to show that these automorphism groups admit a single KMS state.

Keywords: C*-algebras, Cuntz-Krieger algebras, KMS states, gauge action, Ruelle-Perron-Frobenius Theorem.

Mathematical subject classification: 46L55, 37A55.

1 Introduction

In 1978 Olesen and Pedersen [9] showed that the periodic gauge action on the Cuntz algebra \mathcal{O}_n admits a unique KMS state, whose inverse temperature is $\beta = \log n$. Two years later Evans [4: 2.2] extended their result to include, among other things, non-periodic gauge actions, namely one-parameter automorphism groups on \mathcal{O}_n given on the standard generating partial isometries S_j by

$$\gamma_t(S_j) = N_j^{it} S_j, \quad \forall t \in \mathbf{R},$$

where $\{N_j\}_{j=1}^n$ is a collection of real numbers with $N_j > 1$ for all j . See also [3: 3.1]. In 1984 Enomoto, Fujii and Watatani treated the case of the periodic gauge action on the Cuntz-Krieger algebra \mathcal{O}_A for an irreducible matrix A and again arrived at the conclusion that there exists a unique KMS state. The case

Received 21 February 2002.

*Partially supported by CNPq.

of a non-periodic gauge action on \mathcal{O}_A was discussed in [8] in the context of Cuntz-Krieger algebras for infinite matrices but, specializing the conclusions to the finite case, one gets the expected result that if the matrix A is irreducible and the parameters N_j are all greater than 1 then there exists a unique KMS state.

The present work aims to take a new step in the direction of understanding the KMS states on Cuntz-Krieger algebras (over finite matrices) by studying generalized gauge actions on \mathcal{O}_A . In order to describe these actions let Σ_A be the one-sided Markov space for the given matrix A and consider the copy of $C(\Sigma_A)$ within \mathcal{O}_A that is generated by the elements of the form $S_{i_1} \dots S_{i_k} S_{i_k}^* \dots S_{i_1}^*$, where the S_i are the standard generating partial isometries. Fixing an invertible element $U \in C(\Sigma_A)$ it is not hard to see that the correspondence

$$S_j \mapsto US_j$$

extends to give an automorphism of \mathcal{O}_A . Therefore if $H \in C(\Sigma_A)$ is a strictly positive element there exists a unique one-parameter automorphism group $\{\gamma_t\}_{t \in \mathbb{R}}$ of \mathcal{O}_A such that

$$\gamma_t(S_j) = H^{it} S_j.$$

We will refer to γ as the *generalized gauge action*. It is easy to see that this in fact generalizes both the periodic and the non-periodic gauge actions referred to above.

The goal of this paper, as the title suggests, is to study the KMS states for the generalized gauge action on \mathcal{O}_A . Our main result, Theorem 4.4, states that under certain hypotheses there exists a single such KMS state.

The method employed consists of considering \mathcal{O}_A as the crossed product of $C(\Sigma_A)$ by the endomorphism induced by the Markov subshift [6] and applying Theorem 9.6 from [7] to reduce the problem to the search for probability measures on Σ_A which are fixed by Ruelle's transfer operator [12, 13, 1, 2]. This turns out to be closely related to Ruelle's version of the Perron-Frobenius Theorem (see e.g. [2: 1.7]), except that the latter deals with eigenvalues for the transfer operator while we need actual fixed points. With not too much effort we are then able to exploit Ruelle's Theorem in order to understand the required fixed points and thus reach our conclusion.

It should be stressed that Ruelle's Theorem requires two crucial hypotheses, namely that the matrix A be *irreducible and aperiodic* in the sense that there exists a positive integer m such that all entries of A^m are strictly positive (see e.g. [1: Section 1.2]), and that H is Hölder continuous. We are therefore forced to postulate these conditions leaving open the question as to whether one could do without them.

The organization of this paper is as follows: in section (2), the longer and more technical section of this work, we give a brief account of Ruelle's Theorem and draw the conclusions we need with respect to the existence and uniqueness of probability measures that are fixed under the transfer operator.

Section (3) is devoted to reviewing results about crossed products by endomorphisms and in the final section we put all the pieces together proving our main result.

After this article circulated as a preprint we learned of Renault's interesting paper [11] on cocycles for AF-equivalence relations which is closely related to what we do here.

I would finally like to acknowledge helpful conversations with M. Viana who, among other things, brought Ruelle's Theorem to my attention.

2 Ruelle's Perron-Frobenius Theorem

Beyond establishing our notation this section is intended to present Ruelle's Perron-Frobenius Theorem and to develop some further consequences of it to be used in later sections.

Fix, once and for all, an $n \times n$ matrix $A = \{A_{i,j}\}_{1 \leq i,j \leq n}$, with $A_{i,j} \in \{0, 1\}$ for all i and j , such that no row or column of A is identically zero.

Throughout this paper we will be concerned with the associated (one-sided) subshift of finite type, namely the dynamical system (σ, Σ_A) , where Σ_A is the compact topological subspace of the infinite product space $\prod_{i \in \mathbb{N}} \{1, 2, \dots, n\}$ given by

$$\Sigma_A = \left\{ x = (x_0, x_1, x_2, \dots) \in \prod_{i \in \mathbb{N}} \{1, 2, \dots, n\} : A_{x_i, x_{i+1}} = 1 \text{ for all } i \geq 0 \right\},$$

and $\sigma : \Sigma_A \rightarrow \Sigma_A$ is the "left shift", namely the continuous function given by

$$\sigma(x_0, x_1, x_2, \dots) = (x_1, x_2, x_3, \dots).$$

From the assumption that no column of A is identically zero it follows that σ is surjective.

Given a real number $\beta \in (0, 1)$ define a metric d on Σ_A by setting

$$d(x, y) = \beta^{N(x,y)}, \quad \forall x, y \in \Sigma_A,$$

where $N(x, y)$ is the largest integer N such that $x_i = y_i$ for all $i < N$. In the special case in which $x = y$ we set $N(x, y) = +\infty$ and interpret $\beta^{N(x,y)}$ as being zero. It is easy to see that this metric is compatible with the product topology.

Let $C(\Sigma_A)$ denote the C^* -algebra of all continuous complex functions on Σ_A . We will consider the operator

$$\mathcal{L} : C(\Sigma_A) \rightarrow C(\Sigma_A)$$

given by

$$\mathcal{L}(f)|_x = \sum_{y \in \sigma^{-1}(\{x\})} f(y), \quad \forall f \in C(\Sigma_A), \quad \forall x \in \Sigma_A. \quad (2.1)$$

Since σ is surjective one has that $\sigma^{-1}(\{x\})$ is never empty. It is also clear that $\sigma^{-1}(\{x\})$ has at most n elements so that the above sum is finite for every x . One checks that $\mathcal{L}(f)$ is indeed a continuous function and hence that \mathcal{L} is a well defined linear operator on $C(\Sigma_A)$, which is moreover positive and bounded.

Given a real continuous function ϕ on Σ_A the operator

$$\mathcal{L}_\phi : C(\Sigma_A) \rightarrow C(\Sigma_A)$$

given by $\mathcal{L}_\phi(f) = \mathcal{L}(e^\phi f)$ was introduced by Ruelle in [12: 2.3] (see also [13], [2], and [1]) and it is usually referred to as *Ruelle's transfer operator*.

Most of the time we will assume that ϕ is Hölder continuous with respect to the metric d above: recall that a complex function ϕ on a metric space M is said to be *Hölder continuous* when one can find positive constants K and α such that $|\phi(x) - \phi(y)| \leq K d(x, y)^\alpha$, for all x and y in M .

The most important technical tool to be used in this work is the celebrated Ruelle-Perron-Frobenius Theorem which we now state for the convenience of the reader.

Theorem 2.2. (D. Ruelle) *Let A be an $n \times n$ zero-one matrix and let ϕ be a real function defined on Σ_A . Suppose that:*

- (a) *There exists a positive integer m such that $A^m > 0$ (in the sense that all entries are > 0), and*
- (b) *ϕ is Hölder continuous.*

Then there are: a strictly positive function $h \in C(\Sigma_A)$, a Borel probability measure ν on Σ_A , and a real number $\lambda > 0$, such that

- (i) $\mathcal{L}_\phi(h) = \lambda h$,
- (ii) $\mathcal{L}_\phi^*(\nu) = \lambda \nu$, where \mathcal{L}_ϕ^* is the adjoint operator acting on the dual of $C(\Sigma_A)$, and
- (iii) *for every $g \in C(\Sigma_A)$ one has that $\lim_{k \rightarrow \infty} \|\lambda^{-k} \mathcal{L}_\phi^k(g) - \nu(g)h\| = 0$.*

Proof. See e.g. [2: 1.7].

Proposition 2.3. *Under the hypotheses of (2.2) there exists a unique pair (λ_1, ν_1) such that λ_1 is a complex number, ν_1 is a probability measure on Σ_A , and $\mathcal{L}_\phi^*(\nu_1) = \lambda_1 \nu_1$.*

Proof. The existence obviously follows from (2.2.ii). As for uniqueness let (λ_1, ν_1) be such a pair and let (λ, ν) be as in (2.2). For all $g \in C(\Sigma_A)$ we have

$$\lim_{k \rightarrow \infty} \left(\frac{\lambda_1}{\lambda} \right)^k \nu_1(g) = \lim_{k \rightarrow \infty} \nu_1(\lambda^{-k} \mathcal{L}_\phi^k(g)) = \nu(g) \nu_1(h),$$

by (2.2.iii). Plugging $g = 1$ above we conclude that the sequence $\left(\frac{\lambda_1}{\lambda}\right)^k$ converges to the nonzero value $\nu_1(h)$ but this is only possible if $\lambda_1 = \lambda$. For every g we then have that $\nu_1(g) = \nu(g) \nu_1(h)$, so ν_1 is proportional to ν . But since these are probability measures we must have $\nu_1 = \nu$. \square

In particular it follows that both the λ and the ν in the conclusion of (2.2) are uniquely determined. In the following we give an explicit way to compute λ in terms of \mathcal{L}_ϕ (see [1: 1.39]).

Proposition 2.4. *Under the hypotheses of (2.2) one has that*

$$\lambda = \lim_{k \rightarrow \infty} \|\mathcal{L}_\phi^k(1)\|^{1/k}.$$

Proof. Plugging $g = 1$ in (2.2.iii) we conclude that

$$\lim_{k \rightarrow \infty} \lambda^{-k} \|\mathcal{L}_\phi^k(1)\| = \|h\| > 0.$$

So we may choose $n_0 \in \mathbf{N}$ such that for all $n \geq n_0$

$$\frac{\|h\|}{2} < \lambda^{-k} \|\mathcal{L}_\phi^k(1)\| < 2\|h\|.$$

Taking k^{th} roots and then the limit as $k \rightarrow \infty$ we get the conclusion. \square

In the application of Ruelle's Theorem that we have in mind we will take

$$\phi = \phi_\beta = -\beta \log(H), \quad (2.5)$$

where H is a strictly positive continuous function on Σ_A and $\beta > 0$ is a real number.

Observe that if H is Hölder continuous then so is ϕ_β for every real β (this is because “log” is Lipschitz on every compact subset of $(0, +\infty)$, e.g. the range of H). In this case Ruelle’s Theorem gives a correspondence $\beta \rightarrow \lambda$ which we would like to explore more closely in what follows.

Proposition 2.6. *Let A be an $n \times n$ zero-one matrix satisfying (2.2.a) and suppose that H is a Hölder continuous function on Σ_A such that*

$$H(y) > 1, \quad \forall y \in \Sigma_A.$$

For every $\beta \geq 0$ let ϕ_β be as in (2.5) and denote by $\lambda(\beta)$ the unique λ satisfying the conditions of (2.2) for $\phi = \phi_\beta$. Then one has that

- (i) $\lambda(0) > 1$,
- (ii) $\lim_{\beta \rightarrow \infty} \lambda(\beta) = 0$, and
- (iii) λ is a strictly decreasing continuous function of β .

Proof. Observe that

$$\mathcal{L}_{\phi_\beta}(f) = \mathcal{L}(e^{\phi_\beta} f) = \mathcal{L}(e^{-\beta \log(H)} f) = \mathcal{L}(H^{-\beta} f).$$

Let m and M be the supremum and infimum of H on Σ_A , respectively. For every $\beta \geq 0$ and $y \in \Sigma_A$ one therefore has that

$$M^{-\beta} \leq H(y)^{-\beta} \leq m^{-\beta},$$

so that if $f \in C(\Sigma_A)$ is nonnegative we have

$$M^{-\beta} \mathcal{L}(f) \leq \mathcal{L}_{\phi_\beta}(f) \leq m^{-\beta} \mathcal{L}(f).$$

By induction it is easy to see that for all $k \in \mathbf{N}$

$$M^{-k\beta} \mathcal{L}^k(f) \leq \mathcal{L}_{\phi_\beta}^k(f) \leq m^{-k\beta} \mathcal{L}^k(f).$$

Taking norms and k^{th} roots we conclude that

$$M^{-\beta} \|\mathcal{L}^k(f)\|^{1/k} \leq \|\mathcal{L}_{\phi_\beta}^k(f)\|^{1/k} \leq m^{-\beta} \|\mathcal{L}^k(f)\|^{1/k}.$$

Plugging $f = 1$ above and observing that $1 \leq \|\mathcal{L}^k(1)\| \leq \|\mathcal{L}\|^k$ we obtain

$$M^{-\beta} \leq \|\mathcal{L}_{\phi_\beta}^k(1)\|^{1/k} \leq m^{-\beta} \|\mathcal{L}\|,$$

and hence (2.4) yields

$$M^{-\beta} \leq \lambda(\beta) \leq m^{-\beta} \|\mathcal{L}\|.$$

Observing that $H > 1$, and hence that $m > 1$, we deduce (ii). It is also clear from the above that $\lambda(0) \geq 1$ so it is enough to show that $\lambda(0) \neq 1$ in order to obtain (i).

Arguing by contradiction suppose that $\lambda(0) = 1$. Let $h > 0$ be given by (2.2) so that $\mathcal{L}_{\phi_0}(h) = \mathcal{L}(h) = h$. Choose $x^0 \in \Sigma_A$ such that $h(x^0) = \inf_{y \in \Sigma_A} h(y)$, and observe that, since

$$h(x^0) = \sum_{y \in \sigma^{-1}(\{x^0\})} h(y),$$

there exists a unique y in $\sigma^{-1}(\{x^0\})$ which moreover satisfies $h(y) = h(x^0)$. Repeating this process one obtains a sequence $\{x^k\}_{k \in \mathbb{N}}$ in Σ_A such that $\sigma^{-1}(\{x^k\}) = \{x^{k+1}\}$ for all k . Letting $x_k = x_0^k$ (the zeroth coordinate of x^k) we have that $A_{x_{k+1}, x_k} = 1$ and also that this is the only nonzero entry of A in the column x_k . Since A is a finite matrix the sequence $\{x_k\}$ must be periodic. Assuming without loss of generality that the first period of this sequence is $\{1, \dots, m\}$, where $m \leq n$, we see that A has the form

$$A = \begin{bmatrix} S_m & B \\ 0 & C \end{bmatrix},$$

where S_m is the matrix of the forward permutation of m elements. However this is easily seen to contradict (2.2.a) both when $m < n$ (because the zero block in the lower left corner will appear in any power of A) and when $m = n$ (because S_m definitely fails to satisfy (2.2.a)).

In order to prove (iii) let $\delta > 0$ so that $m^\delta \leq H(y)^\delta \leq M^\delta$ for all y in Σ_A . Given $\beta \in \mathbf{R}$ we then have that

$$m^\delta H(y)^{-\beta} \leq H(y)^{-(\beta-\delta)} \leq M^\delta H(y)^{-\beta}.$$

For every nonnegative continuous function f it follows that

$$m^\delta \mathcal{L}_{\phi_\beta}(f) \leq \mathcal{L}_{\phi_{\beta-\delta}}(f) \leq M^\delta \mathcal{L}_{\phi_\beta}(f),$$

and if $k \in \mathbf{N}$ one has

$$m^{k\delta} \mathcal{L}_{\phi_\beta}^k(f) \leq \mathcal{L}_{\phi_{\beta-\delta}}^k(f) \leq M^{k\delta} \mathcal{L}_{\phi_\beta}^k(f).$$

Taking norms and k^{th} roots we conclude that

$$m^\delta \|\mathcal{L}_{\phi_\beta}^k(f)\|^{1/k} \leq \|\mathcal{L}_{\phi_{\beta-\delta}}^k(f)\|^{1/k} \leq M^\delta \|\mathcal{L}_{\phi_\beta}^k(f)\|^{1/k}.$$

With $f = 1$ and taking the limit as $k \rightarrow \infty$, we get by (2.4) that

$$m^\delta \lambda(\beta) \leq \lambda(\beta - \delta) \leq M^\delta \lambda(\beta). \quad (2.7)$$

Substituting $\beta + \delta$ for β above leads to

$$M^{-\delta} \lambda(\beta) \leq \lambda(\beta + \delta) \leq m^{-\delta} \lambda(\beta). \quad (2.8)$$

By (2.7) and (2.8) one sees that λ is a continuous function of β . Since $m > 1$ by hypothesis the rightmost inequality in (2.8) gives $\lambda(\beta + \delta) < \lambda(\beta)$ and hence that λ is strictly decreasing. \square

Corollary 2.9. *Under the hypotheses of (2.6) there exists a unique $\beta > 0$ such that $\lambda(\beta) = 1$.*

3 Preliminaries on Crossed Products

Define the map $\alpha : C(\Sigma_A) \rightarrow C(\Sigma_A)$ by the formula

$$\alpha(f) = f \circ \sigma, \quad \forall f \in C(\Sigma_A).$$

It is easy to see that α is a C^* -algebra endomorphism of $C(\Sigma_A)$. Since σ is surjective one has that α is injective. We should also notice that $\alpha(1) = 1$.

For $x \in \Sigma_A$ let

$$Q(x) = \#\{y \in X : \sigma(y) = x\},$$

(“#” meaning number of elements). Alternatively $Q(x)$ may be defined as the number of “ones” in the column of A indexed by x_0 . Therefore $1 \leq Q(x) \leq n$ for all $x \in \Sigma_A$ so that in particular Q is invertible as an element of $C(\Sigma_A)$. Define the operator

$$\mathcal{L} : C(\Sigma_A) \rightarrow C(\Sigma_A)$$

by $\mathcal{L}(f) = Q^{-1} \mathcal{L}(f)$, where \mathcal{L} is defined in (2.1). It is easy to see that $Q = \mathcal{L}(1)$ and hence that $\mathcal{L}(1) = 1$. Moreover

$$\mathcal{L}(\alpha(f)g) = f \mathcal{L}(g), \quad \forall f, g \in C(\Sigma_A),$$

which tells us that \mathcal{L} is a *transfer operator* for the pair $(C(\Sigma_A), \alpha)$ according to Definition (2.1) in [6]. One may therefore construct the crossed product algebra

$$C(\Sigma_A) \rtimes_{\alpha, \mathcal{L}} \mathbf{N},$$

or $C(\Sigma_A) \rtimes \mathbf{N}$, for short, as in [6:3.7], which turns out to be a C^* -algebra generated by a copy of $C(\Sigma_A)$ and an extra element S which, among other things, satisfies

- $S^*S = 1$,
- $Sf = \alpha(f)S$, and
- $S^*fS = \mathcal{L}(f)$,

for all $f \in C(\Sigma_A)$. See [6] for the precise definition of $C(\Sigma_A) \rtimes \mathbf{N}$.

In [6:6.2] it is proved that $C(\Sigma_A) \rtimes \mathbf{N}$ is isomorphic to the Cuntz-Krieger algebra \mathcal{O}_A . It will be convenient for us to bear in mind the isomorphism between \mathcal{O}_A and $C(\Sigma_A) \rtimes \mathbf{N}$ given in [6], which we next describe. For this consider for each $j = 1, \dots, n$, the clopen subset Σ^j of Σ_A given by

$$\Sigma^j = \{x \in \Sigma_A : x_0 = j\}.$$

These are precisely the sets forming the standard Markov partition of Σ_A . Also let P_j be the characteristic function of Σ^j . According to [6] there exists an isomorphism

$$\Psi : \mathcal{O}_A \rightarrow C(\Sigma_A) \rtimes \mathbf{N}$$

which is determined by the fact that the canonical generating partial isometries $S_j \in \mathcal{O}_A$ are mapped under Ψ as follows:

$$\Psi(S_j) = P_j S Q^{1/2}.$$

We would next like to review the definition of the generalized gauge action on \mathcal{O}_A . For this fix a strictly positive element $H \in C(\Sigma_A)$. According to [7:6.2] there exists a unique one parameter automorphism group γ of $C(\Sigma_A) \rtimes \mathbf{N}$ such that for all $t \in \mathbf{R}$,

$$\gamma_t(S) = H^{it} S, \quad \text{and} \quad \gamma_t(f) = f, \quad \forall f \in C(\Sigma_A).$$

Transferring γ to \mathcal{O}_A via the isomorphism Ψ described above one gets an automorphism group on \mathcal{O}_A which is characterized by the fact that

$$\gamma_t(S_j) = H^{it} S_j, \quad \forall j = 1, \dots, n.$$

Observe that in case H is a constant function, say everywhere equal to Neper's number e , and $A_{ij} \equiv 1$, then \mathcal{O}_A coincides with the Cuntz algebra \mathcal{O}_n and one recovers the action over \mathcal{O}_n considered in [9]. We shall refer to this as the *scalar gauge action*.

For a slightly more general example suppose that H is constant on each Σ^j , taking the value N_j there. Then

$$\gamma_t(S_j) = H^{it} S_j = H^{it} P_j S_j = N_j^{it} S_j,$$

and we obtain special cases of actions studied in [4] or [8].

Observe that the composition $E = \alpha \circ \mathcal{L}$ is a conditional expectation from $C(\Sigma_A)$ onto the range of α . By [7: Section 11], using the set $\{P_1, \dots, P_n\}$, we see that E is of index-finite type. It therefore follows from [7:8.9] that there exists a unique conditional expectation

$$G : \mathcal{O}_A \rightarrow C(\Sigma_A)$$

which is invariant under the scalar gauge action. This conditional expectation must therefore coincide with the conditional expectation given by [5:2.9] for the Cuntz-Krieger bundle (see [10] and [5]).

Let us now give a concrete description of G based on the well known fact that \mathcal{O}_A is linearly spanned by the set of all $S_\mu S_\nu^*$, where μ and ν are finite words in the alphabet $\{1, \dots, n\}$, and we let $S_\mu = S_{\mu_0} \dots S_{\mu_k}$ whenever $\mu = \mu_0 \dots \mu_k$.

For any such μ and ν we have by [5] that

$$G(S_\mu S_\nu^*) = \begin{cases} S_\mu S_\nu^* & , \text{ if } \mu = \nu, \\ 0 & , \text{ if } \mu \neq \nu \end{cases} \quad (3.1)$$

4 KMS states

It is our main goal to describe the KMS states on \mathcal{O}_A for the gauge action γ determined by a given H as above. Recall from [7:9.6] that for every $\beta > 0$ the correspondence

$$\psi \mapsto \nu = \psi|_{C(\Sigma_A)} \quad (4.1)$$

is a bijection from the set of KMS_β states ψ on $C(\Sigma_A) \rtimes \mathbb{N}$ and the set of probability measures¹ ν on Σ_A such that

$$\nu(f) = \nu\left(\mathcal{L}\left(H^{-\beta} \text{ind}(E)f\right)\right), \quad \forall f \in C(\Sigma_A), \quad (4.2)$$

¹By the Riesz Representation Theorem we identify probability measures and states as usual.

where $\text{ind}(E)$ is the Jones-Kosaki-Watatani index of E . See [7] for details. As observed in [7:Section 11] the right hand side of (4.2) coincides with $\nu(\mathcal{L}_{\phi_\beta}(f))$, where ϕ_β is as in (2.5), so that (4.2) is equivalent to

$$\mathcal{L}_{\phi_\beta}^*(\nu) = \nu. \quad (4.3)$$

We now arrive at our main result.

Theorem 4.4. *Let A be an $n \times n$ zero-one matrix satisfying (2.2.a) and let H be a Hölder continuous function on Σ_A such that $H(y) > 1$ for all y in Σ_A . Let γ be the unique one-parameter automorphism group of \mathcal{O}_A such that*

$$\gamma_t(S_j) = H^{it} S_j, \quad \forall j = 1, \dots, n,$$

where the S_j are the canonical partial isometries generating \mathcal{O}_A . Then \mathcal{O}_A admits a unique KMS state ψ for γ . The inverse temperature at which this state occurs is the unique value of β for which $\lambda(\beta) = 1$ (see 2.9). In addition ψ is given by

$$\psi = \nu \circ G,$$

where G is the conditional expectation described in (3.1) and ν is the unique measure on Σ_A satisfying the Ruelle-Perron-Frobenius Theorem for $\phi = -\beta \log(H)$. Finally there are no ground states for γ .

Proof. By (2.9) let $\beta > 0$ be such that $\lambda(\beta) = 1$. Applying (2.2) for $\phi = \phi_\beta = -\beta \log(H)$, let ν be the unique probability measure on Σ_A satisfying

$$\mathcal{L}_{\phi_\beta}^*(\nu) = \lambda(\beta)\nu = \nu.$$

Then (4.3) holds and hence by [7:9.6] the composition $\psi = \nu \circ G$ is a KMS_β state for γ .

Suppose now that $\beta_1 > 0$ and let ψ_1 be a KMS_{β_1} state for γ . Set $\nu_1 = \psi_1|_{C(\Sigma_A)}$ and observe that, again by [7:9.6], one has that ν_1 satisfies (4.3) for ϕ_{β_1} . So the pair $(1, \nu_1)$ satisfies the conditions of (2.3) and hence $\lambda(\beta_1) = 1$ so that $\beta_1 = \beta$ by (2.9). Also by (2.3) ν_1 must coincide with ν and hence $\psi_1 = \psi$ because the correspondence in (4.1) is bijective.

That no ground states exist follows from [7:10.1]. □

References

- [1] V. Baladi. Positive transfer operators and decay of correlations. Advanced Series in Nonlinear Dynamics vol. 16, World Scientific, 2000.
- [2] R. Bowen. Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms. Lecture Notes in Mathematics vol. 470, Springer-Verlag, 1975.
- [3] O. Bratteli, P. E. T. Jorgensen and V. Ostrovskiy. Representation Theory and Numerical AF-invariants: The representations and centralizers of certain states on O_d , preprint, 1999, arXiv:math.OA/9907036.
- [4] D. E. Evans. On O_n , *Publ. Res. Inst. Math. Sci., Kyoto Univ.* **16**: (1980), 915–927.
- [5] R. Exel. Amenability for Fell Bundles, *J. reine angew. Math.*, **492**: (1997), 41–73, arXiv:funct-an/9604009.
- [6] R. Exel. A New Look at The Crossed-Product of a C^* -algebra by an Endomorphism, to appear in *Ergodic Theory Dynam. Systems*.
- [7] R. Exel. Crossed-products by finite index endomorphisms and KMS states, *J. Funct. Analysis*, **199**: (2003), 153–188.
- [8] R. Exel and M. Laca. Partial dynamical systems and the KMS condition, *Commun. Math. Phys.*, **232**: (2003), 223–277, [arXiv:math.OA/0006169].
- [9] D. Olesen and G. K. Pedersen. Some C^* -dynamical systems with a single KMS state, *Math. Scand.* **42**: (1978), 111–118.
- [10] J. Quigg and I. Raeburn. Characterizations of crossed products by partial actions, *J. Oper. Theory*, **37**: (1997), 311–340.
- [11] J. Renault. AF-equivalence relations and their cocycles, Talk at 4th International Conference on Operator Algebras, July 2-7 2001, Constanza, Romania, [arXiv:math.OA/0111182].
- [12] D. Ruelle. Statistical mechanics of a one-dimensional lattice gas, *Commun. Math. Phys.* **9**: (1968), 267–278.
- [13] D. Ruelle. The thermodynamic formalism for expanding maps, *Commun. Math. Phys.* **125**: (1989), 239–262.

Ruy Exel

Departamento de Matemática
 Universidade Federal de Santa Catarina
 88040-900 Florianópolis SC
 BRAZIL

E-mail: exel@mtm.ufsc.br