GEOMETRIC AND ARITHMETIC SUBGROUPS OF THE GROTHENDIECK-TEICHMÜLLER GROUP

HIDEKAZU FURUSHO

Abstract. We compare two geometrically constructed subgroups $\mathbb{\Gamma}$ and GTK of the Grothendieck-Teichmüller group $\widehat{GT}$ with an arithmetically constructed subgroup GTA. We show that the intersection of $\mathbb{\Gamma}$ and GTK is contained in a certain modification of GTA.

0. Introduction

The Grothendieck-Teichmüller group $\widehat{GT}$ is a subgroup of the automorphism group $Aut\widehat{F_2}$ of the free pro-finite group of $\widehat{F_2}$ of rank 2. It is parameterized by elements of $\widehat{\mathbf{Z}}^\times \times [\widehat{F_2}, \widehat{F_2}]$ and is defined by three relations (I), (II) and (III) (see §1). It is a pro-finite group version of the pro-algebraic group $\underline{GT}$ [Dr]. In his study of Galois representations on fundamental groups, Y. Ihara showed that the absolute Galois group $G_{\mathbf{Q}} = Gal(\overline{\mathbf{Q}}/\mathbf{Q})$ can be embedded into $\widehat{GT}$ in [Ih94]. It is still open whether $G_{\mathbf{Q}}$ is equal to $\widehat{GT}$ or not. But recently there have been discovered other several new-type relations satisfied by $G_{\mathbf{Q}}$ in $\widehat{GT}$ (see [Ih00], [LNS] and [NT]). In this article, we show a certain relationship among them. In §1, we shall recall the definition of $\widehat{GT}$ and make a small remark (*Proposition* 1.3) which has been possibly unknown so far. §2 is a review of the definitions of three variants $\mathbb{\Gamma}$ ([LNS]), GTK ([Ih00]) and GTA ([Ih00]) of $\widehat{GT}$. In §3, we introduce main results of the author's master's thesis [F], concerning on a relationship among defining equations of $\mathbb{\Gamma}$, GTK and GTA. In §4, we give many lemmas to complete their proof.

Acknowledgments . The author is profoundly grateful to my master thesis advisor Professor Y. Ihara for warmful encouragements. He also express his special thanks to H. Nakamura and H. Tsunogai for useful discussion and their support and to Professor A. Tamagawa for his kind advises.

2000 *Mathematics Subject Classification*: Primary 14G32, Secondary 11S80.

1. Review of the definition of $\widehat{GT}$

Let $\widehat{F_2}$ be the pro-finite free group of rank 2 with generators x and y. We define $\underline{\widehat{GT}}$ to be the set of pairs $(\lambda, f) \in \widehat{\mathbf{Z}}^\times \times \widehat{F_2}'$ (where $\widehat{F_2}'$ means the topological commutator subgroup $[\widehat{F_2}, \widehat{F_2}]$ of $\widehat{F_2}$) satisfying the following three relations:

$$\begin{cases} \text{(I)} \quad f(x,y)f(y,x) = 1 & \text{((2-cycle relation))} \\ \text{(II)} \quad f(z,x)z^m f(y,z)y^m f(x,y)x^m = 1 & \\ \qquad\qquad \text{for } m = \frac{1}{2}(\lambda - 1),\ xyz = 1 & \text{((3-cycle relation))} \\ \text{(III)} \ f(x_{12},x_{23})f(x_{34},x_{45})f(x_{51},x_{12})f(x_{23},x_{34})f(x_{45},x_{51}) = 1 \quad \text{in } \hat{P}_5^* & \\ \qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{((5-cycle relation))}. \end{cases}$$

Here $\hat{P}_n^*$ is the pro-finite pure sphere braid group with n strings and $x_{i,j} := x_{i,j}^{(n)}$ $(1 \leqslant i,j \leqslant n)$ are its standard generators [Ih94]. For $f \in \widehat{F_2}$ and elements α, β of a pro-finite group H, $f(\alpha,\beta)$ stands for the image of f by the homomorphism $\phi : \widehat{F_2} \to H$ defined by $\phi(x) = \alpha$, $\phi(y) = \beta$. An element $\sigma = (\lambda, f) \in \underline{\widehat{GT}}$ induces an endomorphism of $\widehat{F_2}$ by $\sigma(x) = x^\lambda$, $\sigma(y) = f^{-1}y^\lambda f$, from which we get an embedding $\underline{\widehat{GT}} \hookrightarrow End\widehat{F_2}$ and in fact we can regard $\underline{\widehat{GT}}$ as a sub-monoid of $End\widehat{F_2}$ ([Dr]).

Definition 1.1 ([Dr])**.** The *Grothendieck-Teichmüller group* $\widehat{GT}$ is the group of invertible elements of $\underline{\widehat{GT}}$:

$$\widehat{GT} := \left\{ \sigma = (\lambda, f) \in Aut\ \widehat{F_2} \ \mid (\lambda, f) \text{ satisfies (I)} \sim \text{(III)}. \right\}.$$

Remark 1.2. The above 5-cycle relation (III) is different from the original relation of $\widehat{GT}$ ([Dr])

$$\begin{aligned} \text{(III)}_{\text{DR}} \quad & f(x_{12}^{(4)}, x_{23}^{(4)}x_{24}^{(4)})f(x_{13}^{(4)}x_{23}^{(4)}, x_{34}^{(4)}) \\ & = f(x_{23}^{(4)}, x_{34}^{(4)})f(x_{12}^{(4)}x_{13}^{(4)}, x_{24}^{(4)}x_{34}^{(4)})f(x_{12}^{(4)}, x_{23}^{(4)}) \quad \text{in } \hat{B}_4 \end{aligned}$$

which appeared in [NT], where $\hat{B}_n$ $(n \in \mathbf{N})$ stands for the pro-finite braid group with n-strings. But we can easily show that (I)+(III) is equivalent to (I)+$(\text{III})_{\text{DR}}$.

Proposition 1.3. *Relation* (III) *implies relation* (I) .

Proof . Recall that we have a basic projection $p : \hat{P}_5^* \to \hat{P}_4^*$ which sends $x_{i,j} = x_{i,j}^{(5)} \in \hat{P}_5^*$ to $x_{i,j}^{(4)} \in \hat{P}_4^*$ $(1 \leqslant i,j \leqslant 4)$ and $x_{i,5} \in \hat{P}_5^*$ to

$1 \in \hat{P}_4^*$ $(1 \leqslant i \leqslant 5)$. It is immediate to see that (III) implies (I) because

$$p\big(f(x_{12},x_{23})f(x_{34},x_{45})f(x_{51},x_{12})f(x_{23},x_{34})f(x_{45},x_{51})\big)$$
$$= f(x_{12}^{(4)},x_{23}^{(4)})f(x_{23}^{(4)},x_{34}^{(4)}) = f(x_{12}^{(4)},x_{23}^{(4)})f(x_{23}^{(4)},x_{12}^{(4)})$$

and $\hat{P}_4^*$ is a free pro-finite group of rank 2 with generators $x_{12}^{(4)}$ and $x_{23}^{(4)}$. □

With the $G_{\mathbf{Q}}$-action on the geometric fundamental group $\pi_1(\mathbf{P}^1_{\overline{\mathbf{Q}}} - \{0,1,\infty\}, \overrightarrow{01})$ of the projective line minus 3 points (with respect to a certain tangential base point $\overrightarrow{01}$), we associate a pro-finite group homomorphism $\varphi : G_{\mathbf{Q}} \to Aut\widehat{F_2}$ (recall that $\widehat{F_2} \simeq \pi_1(\mathbf{P}^1_{\overline{\mathbf{Q}}} - \{0,1,\infty\}, \overrightarrow{01})$). It follows from Belyĭ's theorem [Be] that φ is injective. In [Ih90] and [Ih94], it was shown that $\varphi(G_{\mathbf{Q}})$ lies in $\widehat{GT}$. Now to determine whether $G_{\mathbf{Q}}$ is equal to $\widehat{GT}$ or not is a basic open problem, which is also related to a project posed by A. Grothendieck in [Gr]. Recently there have been discovered other new-type relations which $G_{\mathbf{Q}}$ satisfies in $\widehat{GT}$, for example (I′), (II′), (III′), (IV), (IV′), (V), (A_n), (K_n) for $n = 1, 2, 3, \ldots$ (see [Ih00], [LNS], [NS] and [NT]). Although they do not seem to be deduced from defining relations (I), (II) and (III) of $\widehat{GT}$, it has not been shown yet whether they are really new ones and whether they are enough to characterize $G_{\mathbf{Q}}$ as a subgroup of $\widehat{GT}$.

2. Review of the definitions of $\mathbb{\Gamma}$, GTK and GTA

Three subgroups $\mathbb{\Gamma}$, GTK and GTA of $\widehat{GT}$ were introduced in [Ih00] and [LNS]. The subgroup $\mathbb{\Gamma}$ (resp. GTK) was geometrically constructed by P. Lochak, H. Nakamura, and L. Schneps in [LNS] (resp. by Y. Ihara in [Ih00]). On the other hand, GTA was arithmetically constructed by Y. Ihara in [Ih00]. They all contain $G_{\mathbf{Q}}$, but it has not been known whether they are really proper subgroups of $\widehat{GT}$ and whether they are equal to $G_{\mathbf{Q}}$.

2.1. $\mathbb{\Gamma}$.

Definition 2.1 ([LNS]). The *new Grothendieck-Teichmüller group* $\mathbb{\Gamma}$ is the subset of $\widehat{GT}$ defined as follows:

$$\mathbb{\Gamma} := \Big\{\sigma = (\lambda, f) \in \widehat{GT} \mid (\lambda, f) \text{ satisfies (III}') \text{ and (IV) below.}\Big\}.$$

$$\begin{cases} \text{(III}') & g(x_{45},x_{51})f(x_{12},x_{23})f(x_{34},x_{45}) = f(\sigma_1\sigma_3,\sigma_2^2) \qquad \text{in } \hat{B}_5 \\ \text{(IV)} & f(\sigma_1,\sigma_2^4) = \sigma_2^{-8\Psi_2^{(0)}(\sigma)} f(\sigma_1^2,\sigma_2^2)\sigma_1^{-4\Psi_2^{(0)}(\sigma)}(\sigma_2\sigma_1)^{6\Psi_2^{(0)}(\sigma)} \text{ in } \hat{B}_3. \end{cases}$$

Here $g(x,y) \in \widehat{F_2}$ is the auxiliary parameter (depending on $\sigma \in \widehat{GT}$) satisfying $f(x,y) = g(y,x)^{-1}g(x,y)$, which was introduced in [LS]. For $n \in \mathbf{N}$, σ_i $(1 \leqslant i < n)$ stand for standard generators of $\hat{B}_n$ ([LNS]). For the definition of $\Psi_2^{(0)}(\sigma)^{\dagger}$, see §§2.2. In [LNS] and [NS], it was shown that actually $\mathbb{\Gamma}$ forms a subgroup of $\widehat{GT}$. Note that (III$'$) implies (III) ([NS]). These two relations (III$'$) and (IV) just describe the condition for elements of $\widehat{GT}$ to act (as $G_{\mathbf{Q}}$ does) on all types of pro-finite Teichmüller modular groups in a certain consistent way (for more details, see [LNS]).

2.2. *GTK*. For a natural number n, let H_n be the index n normal subgroup of $\widehat{F_2}$ which is freely generated by $n+1$ elements x^n, y, $x^{-1}yx, \cdots, x^{-(n-1)}yx^{n-1}$. For $\sigma = (\lambda, f) \in \widehat{GT}$, f belongs to H_n, since $H_n \supset [\widehat{F_2}, \widehat{F_2}]$. In [Ih99], Ihara constructed the extended Kummer 1-cocycle $\Psi_n^{(0)}(\sigma)$ for $\sigma \in \widehat{GT}$, which is the image of f by the continuous group homomorphism $H_n \to \widehat{\mathbf{Z}}$ defined by $x^n \mapsto 0$, $y \mapsto 1$, $x^{-j}yx^j \mapsto 0$ $(1 \leqslant j < n)$. We remark that especially for $\sigma \in G_{\mathbf{Q}}$, $\Psi_n^{(0)}(\sigma)$ is the Kummer 1-cocycle which is characterized by $\sigma(\sqrt[k]{n}) = \sqrt[k]{n}\zeta_k^{-\Psi_n^{(0)}(\sigma)}$ for $k \in \mathbf{N}$, where $\zeta_k = exp(\frac{2\pi i}{k})$. Suppose that $\varphi_n : H_n \to \widehat{F_2}$ is the continuous group homomorphism defined by $x^n \mapsto x$, $y \mapsto y$, $x^{-j}yx^j \mapsto 1$ $(1 \leqslant j < n)$.

Definition 2.2 ([Ih00]). The *Grothendieck-Teichmüller-Kummer group* GTK is the subset of $\widehat{GT}$ defined as follows:

$$GTK_n := \left\{ \sigma = (\lambda, f) \in \widehat{GT} \mid (\lambda, f) \text{ satisfies } (\mathrm{K}_n) \text{ below.} \right\}$$

$$GTK := \bigcap_{n \in \mathbf{N}} GTK_n$$

$$(\mathrm{K}_n) \qquad \varphi_n(f) = y^{\Psi_n^{(0)}(\sigma)} f \qquad .$$

It follows immediately from [Ih00] *Proposition 1* that GTK_n and GTK actually form subgroups of $\widehat{GT}$. Relation (K_n) just describes the condition for elements of $\widehat{GT}$ to act (as $G_{\mathbf{Q}}$ does) on H_n and $\widehat{F_2}$ consistently with two algebraic morphisms, the Kummer covering $\mathbf{P}^1_{\overline{\mathbf{Q}}} - \{0, \mu_n, \infty\} \twoheadrightarrow \mathbf{P}^1_{\overline{\mathbf{Q}}} - \{0, 1, \infty\}$ defined by $t \mapsto t^n$ and the natural inclusion $\mathbf{P}^1_{\overline{\mathbf{Q}}} - \{0, \mu_n, \infty\} \hookrightarrow \mathbf{P}^1_{\overline{\mathbf{Q}}} - \{0, 1, \infty\}$ defined by $t \mapsto t$ (for more details, see [Ih00]).

---

†The 1-cocycles $\rho_2(\sigma)$ and $\rho_3(\sigma)$ studied in [LNS], [LS], [NS] and [NT] are equal to $-\Psi_2^{(0)}(\sigma)$ and $-\Psi_3^{(0)}(\sigma)$ respectively (see [NT]).

2.3. *GTA*. By the $\widehat{F_2}^{ab}(:= \widehat{F_2}/\widehat{F_2}')$-action on $(\widehat{F_2}')^{ab} := \widehat{F_2}'/[\widehat{F_2}', \widehat{F_2}']$ induced from the conjugation $n \mapsto fnf^{-1}$ for $n \in \widehat{F_2}'$ and $f \in \widehat{F_2}$, we can regard $(\widehat{F_2}')^{ab}$ as a free A_2 $(:= \widehat{\mathbf{Z}}[[\widehat{F_2}ab]])$-module of rank 1, generated by the class $\overline{[x,y]} \in (\widehat{F_2}')^{ab}$ of $[x,y] := xyx^{-1}y^{-1} \in \widehat{F_2}'$ (for more details, see [Ih99]). Thus the action of $\sigma = (\lambda, f) \in \widehat{GT}$ on $(\widehat{F_2}')^{ab}$ induced from that on $\widehat{F_2}$ is expressed as $\sigma(\overline{[x,y]}) = B'_\sigma \cdot \overline{[x,y]}$, where $B'_\sigma \in A_2^\times$. The *adelic beta function* $B_\sigma \in A_2^\times$ was defined by $B'_\sigma = \frac{x^\lambda - 1}{x-1}\frac{y^\lambda - 1}{y-1}B_\sigma$ in [A] (for $\sigma \in G_{\mathbf{Q}}$) and [Ih99] (for $\sigma \in \widehat{GT}$). By the embedding constructed by G. W. Anderson in [A], B_σ can be regarded as a function on $(\mathbf{Q}/\mathbf{Z})^{\oplus 2}$ valued in $\widehat{\mathbf{Z}} \otimes \mathbf{Q}^{ab}$, where $\mathbf{Q}^{ab}$ stands for the maximal abelian extension field over $\mathbf{Q}$. In [Ih99] *Proposition 1.6.1*, it was shown that the adelic beta function has much analogy with the classical beta function. Especially it is remarkable that by using the 5-cycle relation (III) Ihara showed that the adelic beta function $B_\sigma(s_1, s_2)$ $(\sigma \in \widehat{GT},\ (s_1, s_2) \in (\mathbf{Q}/\mathbf{Z})^{\oplus 2})$ can be split (but not uniquely) into the following product: $B_\sigma(s_1, s_2) = \frac{\Gamma_\sigma(s_1)\Gamma_\sigma(s_2)}{\Gamma_\sigma(s_1+s_2)}$. Here the *adelic gamma function* Γ_σ is a function (uniquely determined up to a certain ambiguity) which is defined on $\mathbf{Q}/\mathbf{Z}$ and is valued in the product $\prod_{p:\text{prime}} \mathbb{W}(\overline{\mathbf{F}_p})$ of the Witt vector ring $\mathbb{W}(\overline{\mathbf{F}_p})$ of the algebraic closure of a finite field of characteristic p. In [A](i) *Corollary 8.6.3*, Anderson showed, as an analogy of Gauß' n-th multiplication formula of the classical gamma function, the n-th multiplication formula of the adelic gamma function Γ_σ (*for elements* σ *of* $G_{\mathbf{Q}}$):

$$(\mathrm{A}_n) \qquad \prod_{nc=0} \frac{\Gamma_\sigma(s+c)}{\Gamma_\sigma(c)} \cdot \frac{1}{\Gamma_\sigma(ns)} = 1 \otimes exp\big[2\pi i \cdot n\Psi_n^{(0)}(\sigma)s\big]$$

by using Deligne's theory of absolute Hodge cycles. Ihara suggested that this arithmetic relation (A_n) could be a key condition to distinguish $G_{\mathbf{Q}}$ from $\widehat{GT}$ and considered the following new subgroup of $\widehat{GT}$ containing $G_{\mathbf{Q}}$.

Definition 2.3 ([Ih00]). The *Grothendieck-Teichmüller-Anderson group* GTA is the subset of $\widehat{GT}$ defined as follows:

$$GTA_n := \Big\{\sigma = (\lambda, f) \in \widehat{GT} \ \mid \sigma \text{ satisfies } (\mathrm{A}_n).\Big\}$$
$$GTA := \bigcap_{n \in \mathbf{N}} GTA_n \ .$$

Remark 2.4. To state (A_n) independently from (I)$\sim$(III), it is better to re-formulate it as follows:

$$(\mathrm{A}_n^0) \qquad \prod_{0\leqslant k\leqslant n-1} B_\sigma(s,ks) \Big/ \prod_{nc=0} B_\sigma(c,s) = 1\otimes exp\big[2\pi i\cdot n\Psi_n^{(0)}(\sigma)s\big]$$

because the existence of Γ_σ depends on (I)$\sim$(III). Here we use $\Gamma_\sigma(0) = 1\otimes 1$ ([Ih99]Proposition 1.7.1.(i)).

It can be checked directly from the definitions of $\Psi_n^{(0)}(\sigma)$ and $\Gamma_\sigma(s)$ that GTA and GTA_n actually form subgroups of $\widehat{GT}$.

The relationship among the above three subgroups $\mathbb{I}\Gamma$, GTK and GTA has not been fully understood yet. But we remark that it was shown in [Ih99] that relation (K_n) implies $(Dlog\mathrm{A}_n)$, the logarithmic derivative of equation (A_n).

3. Main results

Theorem 3.1. *The combination of relations* (I) , (II), (IV) *and* (K_2) *imply relation* (A_2^0).

Proof . At first, we introduce a new parameter f_+ in the rank 3 free group $\widehat{F_3}$ with generators W, X and Y. Recall that the index 2 subgroup H_2 (§§2.2) of $\widehat{F_2}$ generated by xyx^{-1}, x^2 and y can be identified with $\widehat{F_3}$ by sending xyx^{-1}, x^2, y into W, X, Y respectively. Since $y^{-2\Psi_2^{(0)}(\sigma)}f$ lies on H_2, we obtain a unique element $f_+(W,X,Y)$ of $\widehat{F_3}$ such that

$$y^{-2\Psi_2^{(0)}(\sigma)}f(x,y) = f_+(xyx^{-1},x^2,y). \tag{1}$$

We note that $W^{\Psi_2^{(0)}(\sigma)}Y^{\Psi_2^{(0)}(\sigma)}f_+ \in [\widehat{F_3},\widehat{F_3}]$. It is immediate that relation (K_2) is equivalent to

$$y^{\Psi_2^{(0)}(\sigma)}f_+(xyx^{-1},x^2,y) = f_+(1,x,y)\ . \tag{2}$$

Here 1 stands for the unit element of $\widehat{F_2}$. In Lemma 4.1, we shall see that (IV) is re-expressed in terms of $f_+\in\widehat{F_3}$ as follows:

$$f_+(W,X,Y)f_+(X^{-1}W^{-1}Y^{-1},Y,X) = 1\ . \tag{3}$$

In Lemma 4.2, we deduce the following equation from (I), (II) and (3):

$$Y^m f_+(W,X,Y)X^m f_+(Z,W,X)W^m f_+(Y,Z,W) \\ Z^m f_+(X,Y,Z) = 1 \qquad \text{for} \quad WXYZ = 1 \tag{4}$$

Translating (A_2^0) into the terms of $A_2 = \widehat{\mathbf{Z}}[[\widehat{F_2}^{ab}]]$ via Anderson's embedding (see §2.3), we get

$$(\mathrm{A}_2^0) \quad \frac{B_\sigma(s,s)}{B_\sigma(\frac{1}{2},s)} = 1 \otimes exp\big[2\pi i \cdot \Psi_2^{(0)}(\sigma)s\big]$$
$$\Longleftrightarrow B_\sigma\{\mathbf{x},\mathbf{x}\} = \mathbf{x}^{-2\Psi_2^{(0)}(\sigma)} B_\sigma\{-1,\mathbf{x}\}.$$

Here $B_\sigma\{\mathbf{x},\mathbf{x}\}$ (resp. $B_\sigma\{-1,\mathbf{x}\}$) stands for the image of $B_\sigma \in A_2$ by the map $A_2 \to A_2$ induced from $\mathbf{x} \to \mathbf{x}$, $\mathbf{y} \to \mathbf{x}$ (resp. $\mathbf{x} \to -1$, $\mathbf{y} \to \mathbf{x}$), where boldfaces $\mathbf{x}$, $\mathbf{y}$ stand for the images in A_2 of standard generators x, y of $\widehat{F_2}$. By properties [II]$\sim$[IV] of [Ih99]§2.3 which were deduced from (I) and (II), we get

$$(\mathrm{A}_2^0) \Longleftrightarrow \frac{\mathbf{x}^\lambda + 1}{\mathbf{x}^m(\mathbf{x}+1)} B_\sigma\{\mathbf{x}, \frac{1}{\mathbf{x}^2}\} = \mathbf{x}^{-2\Psi_2^{(0)}(\sigma)} B_\sigma\{-1,\mathbf{x}\}.$$

By [Ih99]Proposition 2.2.2,

$$(\mathrm{A}_2^0) \Longleftrightarrow \frac{\mathbf{x}^\lambda + 1}{\mathbf{x}^m(\mathbf{x}+1)} \Big\{1 - (\mathbf{x}-1)\frac{df}{dx}(\mathbf{x}, \frac{1}{\mathbf{x}^2})\Big\} = \mathbf{x}^{-2\Psi_2^{(0)}(\sigma)} \Big\{1 + 2\frac{df}{dx}(-1,\mathbf{x})\Big\}.$$

Here we denote the image of $\frac{df}{dx} \in \Lambda_2 := \widehat{\mathbf{Z}}[[\widehat{F_2}]]$ by the map $\Lambda_2 \to A_2$ induced from $\mathbf{x} \mapsto -1$, $\mathbf{y} \mapsto \mathbf{x}$ by $\frac{df}{dx}(-1,\mathbf{x})$ (for the definition of $\frac{df}{dx} \in \Lambda_2$, see §4). For $\alpha \in \Lambda_2$ and $\mathbf{u}$, $\mathbf{v} \in \widehat{F_2}^{ab}$, we denote the image of α by the map $\Lambda_2 \to A_2$ induced from $x \to \mathbf{u}$, $y \to \mathbf{v}$ by $\alpha(\mathbf{u},\mathbf{v}) \in A_2$. By Lemma 4.3, which will be deduced from (1), we get

$$\begin{aligned}(\mathrm{A}_2^0) \Longleftrightarrow & \frac{\mathbf{x}^\lambda + 1}{\mathbf{x}^m(\mathbf{x}+1)} \Big[1 - (\mathbf{x}-1)\Big\{\mathbf{x}^{-2\Psi_2^{(0)}(\sigma)} \frac{df_+}{dW}(\frac{1}{\mathbf{x}^2}, \mathbf{x}^2, \frac{1}{\mathbf{x}^2}) \cdot (1 - \frac{1}{\mathbf{x}^2}) \\ & + \frac{df_+}{dX}(\frac{1}{\mathbf{x}^2}, \mathbf{x}^2, \frac{1}{\mathbf{x}^2}) \cdot 2\mathbf{x}\Big\}\Big] \\ & - \mathbf{x}^{-2\Psi_2^{(0)}(\sigma)} \Big[1 + 2\mathbf{x}^{2\Psi_2^{(0)}(\sigma)} \Big\{(1-\mathbf{x})\frac{df_+}{dW}(\mathbf{x},1,\mathbf{x}) - 2\frac{df_+}{dX}(\mathbf{x},1,\mathbf{x})\Big\}\Big] = 0\ .\end{aligned}$$

For $\frac{df_+}{dW}$ and $\frac{df_+}{dX}$, see §4. By Lemma 4.4, which will be deduced from (2) (namely (K_2)), we get

$(\mathrm{A}_2^0) \Longleftrightarrow$

(5)
$$\frac{\mathbf{x}^\lambda+1}{\mathbf{x}^m(\mathbf{x}+1)}\Big\{1-\mathbf{x}^{-2\Psi_2^{(0)}(\sigma)}\cdot(\mathbf{x}-1)\frac{df_+}{dX}(1,\mathbf{x},\frac{1}{\mathbf{x}^2})\Big\}$$
$$-\mathbf{x}^{-2\Psi_2^{(0)}(\sigma)}\Big[1+2\mathbf{x}^{2\Psi_2^{(0)}(\sigma)}\Big\{(1-\mathbf{x})\frac{df_+}{dW}(\mathbf{x},1,\mathbf{x})-2\frac{df_+}{dX}(\mathbf{x},1,\mathbf{x})\Big\}\Big]=0\ .$$

By formulae (6)$\sim$(14) in §4, which will be deduced from (I), (II), (III) and (K_2), we get the following

$$\begin{aligned}
\frac{dF_+}{dW}(\mathbf{x},1,\mathbf{x}) =&\mathbf{x}^{-2\Psi_2^{(0)}-1}\cdot\frac{dF_+}{dW}(\frac{1}{\mathbf{x}^2},\mathbf{x},1)\\
\frac{dF_+}{dX}(\mathbf{x},1,\mathbf{x}) =&0\\
\frac{dF_+}{dX}(\frac{1}{\mathbf{x}^2},\mathbf{x},1) =&\frac{\mathbf{x}+1}{\mathbf{x}^2}\frac{dF_+}{dW}(\frac{1}{\mathbf{x}^2},\mathbf{x},1)+\frac{\mathbf{x}^{2\Psi_2^{(0)}(\sigma)}-1}{\mathbf{x}-1}\\
\frac{dF_+}{dY}(\mathbf{x},\frac{1}{\mathbf{x}^2},\mathbf{x}) =&-\mathbf{x}^{1-2\Psi_2^{(0)}(\sigma)}\frac{dF_+}{dW}(1,\mathbf{x},\frac{1}{\mathbf{x}^2})-\frac{\mathbf{x}^{2\Psi_2^{(0)}(\sigma)}-1}{\mathbf{x}^{2\Psi_2^{(0)}(\sigma)}(\mathbf{x}-1)}\\
\frac{dF_+}{dY}(\frac{1}{\mathbf{x}^2},\mathbf{x},1) =&\frac{dF_+}{dW}(\frac{1}{\mathbf{x}^2},\mathbf{x},1)
\end{aligned}$$

By substituting $(W,X,Y)=(1,\mathbf{x},\frac{1}{\mathbf{x}^2})$ and above formulae to (15) in Lemma 4.6, we obtain equation (5), which implies the validity of the Anderson duplication formulae (A_2^0). □

Therefore the combination of geometric relations (I), (II), (IV) and (K_2) implies arithmetic relation (A_2^0). Next we will consider three subgroups $\mathbb{\Gamma}$, GTK and GTA.

Lemma 3.2. *For $n,m\in\mathbf{N}$, the combination of relations* (A_n) *and* (A_m) *implies* (A_{nm}).

Proof . Let $n\in\mathbf{N}$. By the slightly more detailed discussion on [Ih00], we find that relation (A_n) implies

$$\Psi_{na}^{(0)}(\sigma)=\Psi_n^{(0)}(\sigma)+\Psi_a^{(0)}(\sigma)\qquad\text{for }\ \forall a\in\mathbf{N}.$$

Therefore the combination of (A_n) and (A_m) implies the following:

$$\begin{aligned}\frac{\prod_{k=0}^{nm-1}\Gamma_\sigma(s+\frac{k}{nm})}{\Gamma_\sigma(nms)\prod_{k=0}^{nm-1}\Gamma_\sigma(\frac{k}{nm})} &= 1\otimes exp\big[2\pi i\cdot nm\{\Psi_n^{(0)}(\sigma)+\Psi_m^{(0)}(\sigma)\}s\big]\\ &= 1\otimes exp\big[2\pi i\cdot nm\Psi_{nm}^{(0)}(\sigma)s\big] \quad \square\end{aligned}$$

We also remark the following formulae, but it is not required to prove Theorem 3.4 below.

Lemma 3.3. *For $n,m\in\mathbf{N}$, the combination of relations* (K_n) *and* (K_m) *implies* (K_{nm}).

Proof . Let $n\in\mathbf{N}$. As in the same way of the case of Lemma 3.2, we get that relation (K_n) implies

$$\Psi_{na}^{(0)}(\sigma)=\Psi_n^{(0)}(\sigma)+\Psi_a^{(0)}(\sigma) \qquad \text{for } \forall a\in\mathbf{N}.$$

Therefore the combination of (K_n) and (K_m) implies the following:

$$\varphi_{nm}(f)=\varphi_n(y^{\Psi_m^{(0)}(\sigma)})\varphi_n(f)=y^{\Psi_n^{(0)}(\sigma)+\Psi_m^{(0)}(\sigma)}f=y^{\Psi_{nm}^{(0)}(\sigma)}f. \quad \square$$

Theorem 3.4. *$GTK\cap\mathbb{\Gamma}\subseteq GTA_{2^\infty}$, where $GTA_{2^\infty}:=\bigcap_{n\in\mathbf{N}}GTA_{2^n}$.*

Proof . We note that one relation (A_2) implies infinite ones (A_{2^n}) for $n=1,2,3,\cdots$. Therefore, by combining Lemma 3.2 with Theorem 3.1, we get the claim. $\square$

Thus we get a relationship among the arithmetic subgroup GTA_{2^∞} and the geometric subgroups GTK and $\mathbb{\Gamma}$. Relations (3) and (4) describe the D_4 (not D_2) symmetry of $\mathbf{P}^1-\{0,\pm1,\infty\}$. Since they were essential in the proof of Theorem 3.1, it does not look possible, at least to the author, to deduce Anderson's n-th multiplication formula ($n\geqslant 5$) from the D_n-symmetry of $\mathbf{P}^1-\{0,\mu_n,\infty\}$. H. Tsunogai suggest to the author the possibility of deducing (A_3) from the $\mathfrak{S}_4$-symmetry of $\mathbf{P}^1-\{0,\mu_3,\infty\}$. Still we would expect the geometric interpretation of arithmetic Anderson's multiplication formulae which is originally of arithmetic nature to be a key to distinguish $G_{\mathbf{Q}}$ from $\widehat{GT}$.

4. Miscellaneous lemmas

We present auxiliary lemmas which were required in the proof of Theorem 3.1. Here we follow the notation in [Ih99].

Lemma 4.1. *Relation* (IV) *is equivalent to* (3).

Proof . Let $\widehat{F_2} \,/\, \langle\langle z^2\rangle\rangle$ denote the quotient pro-finite group of $\widehat{F_2}$ by the normal closure $\langle\langle z^2\rangle\rangle$ of $z^2 = (xy)^{-2}$. By [NS] Theorem 2.2.(22),

$$\begin{aligned}
\text{(IV)} &\Longleftrightarrow y^{4\Psi_2^{(0)}(\sigma)} f(x,y^2) x^{4\Psi_2^{(0)}(\sigma)} f(y,x^2) \equiv 1 && \text{in} \quad \widehat{F_2} \,/\, \langle\langle z^2\rangle\rangle \\
&\Longleftrightarrow f_+(xy^2x^{-1},x^2,y^2) f_+(yx^2y^{-1},y^2,x^2) \equiv 1 && \text{in} \quad \widehat{F_2} \,/\, \langle\langle z^2\rangle\rangle \\
&\Longleftrightarrow f_+(xy^2x^{-1},x^2,y^2) f_+(x^{-1}y^{-2}x^{-1}y^{-2},y^2,x^2) \equiv 1 && \text{in} \quad \widehat{F_2} \,/\, \langle\langle z^2\rangle\rangle.
\end{aligned}$$

On the other hand, since xy^2x^{-1},x^2 and y^2 generate a free pro-finite subgroup of rank 3 in $\widehat{F_2}$,

$$(3) \Longleftrightarrow f_+(xy^2x^{-1},x^2,y^2) f_+(x^{-1}y^{-2}x^{-1}y^{-2},y^2,x^2) = 1 \qquad \text{in} \quad \widehat{F_2}.$$

From the argument in the proof of [NS] Theorem 2.2, the quotient classes of x, y^2 in $\widehat{F_2} \,/\, \langle\langle z^2\rangle\rangle$ generate a free pro-finite subgroup of rank 2. Thus the images of xy^2x^{-1}, x^2, y^2 in $\widehat{F_2} \,/\, \langle\langle z^2\rangle\rangle$ generate a free pro-finite subgroup of rank 3, from which it follows that (IV) is equivalent to (3). □

Lemma 4.2. *The combination of relations* (I), (II) *and* (3) *implies* (4).

Proof . By the permutation $x \mapsto y$, $y \mapsto x$, $z \mapsto x^{-1}y^{-1}$, (II) is re-expressed as follows:

$$f(y,x)y^{m+1}f(z,y)z^{m+1}f(x,z)x^{m+1} = 1.$$

By combining this with (II) and applying (I), we get

$$x^{2m+1}f(x,z)^{-1}z^m f(y,z)y^{2m+1}f(y,z)^{-1}z^{m+1}f(x,z) = 1.$$

Taking the image of the last equation by the continuous homomorphism induced from $x \mapsto y^{-1}x^{-1}$, $y \mapsto x$, $z \mapsto y$, we get

$$\begin{aligned}
&(y^{-1}x^{-1})^{2m+1} f_+(y^{-1}x^{-1}yxy, y^{-1}x^{-1}y^{-1}x^{-1}, y)^{-1} y^m f_+(xyx^{-1},x^2,y) x^{2m+1} \\
&\quad f_+(xyx^{-1},x^2,y)^{-1} y^{m+1} f_+(y^{-1}x^{-1}yxy, y^{-1}x^{-1}y^{-1}x^{-1}, y) = 1.
\end{aligned}$$

By (3),

$$\begin{aligned}
&(y^{-1}x^{-1}y^{-1}x^{-1})^m \cdot f_+(x^2,y,y^{-1}x^{-1}y^{-1}x^{-1}) \cdot y^m \cdot f_+(xyx^{-1},x^2,y) \cdot (x^2)^m \\
&\quad f_+(y^{-1}x^{-1}y^{-1}x^{-1}, xyx^{-1}, x^2) \cdot (xyx^{-1})^m \cdot f_+(y, y^{-1}x^{-1}y^{-1}x^{-1}, xyx^{-1}) = 1.
\end{aligned}$$

Since the subgroup generated by x^2, y and $y^{-1}x^{-1}y^{-1}x^{-1}$ is equal to the one generated by x^2, y and xyx^{-1}, it is a free pro-finite subgroup of rank 3 in $\widehat{F_2}$, which implies (4). □

By Anderson's theorem (see [Ih99] Theorem A.1.), the element $f \in \widehat{F_2}$ can be written uniquely in $\Lambda_2 := \widehat{\mathbf{Z}}[[\widehat{F_2}]]$ as follows:

$$f(x,y) = 1 + \frac{df}{dx} \cdot (x-1) + \frac{df}{dy} \cdot (y-1). \tag{6}$$

Put $\Lambda_3 := \widehat{\mathbf{Z}}[[\widehat{F_3}]]$. Similarly the element $f_+ \in \widehat{F_3} = \langle W, X, Y \rangle^\wedge$ in §3 can be written uniquely in Λ_3 as follows:

$$f_+(W,X,Y) = 1 + \frac{df_+}{dW} \cdot (W-1) + \frac{df_+}{dX} \cdot (X-1) + \frac{df_+}{dY} \cdot (Y-1), \tag{7}$$

where $\frac{df_+}{dW}, \frac{df_+}{dX}, \frac{df_+}{dY} \in \Lambda_3$.

Lemma 4.3.

$$\begin{aligned} \frac{df}{dx}(x,y) =& y^{2\Psi_2^{(0)}(\sigma)} \frac{df_+}{dW}(xyx^{-1}, x^2, y) \cdot (1 - xyx^{-1}) \\ &+ \frac{df_+}{dX}(xyx^{-1}, x^2, y) \cdot 2x. \end{aligned} \tag{8}$$

$$\begin{aligned} \frac{df}{dy}(x,y) =& \frac{y^{2\Psi_2^{(0)}(\sigma)} - 1}{y-1} + y^{2\Psi_2^{(0)}(\sigma)} \Big\{ \frac{df_+}{dW}(xyx^{-1}, x^2, y) \cdot x \\ &+ \frac{df_+}{dY}(xyx^{-1}, x^2, y) \Big\}. \end{aligned} \tag{9}$$

Here for $\beta \in \Lambda_3$ and $a, b, c \in \widehat{F_2}$, we denote the image of β by the map $\Lambda_3 \to \Lambda_2$ induced from $W \mapsto a$, $X \mapsto b$, $Y \mapsto c$ by $\beta(a,b,c) \in \Lambda_2$.

Proof . It follows from (1) by a direct calculation. □

Lemma 4.4.

$$\begin{aligned} \frac{df_+}{dX}(1,x,y) =& y^{\Psi_2^{(0)}(\sigma)} \frac{df_+}{dW}(xyx^{-1}, x^2, y) \cdot (1 - xyx^{-1}) \\ &+ y^{-\Psi_2^{(0)}(\sigma)} \frac{df_+}{dX}(xyx^{-1}, x^2, y) \cdot 2x \quad . \end{aligned} \tag{10}$$

$$\begin{aligned} \frac{df_+}{dY}(1,x,y) =& \frac{y^{\Psi_2^{(0)}(\sigma)} - 1}{y-1} + y^{\Psi_2^{(0)}(\sigma)} \Big\{ \frac{df_+}{dW}(xyx^{-1}, x^2, y) \cdot x \\ &+ \frac{df_+}{dY}(xyx^{-1}, x^2, y) \Big\}. \end{aligned} \tag{11}$$

Proof . It follows from (2) by a direct calculation. □

Lemma 4.5.

$$\frac{df_+}{dW}(W,X,Y) = f_+(W,X,Y)\frac{df_+}{dW}(X^{-1}W^{-1}Y^{-1},Y,X)X^{-1}W^{-1}. \tag{12}$$

$$\begin{aligned}\frac{df_+}{dX}(W,X,Y) = f_+(W,X,Y)\Big\{&\frac{df_+}{dW}(X^{-1}W^{-1}Y^{-1},Y,X)\cdot X^{-1}\\ &-\frac{df_+}{dY}(X^{-1}W^{-1}Y^{-1},Y,X)\Big\}.\end{aligned} \tag{13}$$

$$\begin{aligned}\frac{df_+}{dY}(W,X,Y) = f_+(W,X,Y)\Big\{&\frac{df_+}{dW}(X^{-1}W^{-1}Y^{-1},Y,X)\cdot\\ &X^{-1}W^{-1}Y^{-1}+\frac{df_+}{dX}(X^{-1}W^{-1}Y^{-1},Y,X)\Big\}.\end{aligned} \tag{14}$$

Proof . It follows from (3) by a direct calculation. □

Lemma 4.6.

(15)

$$\begin{aligned}&Y^m\frac{df_+}{dW}(W,X,Y)\\ &\quad+Y^m f_+(W,X,Y)X^m\Big\{\frac{df_+}{dW}(Z,W,X)\cdot(-Z)+\frac{df_+}{dX}(Z,W,X)\Big\}\\ &\quad+Y^m f_+(W,X,Y)X^m f_+(Z,W,X)\frac{W^m-1}{W-1}\\ &\quad+Y^m f_+(W,X,Y)X^m f_+(Z,W,X)W^m\Big\{\frac{df_+}{dX}(Y,Z,W)\cdot(-Z)+\frac{df_+}{dY}(Y,Z,W)\Big\}\\ &\quad+Y^m f_+(W,X,Y)X^m f_+(Z,W,X)W^m f_+(Y,Z,W)\frac{Z^m-1}{Z-1}\cdot(-Z)\\ &\quad+Y^m f_+(W,X,Y)X^m f_+(Z,W,X)W^m f_+(Y,Z,W)Z^m\cdot\\ &\qquad\Big\{\frac{df_+}{dY}(X,Y,Z)\cdot(-Z)\Big\}=0 \qquad \textit{for} \qquad WXYZ=1.\end{aligned}$$

Proof . It follows from (4) by a direct calculation. □

References

[A] Anderson, G. W., (i) The hyperadelic gamma function: a précis, Galois representations and arithmetic algebraic geometry (Kyoto, 1985/Tokyo, 1986), 1–19, Adv. Stud. Pure Math., 12, North-Holland, Amsterdam, 1987,
(ii) The hyperadelic gamma function, Invent. Math. 95 (1989), no. 1, 63–131.

[Be] Belyĭ, G. V., Galois extensions of a maximal cyclotomic field, (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 43 (1979), no. 2, 267–276, 479.

[Dr] Drinfel'd, V. G., On quasitriangular quasi-Hopf algebras and a group closely connected with $\mathrm{Gal}(\overline{Q}/Q)$, Leningrad Math. J. 2 (1991), no. 4, 829–860.

[F] Furusho, H., Geometric and arithmetic subgroups of the Grothendieck-Teichmüller group, Master's thesis (in Japanese), RIMS, Kyoto University, March 2000.

[Gr] Grothendieck, A., Esquisse d'un programme, in Geometric Galois actions, 1, 5–48, edited by Leila Schneps and Pierre Lochak, London Mathematical Society Lecture Note Series, 242, Cambridge University Press, Cambridge, 1997.

[Ih90] Ihara, Y., Braids, Galois groups, and some arithmetic functions, Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990), 99–120, Math. Soc. Japan, Tokyo, 1991.

[Ih91] ——— Automorphisms of pure sphere braid groups and Galois representations, The Grothendieck Festschrift, Vol. II, 353–373, Progr. Math., 87, Birkhäuser Boston, Boston, MA, 1990.

[Ih94] ——— On the embedding of $\mathrm{Gal}(\overline{Q}/Q)$ into $\widehat{\mathrm{GT}}$, London Math. Soc. Lecture Note Ser., 200, The Grothendieck theory of dessins d'enfants (Luminy, 1993), 289–321, Cambridge Univ. Press, Cambridge, 1994.

[Ih99] ——— On beta and gamma functions associated with the Grothendieck-Teichmüller groups, Aspects of Galois theory, 144–179, London Math. Soc. Lecture Note Ser., 256, Cambridge Univ. Press, Cambridge, 1999.

[Ih00] ——— On beta and gamma functions associated with the Grothendieck-Teichmüller group II, J. Reine Angew. Math.527 (2000), 1–11.

[LNS] Lochak, P., Nakamura, H., Schneps, L., On a new version of the Grothendieck-Teichmüller group, C. R. Acad. Sci. Paris Sér. I Math. 325 (1997), no. 1, 11–16.

[LS] Lochak, P., Schneps, L., A cohomological interpretation of the Grothendieck-Teichmüller group, With an appendix by C. Scheiderer, Invent. Math. 127 (1997), no. 3, 571–600.

[NS] Nakamura, H., Schneps, L., On a subgroup of the Grothendieck-Teichmüller group acting on the tower of profinite Teichmüller modular groups, Invent. Math. 141 (2000), no. 3, 503–560.

[NT] Nakamura, H., Tsunogai, H., Harmonic and equianharmonic equations in the Grothendieck-Teichmüller group, Forum Math. (to appear).

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502 Japan

E-mail address: `furusho@kurims.kyoto-u.ac.jp`