

RIMS-1675

Folder complexes and multiflow combinatorial dualities

By

Hiroshi HIRAI

July 2009



京都大学 数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

Folder complexes and multiflow combinatorial dualities

Hiroshi HIRAI

Research Institute for Mathematical Sciences,
Kyoto University, Kyoto 606-8502, Japan
hirai@kurims.kyoto-u.ac.jp

July 2009

Abstract

In multiflow maximization problems, there are several combinatorial duality relations, such as Ford-Fulkerson's max-flow min-cut theorem for single commodity flows, Hu's max-biflow min-cut theorem for two-commodity flows, Lovász-Cherkassky duality theorem for free multiflows, and so on. In this paper, we provide a unified framework for such multiflow combinatorial dualities by using the notion of a *folder complex*, which is a certain 2-dimensional polyhedral complex introduced by Chepoi. We show that for a nonnegative weight μ on terminal set, μ -weighted maximum multiflow problem admits a combinatorial duality relation if and only if μ is represented by distances among certain subsets in a folder complex, and show that the corresponding combinatorial dual problem is a discrete location problem on the graph of the folder complex. This extends a result of Karzanov for the case of metric weights. Also we give a special optimality criterion to such combinatorial dual problems and give a characterization of such weights in terms of the combinatorial dimension.

1 Introduction

Let G be an undirected graph with nonnegative edge capacity $c : EG \rightarrow \mathbf{R}_+$. Let $S \subseteq VG$ be a set of terminals. A path P in G is called an S -path if its ends belong to distinct terminals. A *multiflow* is a pair (\mathcal{P}, λ) of a set \mathcal{P} of S -paths and its nonnegative flow-value function $\lambda : \mathcal{P} \rightarrow \mathbf{R}_+$ satisfying capacity constraint $\sum_{P \in \mathcal{P}: e \in P} \lambda(P) \leq c(e)$ for $e \in EG$. For a nonnegative weight $\mu : \binom{S}{2} \rightarrow \mathbf{R}_+$ on the set of pairs of terminals, the μ -weighted maximum multiflow problem (μ -problem for short) is formulated as

$$(1.1) \quad \begin{aligned} & \text{Maximize} && \sum_{P \in \mathcal{P}} \mu(s_P, t_P) \lambda(P) \\ & \text{subject to} && (\mathcal{P}, \lambda) : \text{multiflow for } (G, c; S), \end{aligned}$$

where s_P, t_P denote the ends of P . The μ -problem (1.1) is a linear problem. So there is a linearly programming duality relation. However, for a special weight μ , the μ -problem admits a *combinatorial duality relation*. For example, suppose that S is a 2-set $\{s, t\}$. Then the μ -problem is the maximum flow problem. The max-flow min-cut theorem, due to Ford-Fulkerson [8], says that the maximum flow value is equal to the minimum s - t cut value. Suppose that S is a four-set $\{s, t, s', t'\}$, and $\mu(s, t) = \mu(s', t') = 1$ and other weights are zero. Then the μ -problem is two-commodity flow maximization. Hu [14] proved that the maximum multiflow value is equal to the minimum of ss' - tt' mincut value and st' - ts' mincut value. Suppose that μ is all-one weight. Then Lovász [23] and Cherkassky [4] proved that the maximum multiflow value is equal to the half of the sum of t - $S \setminus t$ mincut values over $t \in S$.

A remarkable result by Karzanov [17] completely characterized *metric*-weights μ admitting such a combinatorial duality relation. We shall describe it. Here μ is called a *metric* if μ satisfies the triangle inequality. An undirected graph Γ is called a *frame* if

- (i) Γ is bipartite,
- (ii) Γ has no isometric cycle of length greater than four, and
- (iii) Γ is *orientable* in the sense that Γ has an orientation with the property that, for each 4-cycle C , nonadjacent edges have opposite directions with respect to a cyclic ordering of C ; see the left of Figure 4.

We denote the shortest path metric of Γ by d_Γ .

Theorem 1.1 ([17]). *Let μ be a metric on a finite set S . Suppose that there exist a frame Γ and a map $\phi : S \rightarrow V\Gamma$ such that*

$$\mu(s, t) = d_\Gamma(\phi(s), \phi(t)) \quad (s, t \in S).$$

Then, for any capacitated graph (G, c) having S as terminal set, the maximum value of the μ -problem (1.1) is equal to the minimum value of the following discrete location problem on Γ :

$$(1.2) \quad \begin{array}{ll} \text{Minimize} & \sum_{xy \in EG} c(xy) d_\Gamma(\rho(x), \rho(y)) \\ \text{subject to} & \rho : VG \rightarrow V\Gamma, \\ & \rho(s) = \phi(s) \quad (s \in S). \end{array}$$

(Karzanov [17] presented this theorem in the dual form, i.e., the *minimum 0-extension problem* on a frame is solvable by its *metric relaxation*, which is the LP-dual to the metric-weighted maximum multiflow problem.) Moreover, if a rational metric μ cannot be represented by (a dilation of) a submetric of a frame, then there is no such a combinatorial duality relation [18]. From the point of the view in Theorem 1.1, Lovász-Cherkassky duality relation can be understood as follows. Let μ be the all-one distance on S . Then 2μ is represented as the distance among the leaves of a star consisting of $\#S$ edges. A star is clearly a frame. Then (1.2) coincides with the problem of finding t - $S \setminus t$ mincut for each $s \in S$.

Unfortunately, this theorem cannot be applicable to μ -problems for *nonmetric* weights μ , such as two-commodity flows. Our previous paper [11] partially extended Theorems 1.1 to a nonmetric version. However this result needs tedious calculations for the tight span and its subdivision, which cause difficulty and inconvenience for a further combinatorial study to μ -problems (1.1). The main contribution (Theorem 2.1) of this paper is a natural extension of Theorem 1.1 for general weights, which provides a useful and flexible framework for multiflow combinatorial dualities. Also this result is an important step of the complete classification of weights μ having finite fractionality in the next paper [13].

To describe our result, we employ the notion of a *folder complex* (an *F-complex* for short) introduced by Chepoi [3, Section 7]. Roughly speaking, an F-complex is a 2-dimensional polyhedral complex obtained by *filling a 2-dimensional cell into each 4-cycle of a frame*. In Section 2, we describe basic definitions and the main result. A key concept that we newly introduce in this paper is the notion of a *normal set* in an F-complex. Our main result (Theorem 2.1) says that if weight μ can be realized by the distances among normal sets in an F-complex under l_1 -metrization, then the μ -problem

has a combinatorial dual problem similar to (1.2). We also give illustrative examples for multiflow combinatorial dualities by F-complexes, including Karzanov-Lomonosov duality relation for anticlique-bipartite commodity flows and seemingly new combinatorial duality relation for a *weighted version* of two-commodity flows.

Our main result immediately follows from two properties: one is the Helly property of normal sets, and the other one is a decomposition property of finite submetrics on an F-complex. This connection to Helly property sheds a new insight on the multiflow duality. To prove them, in Section 3, we reveal several intriguing properties of shortest paths connecting normal sets in an F-complex, which are previously known for shortest paths connecting vertices of a frame. These properties will also play key roles in the next paper [13]. In Section 4.2, we give a special optimality criterion to combinatorial dual problems, which extends one for the classical tree location problems. Also we verify that weight μ has such an F-complex realization if and only if μ has combinatorial dimension most 2. This result has already been suggested by the previous paper [11].

Notation. Let \mathbf{R} and \mathbf{R}_+ be the sets of reals and nonnegative reals, respectively. The set of functions from a set V to \mathbf{R} is denoted by \mathbf{R}^V . For $p, q \in \mathbf{R}^V$, $p \leq q$ means $p(x) \leq q(x)$ for all $x \in V$.

A function $\mu : S \times S \rightarrow \mathbf{R}$ is called a *distance* on S if $\mu(s, t) = \mu(t, s) \geq \mu(s, s) = 0$ for $s, t \in S$. We regard a nonnegative weight on terminals as a distance. A distance d on S is called a *metric* if d satisfies the triangle inequality $d(x, z) \leq d(x, y) + d(y, z)$ ($x, y, z \in S$). For a metric d on S , the minimum distance between two subsets A and B is denoted by $d(A, B)$, i.e.,

$$d(A, B) = \inf_{x \in A, y \in B} d(x, y).$$

We simply denote $d(\{x\}, A)$ by $d(x, A)$. For a subset $A \subseteq S$ and a nonnegative real r , the *ball* $B(A, r)$ around A of radius r is defined as

$$B(A, r) = \{x \in S \mid d(A, x) \leq r\}.$$

For an undirected graph G , the vertex set and the edge set are denoted by VG and EG , respectively. An edge joining $x, y \in VG$ is simply denoted by xy or yx . Let d_G denote the shortest path metric on VG with some specified edge-length. If no edge length is specified, then d_G means the shortest path metric by unit edge-length. A subgraph G' of G is called *isometric* if $d_{G'}(x, y) = d_G(x, y)$ for $x, y \in VG'$.

Let $\mathcal{F} \subseteq 2^S$ be a set of subsets. We say “ \mathcal{F} has the Helly property” if every subset $\mathcal{F}' \subseteq \mathcal{F}$ having pairwise nonempty intersection

$$A \cap B \neq \emptyset \quad (A, B \in \mathcal{F}')$$

has nonempty intersection

$$\bigcap_{A \in \mathcal{F}'} A \neq \emptyset.$$

A *polyhedral complex* Δ is a set of polyhedra in some Euclidean space such that every face of $\sigma \in \Delta$ belongs to Δ , and the nonempty intersection of $\sigma, \sigma' \in \Delta$ is a face of both σ and σ' . Let $|\Delta|$ denote the underlying set $\bigcup_{\sigma \in \Delta} \sigma$. A member in Δ is called a *cell*. A k -dimensional cell is simply called a *k-cell*. A 1-cell is also called an *edge*. If an edge has ends p, q , then it is also denoted by pq . A k -dimensional polyhedral complex is simply called a *k-complex*. For our applications to the multiflow theory, we always assume that Δ is *finite*.




Figure 1: A square and a triangle equipped with the l_1 -distance




Figure 2: A folder

2 Definitions, results, and examples

Let Δ be a (finite) 2-complex satisfying the following property:

(2.1) (i) For some positive rational κ , each maximal cell σ has an injective continuous map $\varphi_\sigma : \sigma \rightarrow \mathbf{R}^2$ such that the image $\varphi_\sigma(\sigma)$ is

- (a) the convex hull of $\{(0, 0), (\kappa, 0), (0, \kappa), (\kappa, \kappa)\}$,
- (b) the convex hull of $\{(0, 0), (\kappa, 0), (\kappa, \kappa)\}$, or
- (c) the segment $[(0, 0), (\kappa, 0)]$, and

the image of a face of σ is a face of $\varphi_\sigma(\sigma)$.

(ii) If $\sigma \cap \sigma' \neq \emptyset$ for two maximal cells σ, σ' , then $\varphi_{\sigma'} \circ \varphi_\sigma^{-1}$ is an isometry from $\varphi_\sigma(\sigma \cap \sigma')$ to $\varphi_{\sigma'}(\sigma \cap \sigma')$ in the Euclidean distance.

By condition (ii), if 2-cells σ, σ' share a common edge e , then $\varphi_\sigma(e) = [(0, 0), (\kappa, \kappa)]$ implies $\varphi_{\sigma'}(e) = [(0, 0), (\kappa, \kappa)]$. An edge e of Δ is called a *hypotenuse* if $\varphi_\sigma(e) = [(0, 0), (\kappa, \kappa)]$ for a 2-cell σ containing e . Other edge is called a *leg*. In particular, a maximal 1-cell is a leg. κ is called the *leg-length*. A 2-cell σ is called a *square* if its image by φ_σ is (a), and is called a *triangle* if its image by φ_σ is (b). Namely Δ is obtained by gluing squares and isosceles right triangles along same type of edges. A *folder* of Δ is a square or the union of all triangles sharing one common hypotenuse; see Figure 2.

A more combinatorial and abstract approach is often useful. We can regard Δ as a pair of a simple graph Γ_0 and a subset \mathcal{C} of its chordless cycles such that




Figure 3: (a) violation of the flag condition, and (b) violation of the local convexity

- (2.2) (i) the edge set of Γ_0 is partitioned into two types of edges, called *legs* and *hypotenuses*, and
- (ii) each member of \mathcal{C} is
- (a) a 3-cycle (a *triangle*) consisting of one hypotenuse and two legs, or
 - (b) a 4-cycle (a *square*) consisting of four legs,
- and every hypotenuse belongs to some triangle.

Indeed, we can take Γ_0 as the 1-skeleton graph of Δ , and take \mathcal{C} as boundary cycles of 2-cells. Conversely, from $(\Gamma_0; \mathcal{C})$ with property (2.2) and a positive rational κ , we can construct a 2-complex Δ with property (2.1).

We can endow $|\Delta|$ with the l_p -metric by the following way. For each cell σ and a path P in σ , the l_p -length of P is defined by the l_p -length of $\varphi_{\sigma'}(P)$ in (\mathbf{R}^2, l_p) for a maximal cell σ' containing σ (well-defined by (ii)). Then we can define the l_p -length of a path P in $|\Delta|$ by the sum of the l_p -length of $P \cap \sigma^\circ$ over all cells σ , where σ° is the relative interior of σ . Thus we can define the metric d_{Δ, l_p} on $|\Delta|$ by defining $d_{\Delta, l_p}(p, q)$ to be the infimum of the lengths of all paths connecting p, q in $|\Delta|$. In this paper, we are mainly interested in the l_1 -metrization d_{Δ, l_1} , which is simply denoted by d_Δ .

A simply-connected 2-complex Δ with property (2.1) is called a *folder complex* (an *F-complex* for short) [3, Section 7] if

- (2.3) (i) the intersection of any two folders does not contain incident legs, and
- (ii) there are no three folders F_i ($i = 1, 2, 3$) and three distinct legs e_i ($i = 1, 2, 3$) sharing a common vertex such that e_i belongs to F_j exactly when $i \neq j$.

In fact, this is equivalent to the condition that $|\Delta|$ is a CAT(0) space under the l_2 -metrization [3, Theorem 7.1]. We particularly call (ii) the *flag condition*; see Figure 3 (a).

Next we introduce certain subsets of $|\Delta|$. A subset R of $|\Delta|$ is called *normal* if it satisfies the following property:

- (2.4) (i) R coincides with $|\mathcal{K}|$ for a connected subcomplex \mathcal{K} of Δ with the property that if \mathcal{K} contains a leg e , then every cell containing e belongs to \mathcal{K} .
- (ii) there are no two triangles σ and σ' sharing a leg and a right angle such that $(\sigma \cup \sigma') \cap R$ coincides with the union of hypotenuses of σ and σ' .




Figure 4: Orientation

We particularly call (ii) the *local convexity condition*; see Figure 3 (b). Let $\Gamma = \Gamma^\Delta$ be the graph consisting of all legs of Δ , called the *leg-graph* of Δ . Equivalently, Γ is the graph obtained by deleting all hypotenuses from the 1-skeleton graph of Δ . The edge-length of Γ is defined by κ uniformly. An F-complex Δ is said to be *orientable* if the leg-graph Γ has an orientation such that

- (2.5) (i) in each square, its diagonal edges have same direction. More precisely, orientations of two nonincident edges are opposite each other in a cyclic ordering of the corresponding 4-cycle, and
- (ii) in each triangle, its right angle is neither a source nor a sink.

See Figure 4. In fact, the leg-graph of an orientable F-complex is a frame; see Theorem 3.2 in Section 3.1.

For a rational distance μ on set S , a pair $(\Delta; \{R_s\}_{s \in S})$ of an F-complex Δ and a set $\{R_s\}_{s \in S}$ of normal sets is called an *F-complex realization* of μ if it satisfies

$$\mu(s, t) = d_\Delta(R_s, R_t) \quad (s, t \in S).$$

Namely, μ is realized by the l_1 -distances among normal sets R_s . Consider the μ -problem (1.1) for a capacitated graph (G, c) with $S \subseteq VG$, and consider the following continuous and discrete location problems on Δ :

$$(2.6) \quad \begin{array}{ll} \text{Minimize} & \sum_{xy \in EG} c(xy) d_\Delta(\rho(x), \rho(y)) \\ \text{subject to} & \rho : VG \rightarrow |\Delta|, \\ & \rho(s) \in R_s \quad (s \in S). \end{array}$$

$$(2.7) \quad \begin{array}{ll} \text{Minimize} & \sum_{xy \in EG} c(xy) d_\Gamma(\rho(x), \rho(y)) \\ \text{subject to} & \rho : VG \rightarrow V\Gamma, \\ & \rho(s) \in R_s \cap V\Gamma \quad (s \in S). \end{array}$$

Now we are ready to describe our main result.

Theorem 2.1. *Let μ be a rational distance on S . Suppose that μ has an F-complex realization $(\Delta; \{R_s\}_{s \in S})$. Then, for any capacitated graph (G, c) with $S \subseteq VG$, the following holds:*

- (1) *The maximum value of the μ -problem (1.1) is equal to the minimum value of the continuous location problem (2.6).*
- (2) *In addition, if Δ is orientable, then the maximum value of the μ -problem (1.1) is equal to the minimum value of the discrete location problem (2.7).*




Figure 5: 4-subdivision




Figure 6: Two-commodity F-complexes

An F-complex Δ may be not orientable. By the subdivision operation, we can always obtain an orientable F-complex as follows. For a positive integer $k \geq 2$, the k -subdivision Δ^k is obtained from Δ by subdividing each square into k^2 squares, subdividing each triangle into $k(k-1)/2$ squares and k triangles, and subdividing each maximal 1-cell(edge) into k edges as in Figure 5. The leg-length is set to be κ/k . One can easily see:

- (2.8) (i) The k -subdivision Δ^k is an F-complex for any k .
(ii) The 2-subdivision Δ^2 is always orientable.

Next we describe illustrative examples of combinatorial duality relations for μ -problems based on Theorem 2.1.

Two-commodity flows. First consider the case where terminal set S is a four-set $\{s, t, s', t'\}$ and distance μ on S is given as $\mu(s, t) = \mu(s', t') = 1$ and others are zero. Then the μ -problem (1.1) is the two-commodity flow maximization problem [14]. We can realize μ by the distance on an F-complex as follow. Consider four triangles, glue them around a common right angle, and define the leg-length κ to be $1/2$, as in Figure 6 (a). The resulting 2-complex is clearly an orientable F-complex. Moreover Δ realizes μ as the distance among four hypotenuses, which are normal. Theorem 2.1 (ii) gives a combinatorial duality relation for the two-commodity flow maximization.

Weighted two-commodity flows. Also we can consider a *weighted* version of two-commodity flows. Let μ be defined as $(\mu(s, t), \mu(s', t')) = (p, q)$ for relatively prime positive integers p, q , and other distances are zero. Then consider an F-complex Δ as in Figure 6 (b), which is a subdivision of a rectangle of edge-length ratio $(p : q)$. Let $\kappa = 1/2$. Clearly Δ is a F-complex. Again μ is realized by the distance of the

four edges of the rectangle, which are normal. Also one can see that Δ is orientable. Then Theorem 2.1 (ii) yields a combinatorial duality relation, which seems new in the literature.

Anticlique-bipartite commodity flows. A 0-1 distance μ can be identified with a *commodity graph* H on S by $st \in EH \Leftrightarrow \mu(s, t) = 1$. We may assume that H has no isolated vertices. The corresponding μ -problem for (G, c) is the maximization of the total sum of flow-values of multiflows connecting pairs of terminals specified by edges of H . For a multifold f , the total flow-value with respect to H is denoted by $\|f\|_H$. A bipartition (X, Y) of vertex set VG is called a *cut*, and its *cut capacity* $c(X, Y)$ is defined to be the sum of the capacity of edges with exactly one end belonging to X . A *multicut* $\mathcal{X} = (X_1, X_2, \dots, X_k)$ ($k \geq 2$) with respect to H is a partition of VG with the property that ends of each edge of H belong to distinct parts of \mathcal{X} , and its capacity $c(\mathcal{X})$ is the sum of the capacity of edges whose ends belong to distinct parts of \mathcal{X} . For a multifold f and a multicut \mathcal{X} (w.r.t. H), the following obvious weak duality relation holds:

$$(2.9) \quad \|f\|_H \leq c(\mathcal{X}).$$

In general, the duality gap is strict (except single- and two-commodity flows). Consider the following *relaxation* of multicuts. A *semi-multicut* $\mathcal{X} = (X_1, X_2, \dots, X_k)$ ($k \geq 2$) with respect to H is a *set of disjoint subsets* of VG with the property that ends of each edge of H belong to distinct sets in \mathcal{X} , and its capacity $c(\mathcal{X})$ is defined as

$$c(\mathcal{X}) = \frac{1}{2} \sum_{i=1}^k c(X_i, \overline{X_i}).$$

Again the weak duality (2.9) holds for any multifold and any semi-multicut, and the duality gap is still strict. However, the class of commodity graphs H admitting strong duality (for semi-multicuts) is not so narrow; recall Lovász-Cherkassky theorem for the case where H is a complete graph. Karzanov and Lomonosov [20] characterized such a class of commodity graphs H as follows.

Theorem 2.2 ([20]). *Suppose that the set \mathcal{A} of maximal stable sets of commodity graph H has the following properties:*

- (1) *For every triple $A, B, C \in \mathcal{A}$ of maximal stable sets, at least one of $A \cap B$, $B \cap C$, and $C \cap A$ is empty.*
- (2) *The intersection graph of \mathcal{A} is bipartite.*

Then, for any capacitated graph (G, c) having H as commodity graph, the following holds:

$$\max\{\|f\|_H \mid \text{multifold } f\} = \min\{c(\mathcal{X}) \mid \text{semi-multicut } \mathcal{X}\}.$$

See [22], and also see [9] for a polymatroidal approach to this result. A commodity graph with condition (1-2) is called *anticlique-bipartite*. Here we give an interpretation of this results in terms of an F-complex. Let us construct an F-complex realization for 0-1 distance μ corresponding to a commodity graph H with property (1). Let $\Omega_{\mathcal{A}}$ be the intersection graph of \mathcal{A} . From $\Omega_{\mathcal{A}}$, we can construct an abstract 2-complex of (2.2). All edges of $\Omega_{\mathcal{A}}$ are supposed be hypotenuses. Add a new vertex O , and add a leg OA for each $A \in \mathcal{A}$. Let Γ_0 be the resulting graph. 2-cells \mathcal{C} consists of 3-cycles of one hypotenuse AB and two legs OA, OB over all intersecting pairs $A, B \in \mathcal{A}$. Define the leg-length κ to be $1/2$. Let Δ be the resulting 2-complex, which is the join of O and




Figure 7: (a) an anticlique bipartite commodity graph and (b) its F-complex realization




Figure 8: F-complexes for $H = C_5$

$\Omega_{\mathcal{A}}$. By the property (1), the girth of $\Omega_{\mathcal{A}}$ is at least four. This fact implies that Δ is an F-complex; the violation of the flag condition at O implies the existence of a chordless cycle in $\Omega_{\mathcal{A}}$ of length at most three. For $s \in S$, define a normal set R_s by

$$R_s = \begin{cases} \text{hypotenuse } AB & \text{if } s \text{ belongs to exactly two } A, B \in \mathcal{A}, \\ \text{vertex } A & \text{if } s \text{ belongs to exactly one } A \in \mathcal{A}. \end{cases}$$

Note that every terminal s belongs to at most two maximal stable sets by (1) and the assumption that H has no isolated vertices. Then $(\Delta; \{R_s\}_{s \in S})$ is an F-complex realization. Indeed, $d_{\Delta}(R_s, R_t) = 0 \Leftrightarrow R_s \cap R_t \neq \emptyset \Leftrightarrow R_s \cap R_t = \{A\}$ for $A \in \mathcal{A}$ with $s, t \in A \Leftrightarrow \mu(s, t) = 0$, and $R_s \cap R_t = \emptyset$ implies that $d_{\Delta}(R_s, R_t) = 1$. Furthermore Δ is orientable if and only if the condition (2) is fulfilled. Consider the discrete location problem (2.7) for $(\Delta; \{R_s\}_{s \in S})$. For a map $\rho : VG \rightarrow V\Gamma$ feasible to (2.7), the set $\mathcal{X} := (\rho^{-1}(A) \mid A \in \mathcal{A})$ is a semi-multicut, and $\sum_{xy \in EG} c(xy) d_{\Delta}(\rho(x), \rho(y))$ coincides with $c(\mathcal{X})$. Thus Theorem 2.1 implies Karzanov-Lomonosov duality relation above.

Figure 7 illustrates (a) an anticlique-bipartite commodity graph H of $K_{3,3}$ minus one perfect matching, and (b) its F-complex realization Δ . In this case, Δ is obtained by gluing six triangles of leg-length $1/2$, and each R_s is a hypotenuse.

Also for nonorientable case, i.e., H violates (2), we obtain a combinatorial duality relation by subdividing Δ into Δ^2 , which coincides with the one given in [16]. Figure 8 (a) illustrates an F-complex corresponding to the commodity graph of five-cycle, which is nonorientable, and (b) illustrates its 2-subdivision. See the next paper [13] for further examples.

3 Folder complexes and normal sets

Our main Theorem 2.1 is a consequence of the following two properties of an F-complex Δ ; see the paragraph below.

- (3.1) (i) For any collection of normal sets N_1, N_2, \dots, N_k and nonnegative rationals r_1, r_2, \dots, r_k , the collection of balls $\{B(N_i, r_i)\}_{i=1}^k$ has the Helly property.
- (ii) Suppose that Δ is orientable. For a map $\rho : X \rightarrow |\Delta|$ on a finite set X , define metric d^ρ on X by

$$d^\rho(x, y) = d_\Delta(\rho(x), \rho(y)) \quad (x, y \in X).$$

Then there exist maps $\rho_i : X \rightarrow V\Gamma$ ($i \in I$) such that d^ρ is a convex combination of d^{ρ_i} and, for any normal set R , $\rho(x) \in R$ implies $\rho_i(x) \in R$ ($i \in I$).

(i) is an extension of [3, Theorem 7.8] for the case where each N_i is a singleton set; see Section 4.2. The goal of this section is prove (3.1). In Section 3.1, we introduce some of basic notation. In Section 3.2, we study shortest paths connecting normal sets. By using them, in Section 3.3 we prove (3.1) (i). In Section 3.4, we show a certain decomposition property of shortest paths, and prove (3.1) (ii).

Proof of Theorem 2.1. Here we show Theorem 2.1 by assuming (3.1). The second statement (2) of Theorem 2.1 immediately follows from the first (1) and (3.1) (ii) (and (3.7)). So we show the first statement. Let μ be a rational distance on finite set S . For a capacitated graph (G, c) with terminal set S , consider the μ -problem (1.1). As is well-known [22], its LP-dual problem is given by

$$(3.2) \quad \begin{array}{ll} \text{Minimize} & \sum_{xy \in EG} c(xy)m(x, y) \\ \text{Subject to} & m : \text{metric on } VG, \\ & m(s, t) \geq \mu(s, t) \quad (s, t \in S). \end{array}$$

Let $(\Delta; \{R_s\}_{s \in S})$ be an F-complex realization of μ . For a map ρ feasible to (2.6), define metric d^ρ on VG as in (3.1) (ii). Then condition $\rho(s) \in R_s$ implies

$$d^\rho(s, t) = d_\Delta(\rho(s), \rho(t)) \geq d_\Delta(R_s, R_t) = \mu(s, t),$$

and thus d^ρ is feasible. Hence it suffices to show:

- (3.3) For every rational metric m feasible to (3.2), there exists a map $\rho : VG \rightarrow |\Delta|$ such that $d^\rho \leq m$ and $\rho(s) \in R_s$ for $s \in S$.

The following argument is based on the idea used in [1, p.421]. Let m be a feasible rational metric. Let $VG = \{x_1, x_2, \dots, x_n\}$. Define a map $\rho : VG \rightarrow |\Delta|$ recursively by $\rho(x_k)$ being an arbitrary point in

$$(3.4) \quad \bigcap_{s \in S} B(R_s, m(s, x_k)) \cap \bigcap_{i=1}^{k-1} B(\rho(x_i), m(x_i, x_k)) \quad (k = 1, 2, \dots, m).$$

We show that the set (3.4) is nonempty for all k , which implies that ρ is well-defined and satisfies $\rho(s) \in R_s$ and $d^\rho \leq m$. Indeed, $\rho(s) \in B(R_s, m(s, s)) = R_s$, and $\rho(x_j) \in B(\rho(x_i), m(x_i, x_j))$ ($i < j$) implies $d_\Delta(\rho(x_i), \rho(x_j)) \leq m(x_i, x_j)$.

We use the induction on k . Suppose that (3.4) is nonempty for $i < k$. By rationality, we can take $\rho(x_i)$ as a vertex of l -subdivision Δ^l for large l . Then each singleton set $\{\rho(x_i)\}$ is normal in Δ^l . Of course, R_s is also normal in Δ^l . By the Helly property (3.1) (i), it suffices to show that each pairwise intersection is nonempty. Note that two balls $B(R, r)$ and $B(R', r')$ intersect if and only if $d_\Delta(R, R') \leq r + r'$. For $s, t \in S$, the nonemptiness of $B(R_s, m(s, x_k)) \cap B(R_t, m(t, x_k))$ follows from $m(s, x_k) + m(x_k, t) \geq m(s, t) \geq \mu(s, t) = d_\Delta(R_s, R_t)$. For $s \in S$ and $i < k$, the nonemptiness of $B(R_s, m(s, x_k)) \cap B(\rho(x_i), m(x_i, x_k))$ follows from $m(s, x_k) + m(x_k, x_i) \geq m(s, x_i) \geq d_\Delta(R_s, \rho(x_i))$, where the last inequality follows from $\rho(x_i) \in B(R_s, m(s, x_i))$ for $i < k$ (by induction). For $1 \leq i < j < k$, the nonemptiness of $B(\rho(x_i), m(x_i, x_k)) \cap B(\rho(x_j), m(x_j, x_k))$ follows from $m(x_i, x_k) + m(x_k, x_j) \geq m(x_i, x_j) \geq d_\Delta(\rho(x_i), \rho(x_j))$. Thus we are done.

3.1 Preliminaries

Let Γ be an undirected graph. The *interval* $I_\Gamma(x, y)$ between $x, y \in V\Gamma$ is defined to be the set $\{z \in V\Gamma \mid d_\Gamma(x, y) = d_\Gamma(x, z) + d_\Gamma(z, y)\}$ (for some specified edge length). An undirected graph Γ (with unit edge length) is called *modular* if for each triple x, y, z of vertices the intersection $I_\Gamma(x, y) \cap I_\Gamma(y, z) \cap I_\Gamma(z, y)$ is nonempty. An element in $I_\Gamma(x, y) \cap I_\Gamma(y, z) \cap I_\Gamma(z, y)$ is called a *median* of x, y, z . A modular graph Γ satisfies the so-called *quadrangle condition*:

$$(3.5) \quad \text{For any four vertices } x, u, v, y \text{ with } d_\Gamma(x, u) = d_\Gamma(x, v) = 1 \text{ and } d_\Gamma(u, y) = d_\Gamma(v, y) = d_\Gamma(x, y) - 1, \text{ there exists a common neighbor } w \text{ of } u \text{ and } v \text{ such that } d_\Gamma(w, y) = d_\Gamma(x, y) - 2.$$

Indeed, we can take w as a median of u, v, y .

A graph is called *hereditary modular* if all isometric subgraph is modular. Bandelt [2] gave an elegant characterization of hereditary modular graphs as follows.

Theorem 3.1 ([2]). *A graph is hereditary modular if and only if it is bipartite and has no isometric cycles of length greater than four.*

Therefore a frame is just an orientable hereditary modular graph. Chepoi [3] established a relation between hereditary modular graphs and F-complexes. Here a folder consisting of a single triangle is called a *triangle-folder*.

Theorem 3.2 ([3, Theorem 7.1]). *For a simply-connected 2-complex Δ satisfying (2.1) without triangle-folders, the following conditions are equivalent:*

- (i) Δ is an F-complex.
- (ii) its leg-graph Γ is a hereditary modular graph without $K_{3,3}$ and $K_{3,3}^-$ as induced subgraphs.

In addition, (i) \Rightarrow (ii) holds for any simply-connected 2-complex Δ satisfying (2.1) (possibly including triangle-folders).

$K_{3,3}^-$ is the graph obtained from $K_{3,3}$ by deleting one edge. Since both $K_{3,3}$ and $K_{3,3}^-$ are nonorientable, the leg-graph of an orientable F-complex is a frame. We can partially reconstruct an F-complex Δ from its leg-graph Γ based on the following property:

- $$(3.6) \quad \begin{aligned} & \text{(i) every 4-cycle of the leg-graph } \Gamma \text{ belongs to a (unique) folder.} \\ & \text{(ii) every 3-cycle of the 1-skeleton graph of } \Delta, \text{ consisting of two legs and one hypotenuse, is the boundary of a triangle.} \end{aligned}$$

(i) follows from the argument in [3, Section 7]; this is not so obvious, which uses the simply-connectedness, CAT(0) property of Δ under the l_2 -metrization, and the Gauss-Bonnet Theorem. (ii) follows from (i). Indeed, the hypotenuse of 3-cycle belongs to some triangle. Use (i) for a possibly appearing 4-cycle of legs.

Therefore, if there is no folder consisting of at most two triangles, then Δ is completely reconstructed from its leg-graph; there are two ways splitting a square to two triangles. Moreover, for an arbitrary hereditary modular graph without $K_{3,3}$ and $K_{3,3}^-$ as induced subgraphs, by filling a folder to each maximal complete bipartite graph $K_{2,n}$ ($n \geq 2$), we obtain an F-complex without triangle-folders [3].

3.2 Geodesics

Here we investigate shortest paths connecting normal sets. Several important properties previously known for shortest paths connecting *vertices* can be naturally extended for shortest paths connecting *normal sets*.

Let Δ be an F-complex. In this subsection, *we assume that the leg-length κ is equal to 1 for simplicity*. We begin with:

$$(3.7) \quad \text{For normal sets } N \text{ and } M, \text{ we have } d_\Delta(N, M) = d_\Gamma(N \cap V\Gamma, M \cap V\Gamma).$$

Indeed, we can easily modify a (polygonal) geodesic between N and M so that it lies on the union of legs (as in the proof of [11, Proposition 4.2]). For notational simplicity, in the sequel we use the same d for d_Δ and d_Γ , and we denote $I_\Gamma(x, y)$ simply by $I(x, y)$. For a normal set R , let Γ_R be the 1-skeleton graph of the subcomplex of Δ induced by R . The edge length of e of Γ_R is defined to be 1 if e is a leg and 2 if e is a hypotenuse.

Lemma 3.3. *For a normal set R , we have $d_{\Gamma_R}(x, y) = d(x, y)$ for $x, y \in V\Gamma_R$.*

Proof. Suppose to the contrary that there are $x, y \in V\Gamma_R$ with $d_{\Gamma_R}(x, y) > d(x, y)$. Take such x, y with $d_{\Gamma_R}(x, y)$ minimum. Then $d_{\Gamma_R}(x, y) \geq 4$. Indeed, $d_{\Gamma_R}(x, y) \leq 2$ is clearly impossible. Suppose $d_{\Gamma_R}(x, y) = 3$. Since Γ is bipartite, we have $d(x, y) = 1$ and therefore x and y are incident in Γ . Take a shortest path P connecting x, y in Γ_R . $P \cup \{xy\}$ is a 3-cycle or a 4-cycle in 1-skeleton graph of Δ . In both cases, by (3.6) there is a folder F containing $P \cup \{xy\}$. Then R violates the normality (i) at F by $xy \not\subseteq R$. A contradiction.

Take $z \in I_{\Gamma_R}(x, y)$ with $\min\{d(x, z), d(z, y)\} \geq 2$. By minimality assumption, we have $d_{\Gamma_R}(x, z) = d(x, z)$ and $d_{\Gamma_R}(y, z) = d(y, z)$. Take a median w of x, y, z in Γ . Then $w \notin R$. Take $w' \in I(z, w)$ incident to z (by a leg zw'). We show that there exists a triangle σ of vertices z, w', u and hypotenuse zu such that $u \in I_{\Gamma_R}(z, x)$ and $\sigma \cap R = zu$. Then there also exists such a triangle σ' for $I_{\Gamma_R}(z, y)$, and R violates the local convexity in $\sigma \cup \sigma'$. Take $x' \in I_{\Gamma_R}(z, x)$ incident to z . Suppose that $x'z$ is a leg. Then by the quadrangle condition (3.5) for z, x', w', x there exists x'' incident to x' and w' with $d(z, x) = d(z, x'') - 2$ and therefore there is a folder containing z, x', x'', w' . By the normality (i), x'' must belong to R , and thus zx'' is a hypotenuse, and there is a required triangle σ having z, w', x'' . Suppose that $x'z$ is the hypotenuse of some triangle $\tilde{\sigma}$ having vertices x', z, x_0 . By the quadrangle condition for z, w', x_0, x , there are a common neighbor x'' of x_0 and w' with $d(z, x'') = d(z, x) - 2$, and a folder F containing z, w', x_0, x'' . If $x'' = x'$, then $\tilde{\sigma}$ is a required triangle. So $x'' \neq x'$. By the quadrangle condition for x_0, x', x'', x we can find a folder F' violating the flag condition at x_0 with $\tilde{\sigma}$ and F . A contradiction. \square

We will often use the following *bipartite property*:

Lemma 3.4. For a normal set R and a leg $e = xy$, if $e \not\subseteq R$, then we have

$$d(R, x) - d(R, y) \in \{-1, 1\}.$$

Proof. Let T be the closure of a connected component of $|\Delta| \setminus R$. Since Δ is simply-connected, $T \cap R$ is connected. By the normality, $T \cap R$ consists of hypotenuses (or a single point). Therefore $T \cap R \cap V\Gamma$ is contained by one color class of bipartite graph Γ . The statement immediately follows from this fact. \square

The next lemma is an extension of the quadrangle condition (3.5). We also call it the *quadrangle condition*.

Lemma 3.5. For a triple x, u, v of vertices and a normal set R , suppose that $d(x, u) = d(x, v) = 1$ and $d(u, R) = d(v, R) = d(x, R) - 1$.

- (1) If $d(x, R) = 1$, then there exists a triangle σ of vertices x, u, v such that $R \cap \sigma = uv$ (uv is a hypotenuse).
- (2) If $d(x, R) \geq 2$, then there exists a common neighbor w of u and v such that $d(w, R) = d(x, R) - 2$.

Proof. (1). By Lemma 3.3, x and y are joined by a path P of length 2 in Γ_R . Apply (3.6) for cycle $P \cup \{ux, uv\}$. (2). So suppose that $d(x, R) = k \geq 2$. We use the induction on k . Take $u', v' \in R$ satisfying $d(u, u') = d(v, v') = k - 1$ with $d(u', v') (\leq 2k)$ minimum. Suppose $d(u', v') = 2k$. Then $x \in I(u', v')$. Let u'_0 be a neighbor of u' in $I(u', x)$. By Lemma 3.3, we can take a vertex $u'_1 \in I_{\Gamma_R}(u', v')$ joined to u' by a leg or a hypotenuse in R . By repeated applications of the quadrangle condition (for u', u'_0, \cdot, v' and for u'_0, \cdot, v'), as in the last part of the proof of Lemma 3.3, we can find a triangle σ of vertices u', u'_0, u'_1 with $R \cap \sigma = u'u'_1$ and $d(v', u'_1) = d(v', u') - 2$. Then $d(u'_1, u) = k - 1$, and this contradicts to the minimality assumption of (u', v') .

So suppose $d(u', v') < 2k$. Take a median $w^* (\neq x)$ of u', v', x and take a neighbor w of x from $I(x, w^*)$. If $w = v$, then by the quadrangle condition for x, u, v, u' we can find a required vertex. Therefore $w \neq u, v$. By the quadrangle condition for x, u, w, u' , we can find a common neighbor u^* of u, w with $d(u^*, u') = d(u^*, R) = k - 2$. Similarly we can find a common neighbor v^* of w, v with $d(v^*, v') = d(v^*, R) = k - 2$. If $u^* = v^*$, then we are done. So $u^* \neq v^*$. Let F_1 and F_2 be the folders containing x, u, w, u^* and x, v, w, v^* , respectively. Apply the induction to quadrangle w, u^*, v^*, R . Then we can find a folder F_3 violating the flag condition at w with F_1 and F_2 . A contradiction. \square

The next lemma is a certain convexity property of normal sets.

Lemma 3.6. Let M, N be two normal sets. For a vertex $x \in M$, if $d(x, N) > d(M, N)$, then there exists an edge (leg or hypotenuse) $xy \subseteq M$ such that $d(y, N) < d(x, N)$.

Proof. Take a vertex w in M with $d(w, N) = d(M, N)$. Since M is connected, we can take a path P in Γ_M connecting x and w . Among such paths, take P with the following properties:

- (a) the integration I of $d(\cdot, N)$ along P , i.e.,

$$I = \sum_{uv: \text{leg in } P} \frac{d(u, N) + d(v, N)}{2} + \sum_{uv: \text{hypotenuse in } P} d(u, N) + d(v, N)$$

is minimum.

(b) P contains legs as much as possible (under property (a)).

Let $P = (x = x_0, x_1, \dots, x_m = w)$. Define a sequence $\{a_i\}_{i=0, \dots, m}$ by $a_i = d(x_i, N)$. We show that $\{a_i\}_{i=0, \dots, m}$ has no triple a_{i-1}, a_i, a_{i+1} with the property (i) $a_{i-1} = a_i > a_{i+1}$ or (ii) $a_{i-1} < a_i > a_{i+1}$. This immediately implies the statement.

Case (i). Suppose to the contrary that $a_{i-1} = a_i > a_{i+1}$ holds for some i . By the bipartite property, edge $x_{i-1}x_i$ is necessarily a hypotenuse. By construction, there is a triangle σ of vertices x_{i-1}, x_i, u such that $u \notin M$. Also $u \notin N$. Suppose that x_ix_{i+1} is a leg. Then by the quadrangle condition for x_i, x_{i+1}, u, N , there are a common neighbor v of x_{i+1}, u and a folder F containing x_i, x_{i+1}, u, v . By normality, we have $v \in M$. Then there is a triangle σ' of vertices u, v, x_i in F such that M violates the local convexity at $\sigma \cup \sigma'$. A contradiction. Suppose that x_ix_{i+1} is a hypotenuse. There is a triangle σ of vertices x_i, x_{i+1}, v such that $v \notin M$ (by construction). If $v = u$, then M violates the local convexity on $\sigma \cup \sigma'$. So $v \neq u$. By the quadrangle condition for x_i, v, u, N , there is a common neighbor $z (\neq x_i)$ of v, u and a folder F supported by x_i, v, u, z . Again, by the quadrangle condition for v, z, x_{i+1}, N , there is a folder F' violating the flag condition at v with σ' and F . A contradiction.

Case (ii). Suppose to the contrary that $a_{i-1} < a_i > a_{i+1}$ holds for some i . Then at least one of $x_{i-1}x_i$ and x_ix_{i+1} is a hypotenuse. Otherwise, by the quadrangle condition and the normality, the segment $x_{i-1}x_{i+1}$ belongs to M . This contradicts to the construction. Suppose that both $x_{i-1}x_i$ and x_ix_{i+1} are hypotenuses. Then there are triangles σ, σ' of vertices x_{i-1}, x_i, u and x_{i-1}, x_i, v (resp.) such that $u, v \notin M$. If $u = v$, then M violates the local convexity on $\sigma \cup \sigma'$. Therefore $u \neq v$. Also in this case, we can find a triple of folders violating the flag condition at u (or v) by repeated applications of the quadrangle condition, as above. Suppose that $x_{i-1}x_i$ is a hypotenuse and x_ix_{i+1} is a leg. Then there is a triangle σ of vertices x_{i-1}, x_i, u with $u \notin M$. Again repeated applications of the quadrangle condition imply the existence of a triple of folders violating the flag condition at u . A contradiction. \square

From the proof, we obtain the following:

Lemma 3.7. *Let M, N be two normal sets. For two distinct vertices $x, y \in M$ with $d(x, N) = d(y, N) = d(M, N)$, there is a hypotenuse xz in M with $d(z, y) = d(x, y) - 2$ and $d(z, N) = d(M, N)$.*

Next we discuss the extendability of a path to a shortest path connecting normal sets. Although our goal is Lemma 3.11 below, we derive it from a more general version (Lemma 3.9) for future applications in [13].

An F-complex Δ is called *star-shaped* if there is a vertex p such that every folder contains p and no triangle contains p as its right angle. For a vertex p in Δ , let Δ_p be the subcomplex of Δ consisting of cells containing p and their faces. Clearly Δ_p is also an F-complex. Although Δ_p may not be star-shaped, $(\Delta^2)_p$ is always star-shaped.

Lemma 3.8. *Let p be a vertex, and let R be a normal set not containing p . Suppose that Δ_p is star-shaped. Then a point x in $|\Delta_p|$ at minimum distance from R is a vertex and uniquely determined.*

We call this vertex x the *gate* of R in Δ_p ; this definition is compatible to that in [7].

Proof. By $p \notin R$ and (2.4) (i), R does not contain a leg in Δ_p . Since Δ_p is star-shaped, the boundary (relative to $|\Delta|$) consists of legs. By Lemma 3.4, the map $x \mapsto d(x, R)$ on a leg is monotone decreasing or increasing. Hence the minimum attains at a vertex. Suppose that two vertices x, y have the minimum distance from R . Consider the case

where x and y are not incident to p and have a common neighbor v ($\in \Delta_p$). Then x, v, p and y, v, p belong to folders F_1 and F_2 , respectively. By the quadrangle condition for v, x, y, R , we obtain a folder F_3 violating the flag condition with F_1 and F_2 . A contradiction. Suppose other cases. Take a path P connecting x and y in Δ_p passing p . By applying the quadrangle condition for three consecutive vertices in P and R , we find a vertex u in Δ_p with $d(u, R) < d(x, R)$, or a vertex z in Δ_p with $d(z, R) = d(x, R)$ such that z and x have a common neighbor. The first case is clearly impossible, and the second case reduces to the case above for (x, z) . \square

The extendability of a shortest path passing a vertex p can be locally characterized by the gates in Δ_p (or $(\Delta^2)_p$ when Δ_p is not star-shaped).

Lemma 3.9. *Let p be a vertex, and let M, N be normal sets not containing p . Suppose that Δ_p is star-shaped. Let x and y be the gates of M and N in Δ_p , respectively. Then the following two conditions are equivalent:*

- (1) $d(M, N) = d(M, x) + d(x, p) + d(p, y) + d(y, N)$.
- (2) $d(x, y) = d(x, p) + d(p, y)$.

Proof. (1) \Rightarrow (2) is obvious. We show the converse. We first consider the case where M is a singleton $\{w\}$. Our proof is based on that of [3, Lemma 7.6]. Suppose to the contrary that $d(w, N) < d(w, x) + d(x, p) + d(p, y) + d(y, N)$. We take such $\{w\}$ with $k = d(w, x)$ minimum. Note that $d(p, N) = d(p, y) + d(y, N)$ necessarily holds. Suppose the case $k = 0$ ($x = w$). Suppose that x is incident to p . Then $d(x, N) = d(p, N) - 1$. Take a neighbor z of p from $I(p, y)$. Apply the quadrangle condition for p, x, z, N . Then we obtain a common neighbor u of x, z with $d(u, N) = d(z, N) - 1$. Since u belongs to Δ_p , we have $z \neq y$ and $d(x, y) = 3$. Thus $d(u, N) = d(y, N)$ and $u \neq y$, which contradicts to Lemma 3.8. Suppose that x is not incident to p ($d(p, x) = 2$). Let z be a common neighbor of x and p . Then $d(z, N) = d(z, p) + d(p, y) + d(y, N)$ by above. Thus $d(x, N) = d(z, N) - 1$ holds. Apply the quadrangle condition for z, x, p, N . Then we obtain a common neighbor u of x and p with $d(u, N) = d(p, N) - 1$. Then y is not incident to p , i.e., $d(x, y) = 4$. Take a common neighbor v of p and y . Apply the quadrangle condition for p, v, u, N . Then we obtain a common neighbor q of v, u with $d(q, N) = d(y, N)$. However $q \neq y$ by $d(x, y) = 4$. A contradiction to Lemma 3.8.

Suppose $k = 1$, i.e., w is incident to x . By the assumption and the bipartite property, we have $d(w, N) = d(x, N) - 1$. If x is incident to p , then by the quadrangle condition for x, w, p, N there is a folder including w, p, x , and thus w belongs to Δ_p . A contradiction. Suppose that x is not incident to p . Take a common neighbor z of x and p . Then z, x, p belong to a common folder F_1 . Apply the quadrangle condition for x, w, z, N . We obtain a common neighbor u of w and z such that $d(u, N) = d(w, N) - 1$. Then u, w, z, x belong to a common folder F_2 ($\neq F_1$). Apply the quadrangle condition for z, u, p, N . Then we obtain a folder F_3 violating the flag condition with F_1 and F_2 . A contradiction.

So suppose $k \geq 2$. Take a neighbor z of w from $I(w, x)$, and take a neighbor u of z from $I(z, x)$. Both gates of z and u in Δ_p are x . By minimality assumption, $d(z, N) = d(z, x) + d(x, p) + d(p, y) + d(y, N)$. Therefore $d(w, N) = d(z, N) - 1$. By the quadrangle condition for z, w, u, N , there is a common neighbor r of w, u such that $d(r, N) = d(z, N) - 2$. Also the gate of r in Δ_p is x and $d(r, x) = d(w, x) - 1$. Thus $d(r, N) < d(r, x) + d(x, p) + d(p, y) + d(y, N)$. A contradiction to the minimality of w .

Next we show that M is general. Suppose that (1) fails for $k = d(M, x) = 0$ ($x \in M$). Suppose that x is incident to p . By Lemma 3.6 and the quadrangle condition, we can find a triangle such that its hypotenuse belongs to M and its right angle is p . This

contradicts to the fact that Δ_p is star-shaped. Suppose that x is not incident to p . Let z be a common neighbor of x and p . Let F be a folder containing z, x, p . By Lemma 3.6 and the quadrangle condition, we can find a triangle σ of vertices x, z, u such that $xu = M \cap \sigma$ and $d(u, N) = d(x, N) - 2$. By the quadrangle condition for z, p, u, N , we can find a folder F' violating the flag condition with F and σ . Suppose that (1) fails for $k = d(M, x) > 0$. Take $w \in M$ with $d(w, x) = d(M, x)$ such that $d(w, N)$ is minimum. By the singleton case above, $d(w, N) = d(w, x) + d(x, p) + d(p, y) + d(y, N)$ holds, and thus $d(w, N) > d(N, M)$. By Lemma 3.6, there is $z \in M$ incident to w by a leg or a hypotenuse such that $d(z, N) < d(w, N)$. Let $w' \in I(w, x)$ be a neighbor of w . By repeated applications of the quadrangle condition, we can find a triangle σ of vertices w, w', z' such that $w'z' = \sigma \cap M$ and $d(z', N) = d(w, N) - 2$. Consider the distance $d(z', x)$. Then $d(z', x) < d(w, x)$ contradicts to the definition of w . Therefore $d(z', x) = d(w, x) = d(M, x)$. However this contradicts to the minimality of $d(w, N)$. \square

By the same idea of the proof of Lemma 3.8, we have:

Lemma 3.10. *For a folder F and a normal set R not meeting the interior of F , a point x in F at minimum distance from R is a vertex and uniquely determined.*

Also we call x the *gate* of R in F . Since $F = |(\Delta^2)_p|$ for the center vertex p of the 2-subdivision of F , as a corollary of Lemma 3.9, we obtain the following fact, which is an extension of [17, p.87, Claim 3] for the case where N, M are singleton sets.

Lemma 3.11. *Let F be a folder, and M, N normal sets not meeting the interior of F . Let x and y be the gates of M and N in F , respectively. If x and y are not adjacent, then we have*

$$d(M, N) = d(M, x) + 2 + d(y, N).$$

We also give a variant of Lemma 3.11; a proof is similar.

Lemma 3.12. *Let M, N be normal sets, and let σ be a triangle with vertices x, y, z and hypotenuse yz . Suppose that $M \cap \sigma = yz$ and x is the gate of N in the folder containing σ . Then we have*

$$d(M, N) = 1 + d(x, N).$$

3.3 Helly property of normal sets

The goal of this subsection is to prove:

Theorem 3.13. *For an F -complex, the collection of normal sets has the Helly property.*

In fact, (3.1) (i) is a simple corollary of it. For a ball $B(N, r)$ for normal set N and rational radius r , we can subdivide Δ into Δ^k and further subdivide Δ^k into $\tilde{\Delta}^k$ by splitting squares into two triangles so that $B(N, r)$ is the union of a subcomplex of $\tilde{\Delta}^k$.

$$(3.8) \quad B(N, r) \text{ is normal in } \tilde{\Delta}^k.$$

Proof. Note that N is also normal in $\tilde{\Delta}^k$. Then (2.4) (i) immediately follows from the bipartite property. Suppose to the contrary that there are adjacent two triangles σ, σ' violating (2.4) (ii) with $B(N, r)$. Let x be a common right angle of σ and σ' . Then $d(x, N) = r + 1/k$. Let y and z be distinct neighbors of x belonging to σ and σ' , respectively. Then $d(y, N) = d(z, N) = r$. By the quadrangle condition for x, y, z, N , we can find a folder F violating the flag condition at x with σ and σ' . A contradiction. \square

We prove Theorem 3.13 by extending Chepoi's shelling argument in [3, p.157]. As Chepoi did, we will remove cells at the maximum distance from some vertex x . So we begin with studying the set

$$M_x = \left\{ y \in |\Delta| \mid d(y, x) = \max_{u \in |\Delta|} d(u, x) \right\}.$$

It is easy to see:

(3.9) M_x is the union of isolated vertices and hypotenuses of triangle-folders.

M_x itself may not be normal.

(3.10) Any normal set M in M_x is a vertex or a tree consisting of hypotenuses.

Proof. We can regard M as a graph of hypotenuses. Suppose that M has a cycle C whose vertices are ordered cyclically as $(y_0, y_1, \dots, y_{m-1})$. Each edge $y_i y_{i+1}$ is the hypotenuse of the unique triangle-folder σ_i . Here the indices are taken modulo m . Let y_i^1 be the right angle of σ_i . By normality, $y_i^1 \neq y_{i+1}^1$ for each i . By the quadrangle condition for y_i, y_{i-1}^1, y_i^1, x , there is a common neighbor y_i^2 of y_{i-1}^1, y_i^1 with $d(y_i^2, x) = d(y_i, x) - 2$ for each i . If $y_i^2 = y_{i-1}^2$, then we obtain a triple of folders violating the flag condition at y_i^1 . Therefore $y_i^2 \neq y_{i-1}^2$ for each i . Again by the quadrangle condition for $y_i^1, y_i^2, y_{i+1}^1, x$, there is a common neighbor y_i^3 of y_i^1, y_{i+1}^1 with $d(y_i^3, x) = d(y_i^1, x) - 2$ for each i . Similarly $y_i^3 \neq y_{i-1}^3$ holds for each i . Repeat this procedure. Then $d(y_i^j, x)$ strictly decreases. In the final step $k = d(x, M)$, we have $y_i^k = x$ for each i . Thus we obtain a triple of folders violating the flag condition at y_i^{k-1} . A contradiction. \square

We are ready to prove Theorem 3.13. We use the induction on the size of Δ . Take some vertex x . Consider M_x and its connected components.

Case 1: M_x has a connected component of an isolated vertex y . By the construction of M_x , all neighbors y_1, y_2, \dots, y_m of y belong to $I_\Gamma(y, x)$. There are three cases:

- (1) y belongs to a unique maximal 1-cell yy_1 ($m = 1$).
- (2) y, y_1, y_2 are vertices of a triangle σ having y as a right angle ($m = 2$).
- (3) y belongs to a folder F such that F is a square, or y is one end of the hypotenuse of F .

We may assume that $N_i \neq \{y\}$ for all i . Consider 2-complex Δ' by deleting all cells containing y from Δ . Clearly Δ' is also simply-connected, and thus an F-complex. Let $N'_i = N_i \cap |\Delta'|$. Then each N'_i is normal in Δ' . We show that each pair N'_i, N'_j has nonempty intersection on $|\Delta'|$, and then we can apply induction. It suffices to verify it for a pair i, j such that N_i and N_j have nonempty intersection on the union of cells containing y . Suppose (1). Clearly $N_i \cap N_j$ necessarily contains y_1 , and so does $N'_i \cap N'_j$. Suppose (2). By normality, $N_i \cap N_j$ includes the hypotenuse $y_1 y_2$, and so does $N'_i \cap N'_j$. Suppose (3). If F is a square, then $N_i \cap N_j \cap F = F$, and $N'_i \cap N'_j$ contains a vertex w in F diagonal to y . If y belongs to the hypotenuse yw of F , then $N_i \cap N_j \cap F$ contains the hypotenuse yw , and $N'_i \cap N'_j$ contains w .

Case 2: M_x has no connected component of an isolated vertex. Take a maximal normal set $M \subseteq M_x$. Let y be a degree-one vertex of tree M , and let y_1 be a unique vertex incident to y in M . Let σ be a unique triangle-folder of hypotenuse yy_1 with right angle u .

(3.11) If there is another hypotenuse e joined to y , then a triangle σ' having e shares a leg yu with σ (in particular $e \subseteq M_x$).

Proof. Let $e = yy'$. Let σ' be a triangle containing e and let u' be the right angle of σ' . By $y \in M_x$, we have $d(y, x) - d(y', x) \in \{0, 2\}$. Suppose $d(y, x) = d(y', x)$. Then e belongs to M_x , and thus belongs to the unique triangle-folder σ' . Then $y' \notin M$ since $y' \in M$ implies $e \in M$ by Lemma 3.3. If $u' \neq u$, then $M \cup e$ is also normal, and this is a contradiction to the maximality of M . Suppose $d(y, x) = d(y', x) + 2$. Suppose $u' \neq u$. Apply the quadrangle condition for y, u, u', x , and again apply it at u' . Then we obtain a violating triple of folders. A contradiction. Thus $u' = u$. \square

Consider Δ' by deleting σ and yy_1 from Δ . Clearly Δ' is an F-complex. Suppose that there is no N_i with $N_i \cap \sigma = yy_1$. Let $N'_i = N_i \cap |\Delta'|$ for each i . Then each N'_i is normal and $\{N'_i\}$ has pairwise nonempty intersection on $|\Delta'|$. Apply the induction. Therefore we assume that there is N_i with $N_i \cap \sigma = yy_1$. Suppose that there is N_j with $N_i \cap N_j = \{y\}$. Otherwise, consider Δ' as above. Let $N'_k = N_k \setminus (yy_1 \setminus y_1)$ if $N_k \cap \sigma = yy_1$, and $N'_k = N_k \cap |\Delta'|$ otherwise. Then each N'_k with $N_k \cap \sigma = yy_1$ is connected and thus normal since y is necessarily a degree-one vertex in Γ_{N_k} from (3.11). Apply the induction. We show that all N_k satisfy $N_i \cap N_k \supseteq \{y\}$; y is a desired common point. Suppose that N_k contains neither yy_1 nor y . If y_1 is the gate of N_k in σ , then by Lemma 3.11 we have $0 = d(N_j, N_k) = d(N_j, y) + 2 + d(y_1, N_k) > 0$ (the gate of N_j is y), and this is a contradiction. If u is the gate of N_k in σ , then, by Lemma 3.12, we have $0 = d(N_i, N_k) = 1 + d(u, N_k) > 0$, and this is a contradiction. So y is the gate of N_k . By $d(y, N_k) > d(N_i, N_k) = 0$ and Lemma 3.6, y is joined to a vertex y' by a leg or a hypotenuse in N_i with $d(y, N_k) > d(y', N_k)$. If yy' is a hypotenuse, then N_i violates the local convexity by (3.11). A contradiction. If yy' is a leg, then by the quadrangle condition for y, y', u, x there is a hypotenuse $e \subseteq N_i$ joined to y . Again (3.11) leads a contradiction. Thus we complete the proof of Theorem 3.13.

3.4 Orbits

We recall the notion of orbits [17, 18] with a slight modification by [11]. Edges $e, e' \in E\Gamma$ are called *mates* if there exists a square σ in Δ containing e, e' as nonincident edges or there exists a folder F consisting of triangles such that F contains both e, e' as legs. Edges e, e' are *projective* if there is a sequence $e = e_0, e_1, \dots, e_m = e'$ such that e_i and e_{i+1} are mates. The projectiveness defines an equivalence relation on $E\Gamma$. An equivalence class is called an *orbit*. The set of all orbits is denoted by $\mathcal{O} = \mathcal{O}^\Delta$. The following property is an extension of [18, Statement 2.2].

Proposition 3.14. *Let M, N be normal sets. Let P, Q be paths in Γ connecting M and N . If P is shortest, then $\#(P \cap O) \leq \#(Q \cap O)$ for each orbit $O \in \mathcal{O}$.*

Proof. We first consider the case where M is a singleton set $\{x\}$. We use the induction on the length k of Q . We may assume that $k \geq 1$ and the length of P is also positive, and assume that vertices u and v following x in P and Q (resp.) are distinct. By the bipartite property, $d(x, N) - d(v, N) \in \{1, -1\}$. Suppose $d(x, N) - d(v, N) = -1$. Then $P \cup \{xv\}$ is a shortest path between v and N , and $Q \setminus \{xv\}$ is a path between v and N of length $k - 1$. By induction we have $\#(Q \cap O) \geq \#((Q \setminus \{xv\}) \cap O) \geq \#((P \cup \{xv\}) \cap O) \geq \#(P \cap O)$. Suppose $d(x, N) - d(v, N) = 1$. Suppose $\#P = 1$. By the quadrangle condition, there is a triangle σ of vertices x, u, v with $N \cap \sigma = uv$. Therefore xu and xv belong to the same orbit, and $\#(P \cap O) \leq \#(Q \cap O)$ is obvious. Suppose $\#Q \geq \#P \geq 2$. By the quadrangle condition for x, u, v, N , we can find a

common neighbor w of u, v with $d(w, N) = d(x, N) - 2$. We can take a shortest path P^* connecting w and N . Clearly $P^* \cup \{uw\}$ is a shortest path connecting u and N . By the induction, $\#((P \setminus \{xu\}) \cap O) = \#((P^* \cup \{uw\}) \cap O)$. Similarly, $\#((Q \setminus \{xv\}) \cap O) \geq \#((P^* \cup \{vw\}) \cap O)$. Since xu and vw are mates and xv and uw are mates, we have $\#(P \cap O) = \#((P^* \cup \{wu, ux\}) \cap O) = \#((P^* \cup \{wv, vx\}) \cap O) \leq \#(Q \cap O)$.

Suppose that M is not a singleton set. Let x and y be the ends of P and Q belonging to M , respectively. By Lemma 3.6 and the singleton case above, it suffices to show the equality $\#(Q \cap O) = \#(P \cap O)$ for the case where Q is also shortest. By above, we may assume $x \neq y$. We use the induction on $d(x, y)$. By Lemma 3.7, we can take $x' \in M$ joined to x by a hypotenuse with $d(x', N) = d(x, N)$ and $d(x', y) = d(x, y) - 2$. There is a triangle of hypotenuse xx' and right angle w with $d(w, N) = d(x, N) - 1$. Take any shortest path \tilde{P} connecting x' and M . Then we have $\#(P \cap O) = \#((\tilde{P} \cup \{xw\}) \cap O) = \#((\tilde{P} \cup \{x'w\}) \cap O) = \#(Q \cap O)$, where the last equality follows from the induction. \square

For a (disjoint) union U of orbits, we can construct a new 2-complex Δ^U from Δ by identifying the ends of all edges not belonging to U . Namely regard Δ as an abstract 2-complex $(\Gamma_0; \mathcal{C})$ with property (2.2). Contract all edges not in U in Γ_0 and delete parallel edges and loops (of hypotenuses) appeared, and also delete 1-cycles (loops) and 2-cycles from \mathcal{C} . Again the resulting 2-complex Δ^U satisfies (2.1). The leg-graph of Δ^U is denoted by Γ^U . This construction naturally induces the map $(\cdot)^U : V\Gamma \rightarrow V\Gamma^U$ by defining x^U to be the contracted vertex. By extending linearly, we obtain map $(\cdot)^U : |\Delta| \rightarrow |\Delta^U|$.

Proposition 3.15. *Let Δ be an orientable F -complex, and let U be the union of some orbits, and let $W = E\Gamma \setminus U$.*

- (1) Δ^U is an orientable F -complex.
- (2) For a normal set R in Δ , R^U is also normal in Δ^U .
- (3) For a shortest path P in Γ connecting normal sets M and N , P^U is also a shortest path in Γ^U connecting M^U and N^U . In particular, we have

$$d_\Delta(M, N) = d_{\Delta^U}(M^U, N^U) + d_{\Delta^W}(M^W, N^W).$$

Proof. If Δ has no triangle-folder, then (1) follows from [19, Statement 3.2]. Suppose that Δ has triangle-folders. Gluing a new triangle to each triangle-folder along the hypotenuse. Then the resulting 2-complex Δ' is also an orientable F -complex without triangle-folders. U is naturally extended to the union U' of orbits in Δ' . Apply the case above, and delete attached triangles from $(\Delta')^{U'}$.

(3). Suppose that P^U is not shortest. Then there is a shorter path P' in Γ^U . By adding edges not in U to P' , we obtain a (not necessarily shortest) path P^* in Γ connecting M and N . Then there is an orbit $O \subseteq U$ such that $\#(O \cap P^*) < \#(O \cap P)$. By Proposition 3.14, P is not shortest. A contradiction.

(2). It suffices to verify the local convexity condition. Suppose that there are a normal set R and two triangles σ_1, σ_2 such that R^U violates the local convexity on the union of (bijective images) σ_1^U and σ_2^U . Suppose that σ_i has vertices x_i, y_i, r_i with hypotenuse $x_i y_i = R \cap \sigma_i$ ($i = 1, 2$), and that $(r_1^U, x_1^U) = (r_2^U, x_2^U)$. Consider the gate g of r_1 in the folder containing σ_2 . If $g = r_2$, then by Lemma 3.12 we have $d(r_1, R) = d(r_1, g) + 1 > 1 = d(r_1, R)$. Therefore $g \neq r_2$. Take a shortest path P from r_1 to r_2 including g , and consider its image P^U , which is a shortest path connecting r_1^U and r_2^U in Γ^U by (3). However its length is at least 1. A contradiction to $r_1^U = r_2^U$. \square




Figure 9: the construction of U_i ; the bold lines represent U_2

Note that (2) and (3) holds without orientability condition. Now we are ready to prove (3.1) (ii). For a map $\rho : X \rightarrow |\Delta|$, we define metric d^ρ on X by

$$(3.12) \quad d^\rho(x, y) = d_\Delta(\rho(x), \rho(y)) \quad (x, y \in X).$$

Proposition 3.16. *Let Δ be an orientable F -complex, and let Γ be its leg-graph. For any map $\rho : X \rightarrow |\Delta|$ on a finite set X , there exist maps $\rho_i : X \rightarrow V\Gamma$ ($i \in I$) such that d^ρ is a convex combination of d^{ρ_i} , and for any normal set R , $\rho(x) \in R$ implies $\rho_i(x) \in R$ ($i \in I$).*

We may consider the case where $\rho(X) \subseteq V\Gamma^k$ for sufficiently large k . Each edge e of Γ is subdivided into k edges e_1, e_2, \dots, e_k . Fix an orientation of Γ with property (2.5). Suppose that (e_1, e_2, \dots, e_k) is arranged from the source to the sink in the orientation of e . Let U_i be the set of edges projective to i -th edge e_i . By construction, $U_i \cap U_j = \emptyset$ for $i \neq j$, and U_i is the union of orbits of Δ^k ; see Figure 9. By Proposition 3.15 (3), we have

$$d_\Gamma(p, q) = \frac{1}{k} \sum_{i=1}^k d_{(\Gamma^k)U_i}(p^{U_i}, q^{U_i}) \quad (p, q \in V\Gamma).$$

Each Δ^{U_i} is isomorphic to Δ up to dilation factor $1/k$. Therefore $p \mapsto p^{U_i}$ induces the map $\phi_i : |\Delta| \rightarrow V\Gamma$. Then $\phi_i \circ \rho$ ($i = 1, 2, \dots, k$) are desired maps. Also, by construction, $\rho(x) \in \sigma$ implies $\rho_i(x) \in \sigma$ for any cell $\sigma \in \Delta$. This implies the latter statement.

4 Related results

In this section, we present two results related to our argument. The first one is an optimal criterion of a combinatorial dual problem (2.7), which roughly characterizes the adjacency of extreme points of the feasible region in LP-dual (3.2). This result will play a crucial role in the splitting-off argument [13]. The second one is a characterization of distances of combinatorial dimension at most 2 in terms of F -complex realizations.

4.1 An optimality criterion for combinatorial dual problems

Here we show that (2.7) has an optimality criterion of the following type:

If ρ is not optimal, then there exists another ρ' close to ρ having the smaller objective value.

We will characterize the closedness in terms of orientations of Γ . An orientation of Γ with property (2.5) is said to be *admissible*. Admissible orientations are obtained by orienting orbits independently. Such an orientation of an orbit is also said to be

admissible. Each orbit has exactly two admissible orientations; one is the reverse of the other one. A map ρ feasible to (2.7) is called a *potential*.

Let \vec{O} be an oriented orbit by an admissible orientation. A potential ρ' is called a *neighbor* of a potential ρ with respect to \vec{O} if for each $x \in VG$ with $\rho'(x) \neq \rho(x)$

- (4.1) (i) there is an oriented edge $\vec{pq} \in \vec{O}$ such that $\rho(x) = p$ and $\rho'(x) = q$, or
- (ii) there are two oriented edges $\vec{pq}, \vec{qr} \in \vec{O}$ belonging to a common 2-cell in Δ such that $\rho(x) = p$ and $\rho'(x) = r$.

Main theorem here is:

Theorem 4.1. *If a potential ρ is not optimal, then there exists a neighbor ρ' of ρ having the smaller objective value.*

Remark 4.2. The combinatorial dual problem (2.7) is a variant of *multifacility location problem* on graph Γ [24]. From the point of the view, Theorem 4.1 can be understood as a generalization of a result of Dearing, Francis, and Lowe [5] for tree location problems (the case where Γ is a tree); see also [21, Chapter 3].

The proof is based on the idea used in [12]. A map ρ feasible to (2.6) is called a *continuous potential*. We denote the distance among points $\rho(x)$ by d^ρ as (3.12), and also denote the corresponding objective value $\sum_{xy \in EG} c(xy) d_\Gamma(\rho(x), \rho(y))$ simply by $\langle c, d^\rho \rangle$.

Recall the correspondence between metrics m feasible to (3.2) and a continuous potentials ρ , discussed in the beginning of Section 3. Namely, for a minimal feasible metric m , there is a continuous potential ρ satisfying $m = d^\rho$. In general, this correspondence $m \mapsto \rho$ is not one-to-one. However one can establish the one-to-one correspondence by the following trick. Add isolated terminals to S and extend μ so that $\{R_s\}_{s \in S}$ contains all singleton sets of $V\Gamma$. In this case, the correspondence $m \mapsto \rho$ is one-to-one, and is continuous. Suppose that for a minimal feasible metric m there are two continuous potentials ρ, ρ' with $m = d^\rho = d^{\rho'}$. Suppose $\rho(x) \neq \rho'(x)$ for some $x \in VG$. Take minimal cells σ and σ' of Δ containing $\rho(x)$ and $\rho'(x)$, respectively. Suppose that σ and σ' are distinct. Then there is a vertex v in σ such that $d_\Delta(v, \rho(x)) \neq d_\Delta(v, \rho'(x))$. By the hypothesis, there is a terminal s with $R_s = \{v\}$. Then $\rho(s) = \rho'(s) = v$ and $d^\rho(s, x) = d_\Gamma(v, \rho(x)) \neq d_\Gamma(v, \rho'(x)) = d^{\rho'}(s, x)$. A contradiction. Therefore $\sigma = \sigma'$. Since distances $d_\Delta(v, \rho(x)) (= d_\Delta(v, \rho'(x)))$ from vertices v of σ uniquely determines $\rho(x)$, two maps ρ and ρ' must coincide.

We are ready to prove the statement. We remark that if the image of ρ belongs to $V\Gamma$, then d^ρ is minimal; this also follows from the fact that $\{R_s\}_{s \in S}$ contains all singleton sets. Suppose that a potential ρ is not optimal. Equivalently, d^ρ is not optimal to (3.2). Since d^ρ is minimal and nonoptimal, there is another minimal feasible metric m' sufficiently close to d^ρ such that $\langle c, m' \rangle < \langle c, d^\rho \rangle$. By the minimality, m' is represented by $d^{\rho'}$ for a continuous potential ρ' . Since m' is sufficiently close to d^ρ and $m' \mapsto \rho'$ is continuous, we may assume that

$$(4.2) \quad d_\Delta(\rho(x), \rho'(x)) < \kappa/2 \quad (x \in VG).$$

Recall that κ is the leg-length of Δ . Furthermore we may assume that $\rho(VG) \subseteq V\Gamma^k$ for sufficiently large k . Fix some admissible orientation $\vec{\Gamma}$ of Γ . Decompose $E\Gamma^k$ into U_1, U_2, \dots, U_k and define maps $\phi_1, \phi_2, \dots, \phi_k : V\Gamma \rightarrow V\Gamma$ as in the proof of Proposition 3.16. Then we have $d^{\rho'} = \sum_{i=1}^k d^{\phi_i \circ \rho'} / k$. Therefore there is an index i such that $\langle c, d^{\phi_i \circ \rho'} \rangle < \langle c, d^\rho \rangle$. We may assume that $1 \leq i \leq k/2$ by reversing the orientation if

necessarily. Compare $\phi_i \circ \rho'$ with ρ . By the condition (4.2), if $(\phi_i \circ \rho')(x) \neq \rho(x)$, then

- (4.3) (i) there is an oriented edge $\vec{pq} \in \vec{E\Gamma}$ such that $\rho(x) = p$ and $(\phi_i \circ \rho')(x) = q$,
or
(ii) there are two oriented edges $\vec{pq}, \vec{qr} \in \vec{E\Gamma}$ belonging to a common 2-cell in Δ such that $\rho(x) = p$ and $(\phi_i \circ \rho')(x) = r$.

Indeed, we can take a cell σ in Δ containing both $\rho'(x)$ and $\rho(x)$. Then $(\phi_i \circ \rho')(x) \neq \rho(x)$ if and only if a shortest path P in Γ^k connecting $\rho'(x)$ and $\rho(x)$ crosses U_i ; recall Figure 9. If $i > k/2$, then P never crosses U_i by (4.2). Thus if P crosses U_i , then $p = \rho(x)$ is necessarily a source of an oriented leg of σ . Since P crosses edges in U_i at most twice, the position of $(\phi_i \circ \rho')(x)$ is (i) (P crosses U_i once) or (ii) (P crosses U_i twice).

Let $\mathcal{O} = \{O_1, O_2, \dots, O_m\}$ be the set of orbits of Δ . As in Section 3.4, we define maps $\varphi_j : V\Gamma \rightarrow V\Gamma^{O_j}$ by $p \mapsto p^{O_j}$ ($j = 1, 2, \dots, m$). By Proposition 3.15, we have $d^\rho = \sum_{j=1}^k d^{\varphi_j \circ \rho}$ (for any potential ρ). Therefore we have $\sum_{j=1}^m \langle c, d^{\varphi_j \circ \phi_i \circ \rho'} \rangle < \sum_{j=1}^m \langle c, d^{\varphi_j \circ \rho} \rangle$. Then there is an index j with $\langle c, d^{\varphi_j \circ \phi_i \circ \rho'} \rangle < \langle c, d^{\varphi_j \circ \rho} \rangle$. Let \vec{O}_j be the admissible orientation of O_j induced by $\vec{\Gamma}$. By using \vec{O}_j and $\phi_i \circ \rho'$, we can construct a required neighbor ρ'' of ρ with respect to \vec{O}_j as follows. For each $x \in VG$, if there is $\vec{pq} \in \vec{O}_j$ with $\rho(x) = p$ and $(\varphi_j \circ \phi_i \circ \rho')(x) = q^{O_j}$ or there are $\vec{pr}, \vec{rq} \in \vec{O}_j$ belonging to a common 2-cell with $\rho(x) = p$ and $(\varphi_j \circ \phi_i \circ \rho')(x) = q^{O_j}$, then define $\rho''(x) = q$. For other vertex x , define $\rho''(x) = \rho(x)$. By construction, ρ'' is a neighbor of ρ w.r.t. \vec{O}_j , and we have

$$\varphi_l \circ \rho'' = \begin{cases} \varphi_j \circ \phi_i \circ \rho' & \text{if } l = j, \\ \varphi_l \circ \rho & \text{otherwise,} \end{cases} \quad (l = 1, 2, \dots, m).$$

Then we have $\langle c, d^{\rho''} \rangle = \sum_{l=1}^m \langle c, d^{\varphi_l \circ \rho''} \rangle = \sum_{l \neq j} \langle c, d^{\varphi_l \circ \rho} \rangle + \langle c, d^{\varphi_j \circ \phi_i \circ \rho'} \rangle < \sum_{l=1}^m \langle c, d^{\varphi_l \circ \rho} \rangle = \langle c, d^\rho \rangle$. Thus ρ'' is a required neighbor.

4.2 Distances of combinatorial dimension 2

For a distance μ on a finite set S , define two polyhedral sets P_μ and T_μ in \mathbf{R}^S by

$$\begin{aligned} P_\mu &= \{p \in \mathbf{R}^S \mid p(s) + p(t) \geq \mu(s, t) \quad (s, t \in S)\}, \\ T_\mu &= \text{the set of minimal elements of } P_\mu. \end{aligned}$$

T_μ is called the *tight span* of μ [15, 6]. Its dimension $\dim T_\mu$ is defined to be the largest dimension of its faces, and is called the *combinatorial dimension* of μ [6]. The geometry of T_μ reflects the combinatorial property of μ as follows.

- (4.4) (i) For a metric μ , $\dim T_\mu \leq 1$ if and only if μ is a tree metric [6].
(ii) For a distance μ , $\dim T_\mu \leq 1$ if and only if μ is a distance between subtrees of a tree [10].
(iii) For a rational metric, $\dim T_\mu \leq 2$ if and only if μ is a dilation of a submetric of a frame [18].

The main aim here is to give an extension of (iii) to a nonmetric version as follows.

Theorem 4.3. *Let μ be a rational distance on a finite set S . The following two conditions are equivalent:*

(1) $\dim T_\mu \leq 2$.

(2) μ has an F -complex realization.

The larger part of the proof has already been given in [11]. So we sketch it. First we show (2) \Rightarrow (1). Suppose that μ has an F -complex realization. Then there exists a positive integer k (the denominator of rational leg-length κ) such that LP-dual (3.2) always has a $1/k$ -integral optimal solution. Namely, the *dual fractionality* of μ is finite. Then [11, Theorem 1.1] implies $\dim T_\mu \leq 2$.

Second, we show (1) \Rightarrow (2). Suppose $\dim T_\mu \leq 2$. Then [11] showed that T_μ has a polyhedral subdivision Δ with property (2.1), and (T_μ, l_∞) is isometric to $(|\Delta|, d_\Delta)$. For $s \in S$, define a subset $T_{\mu,s}$ by $\{p \in T_\mu \mid p(s) = 0\}$. We verify that $(\Delta; \{T_{\mu,s}\}_{s \in S})$ is an F -complex realization of μ . It is known that $d_\Delta(T_{\mu,s}, T_{\mu,t}) = \mu(s, t)$ [11, Lemma 3.6]. Hence it suffices to verify that Δ is an F -complex and each $T_{\mu,s}$ is normal. Although one can directly prove that Δ satisfies (2.3), this approach needs full details of [11, Section 3]. Instead, we use Chepoi's characterization [3] of F -complexes in terms of hyperconvexity. Here a metric space is said to be *hyperconvex* (in the sense of [1]) if the family of balls (around points) has the Helly property.

Theorem 4.4 ([3, Theorem 7.8]). *A locally-finite and simply-connected 2-complex Δ satisfying (2.1) without triangle-folders is an F -complex if and only if $(|\Delta|, d_\Delta)$ is hyperconvex.*

In fact, this theorem holds for any finite and simply-connected 2-complex with property (2.1) possibly having triangle-folders. Indeed, the only if-part follows from Theorem 3.13 (with some care for infinite families and irrational radius). Also the if-part follows from a slight change in Chepoi's proof by the following way. Suppose that $(|\Delta|, d_\Delta)$ is hyperconvex, and suppose to the contrary that there are three folders F_1, F_2, F_3 violating (2.3) at a common point p . Some of F_1, F_2, F_3 may be triangle-folders. In this case, consider 2-subdivision Δ^2 . Again we can take folders violating (2.3) at p each of which is not a triangle-folder. Then apply the argument in [3, p. 156] to them, which leads a contradiction. So it suffices to verify the following, which is well-known in the literature for metrics μ .

Lemma 4.5. *(T_μ, l_∞) is hyperconvex.*

This is an easy consequence of the next lemma.

Lemma 4.6 ([6, p.331, (1.9)]). *There exists a map $\phi : P_\mu \rightarrow T_\mu$ such that*

- (1) $\|\phi(p) - \phi(q)\|_\infty \leq \|p - q\|_\infty$ for $p, q \in P_\mu$, and
- (2) $\phi(p) \leq p$ for $p \in P_\mu$, and thus ϕ is identical on T_μ .

(This lemma holds for nonmetric distances μ although it was originally stated for metrics.)

The proof of Lemma 4.5. For $\{p_i\}_{i \in I} \subseteq T_\mu$ and $\{r_i\}_{i \in I} \subseteq \mathbf{R}_+$, suppose that the collection of balls $\{B(p_i, r_i)\}_{i \in I}$ in (\mathbf{R}^S, l_∞) has pairwise intersection at T_μ . Each ball $B(p_i, r_i) \subseteq \mathbf{R}^S$ is a direct product of segments $[p_i(s) - r_i, p_i(s) + r_i]$ and thus $\{B(p_i, r_i)\}_{i \in I}$ has the Helly property on (\mathbf{R}^S, l_∞) . Let p^* be a unique maximal element in (nonempty intersection) $\bigcap_{i \in I} B(p_i, r_i)$. Then one can verify that p^* belongs to P_μ . Take a nonexpansive map ϕ in the previous lemma. Then $\phi(p^*)$ belongs to $\bigcap_{i \in I} T_\mu \cap B(p_i, r_i)$. \square

Finally we verify that $T_{\mu,s}$ is normal. For a face F of P_μ , let K_F be the graph on S with edge set $EK_F = \{st \mid p(s) + p(t) = \mu(s, t) \ (p \in F, s, t \in S)\}$. Namely K_F represents the set of facets of P_μ active at F . Note that K_F has a loop ss exactly when $F \subseteq \{p \in \mathbf{R}^S \mid p(s) = 0\}$. Then F belongs to T_μ if and only if K_F has no isolated vertices. $T_{\mu,s}$ is the union of subcomplex of Δ since $T_{\mu,s}$ is the union of faces F of T_μ whose K_F has a loop at s . The connectivity is obvious. We show that every face containing e belongs to $T_{\mu,s}$, which implies (2.4) (i). A leg coincides with an l_1 -edge in the sense of [11]. Let F be a minimal face of T_μ containing e . In the graph K_F , s belongs to the connected component each of whose vertices has a loop [11, Lemma 3.8]. So every face F' of T_μ containing F necessarily belongs to $T_{\mu,s}$. Indeed, by $EK_{F'} \subseteq EK_F$, $ss \notin EK_{F'}$ implies that s is isolated in $K_{F'}$ ($F' \not\subseteq T_\mu$). Suppose to the contrary that $T_{\mu,s}$ violates the local convexity condition, i.e., there are two triangles $\sigma, \sigma' \in \Delta$ of vertices p, r, v and q, r, v (resp.) such that pr and rq are hypotenuse with $T_{\mu,s} \cap (\sigma \cup \sigma') = pr \cap rq$. Then $\|p - q\|_\infty = 2\kappa$. Consider a geodesic P connecting p and r in $\{p \in P_\mu \mid p(s) = 0\}$. Consider the image $\phi(P)$ of P by a nonexpansive map ϕ in Lemma 4.6. Then $\phi(P)$ is also a geodesic connecting p and q in $T_{\mu,s}$ with length 2κ . By (3.6) and (2.4) (i), Δ must have a triangle of vertices p, q, v and hypotenuse pq , which violates the flag condition at v . A contradiction.

Remark 4.7. A metric space X is called *injective* (or *an absolute retract*) if for every metric space Y containing X as a subspace there exists a nonexpansive retraction from Y to X . Aronszajn and Panitchpakdi [1] showed that X is injective if and only if X is hyperconvex. So two Lemmas 4.5 and 4.6 are essentially equivalent.

Acknowledgement

This work is supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- [1] N. Aronszajn and P. Panitchpakdi, Extensions of uniformly continuous transformations and hyperconvex metric spaces, *Pacific Journal of Mathematics* **6** (1956), 405–439.
- [2] H.-J. Bandelt, Hereditary modular graphs, *Combinatorica* **8** (1988), 149–157.
- [3] V. Chepoi, Graphs of some CAT(0) complexes, *Advances in Applied Mathematics* **24** (2000), 125–179.
- [4] B. V. Cherkassky, Solution of a problem of multiproduct flows in a network, *Ekonomika i Matematicheskie Metody* **13** (1977), 143–151, (in Russian).
- [5] P. M. Dearing, R. L. Francis, and T. J. Lowe, Convex location problems on tree networks. *Operations Research* **24** (1976), 628–642.
- [6] A. W. M. Dress, Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups: a note on combinatorial properties of metric spaces, *Advances in Mathematics* **53** (1984), 321–402.
- [7] A. W. M. Dress and R. Scharlau, Gated sets in metric spaces, *Aequationes Mathematicae* **34** (1987), 112–120.

- [8] L. R. Ford and D. R. Fulkerson, *Flows in networks*, Princeton University Press, Princeton, 1962.
- [9] A. Frank, A. V. Karzanov, and A. Sebö, On integer multiflow maximization, *SIAM Journal on Discrete Mathematics* **10** (1997), 158–170.
- [10] H. Hirai, Characterization of the distance between subtrees of a tree by the associated tight span, *Annals of Combinatorics* **10** (2006), 111–128.
- [11] H. Hirai, Tight spans of distances and the dual fractionality of undirected multiflow problems, *Journal of Combinatorial Theory, Series B*, to appear.
- [12] H. Hirai, Bounded fractionality of the multiflow feasibility problem for demand graph $K_3 + K_3$ and related maximization problems, RIMS-preprint 1645, (2008).
- [13] H. Hirai, The maximum multiflow problems with bounded fractionality, in preparation.
- [14] T. C. Hu, Multi-commodity network flows, *Operations Research* **11** (1963), 344–360.
- [15] J. R. Isbell, Six theorems about injective metric spaces, *Commentarii Mathematici Helvetici* **39** (1964), 65–76.
- [16] A. V. Karzanov, Polyhedra related to undirected multicommodity flows, *Linear Algebra and its Applications* **114/115** (1989), 293–328.
- [17] A. V. Karzanov, Minimum 0-extensions of graph metrics, *European Journal of Combinatorics* **19** (1998), 71–101.
- [18] A. V. Karzanov, Metrics with finite sets of primitive extensions, *Annals of Combinatorics* **2** (1998), 211–241.
- [19] A. V. Karzanov, One more well-solved case of the multifacility location problem, *Discrete Optimization*, **1** (2004), 51–66.
- [20] A. V. Karzanov and M. V. Lomonosov, Systems of flows in undirected networks, in: *Mathematical Programming* (O.I. Larichev, ed.), Institute for System Studies, Moscow, 1978, 59–66 (in Russian).
- [21] A. W. J. Kolen, *Tree network and planar rectilinear location theory*. CWI Tract, 25, Amsterdam, 1986.
- [22] M. V. Lomonosov, Combinatorial approaches to multiflow problems, *Discrete Applied Mathematics* **11** (1985), 93 pp.
- [23] L. Lovász, On some connectivity properties of Eulerian graphs, *Acta Mathematica Academiae Scientiarum Hungaricae* **28** (1976), 129–138.
- [24] B. C. Tansel, R. L. Francis, and T. J. Lowe, Location on networks: a survey. II. Exploiting tree network structure, *Management Science* **29** (1983), 498–511.