

RIMS-1694

**Lie Group-Lie Algebra Correspondences of
Unitary Groups in Finite von Neumann Algebras**

By

Hiroshi ANDO and Yasumichi MATSUZAWA

May 2010



京都大学 数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

Lie Group-Lie Algebra Correspondences of Unitary Groups in Finite von Neumann Algebras

Hiroshi Ando

Research Institute for Mathematical Sciences, Kyoto University
Kyoto, 606-8502, Japan

E-mail: andonuts@kurims.kyoto-u.ac.jp

Yasumichi Matsuzawa^{1,2,*}

¹ Mathematisches Institut, Universität Leipzig
Johannisgasse 26, 04103, Leipzig, Germany

² Department of Mathematics, Hokkaido University
Kita 10, Nishi 8, Kita-ku, Sapporo, 060-0810, Japan

E-mail: matsuzawa@math.sci.hokudai.ac.jp

May 6, 2010

Abstract

We give an affirmative answer to the question whether there exist Lie algebras for suitable closed subgroups of the unitary group $U(\mathcal{H})$ in a Hilbert space \mathcal{H} with $U(\mathcal{H})$ equipped with the strong operator topology. More precisely, for any strongly closed subgroup G of the unitary group $U(\mathfrak{M})$ in a finite von Neumann algebra \mathfrak{M} , we show that the set of all generators of strongly continuous one-parameter subgroups of G forms a complete topological Lie algebra with respect to the strong resolvent topology. We also characterize the algebra $\overline{\mathfrak{M}}$ of all densely defined closed operators affiliated with \mathfrak{M} from the viewpoint of a tensor category.

Keywords. finite von Neumann algebra, unitary group, affiliated operator, measurable operator, strong resolvent topology, tensor category, infinite dimensional Lie group, infinite dimensional Lie algebra.

Mathematics Subject Classification (2000). 22E65, 46L51.

*Supported by Research fellowships of the Japan Society for the Promotion of Science for Young Scientists (Grant No. 2100165000).

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction and Main Theorem | 2 |
| 2 | Preliminaries | 5 |
| 2.1 | von Neumann Algebras | 5 |
| 2.2 | Murray-von Neumann's Result | 7 |
| 2.3 | Converse of Murray-von Neumann's Result | 14 |
| 3 | Topological Structures of $\overline{\mathfrak{M}}$ | 15 |
| 3.1 | Strong Resolvent Topology | 15 |
| 3.2 | Strong Exponential Topology | 19 |
| 3.3 | τ -Measure Topology | 20 |
| 3.4 | Almost Everywhere Convergence | 20 |
| 3.5 | Proof of Lemma 3.11 | 22 |
| 3.6 | Direct Sums of Algebras of Unbounded Operators | 24 |
| 3.7 | Proof of Theorem 3.9 | 29 |
| 3.8 | Local Convexity | 29 |
| 4 | Lie Group-Lie Algebra Correspondences | 32 |
| 4.1 | Existence of Lie Algebra | 32 |
| 4.2 | Closed Subalgebras of $\overline{\mathfrak{M}}$ | 35 |
| 5 | Categorical Characterization of $\overline{\mathfrak{M}}$ | 37 |
| 5.1 | Introduction | 37 |
| 5.2 | fvN and fRng as Tensor Categories | 38 |
| A | Direct Sums of Operators | 49 |
| B | Fundamental Results of SRT | 51 |
| C | Tensor Categories | 52 |

1 Introduction and Main Theorem

Lie groups played important roles in mathematics because of its close relations with the notion of symmetries. They appear in almost all branches of mathematics and have many applications. While Lie groups are usually understood as finite dimensional ones, many infinite dimensional symmetries appear in natural ways: for instance, loop groups $C^\infty(S^1, G)$ [17], current groups $C_c^\infty(M, G)$ [1], diffeomorphism groups $Diff^\infty(M)$ of manifolds [3] and Hilbert-Schmidt groups [5] are among well-known cases. They have been extensively investigated in several concrete ways.

In this context, it would be meaningful to consider a general theory of infinite dimensional Lie groups. One of the most fundamental infinite dimensional groups are Banach-Lie groups. They are modeled on Banach spaces and many

theorems in finite dimensional cases are also applicable to them. Since it has been shown that a Banach-Lie group cannot act transitively and effectively on a compact manifold as a transformation group [16], however, Banach-Lie groups are not sufficient for treating infinite dimensional symmetries. After the birth of Banach-Lie group theory, more general notions of infinite dimensional Lie groups have been scrutinized to date: locally convex Lie groups [12], ILB-Lie groups [15], pro-Lie groups [6, 7] and so on. While there are many interesting and important results about them, we note that not all theorems in finite dimensional cases remain valid in these categories and their treatments are complicated. For example, the exponential map might not be a local homeomorphism and the Baker-Campbell-Hausdorff formula may no longer be true [10].

We understand that the one of the most fundamental class of finite dimensional Lie groups are the unitary groups $U(n)$ in such a sense that any compact Lie group can be realized as a closed subgroup of them. From this viewpoint, it would be important to study the infinite dimensional analogue of it; that is, we like to explicate the Lie theory for the unitary group $U(\mathcal{H})$ of an infinite dimensional Hilbert space \mathcal{H} . One of the most fundamental question is whether $\text{Lie}(G)$ defined as the set of all generators of continuous one-parameter subgroups of a closed subgroup G of $U(\mathcal{H})$ forms a Lie algebra or not. For the infinite dimensional Hilbert space \mathcal{H} , there are at least two topologies on $U(\mathcal{H})$, (a) the norm topology and (b) the strong operator topology. We discuss the above topologies separately. In the case (a), $U(\mathcal{H})$ is a Banach-Lie group and for each closed subgroup the set $\text{Lie}(G)$ forms a Lie algebra. But it is well known that there are not many “nice” continuous unitary representations of groups in \mathcal{H} , and hence, $U(\mathcal{H})$ with the norm topology is very narrow. On the other hand, $U(\mathcal{H})$ with the strong operator topology (b) is important, because there are many “nice” continuous unitary representations of groups in \mathcal{H} —say, diffeomorphism groups of compact manifolds, etc. However, the answer is negative to the question whether there exists a corresponding Lie algebra or not. Indeed, by the Stone theorem, the Lie algebra of $U(\mathcal{H})$ coincides with the set of all (possibly unbounded) skew-adjoint operators on \mathcal{H} , but we cannot define naturally a Lie algebra structure with addition and Lie bracket operations on it. This arises from the problem of the domains of unbounded operators. For two skew-adjoint operators A, B on \mathcal{H} , $\text{dom}(A + B) = \text{dom}(A) \cap \text{dom}(B)$ is not always dense. Even worse, it can be $\{0\}$. Because of this, the Lie theory for $U(\mathcal{H})$ has not been successful, although the group itself is a very natural object. On the other hand, it is possible that even though the whole group $U(\mathcal{H})$ does not have a Lie algebra, some suitable class of closed subgroups of it have ones. Indeed their Lie algebras $\text{Lie}(G)$ are smaller than $\text{Lie}(U(\mathcal{H}))$.

We give an affirmative answer to the last question. Furthermore we prove that for a suitable subgroup G , $\text{Lie}(G)$ is a complete topological Lie algebra with respect to some natural topology. We outline below the essence of our detailed discussions in the text.

First, a group G to be studied in this paper is a closed subgroup of the unitary group $U(\mathfrak{M})$ of some finite von Neumann algebra \mathfrak{M} acting on a Hilbert space \mathcal{H} . Clearly it is also a closed subgroup of $U(\mathcal{H})$. The key proposition is

the following result of Murray-von Neumann (cf. Theorem 2.17):

Theorem 1.1 (Murray-von Neumann). *The set $\overline{\mathfrak{M}}$ of all densely defined closed operators affiliated with a finite von Neumann algebra \mathfrak{M} on \mathcal{H} ,*

$$\overline{\mathfrak{M}} := \left\{ A ; \begin{array}{l} A \text{ is a densely defined closed operator on } \mathcal{H} \\ \text{such that } uAu^* = A \text{ for all } u \in U(\mathfrak{M}'). \end{array} \right\},$$

constitutes a $$ -algebra under the sum $\overline{A+B}$, the scalar multiplication $\overline{\alpha A}$ ($\alpha \in \mathbb{C}$), the product \overline{AB} and the involution A^* , where \overline{X} denotes the closure of a closable operator X .*

The inclusion $G \subset U(\mathfrak{M})$ implies $\text{Lie}(G) \subset \overline{\mathfrak{M}}$ and hence, for arbitrary two elements $A, B \in \text{Lie}(G)$, the sum $\overline{A+B}$, the scalar multiplication $\overline{\alpha A}$, the Lie bracket $[A, B] := \overline{AB - BA}$ are determined as elements of $\overline{\mathfrak{M}}$. We can prove that they are again elements of $\text{Lie}(G)$, which is not trivial. Therefore $\text{Lie}(G)$ indeed forms a Lie algebra which is infinite dimensional in general. Thus if we do not introduce a topology, it is difficult to investigate it. Then, what is the natural topology on $\text{Lie}(G)$? Since $\text{Lie}(G)$ is a Lie algebra, it should be a vector space topology. Furthermore, in view of the correspondences between Lie groups and Lie algebras it is natural to require the continuity of the mapping

$$\exp : \text{Lie}(G) \ni A \longmapsto e^A \in G,$$

where G is equipped with the strong operator topology and e^A is defined by the spectral theorem. Under these assumptions, a necessary condition for a sequence $\{A_n\}_{n=1}^\infty \subset \text{Lie}(G)$ to converge to $A \in \text{Lie}(G)$ is given by

$$\text{s-} \lim_{n \rightarrow \infty} e^{tA_n} = e^{tA}, \quad \text{for all } t \in \mathbb{R}.$$

This condition is equivalent to

$$\text{s-} \lim_{n \rightarrow \infty} (A_n + 1)^{-1} = (A + 1)^{-1}.$$

The latter convergence is well known in the field of (unbounded) operator theory as the convergence with respect to the strong resolvent topology. Therefore it seems natural to consider the strong resolvent topology for $\text{Lie}(G)$. However, there arises, unfortunately, another troublesome question as to whether the vector space operations and the Lie bracket operation are continuous with respect to the strong resolvent topology of $\text{Lie}(G)$. For example, even if sequences $\{A_n\}_{n=1}^\infty, \{B_n\}_{n=1}^\infty$ of skew-adjoint operators converge, respectively to skew-adjoint operators A, B with respect to the strong resolvent topology, the sequences $\{A_n + B_n\}_{n=1}^\infty$ are not guaranteed to converge to $A + B$. We can solve this difficulty by applying the noncommutative integration theory and proving that the Lie algebraic operations are continuous with respect to the strong resolvent topology and that $\text{Lie}(G)$ is complete as a uniform space. Hence $\text{Lie}(G)$ forms a complete topological Lie algebra. Finally, let us remark one point: remarkably, $\text{Lie}(G)$ is not locally convex in general. Most of infinite dimensional

Lie theories assume the locally convexity explicitly, but as soon as we consider such groups as natural infinite dimensional analogues of classical Lie groups, there appear non-locally convex examples.

We shall explain the contents of the paper. §2 is a preliminary section. We recall the basic facts about closed operators affiliated with a finite von Neumann algebra and explain the generalization of the Murray-von Neumann theorem for non-countably decomposable case. In §3, we introduce three topologies on the set $\overline{\mathfrak{M}}$ of all densely defined closed operators affiliated with a finite von Neumann algebra \mathfrak{M} . The first topology originates from (unbounded) operator theory, the second one is Lie theoretical and the last one derives from the noncommutative integration theory. We discuss their topological properties and show that they do coincide on $\overline{\mathfrak{M}}$. The main result of this section is Theorem 3.9 which states that $\overline{\mathfrak{M}}$ forms a complete topological $*$ -algebra with respect to the strong resolvent topology. In §4 constituting the main contents of the paper, we show that $\text{Lie}(G)$ is a complete topological Lie algebra and discuss some aspects of it. The main result is given in Theorem 4.6. In §5, applying the results of §3, we consider the following problem: *What kind of unbounded operator algebras can they be represented in the form of $\overline{\mathfrak{M}}$?* We give their characterization from the viewpoint of a tensor category. We show that \mathcal{R} can be represented as $\overline{\mathfrak{M}}$ if and only if it is an object of the category \mathbf{fRng} (cf. Definition 5.2). The main result is Theorem 5.5, which says that the category \mathbf{fRng} is isomorphic to the category \mathbf{fvN} of finite von Neumann algebras as a tensor category. In Appendix, we list up some fundamental definitions and results of the direct sums, the strong resolvent convergence and the categories.

2 Preliminaries

In this section we review some basic facts about operator algebras and unbounded operators. For the details, see [18, 21]. See also Appendix A for the direct sums.

2.1 von Neumann Algebras

Let \mathcal{H} be a Hilbert space with an inner product $\langle \xi, \eta \rangle$, which is linear with respect to η . We denote the algebra of all bounded operators on \mathcal{H} by $\mathfrak{B}(\mathcal{H})$. Let \mathfrak{M} be a von Neumann algebra acting on \mathcal{H} . The set $\mathfrak{M}' := \{x \in \mathfrak{B}(\mathcal{H}) ; xy = yx, \text{ for all } y \in \mathfrak{M}\}$ is called the *commutant* of \mathfrak{M} . The group of all unitary operators in \mathfrak{M} is denoted by $U(\mathfrak{M})$. The lattice of all projections in \mathfrak{M} is denoted by $P(\mathfrak{M})$. The orthogonal projection onto the closed subspace $\mathcal{K} \subset \mathcal{H}$ is denoted by $P_{\mathcal{K}}$.

Definition 2.1. Let \mathfrak{M} be a von Neumann algebra acting on a Hilbert space \mathcal{H} .

- (1) A von Neumann algebra with no non-unitary isometry is called *finite*.

- (2) A von Neumann algebra is called *countably decomposable* if it admits at most countably many non-zero orthogonal projections.
- (3) A subset \mathcal{D} of \mathcal{H} is called *separating* for \mathfrak{M} if $x\xi = 0$, $x \in \mathfrak{M}$ for all $\xi \in \mathcal{D}$ implies $x = 0$.

It is known that a von Neumann algebra \mathfrak{M} acting on a Hilbert space \mathcal{H} is countably decomposable if and only if there exists a countable separating subset of \mathcal{H} for \mathfrak{M} .

Definition 2.2. Let \mathfrak{M} be a von Neumann algebra.

- (1) A state τ on \mathfrak{M} is called *tracial* if for all $x, y \in \mathfrak{M}$,

$$\tau(xy) = \tau(yx)$$

holds.

- (2) A tracial state τ is called *faithful* if $\tau(x^*x) = 0$ ($x \in \mathfrak{M}$) implies $x = 0$.
- (3) A tracial state τ is called *normal* if it is σ -weakly continuous.

It is known that a von Neumann algebra is countably decomposable and finite if and only if there exists a faithful normal tracial state on it. For more informations about tracial states, see [21].

Let \mathfrak{M} be a von Neumann algebra and $p \in \mathfrak{M} \cup \mathfrak{M}'$ be a projection. Define the set \mathfrak{M}_p of bounded operators on the Hilbert space $\text{ran}(p)$ as

$$\{px|_{\text{ran}(p)} ; x \in \mathfrak{M}\},$$

then \mathfrak{M}_p forms a von Neumann algebra acting on the Hilbert space $\text{ran}(p)$ and $(\mathfrak{M}_p)' = (\mathfrak{M}')_p$ holds.

If $(\mathfrak{M}, \mathcal{H})$ and $(\mathfrak{N}, \mathcal{K})$ are von Neumann algebras and if there exists a unitary operator U of \mathcal{H} onto \mathcal{K} such that

$$U\mathfrak{M}U^* = \mathfrak{N},$$

then $(\mathfrak{M}, \mathcal{H})$ and $(\mathfrak{N}, \mathcal{K})$ are said to be *spatially isomorphic*. The map π of \mathfrak{M} onto \mathfrak{N} defined by

$$\varphi(x) = UxU^*, \quad x \in \mathfrak{M},$$

is called a *spatial isomorphism*. The next is useful.

Lemma 2.3. *Let $(\mathfrak{M}, \mathcal{H})$ be a finite von Neumann algebra. Then there exists a family of countably decomposable finite von Neumann algebras $\{(\mathfrak{M}_\alpha, \mathcal{H}_\alpha)\}_\alpha$ such that $(\mathfrak{M}, \mathcal{H})$ is spatially isomorphic to the direct sum $\left(\bigoplus_\alpha^b \mathfrak{M}_\alpha, \bigoplus_\alpha \mathcal{H}_\alpha\right)$.*

A von Neumann algebra \mathfrak{M} is called *atomic* if each non-zero projection in \mathfrak{M} majorizes a non-zero minimal projection. It is known that a finite von Neumann algebra is atomic if and only if it is spatially isomorphic to the direct sum of

finite dimensional von Neumann algebras $M_n(\mathbb{C})$ ($n \in \mathbb{N}$), where $M_n(\mathbb{C})$ is the algebra of all $n \times n$ complex matrices.

A von Neumann algebra with no non-zero minimal projection is called *diffuse*. It is known that every von Neumann algebra is spatially isomorphic to the direct sum of some atomic von Neumann algebra $\mathfrak{M}_{\text{atomic}}$ and diffuse von Neumann algebra $\mathfrak{M}_{\text{diffuse}}$. These von Neumann algebras $\mathfrak{M}_{\text{atomic}}$ and $\mathfrak{M}_{\text{diffuse}}$ are unique up to spatial isomorphism. We call $\mathfrak{M}_{\text{atomic}}$ and $\mathfrak{M}_{\text{diffuse}}$ the *atomic part* and the *diffuse part* of \mathfrak{M} , respectively.

2.2 Murray-von Neumann's Result

The domain of a linear operator T on \mathcal{H} is written as $\text{dom}(T)$ and the range of it is written as $\text{ran}(T)$. If T is a closable operator, we write \overline{T} for the closure of T .

Definition 2.4. A densely defined closable operator T on \mathcal{H} is said to be *affiliated* with a von Neumann algebra \mathfrak{M} if for any $u \in U(\mathfrak{M}')$, $uTu^* = T$ holds. If T is affiliated with \mathfrak{M} , so is \overline{T} . The set of all densely defined closed operators affiliated with \mathfrak{M} is denoted by \mathfrak{M} .

Note that T is affiliated with \mathfrak{M} if and only if $xT \subset Tx$ for all $x \in \mathfrak{M}'$. Next, we define algebraic structures of unbounded operators in the style of Murray-von Neumann [11].

Let $x_1, y_1, x_2, y_2, \dots$ be (finite or countable infinite number of) indeterminants. A non-commutative monomial with indeterminants $\{x_i, y_i\}_i$ is a formal product $z_1 z_2 \cdots z_n$, where all z_k equal to x_i or y_i . If $n = 0$, we write this monomial as 1. A non-commutative polynomial $p(x_1, y_1, \dots)$ is a formal sum of finite number of monomials. $p(x_1, y_1, \dots)$ has the following form:

$$p(x_1, y_1, \dots) = \begin{cases} \sum_{\rho=1}^q a_\rho \cdot z_1^{(\rho)} \cdots z_{n_\rho}^{(\rho)} & (q = 1, 2, \dots), \\ 0 & (q = 0). \end{cases}$$

Here, $a_\rho \in \mathbb{C}$ and we allow such a term as $0 \cdot z_1 z_2 \cdots z_n$ in this expression. If there is a term with coefficient 0, it cannot be omitted in the representation. Hence x_1 is different from $x_1 + 0 \cdot y_1$ as non-commutative polynomials. If there are two such terms as $a \cdot z_1 \cdots z_n$, $b \cdot z_1 \cdots z_n$, we identify the sum of them with the term $(a + b) \cdot z_1 \cdots z_n$. The sum, the scalar multiplication and the multiplication of non-commutative polynomials are defined naturally, where we do not ignore the terms with 0 coefficients.

Once a non-commutative polynomial $p(x_1, y_1, \dots)$ is given we obtain a new polynomial $p^{(r)}(x_1, y_1, \dots)$ by omitting terms with coefficient $a_\rho = 0$ in the representation of p . We call $p^{(r)}(x_1, y_1, \dots)$ the reduced polynomial of p . We also define the adjoint element by $x_i^\dagger := y_i$, $y_i^\dagger := x_i$. We also define the

conjugate polynomial of p by

$$p(x_1, y_1, \dots)^+ := \begin{cases} \sum_{\rho=1}^q \overline{a_\rho} \cdot (z_{n_\rho}^{(\rho)})^+ \cdots (z_1^{(\rho)})^+ & (q = 1, 2, \dots), \\ 0 & (q = 0). \end{cases}$$

Suppose there is a corresponding sequence $\{X_i\}_i$ of densely defined closed operators on \mathcal{H} . For all i , we assume (x_i, y_i) corresponds to the pair of the closed operators (X_i, X_i^*) . In this case we define a new operator $p(X_1, X_1^*, \dots)$ obtained by substituting each $\{x_i, y_i\}$ in the representation of $p(x_1, y_1, \dots)$ of the pairs (X_i, X_i^*) . More precisely, the domain of $p(X_1, X_1^*, \dots)$ is defined according to the following rules:

- (1) $\text{dom}(0) = \text{dom}(1) = \mathcal{H}$, $0\xi := 0$, $1\xi := \xi$, for all $\xi \in \mathcal{H}$,
- (2) $\text{dom}(aX) := \text{dom}(X)$, $(aX)\xi := a(X\xi)$, for all $\xi \in \text{dom}(aX)$,
- (3) $\text{dom}(X + Y) := \text{dom}(X) \cap \text{dom}(Y)$, $(X + Y)\xi := X\xi + Y\xi$,
for all $\xi \in \text{dom}(X + Y)$,
- (4) $\text{dom}(XY) := \{\xi \in \text{dom}(Y); Y\xi \in \text{dom}(X)\}$, $(XY)\xi := X(Y\xi)$,
for all $\xi \in \text{dom}(XY)$,

where X and Y are densely defined closed operators on \mathcal{H} and $a \in \mathbb{C}$. In general, \mathfrak{M} is not a *-algebra under these operations. This is the reason for the difficulty of constructing Lie theory in infinite dimensions. However, Murray and von Neumann proved, in the pioneering paper [11], that for a finite von Neumann algebra \mathfrak{M} , \mathfrak{M} does constitute a *-algebra of unbounded operators, which we will explain more precisely in the sequel.

Murray-von Neumann proved the following results for a countably decomposable case. Since we need to apply these results for a general finite von Neumann algebra case, we shall offer the generalization of their proofs. First of all, we recall the notion of complete density, which is important for later discussions.

Definition 2.5. A subspace $\mathcal{D} \subset \mathcal{H}$ is said to be *completely dense* for a finite von Neumann algebra \mathfrak{M} if there exists an increasing net $\{p_\alpha\}_\alpha \subset P(\mathfrak{M})$ of projections in \mathfrak{M} such that

- (1) $p_\alpha \nearrow 1$ (strongly).
- (2) $p_\alpha \mathcal{H} \subset \mathcal{D}$ for any α .

It is clear that a completely dense subspace is dense in \mathcal{H} . We often omit the phrase “for \mathfrak{M} ” when the von Neumann algebra in consideration is obvious from the context.

Remark 2.6. In [11], Murray and von Neumann used the term “strongly dense”. However, this terminology is somewhat confusing. Therefore we tentatively use the term “completely dense”.

Lemma 2.7. *Let \mathfrak{M} be a countably decomposable, finite von Neuman algebra on a Hilbert space \mathcal{H} , τ be a faithful normal tracial state on \mathfrak{M} . For a completely dense subspace $\mathcal{D} \subset \mathcal{H}$, the following are equivalent.*

(1) \mathcal{D} is completely dense.

(2) There exists an increasing sequence $\{p_n\}_{n=1}^\infty \subset P(\mathfrak{M})$ such that

$$p_n \nearrow 1 \text{ (strongly)}, \quad \text{ran}(p_n) \subset \mathcal{D}.$$

(3) For every $\varepsilon > 0$, there exists $p \in P(\mathfrak{M})$ such that

$$\tau(p^\perp) < \varepsilon, \quad p\mathcal{H} \subset \mathcal{D}.$$

Proof. It is clear that (2) \Rightarrow (1) \Rightarrow (3) holds. We shall prove (3) \Rightarrow (2). By assumption, for all $n \in \mathbb{N}$, there exists $p_n \in P(\mathfrak{M})$ such that $\tau(p_n^\perp) < 1/2^n$ and $p_n\mathcal{H} \subset \mathcal{D}$. Put

$$q_n := \bigwedge_{k=n}^\infty p_k \in P(\mathfrak{M}).$$

Since $q_n \leq q_{n+1}$, the strong limit $s\text{-}\lim_{n \rightarrow \infty} q_n =: q \in P(\mathfrak{M})$ exists. It holds that

$$\begin{aligned} \tau(q^\perp) &= \lim_{n \rightarrow \infty} \tau(q_n^\perp) = \lim_{n \rightarrow \infty} \tau\left(\bigvee_{k=n}^\infty p_k^\perp\right) \\ &\leq \lim_{n \rightarrow \infty} \sum_{k=n}^\infty \tau(p_k^\perp) \leq \lim_{n \rightarrow \infty} \sum_{k=n}^\infty \frac{1}{2^k} = 0. \end{aligned}$$

Therefore we have $q = 1$. □

Lemma 2.8. *Let $\{(\mathfrak{M}_\lambda, \mathcal{H}_\lambda)\}_{\lambda \in \Lambda}$ be a family of countably decomposable, finite von Neumann algebras. Let*

$$\mathfrak{M} := \bigoplus_{\lambda \in \Lambda}^b \mathfrak{M}_\lambda, \quad \mathcal{H} := \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda.$$

For each $\lambda \in \Lambda$, let $\mathcal{D}_\lambda \subset \mathcal{H}_\lambda$ be a completely dense subspace for \mathfrak{M}_λ . Then $\widehat{\bigoplus_{\lambda \in \Lambda} \mathcal{D}_\lambda} \subset \mathcal{H}$ is a completely dense subspace for \mathfrak{M} .

Proof. By Lemma 2.7, for each $\lambda \in \Lambda$, there exists an increasing sequence $\{p_{\lambda,n}\}_{n=1}^\infty \subset P(\mathfrak{M}_\lambda)$ such that $p_{\lambda,n} \nearrow 1$ (strongly) and $\text{ran}(p_{\lambda,n}) \subset \mathcal{D}_\lambda$. For a finite set $F \subset \Lambda$, define

$$\begin{aligned} p_{F,n} &:= \bigoplus_{\lambda \in F} p_{\lambda,n}^{(\lambda)}, \\ p_{F,n}^{(\lambda)} &:= \begin{cases} p_{\lambda,n} & (\lambda \in F), \\ 0 & (\lambda \notin F). \end{cases} \end{aligned}$$

Then we have $p_{F,n} \in P(\mathfrak{M})$ and $\{p_{F,n}\}_{(F,n)}$ is an increasing net of projections. Here, we define $(F, n) \leq (F', n')$ by $F \subset F'$ and $n \leq n'$. It is clear that $p_{F,n} \nearrow 1$ (strongly) and $\text{ran}(p_{F,n}) \subset \widehat{\bigoplus_{\lambda \in \Lambda} \mathcal{D}_\lambda}$. Hence $\widehat{\bigoplus_{\lambda \in \Lambda} \mathcal{D}_\lambda}$ is completely dense. \square

Remark 2.9. Lemma 2.7 does not hold if \mathfrak{M} is not countably decomposable. We will show a counterexample. Let

$$\mathcal{H} := \bigoplus_{t \in \mathbb{R}} \ell^2(\mathbb{N}), \quad \mathfrak{M} := \bigoplus_{t \in \mathbb{R}}^b \mathfrak{M}_t, \quad \mathcal{D} := \widehat{\bigoplus_{t \in \mathbb{R}} \ell^2(\mathbb{N})}$$

Here, all \mathfrak{M}_t are isomorphic copies of some finite von Neumann algebra on $\ell^2(\mathbb{N})$. By Lemma 2.8, \mathcal{D} is completely dense for \mathfrak{M} . Suppose (2) of Lemma 2.7 holds. Then there exists $p_n \in P(\mathfrak{M})$ such that $\text{ran}(p_n) \subset \mathcal{D}$ and $p_n \nearrow 1$ (strongly). Represent p_n as $\bigoplus_t p_{t,n}$ ($p_{t,n} \in P(\mathfrak{M}_t)$). Then we have

$$\bigoplus_{t \in \mathbb{R}} \text{ran}(p_{t,n}) = \text{ran}(p_n) \subset \mathcal{D} = \widehat{\bigoplus_{t \in \mathbb{R}} \ell^2(\mathbb{N})}.$$

Therefore for each $n \in \mathbb{N}$, there exists a finite set $F_n \subset \mathbb{R}$ such that $p_{t,n} = 0$ for $t \notin F_n$. Since $F := \bigcup_{n=1}^{\infty} F_n \subset \mathbb{R}$ is at most countable, there exists some $t_0 \notin F$. Choose $\xi^{(t_0)} \in \ell^2(\mathbb{N})$ to be a unit vector and $\xi^{(t)} := 0$ ($t \neq t_0$). Then for $\xi = \{\xi^{(t)}\}_{t \in \mathbb{R}} \in \mathcal{H}$, it follows that

$$\begin{aligned} \|p_n \xi - \xi\|^2 &= \sum_{t \in \mathbb{R}} \|p_{t,n} \xi^{(t)} - \xi^{(t)}\|^2 = \underbrace{\|p_{t_0,n} \xi^{(t_0)} - \xi^{(t_0)}\|^2}_{=0} \\ &= \|\xi^{(t_0)}\|^2 = 1. \end{aligned}$$

On the other hand, we have $\|p_n \xi - \xi\|^2 \rightarrow 0$, which is a contradiction.

Proposition 2.10 (Murray-von Neumann [11]). *Let \mathfrak{M} be a finite von Neumann algebra on a Hilbert space \mathcal{H} . Let $\{\mathcal{D}_i\}_{i=1}^{\infty} \subset \mathcal{H}$ be a sequence of completely dense subspaces for \mathfrak{M} . Then the intersection $\bigcap_{i=1}^{\infty} \mathcal{D}_i$ is also completely dense.*

The proof requires some lemmata.

Lemma 2.11. *Proposition 2.10 holds if \mathfrak{M} is countably decomposable.*

Proof. Let τ be a faithful normal tracial state on \mathfrak{M} . By Lemma 2.7, for each $\varepsilon > 0$ and $i \in \mathbb{N}$, there exists $p_i \in P(\mathfrak{M})$ such that $\tau(p_i^\perp) < \varepsilon/2^i$ and $p_i \mathcal{H} \subset \mathcal{D}_i$. Put

$$p := \bigwedge_{i=1}^{\infty} p_i \in P(\mathfrak{M}).$$

Then we have

$$\begin{aligned}\tau(p^\perp) &= \tau\left(\bigvee_{i=1}^{\infty} p_i^\perp\right) \leq \sum_{i=1}^{\infty} \tau(p_i^\perp) \leq \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i} = \varepsilon, \\ p\mathcal{H} &= \bigcap_{i=1}^{\infty} (p_i\mathcal{H}) \subset \bigcap_{i=1}^{\infty} \mathcal{D}_i.\end{aligned}$$

Hence by Lemma 2.7, the intersection $\bigcap_{i=1}^{\infty} \mathcal{D}_i$ is completely dense. \square

Lemma 2.12. *Let $\{(\mathfrak{M}_\lambda, \mathcal{H}_\lambda)\}_{\lambda \in \Lambda}$ be a family of countably decomposable, finite von Neumann algebras. Put*

$$\mathfrak{M} := \bigoplus_{\lambda \in \Lambda}^b \mathfrak{M}_\lambda, \quad \mathcal{H} := \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda.$$

Let $\mathcal{D} \subset \mathcal{H}$ be a completely dense subspace for \mathfrak{M} . Then for each $\lambda \in \Lambda$, there exists some completely dense subspace $\mathcal{D}_\lambda \subset \mathcal{H}_\lambda$ for \mathfrak{M}_λ such that

$$\widehat{\bigoplus_{\lambda \in \Lambda} \mathcal{D}_\lambda} \subset \mathcal{D}.$$

Proof. By the definition, there exists an increasing net $\{p_\alpha\}_{\alpha \in A} \subset P(\mathfrak{M})$ such that $p_\alpha \nearrow 1$ (strongly) and $\text{ran}(p_\alpha) \subset \mathcal{D}$. Let $p_\alpha =: \bigoplus_{\lambda} p_{\lambda, \alpha}$ ($p_{\lambda, \alpha} \in P(\mathfrak{M}_\lambda)$). Then it holds that $p_{\lambda, \alpha} \nearrow 1$ (strongly). Put

$$\mathcal{D}_\lambda := \bigcup_{\alpha \in A} \text{ran}(p_{\lambda, \alpha}) \subset \mathcal{H}_\lambda.$$

We see that \mathcal{D}_λ is completely dense for \mathfrak{M}_λ . It is clear that $\widehat{\bigoplus_{\lambda \in \Lambda} \mathcal{D}_\lambda} \subset \mathcal{D}$ holds. \square

Proof of Proposition 2.10. Since \mathfrak{M} is finite, there exists a family of countably decomposable, finite von Neumann algebras $\{(\mathfrak{M}_\lambda, \mathcal{H}_\lambda)\}_{\lambda \in \Lambda}$ and a unitary operator $U : \mathcal{H} \rightarrow \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda$ such that $U\mathfrak{M}U^* = \bigoplus_{\lambda \in \Lambda} \mathfrak{M}_\lambda$. Put $\mathcal{D}'_i := U\mathcal{D}_i$. To prove the proposition, it suffices to prove that $\bigcap_{i=1}^{\infty} \mathcal{D}'_i$ is completely dense for $\bigoplus_{\lambda \in \Lambda} \mathfrak{M}_\lambda$. By Lemma 2.12, for each $i \in \mathbb{N}$, there exist completely dense subspaces $\mathcal{D}_{\lambda, i} \subset \mathcal{H}_\lambda$ for \mathfrak{M}_λ such that $\mathcal{D}'_i \supset \widehat{\bigoplus_{\lambda \in \Lambda} \mathcal{D}_{\lambda, i}}$. Then it follows that

$$\bigcap_{i=1}^{\infty} \mathcal{D}'_i \supset \bigcap_{i=1}^{\infty} \left(\widehat{\bigoplus_{\lambda \in \Lambda} \mathcal{D}_{\lambda, i}} \right) = \widehat{\bigoplus_{\lambda \in \Lambda} \left(\bigcap_{i=1}^{\infty} \mathcal{D}_{\lambda, i} \right)}.$$

By Lemma 2.11, $\bigcap_{i=1}^{\infty} \mathcal{D}_{\lambda, i}$ is completely dense for \mathfrak{M}_λ . Therefore by Lemma 2.8, $\widehat{\bigoplus_{\lambda \in \Lambda} \left(\bigcap_{i=1}^{\infty} \mathcal{D}_{\lambda, i} \right)}$ is completely dense for $\bigoplus_{\lambda \in \Lambda} \mathfrak{M}_\lambda$, which implies $\bigcap_{i=1}^{\infty} \mathcal{D}'_i$ is also completely dense for $\bigoplus_{\lambda \in \Lambda} \mathfrak{M}_\lambda$. \square

Proposition 2.13 (Murray-von Neumann [11]). *Let \mathfrak{M} be a finite von Neumann algebra. Then for each $X \in \overline{\mathfrak{M}}$ and a completely dense subspace \mathcal{D} for \mathfrak{M} , the subspace*

$$\{\xi \in \text{dom}(X) ; X\xi \in \mathcal{D}\}$$

is also completely dense. In particular, $\text{dom}(X)$ is completely dense for all $X \in \overline{\mathfrak{M}}$.

Proof. See [11]. □

Proposition 2.14 (Murray-von Neumann [11]). *Let \mathfrak{M} be a finite von Neumann algebra.*

- (1) *Every closed symmetric operator in $\overline{\mathfrak{M}}$ is self-adjoint.*
- (2) *There are no proper closed extensions of operators in $\overline{\mathfrak{M}}$. Namely, if $X, Y \in \overline{\mathfrak{M}}$ satisfy $X \subset Y$, then $X = Y$.*
- (3) *Let $\{X_i\}_i$ be a (finite or infinite) sequence in $\overline{\mathfrak{M}}$. The intersection of domains*

$$\mathcal{D}_{\mathcal{P}} := \bigcap_{p \in \mathcal{P}} \text{dom}(p(X_1, X_1^*, X_2, X_2^*, \dots))$$

of all unbounded operators obtained by substituting $\{X_i\}_i$ into the non-commutative polynomial $p(x_1, y_1, \dots)$ is completely dense for $\overline{\mathfrak{M}}$, where \mathcal{P} is the set of all non-commutative polynomials with indefinite elements $\{x_i, y_i\}_i$.

Proof. See [11]. □

Remark 2.15. Murray-von Neumann proved (1) of Proposition 2.14 using Cayley transform, but there is a simpler proof. We record it here.

Proof. Let $A \in \overline{\mathfrak{M}}$ be a symmetric operator. It is easy to see that $A + i$ is injective. Let $A + i = u|A + i|$ be its polar decomposition. From the injectivity, $u^*u = P_{\ker(A+i)^\perp} = 1_{\mathcal{H}}$. Since \mathfrak{M} is finite and $uu^* = P_{\overline{\text{ran}}(A+i)}$, we see that $\overline{\text{ran}}(A+i) = \mathcal{H}$. On the other hand, since A is closed and symmetric, $\text{ran}(A+i)$ is closed. Therefore we obtain $\text{ran}(A+i) = \mathcal{H}$. By the same way, it holds that $\text{ran}(A-i) = \mathcal{H}$, which means A is a self-adjoint operator. □

Similarly, we see that for $X \in \overline{\mathfrak{M}}$ the injectivity of X is equivalent to the density of $\text{ran}(X)$.

Lemma 2.16 (Murray-von Neumann [11]). *Let \mathfrak{M} be a finite von Neumann algebra and $\{X_i\}_i$ be a (finite or infinite) sequence in $\overline{\mathfrak{M}}$. Let*

$$p(x_1, y_1, x_2, y_2, \dots), \quad q(x_1, y_1, x_2, y_2, \dots), \quad r(x_1, y_1, x_2, y_2, \dots)$$

be non-commutative polynomials and $p(X_1, X_1^, X_2, X_2^*, \dots)$ be an operator obtained by substituting (x_i, y_i) by (X_i, X_i^*) .*

(1) $p(X_1, X_1^*, X_2, X_2^*, \dots)$ is a densely defined closable operator on \mathcal{H} , and

$$\overline{p(X_1, X_1^*, X_2, X_2^*, \dots)} \in \overline{\mathfrak{M}}.$$

(2) If $p^{(r)}(x_1, y_1, \dots) = q^{(r)}(x_1, y_1, \dots)$, then

$$\overline{p(X_1, X_1^*, \dots)} = \overline{q(X_1, X_1^*, \dots)}.$$

Namely, the closure of the substitution of operators depends on a reduced polynomial only.

(3) If $p(x_1, y_1, \dots)^+ = q(x_1, y_1, \dots)$, then

$$\left\{ \overline{p(X_1, X_1^*, \dots)} \right\}^* = \overline{q(X_1, X_1^*, \dots)}.$$

(4) If $\alpha p(x_1, y_1, \dots) = q(x_1, y_1, \dots)$ ($\alpha \in \mathbb{C}$), then

$$\overline{\alpha \cdot \left\{ \overline{p(X_1, X_1^*, \dots)} \right\}^*} = \overline{q(X_1, X_1^*, \dots)}.$$

(5) If $p(x_1, y_1, \dots) + q(x_1, y_1, \dots) = r(x_1, y_1, \dots)$, then

$$\overline{\overline{p(X_1, X_1^*, \dots)} + \overline{q(X_1, X_1^*, \dots)}} = \overline{r(X_1, X_1^*, \dots)}.$$

(6) If $p(x_1, y_1, \dots) \cdot q(x_1, y_1, \dots) = r(x_1, y_1, \dots)$, then

$$\overline{\overline{p(X_1, X_1^*, \dots)} \cdot \overline{q(X_1, X_1^*, \dots)}} = \overline{r(X_1, X_1^*, \dots)}.$$

Proof. See [11]. □

In summary, we have the following theorem.

Theorem 2.17 (Murray-von Neumann [11]). *For an arbitrary finite von Neumann algebra \mathfrak{M} , the set $\overline{\mathfrak{M}}$ forms a *-algebra of unbounded operators, where the algebraic operations are defined by¹*

$$\begin{aligned} (X, Y) &\mapsto \overline{X + Y}, & (\alpha, X) &\mapsto \overline{\alpha X}, \\ (X, Y) &\mapsto \overline{XY}, & X &\mapsto X^*. \end{aligned}$$

To conclude these preliminaries, we shall show a simple but useful lemma.

Lemma 2.18. *Let \mathfrak{M} be a finite von Neumann algebra, A be an operator in $\overline{\mathfrak{M}}$. If \mathcal{D} is a completely dense subspace of \mathcal{H} contained in $\text{dom}(A)$, then it is a core of A . That is, $\overline{A|_{\mathcal{D}}} = A$.*

¹ $\overline{\alpha X}$ equals αX when $\alpha \neq 0$. However, $\text{dom}(\overline{0 \cdot X}) = \mathcal{H} \neq \text{dom}(X)$.

Proof. From the complete density of \mathcal{D} , there exists an increasing net of closed subspaces $\{M_\alpha\}_\alpha$ of \mathcal{H} with $P_\alpha := P_{M_\alpha} \in \mathfrak{M}$ such that

$$\mathcal{D}_0 := \bigcup_{\alpha} M_\alpha \subset \mathcal{D}$$

is dense in \mathcal{H} . Define $A_0 := A|_{\mathcal{D}_0}$. Take an arbitrary $u \in U(\mathfrak{M}')$. Let $\xi \in \mathcal{D}_0 = \text{dom}(A_0)$, so that there is some α such that $\xi \in M_\alpha$. Then we have

$$\begin{aligned} uA_0\xi &= uA\xi = Au\xi = AuP_\alpha\xi \\ &= AP_\alpha u\xi = A_0P_\alpha u\xi \\ &= A_0u\xi. \end{aligned}$$

Therefore $uA_0 \subset A_0u$ holds. Since $u \in U(\mathfrak{M}')$ is arbitrary, we have $uA_0u^* = A_0$. Taking the closure of both sides, we see that $\overline{A_0} = u\overline{A_0}u^*$. This means $\overline{A_0} \in \overline{\mathfrak{M}}$. Therefore, it follows that

$$\overline{A_0} = \overline{A|_{\mathcal{D}}} \subset \overline{A} = A$$

Therefore by Proposition 2.14, we have $\overline{A_0} = A$. \square

2.3 Converse of Murray-von Neumann's Result

The converse of Theorem 2.17 is also true. We shall give a proof here.

Lemma 2.19. *Let \mathfrak{M} be a von Neumann algebra acting on a Hilbert space \mathcal{H} . Assume that, for all $A, B \in \mathfrak{M}$, the domains $\text{dom}(A+B)$ and $\text{dom}(AB)$ are dense in \mathcal{H} . Then $A+B$ and AB are densely defined closable operators on \mathcal{H} and the closures $\overline{A+B}$ and \overline{AB} are affiliated with \mathfrak{M} for all $A, B \in \mathfrak{M}$.*

Proof. By the assumption, $A+B$ is densely defined and

$$(A+B)^* \supset A^* + B^*.$$

Since the right hand side is densely defined, $A+B$ is closable. As same as above, we see that AB is closable. Affiliation property is easy. \square

Remark 2.20. Let \mathfrak{M} be a von Neumann algebra. It is easy to check that αA ($\alpha \in \mathbb{C}$, $A \in \overline{\mathfrak{M}}$) is always densely defined closable and its closure $\overline{\alpha A}$ is affiliated with \mathfrak{M} . Moreover $\overline{\mathfrak{M}}$ is closed with respect to the involution $A \mapsto A^*$.

Theorem 2.21. *Let \mathfrak{M} be a von Neumann algebra acting on a Hilbert space \mathcal{H} . Assume that, for all $A, B \in \mathfrak{M}$, the domains $\text{dom}(A+B)$ and $\text{dom}(AB)$ are dense in \mathcal{H} . If the set $\overline{\mathfrak{M}}$ forms a *-algebra with respect to the sum $\overline{A+B}$, the scalar multiplication $\overline{\alpha A}$ ($\alpha \in \mathbb{C}$), the multiplication \overline{AB} and the involution A^* , then \mathfrak{M} is a finite von Neumann algebra.*

Proof. Step 1. We first show that all closed symmetric operators affiliated with \mathfrak{M} are automatically self-adjoint. Let A be a closed symmetric operator affiliated with \mathfrak{M} . Define operators $B \in \overline{\mathfrak{M}}$ and $C \in \overline{\mathfrak{M}}$ as

$$B := \frac{1}{2} \overline{(A + A^*)}, \quad C := \frac{1}{2i} \overline{(A - A^*)},$$

then B and C are self-adjoint and $A = \overline{B + iC}$ holds because $\overline{\mathfrak{M}}$ is a $*$ -algebra. Since A is symmetric, we see that

$$C \supset \frac{1}{2i} (A - A^*) \supset \frac{1}{2i} (A - A) = 0|_{\text{dom}(A)}.$$

By taking the closure, we obtain $C = 0$. Hence $A = B$ is self-adjoint.

Step 2. We shall prove that \mathfrak{M} is finite. Let v be an arbitrary isometry in \mathfrak{M} . By the Wold decomposition, there exists a unique projection $p \in \mathfrak{M}$ such that $\text{ran}(p)$ reduces v , $s := v|_{\text{ran}(p)} \in \mathfrak{M}_p$ is a unilateral shift operator and $u := v|_{\text{ran}(p^\perp)} \in \mathfrak{M}_{p^\perp}$ is unitary. It is easy to see that

$$\ker(1 - s) = \{0\}, \quad \ker(1 - s^*) = \{0\},$$

so that we can define the closed symmetric operator T on $\text{ran}(p)$ as follows:

$$T := i(1 + s)(1 - s)^{-1}.$$

We immediately see that T is affiliated with the von Neumann algebra \mathfrak{M}_p . Define the operator A on $\mathcal{H} = \text{ran}(p) \oplus \text{ran}(p^\perp)$ by

$$A := T \oplus 0_{\text{ran}(p^\perp)},$$

then A is a closed symmetric operator and it is affiliated with \mathfrak{M} . From Step 1, A is self-adjoint, so that T is also self-adjoint. Since the Cayley transform of a self-adjoint operator is always unitary and the Cayley transform of T is s , s is unitary. This implies $p = 0$ because a unilateral shift operator admits no non-zero reducing closed subspace on which it is unitary. Hence $v = u$ is unitary. \square

3 Topological Structures of $\overline{\mathfrak{M}}$

In this section we investigate topological properties of $\overline{\mathfrak{M}}$. We need these results in the next section. We first endow $\overline{\mathfrak{M}}$ with two topologies, called the strong resolvent topology and the strong exponential topology. The former is (unbounded) operator theoretic and the latter is Lie theoretic. To show that these two topologies do coincide and $\overline{\mathfrak{M}}$ forms a complete topological $*$ -algebra with respect to them, we introduce another topology, called the τ -measure topology which originates from the noncommutative integration theory. They seem quite different to each other, but in fact they also coincide. The main topic of the present section is to study correlations between them.

3.1 Strong Resolvent Topology

First of all, we define the topology called the strong resolvent topology on the suitable subset of densely defined closed operators. Let \mathcal{H} be a Hilbert space. We call a densely defined closed operator A on \mathcal{H} belongs to the *resolvent class* $\mathcal{RC}(\mathcal{H})$ if A satisfies the following two conditions:

(RC.1) there exist self-adjoint operators X and Y on \mathcal{H} such that the intersection $\text{dom}(X) \cap \text{dom}(Y)$ is a core of X and Y ,

$$(RC.2) \quad A = \overline{X + iY}, \quad A^* = \overline{X - iY}.$$

Note that (RC.1) implies $\text{dom}(X) \cap \text{dom}(Y)$ is dense, so $X + iY$ and $X - iY$ are closable. Thus $\overline{X + iY}$ and $\overline{X - iY}$ are always defined. Furthermore, we have

$$\frac{1}{2}(A + A^*) = \frac{1}{2}(\overline{X + iY} + \overline{X - iY}) \supset X|_{\text{dom}(X) \cap \text{dom}(Y)}.$$

Since $A + A^*$ is closable and by (RC.1), we get

$$\frac{1}{2}\overline{A + A^*} \supset X.$$

As X is self-adjoint, X has no non-trivial symmetric extension, we have

$$\frac{1}{2}\overline{A + A^*} = X.$$

Therefore, X is uniquely determined. As same as above, Y is also unique and

$$\frac{1}{2i}\overline{A - A^*} = Y.$$

We denote

$$\text{Re}(A) := X = \frac{1}{2}\overline{A + A^*}, \quad \text{Im}(A) := Y = \frac{1}{2i}\overline{A - A^*}.$$

Also note that bounded operators and (possibly unbounded) normal operators belong to $\mathcal{RC}(\mathcal{H})$.

Now we endow $\mathcal{RC}(\mathcal{H})$ with the *strong resolvent topology* (SRT for short), the weakest topology for which the following mappings

$$\mathcal{RC}(\mathcal{H}) \ni A \mapsto \{\text{Re}(A) - i\}^{-1} \in (\mathfrak{B}(\mathcal{H}), SOT)$$

and

$$\mathcal{RC}(\mathcal{H}) \ni A \mapsto \{\text{Im}(A) - i\}^{-1} \in (\mathfrak{B}(\mathcal{H}), SOT)$$

are continuous. Thus a net $\{A_\alpha\}_\alpha$ in $\mathcal{RC}(\mathcal{H})$ converges to $A \in \mathcal{RC}(\mathcal{H})$ with respect to the strong resolvent topology if and only if

$$\{\text{Re}(A_\alpha) - i\}^{-1}\xi \rightarrow \{\text{Re}(A) - i\}^{-1}\xi, \quad \{\text{Im}(A_\alpha) - i\}^{-1}\xi \rightarrow \{\text{Im}(A) - i\}^{-1}\xi,$$

for each $\xi \in \mathcal{H}$. This topology is well-studied in the field of unbounded operator theory and suitable for the operator theoretical study. We denote the system of open sets of the strong resolvent topology by \mathcal{O}_{SRT} .

Let \mathfrak{M} be a finite von Neumann algebra on a Hilbert space \mathcal{H} . We shall show that $\overline{\mathfrak{M}}$ is a closed subset of the resolvent class $\mathcal{RC}(\mathcal{H})$. This fact follows from Proposition 2.10, Theorem 2.17, Lemma 2.18 and the following lemmata.

Lemma 3.1. *Let \mathfrak{M} be a finite von Neumann algebra on a Hilbert space \mathcal{H} , A be in $\overline{\mathfrak{M}}$. Then there exist unique self-adjoint operators B and C in $\overline{\mathfrak{M}}$ such that*

$$A = \overline{B + iC}.$$

Proof. Put

$$B := \frac{1}{2}\overline{A + A^*}, \quad C := \frac{1}{2i}\overline{A - A^*}.$$

Applying Proposition 2.10, $\text{dom}(B)$ and $\text{dom}(C)$ are dense in \mathcal{H} . Hence B and C are closed symmetric operators affiliated with \mathfrak{M} . By Proposition 2.14, in fact, B and C are self-adjoint. As $\overline{\mathfrak{M}}$ is a $*$ -algebra, we have

$$A = \overline{B + iC}.$$

□

Lemma 3.2. *Let \mathfrak{M} be a finite von Neumann algebra. Then $\overline{\mathfrak{M}}$ is closed with respect to the strong resolvent topology.*

Proof. Let $\{A_\alpha\}_\alpha \subset \overline{\mathfrak{M}}$ be a net converging to $A \in \mathcal{RC}(\mathcal{H})$ with respect to the strong resolvent topology. Then, for all $u \in U(\mathfrak{M}')$, we have

$$\begin{aligned} \{u\text{Re}(A)u^* - i\}^{-1} &= u\{\text{Re}(A) - i\}^{-1}u^* = \text{s-lim}_\alpha u\{\text{Re}(A_\alpha) - i\}^{-1}u^* \\ &= \text{s-lim}_\alpha \{u\text{Re}(A_\alpha)u^* - i\}^{-1} = \text{s-lim}_\alpha \{\text{Re}(A_\alpha) - i\}^{-1} \\ &= \{\text{Re}(A) - i\}^{-1}. \end{aligned}$$

This implies $\text{Re}(A)$ belongs to $\overline{\mathfrak{M}}$. As same as above, we obtain $\text{Im}(A) \in \overline{\mathfrak{M}}$. Thus so is $A = \text{Re}(A) + i\text{Im}(A)$. □

The next lemma is important in our discussion.

Lemma 3.3. *Let \mathfrak{M} be a finite von Neumann algebra acting on a Hilbert space \mathcal{H} . Then the following are equivalent:*

- (1) \mathfrak{M} is countably decomposable,
- (2) $(\overline{\mathfrak{M}}, \text{SRT})$ is metrizable as a topological space,
- (3) $(\overline{\mathfrak{M}}, \text{SRT})$ satisfies the first countability axiom.

Proof. (1) \Rightarrow (2). Let $\{\xi_k\}_k$ be a countable separating family of unit vectors in \mathcal{H} for \mathfrak{M} . For each $A, B \in \mathfrak{M}$, we define

$$\begin{aligned} d(A, B) &:= \sum_k \frac{1}{2^k} \|\{\text{Re}(A) - i\}^{-1}\xi_k - \{\text{Re}(B) - i\}^{-1}\xi_k\| \\ &\quad + \sum_k \frac{1}{2^k} \|\{\text{Im}(A) - i\}^{-1}\xi_k - \{\text{Im}(B) - i\}^{-1}\xi_k\|. \end{aligned}$$

It is easy to see that the above d is a distance function on the space $\overline{\mathfrak{M}}$, and the topology induced by the distance function d coincide with the strong resolvent topology on $\overline{\mathfrak{M}}$.

(2) \Rightarrow (3) is trivial.

(3) \Rightarrow (1). Let $S \subset P(\mathfrak{M})$ be a family of mutually orthogonal nonzero projections in \mathfrak{M} . Since $(\overline{\mathfrak{M}}, SRT)$ satisfies the first countability axiom, the origin $0 \in \overline{\mathfrak{M}}$ has a countable fundamental system of neighborhoods $\{V_k\}_k$. Put

$$S_k := \{p \in S ; p \notin V_k\},$$

then $S = \bigcup_k S_k$. This follows from the Hausdorff property of the strong resolvent topology. Next we show that each S_k is a finite set. Suppose S_k is an infinite set, then we can take a countably infinite subset $\{p_n ; n \in \mathbb{N}\}$ of S_k . Define

$$p := s\text{-}\lim_{N \rightarrow \infty} \sum_{n=1}^N p_n.$$

For every $\xi \in \mathcal{H}$ we see that

$$\begin{aligned} \|p_n \xi\| &= \left\| \sum_{i=1}^n p_n \xi - \sum_{i=1}^{n-1} p_n \xi \right\| \\ &\leq \left\| \sum_{i=1}^n p_n \xi - p \xi \right\| + \left\| p \xi - \sum_{i=1}^{n-1} p_n \xi \right\| \\ &\longrightarrow 0. \end{aligned}$$

Thus p_n converges strongly to 0. By Lemma B.1, this implies p_n converges to 0 with respect to the strong resolvent topology. Hence there exists a number $n \in \mathbb{N}$ such that $p_n \in V_k$. This is a contradiction to $p_n \in S_k$. Therefore S_k is a finite set. From the above arguments, we conclude that $S = \bigcup_k S_k$ is at most countable. \square

Remark 3.4. As we see in the sequel, $(\overline{\mathfrak{M}}, SRT)$ is a Hausdorff topological linear space. Thus in the case that \mathfrak{M} satisfies conditions (1), (2) or (3) of Lemma 3.3, $(\overline{\mathfrak{M}}, SRT)$ is metrizable with a translation invariant distance function. In particular, it is also metrizable as a uniform space.

Finally, we state one lemma.

Lemma 3.5. *Let \mathfrak{M} be a finite von Neumann algebra acting on a Hilbert space \mathcal{H} . Then the strong resolvent topology and the strong operator topology coincide on the closed unit ball \mathfrak{M}_1 .*

Proof. Note that if a von Neumann algebra is finite, then the involution is strongly continuous on the closed unit ball. The lemma follows immediately from this fact, Lemma B.1 and Lemma B.5. \square

See Appendix B for more informations of the strong resolvent topology.

3.2 Strong Exponential Topology

Next we introduce a Lie theoretic topology on $\overline{\mathfrak{M}}$. Let \mathcal{H} be a Hilbert space. For each $A \in \mathcal{R}\mathcal{C}(\mathcal{H})$, each SOT-neighborhood V at $1 \in \mathfrak{B}(\mathcal{H})$ and each compact set K of \mathbb{R} , we define $W(A; V, K)$ the subset of $\mathcal{R}\mathcal{C}(\mathcal{H})$ by

$$W(A; V, K) := \left\{ B \in \mathcal{R}\mathcal{C}(\mathcal{H}) ; \begin{array}{l} e^{-it\operatorname{Re}(A)} e^{it\operatorname{Re}(B)} \in V, \\ e^{-it\operatorname{Im}(A)} e^{it\operatorname{Im}(B)} \in V, \forall t \in K. \end{array} \right\},$$

then $\{W(A; V, K)\}_{A, V, K}$ is a fundamental system of neighborhoods on $\mathcal{R}\mathcal{C}(\mathcal{H})$. We denote the system of open sets of the topology induced by this fundamental system of neighborhoods by \mathcal{O}_{SET} , and call this topology the *strong exponential topology* (SET for short). Note that a net $\{A_\lambda\}_{\lambda \in \Lambda}$ in $\mathcal{R}\mathcal{C}(\mathcal{H})$ converges to $A \in \mathcal{R}\mathcal{C}(\mathcal{H})$ in the strong exponential topology if and only if

$$e^{it\operatorname{Re}(A_\lambda)} \xi \longrightarrow e^{it\operatorname{Re}(A)} \xi, \quad e^{it\operatorname{Im}(A_\lambda)} \xi \longrightarrow e^{it\operatorname{Im}(A)} \xi,$$

for each $\xi \in \mathcal{H}$, uniformly for t in any finite interval. This topology is important from the viewpoint of Lie theory. Indeed it can be defined by the unitary group $U(\mathcal{H})$ only. Before stating the main theorem in this section, we study relations between the strong resolvent topology and the strong exponential topology.

Lemma 3.6. *Let \mathfrak{M} be a countably decomposable finite von Neumann algebra acting on a Hilbert space \mathcal{H} . Then $(\overline{\mathfrak{M}}, \text{SET})$ is metrizable as a topological space.*

Proof. Let $\{\xi_n\}_n$ be a countable separating family of unit vectors in \mathcal{H} for \mathfrak{M} . For each $A, B \in \overline{\mathfrak{M}}$ we define

$$\begin{aligned} d(A, B) := & \sum_n \sum_{m=1}^{\infty} \frac{1}{2^{n+m}} \sup_{t \in [-m, m]} \|e^{it\operatorname{Re}(A)} \xi_n - e^{it\operatorname{Re}(B)} \xi_n\| \\ & + \sum_n \sum_{m=1}^{\infty} \frac{1}{2^{n+m}} \sup_{t \in [-m, m]} \|e^{it\operatorname{Im}(A)} \xi_n - e^{it\operatorname{Im}(B)} \xi_n\|. \end{aligned}$$

It is easy to see that the above d is a distance function on the space $\overline{\mathfrak{M}}$, and the topology induced by the distance function d coincide with the strong exponential topology on $\overline{\mathfrak{M}}$. \square

Lemma 3.7. *Let \mathfrak{M} be a countably decomposable finite von Neumann algebra. Then the strong resolvent topology and the strong exponential topology coincide on $\overline{\mathfrak{M}}$.*

Proof. This follows immediately from Lemma 3.3, Lemma 3.6 and Lemma B.2. \square

Remark 3.8. Similar to the above argument, one can prove that the strong resolvent topology and the strong exponential topology coincide on $\mathcal{R}\mathcal{C}(\mathcal{H})$ if the Hilbert space \mathcal{H} is separable. But the authors do not know whether this is true or not if \mathcal{H} is not separable. However we can show the following theorem.

The next is the main theorem in this section.

Theorem 3.9. *Let \mathfrak{M} be a finite von Neumann algebra acting on a Hilbert space \mathcal{H} . Then $\overline{\mathfrak{M}}$ is a complete topological $*$ -algebra with respect to the strong resolvent topology. Moreover the strong resolvent topology and the strong exponential topology coincide on $\overline{\mathfrak{M}}$.*

Throughout this section, we prove the above theorem.

3.3 τ -Measure Topology

We first prove Theorem 3.9 in a countably decomposable von Neumann algebra case. In this case, we can use the noncommutative integration theory thanks to a faithful normal tracial state. We shall introduce the τ -measure topology. Let \mathfrak{M} be a countably decomposable finite von Neumann algebra acting on a Hilbert space \mathcal{H} . Fix a faithful normal tracial state τ on \mathfrak{M} . The τ -measure topology (MT for short) on $\overline{\mathfrak{M}}$ is the linear topology whose fundamental system of neighborhoods at 0 is given by

$$N(\varepsilon, \delta) := \left\{ A \in \overline{\mathfrak{M}} ; \begin{array}{l} \text{there exists a projection } p \in \mathfrak{M} \\ \text{such that } \|Ap\| < \varepsilon, \tau(p^\perp) < \delta \end{array} \right\},$$

where ε and δ run over all strictly positive real numbers. It is known that $\overline{\mathfrak{M}}$ is a complete topological $*$ -algebra with respect to this topology [14]. We denote the system of open sets with respect to the τ -measure topology by \mathcal{O}_τ . Note that the τ -measure topology satisfies the first countability axiom.

Remark 3.10. In this context, the operators in $\overline{\mathfrak{M}}$ are sometimes called τ -measurable operators [4].

Thus there are two topologies on $\overline{\mathfrak{M}}$, the strong resolvent topology and the τ -measure topology. It seems that these two topologies are quite different. However, in fact, they coincide on $\overline{\mathfrak{M}}$, i.e.,

Lemma 3.11. *Let \mathfrak{M} be a countably decomposable finite von Neumann algebra acting on a Hilbert space \mathcal{H} . Then the strong resolvent topology and the τ -measure topology coincide on $\overline{\mathfrak{M}}$. In particular, $\overline{\mathfrak{M}}$ forms a complete topological $*$ -algebra, whose topology is independent of the choice of a faithful normal tracial state τ . Moreover the τ -measure topology is independent of the choice of a faithful normal tracial state τ .*

This lemma is the first step to our goal.

3.4 Almost Everywhere Convergence

To prove Lemma 3.11, we define almost everywhere convergence. Let \mathfrak{M} be a countably decomposable finite von Neumann algebra on a Hilbert space \mathcal{H} .

Definition 3.12. A sequence $\{A_n\}_{n=1}^\infty \subset \overline{\mathfrak{M}}$ converges almost everywhere (with respect to \mathfrak{M}) to $A \in \overline{\mathfrak{M}}$ if there exists a completely dense subspace \mathcal{D} such that

- (i) $\mathcal{D} \subset \bigcap_{n=1}^{\infty} \text{dom}(A_n) \cap \text{dom}(A)$,
- (ii) $A_n \xi$ converges to $A\xi$ for each $\xi \in \mathcal{D}$.

We shall investigate the relations between the almost everywhere convergence and the other topologies.

Lemma 3.13. *Let $\{A_n\}_{n=1}^{\infty} \subset \overline{\mathfrak{M}}$ be a sequence, $A \in \overline{\mathfrak{M}}$. Suppose A_n converges to A in the τ -measure topology, then there exists a subsequence $\{A_{n_k}\}_{k=1}^{\infty}$ of $\{A_n\}_{n=1}^{\infty}$ such that A_{n_k} converges almost everywhere to A .*

Proof. For all $j \in \mathbb{N}$, we can take $n_j \in \mathbb{N}$ and $p_j \in P(\mathfrak{M})$ which satisfy the following conditions:

$$\|(\overline{A_{n_j} - A})p_j\| < 1/2^j, \quad \tau(p_j^{\perp}) < 1/2^j, \quad n_j < n_{j+1}.$$

Put $p := \bigvee_{l=1}^{\infty} \bigwedge_{k=l}^{\infty} p_k \in P(\mathfrak{M})$, then $\text{ran}(p) = \overline{\bigcup_{l=1}^{\infty} \bigcap_{k=l}^{\infty} \text{ran}(p_k)}$. On the other hand,

$$\begin{aligned} \tau(p^{\perp}) &= \lim_{l \rightarrow \infty} \tau \left(\bigvee_{k=l}^{\infty} p_k^{\perp} \right) \leq \lim_{l \rightarrow \infty} \sum_{k=l}^{\infty} \tau(p_k^{\perp}) \\ &\leq \lim_{l \rightarrow \infty} \sum_{k=l}^{\infty} \frac{1}{2^k} = 0. \end{aligned}$$

Therefore, $\mathcal{H} = \text{ran}(p) = \overline{\bigcup_{l=1}^{\infty} \bigcap_{k=l}^{\infty} \text{ran}(p_k)}$. This implies

$$\mathcal{D}_0 := \bigcup_{l=1}^{\infty} \bigcap_{k=l}^{\infty} \text{ran}(p_k)$$

is completely dense. Let \mathcal{D}_1 be the intersection of the domains of all non-commutative polynomials of operators $\{A_{n_k}, A, p_k\}_{k=1}^{\infty}$, where we do not take closure for each non-commutative polynomial of operators. Then \mathcal{D}_1 is also completely dense and so is $\mathcal{D} := \mathcal{D}_0 \cap \mathcal{D}_1$. Take $\xi \in \mathcal{D}$, then there exists $k_0 \in \mathbb{N}$ such that $\xi \in \bigcap_{k=k_0}^{\infty} \text{ran}(p_k)$. Consequently, for all $k \geq k_0$, we get

$$\xi = p_k \xi, \quad p_k \xi \in \text{dom}(A) \cap \text{dom}(A_k), \quad \xi \in \text{dom}(A) \cap \text{dom}(A_k),$$

and

$$\begin{aligned} \|(\overline{A_{n_k} - A})\xi\| &= \|(\overline{A_{n_k} - A})p_j \xi\| \\ &\leq \|(\overline{A_{n_j} - A})p_j\| \cdot \|\xi\| \\ &\leq \frac{1}{2^k} \cdot \|\xi\| \longrightarrow 0. \end{aligned}$$

Thus A_{n_k} converges almost everywhere to A . □

Lemma 3.14. *Let $\{A_n\}_{n=1}^\infty$ be a sequence in $\overline{\mathfrak{M}}$ converging almost everywhere to $A \in \overline{\mathfrak{M}}$. Suppose $\{A_n^*\}_{n=1}^\infty$ also converges almost everywhere to A^* , then $\{A_n\}_{n=1}^\infty$ converges to A in the strong resolvent topology.*

Proof. It is easy to check that $\operatorname{Re}(A_n)$ and $\operatorname{Im}(A_n)$ converge almost every where to $\operatorname{Re}(A)$ and $\operatorname{Im}(A)$, respectively. Applying Lemma B.1 and Lemma 2.18 for $\operatorname{Re}(A_n)$ and $\operatorname{Im}(A_n)$, we see that $\operatorname{Re}(A_n)$ and $\operatorname{Im}(A_n)$ converge to $\operatorname{Re}(A)$ and $\operatorname{Im}(A)$ in the strong resolvent topology, respectively. This implies $\{A_n\}_{n=1}^\infty$ converges to A in the strong resolvent topology. \square

The following is well-known:

Lemma 3.15. *Let X be a metric space, $\{x_n\}_{n=1}^\infty \subset X$ be a sequence, $x \in X$. Suppose for each subsequence $\{x_{n_k}\}_{k=1}^\infty$ of $\{x_n\}_{n=1}^\infty$ has a subsequence $\{x_{n_{k_l}}\}_{l=1}^\infty$ of $\{x_{n_k}\}_{k=1}^\infty$ which converges to x , then x_n converges to x .*

3.5 Proof of Lemma 3.11

We shall start to prove Lemma 3.11. We prove that the system of open sets of the strong resolvent topology \mathcal{O}_{SRT} and the system of open sets of the τ -measure topology \mathcal{O}_τ coincide on $\overline{\mathfrak{M}}$. Let $\{A_n\}_{n=1}^\infty \subset \overline{\mathfrak{M}}$ be a sequence, $A \in \overline{\mathfrak{M}}$.

$\mathcal{O}_{\text{SRT}} \subset \mathcal{O}_\tau$: Suppose that $\{A_n\}_{n=1}^\infty$ converges to A in the τ -measure topology. Let $\{A_{n_k}\}_{k=1}^\infty$ be an arbitrary subsequence of $\{A_n\}_{n=1}^\infty$. By Proposition 3.13, there exists a subsequence $\{A_{n_{k_l}}\}_{l=1}^\infty$ of $\{A_{n_k}\}_{k=1}^\infty$ such that $\{A_{n_{k_l}}\}_{l=1}^\infty$ and $\{A_{n_{k_l}}^*\}_{l=1}^\infty$ converge almost everywhere to A and A^* , respectively. Applying Lemma 3.14, $A_{n_{k_l}}$ converges to A in the strong resolvent topology. This implies A_n converges to A in the strong resolvent topology, by Lemma 3.15. Thus we get $\mathcal{O}_{\text{SRT}} \subset \mathcal{O}_\tau$.

$\mathcal{O}_\tau \subset \mathcal{O}_{\text{SRT}}$: Suppose that $\{A_n\}_{n=1}^\infty$ converges to A with respect to the strong resolvent topology. First we consider the case that A_n and A are self-adjoint. Let $|A_n| =: \int_0^\infty \lambda dE_n(\lambda)$ and $|A| =: \int_0^\infty \lambda dE(\lambda)$ be spectral resolutions of $|A_n|$ and $|A|$, respectively. Fix an arbitrary positive number $\varepsilon > 0$. It is clear that $\text{s-lim}_{\lambda \rightarrow \infty} E([0, \lambda]) = 1$, so there exists a positive number $\Lambda > 0$ such that $\tau(E([0, \Lambda])^\perp) < \varepsilon$, where we can take $\Lambda > 0$ which is not a point spectrum of $|A|$. Indeed, self-adjoint operators have at most countable point spectra, as \mathfrak{M} is countably decomposable. Next we define a continuous function ϕ on \mathbb{R} as follows:

$$\phi(\lambda) := \begin{cases} 0 & \text{if } \lambda \leq -2\Lambda, \\ -\lambda - 2\Lambda & \text{if } -2\Lambda \leq \lambda \leq -\Lambda, \\ \lambda & \text{if } -\Lambda \leq \lambda \leq \Lambda, \\ -\lambda + 2\Lambda & \text{if } \Lambda \leq \lambda \leq 2\Lambda, \\ 0 & \text{if } 2\Lambda \leq \lambda. \end{cases}$$

Let $|\phi(A_n) - \phi(A)| =: \int_0^\infty \lambda dF_n(\lambda)$ be a spectral resolution of $|\phi(A_n) - \phi(A)|$, e be a spectral measure of A . Note that $E([0, \Lambda]) = e((-\Lambda, \Lambda))$. For each $\xi \in \mathcal{H}$,

$$\begin{aligned} \langle \xi, AE([0, \Lambda])\xi \rangle &= \int_{(-\Lambda, \Lambda)} \lambda d\langle \xi, e(\lambda)\xi \rangle \\ &= \int_{(-\Lambda, \Lambda)} \phi(\lambda) d\langle \xi, e(\lambda)\xi \rangle \\ &= \int_{\mathbb{R}} \phi(\lambda) d\langle \xi, e(\lambda)E([0, \Lambda])\xi \rangle \\ &= \langle \xi, \phi(A)E([0, \Lambda])\xi \rangle. \end{aligned}$$

Thus we have $AE([0, \Lambda])\xi = \phi(A)E([0, \Lambda])\xi$. Similar to the above argument, we get $A_n E_n([0, \Lambda])\xi = \phi(A_n)E_n([0, \Lambda])\xi$. Therefore, for all $\xi \in \mathcal{H}$, we see that

$$\begin{aligned} &\|(A_n - A)\{E_n([0, \Lambda]) \wedge E([0, \Lambda]) \wedge F_n([0, \varepsilon])\}\xi\|^2 \\ &= \|\{\phi(A_n) - \phi(A)\}\{E_n([0, \Lambda]) \wedge E([0, \Lambda]) \wedge F_n([0, \varepsilon])\}\xi\|^2 \\ &= \|\phi(A_n) - \phi(A)\|^2 \|\{E_n([0, \Lambda]) \wedge E([0, \Lambda]) \wedge F_n([0, \varepsilon])\}\xi\|^2 \\ &= \int_{[0, \varepsilon]} \lambda^2 d\|F_n(\lambda)\{E_n([0, \Lambda]) \wedge E([0, \Lambda]) \wedge F_n([0, \varepsilon])\}\xi\|^2 \\ &\leq \varepsilon^2 \|\xi\|^2. \end{aligned}$$

This implies

$$\| \overline{(A_n - A)\{E_n([0, \Lambda]) \wedge E([0, \Lambda]) \wedge F_n([0, \varepsilon])\}} \| \leq \varepsilon.$$

On the other hand,

$$\begin{aligned} &\tau(\{E_n([0, \Lambda]) \wedge E([0, \Lambda]) \wedge F_n([0, \varepsilon])\}^\perp) \\ &\leq \tau(E_n([0, \Lambda])^\perp) + \tau(E([0, \Lambda])^\perp) + \tau(F_n([0, \varepsilon])^\perp) \\ &\leq \tau(E_n([0, \Lambda])^\perp) + \varepsilon + \tau(F_n([0, \varepsilon])^\perp). \end{aligned}$$

By Lemma B.4, $|A_n|$ converges to $|A|$ in the strong resolvent topology, as the function

$$\mathbb{R} \ni \lambda \longmapsto (|\lambda| - i)^{-1} \in \mathbb{C}$$

is bounded continuous. By Lemma B.3,

$$\begin{aligned} E_n([0, \Lambda]) &= E_n((-1, \Lambda)) \\ &\xrightarrow{SOT} E((-1, \Lambda)) = E([0, \Lambda]). \end{aligned}$$

Thus for all sufficiently large number $n \in \mathbb{N}$,

$$\begin{aligned} \tau(E_n([0, \Lambda])^\perp) &= \tau(E([0, \Lambda])^\perp) + \tau(E([0, \Lambda]) - E_n([0, \Lambda])) \\ &\leq \varepsilon + \varepsilon = 2\varepsilon. \end{aligned}$$

Furthermore, by Lemma B.4, $\phi(A_n)$ converges strongly to $\phi(A)$. We obtain that for each $\xi \in \mathcal{H}$,

$$\|\phi(A_n) - \phi(A)\xi\| = \|\{\phi(A_n) - \phi(A)\}\xi\| \longrightarrow 0.$$

Applying Lemma B.1 and Lemma B.3 for $|\phi(A_n) - \phi(A)|$, we see that

$$F_n([0, \varepsilon]) = F_n((-1, \varepsilon)) \xrightarrow{SOT} 1.$$

Hence, for all sufficiently large numbers $n \in \mathbb{N}$, $\tau(F_n([0, \varepsilon])^\perp) < \varepsilon$. Thus, for all sufficiently large numbers $n \in \mathbb{N}$, we have

$$\tau(\{E_n([0, \Lambda]) \wedge E([0, \Lambda]) \wedge F_n([0, \varepsilon])\}^\perp) \leq 4\varepsilon.$$

From the above argument, we conclude that A_n converges to A in the τ -measure topology. In a general case, self-adjoint operators $\operatorname{Re}(A_n)$ and $\operatorname{Im}(A_n)$ converge to $\operatorname{Re}(A)$ and $\operatorname{Im}(A)$ in the strong resolvent topology, respectively. By the above argument, we see that $\operatorname{Re}(A_n)$ and $\operatorname{Im}(A_n)$ converge to $\operatorname{Re}(A)$ and $\operatorname{Im}(A)$ in the τ -measure topology, respectively. Since the addition is continuous with respect to the τ -measure topology, A_n converges to A in the τ -measure topology. This implies $\mathcal{O}_\tau \subset \mathcal{O}_{\text{SRT}}$. Hence the proof of Lemma 3.11 is complete.

Remark 3.16. We referred to the proof of Theorem 5.5 of the paper [20] to prove the inclusion $\mathcal{O}_\tau \subset \mathcal{O}_{\text{SRT}}$.

3.6 Direct Sums of Algebras of Unbounded Operators

To prove Theorem 3.9 in a general case, we show some facts about the direct sums of unbounded operators. See Appendix A for the definition of the direct sums of unbounded operators. The next lemma follows immediately from Lemma A.3.

Lemma 3.17. *Let \mathcal{H}_α be a Hilbert space, \mathcal{H} be the direct sum Hilbert space of $\{\mathcal{H}_\alpha\}_\alpha$. For each α , we consider a net $\{A_{\alpha,\lambda}\}_{\lambda \in \Lambda}$ of self-adjoint operators on \mathcal{H}_α and self-adjoint operator A_α on \mathcal{H}_α . Set*

$$A_\lambda := \bigoplus_\alpha A_{\alpha,\lambda},$$

and

$$A := \bigoplus_\alpha A_\alpha,$$

on the Hilbert space \mathcal{H} .

- (1) A_λ converges to A in the strong resolvent topology if and only if each $\{A_{\alpha,\lambda}\}_{\lambda \in \Lambda}$ converges to A_α in the strong resolvent topology.
- (2) A_λ converges to A in the strong exponential topology if and only if each $\{A_{\alpha,\lambda}\}_{\lambda \in \Lambda}$ converges to A_α in the strong exponential topology.

Proof. (1). By Lemma A.3, we have

$$(A_\lambda - i)^{-1} = \oplus_\alpha (A_{\alpha,\lambda} - i)^{-1}, \quad (A - i)^{-1} = \oplus_\alpha (A_\alpha - i)^{-1}.$$

The necessary condition is trivial. On the other hand, it is easy to see that $\{(A_\lambda - i)^{-1}\}_{\lambda \in \Lambda}$ converges to $(A - i)^{-1}$ on $\hat{\oplus}_\alpha \mathcal{H}_\alpha$. Since $\widehat{\oplus}_\alpha \mathcal{H}_\alpha$ is dense in $\oplus_\alpha \mathcal{H}_\alpha$ and $\{(A_\lambda - i)^{-1}\}_{\lambda \in \Lambda}$ is uniformly bounded, the sufficient condition follows.

(2). Similar to the proof (1). \square

The next lemma is the key to prove Theorem 3.9.

Lemma 3.18. *Let \mathfrak{M}_α be a finite von Neumann algebra acting on \mathcal{H}_α , and put*

$$\mathfrak{M} := \bigoplus_\alpha^b \mathfrak{M}_\alpha.$$

Then

$$\overline{\mathfrak{M}} = \bigoplus_\alpha \overline{\mathfrak{M}_\alpha}$$

holds. The sum, the scalar multiplication, the multiplication and the involution are given by

$$\begin{aligned} \overline{(\oplus_\alpha A_\alpha) + (\oplus_\alpha B_\alpha)} &= \oplus_\alpha \overline{(A_\alpha + B_\alpha)}, \\ \overline{\lambda (\oplus_\alpha A_\alpha)} &= \oplus_\alpha \overline{(\lambda A_\alpha)}, \quad \text{for all } \lambda \in \mathbb{C}, \\ \overline{(\oplus_\alpha A_\alpha) (\oplus_\alpha B_\alpha)} &= \oplus_\alpha \overline{(A_\alpha B_\alpha)}, \\ (\oplus_\alpha A_\alpha)^* &= \oplus_\alpha (A_\alpha^*). \end{aligned}$$

In addition, if each $\overline{\mathfrak{M}_\alpha}$ is countably decomposable, then $\overline{\mathfrak{M}}$ is a complete topological *-algebra with respect to the strong resolvent topology, and the strong resolvent topology coincides with the strong exponential topology on $\overline{\mathfrak{M}}$.

Proof. we shall prove this lemma step by step.

Step 1. We first show that $\bigoplus_\alpha \overline{\mathfrak{M}_\alpha} \subset \overline{\mathfrak{M}}$. Indeed let $\oplus_\alpha A_\alpha \in \bigoplus_\alpha \overline{\mathfrak{M}_\alpha}$. By Lemma A.1 and Lemma A.4, each unitary operator $u \in U(\mathfrak{M}')$ can be written as $u = \oplus_\alpha u_\alpha$, where $u_\alpha \in U(\mathfrak{M}'_\alpha)$. Thus we have

$$u (\oplus_\alpha A_\alpha) = \oplus_\alpha (u_\alpha A_\alpha) \subset \oplus_\alpha (A_\alpha u_\alpha) = (\oplus_\alpha A_\alpha) u.$$

This implies $\oplus_\alpha A_\alpha \in \overline{\mathfrak{M}}$.

Step 2. We show that the converse inclusion $\overline{\mathfrak{M}} \subset \bigoplus_\alpha \overline{\mathfrak{M}_\alpha}$. For each β , we put

$$q_\beta := \oplus_\alpha (\delta_{\alpha\beta} 1_{\mathcal{H}_\alpha}) \in \mathfrak{M}',$$

where $\delta_{\alpha\beta}$ is the Kronecker delta and $1_{\mathcal{H}_\alpha}$ is the identity operator on \mathcal{H}_α . From Lemma A.1, q_β is a projection and

$$\text{ran}(q_\beta) = \bigoplus_{\alpha} (\delta_{\alpha\beta} \mathcal{H}_\alpha) =: \tilde{\mathcal{H}}_\beta.$$

Let $A \in \overline{\mathfrak{M}}$ be a self-adjoint operator. We would like to prove $A \in \bigoplus_{\alpha} \overline{\mathfrak{M}}_\alpha$. Since $q_\beta \in \mathfrak{M}'$, we have $q_\beta A \subset A q_\beta$ for all β . This implies that each $\tilde{\mathcal{H}}_\beta$ reduces A . We denote reduced part of A to $\tilde{\mathcal{H}}_\beta$ by $A_{\tilde{\mathcal{H}}_\beta}$. $A_{\tilde{\mathcal{H}}_\beta}$ is obviously self-adjoint. For each β , we consider natural unitary operator $v_\beta : \mathcal{H}_\beta \rightarrow \tilde{\mathcal{H}}_\beta$. Then the operator $A_\beta := v_\beta^* A_{\tilde{\mathcal{H}}_\beta} v_\beta$ is again self-adjoint. To prove $A = \bigoplus_{\alpha} A_\alpha$, we take an arbitrary $\xi \in \widehat{\bigoplus}_{\alpha} \text{dom}(A_\alpha)$. Since $v_\alpha \xi^{(\alpha)} \in \text{dom}(A_{\tilde{\mathcal{H}}_\alpha}) \subset \text{dom}(A)$, we see that

$$\xi = \sum_{\substack{\alpha \\ \text{finite sum}}} v_\alpha \xi^{(\alpha)} \in \text{dom}(A).$$

Therefore we obtain

$$\begin{aligned} (\bigoplus_{\alpha} A_\alpha) \xi &= \left\{ v_\alpha^* A v_\alpha \xi^{(\alpha)} \right\}_{\alpha} = \sum_{\substack{\alpha \\ \text{finite sum}}} A v_\alpha \xi^{(\alpha)} \\ &= A \left(\sum_{\substack{\alpha \\ \text{finite sum}}} v_\alpha \xi^{(\alpha)} \right) = A \xi. \end{aligned}$$

Hence $(\bigoplus_{\alpha} A_\alpha) |_{\widehat{\bigoplus}_{\alpha} \text{dom}(A_\alpha)} \subset A$. By Lemma A.2, we have $\bigoplus_{\alpha} A_\alpha \subset A$. On the other hand, both of $\bigoplus_{\alpha} A_\alpha$ and A are self-adjoint and self-adjoint operators have no non-trivial self-adjoint extension. These facts implies $\bigoplus_{\alpha} A_\alpha = A$. Next we show that each A_α is in $\overline{\mathfrak{M}}_\alpha$. Taking arbitrary unitary operators $u_\alpha \in U(\mathfrak{M}'_\alpha)$ and putting $u := \bigoplus_{\alpha} u_\alpha$, then by Lemma A.4, u is a unitary operator in \mathfrak{M}' . Since $A \in \overline{\mathfrak{M}}$, we see that

$$(\bigoplus_{\alpha} u_\alpha) (\bigoplus_{\alpha} A_\alpha) = u A \subset A u = (\bigoplus_{\alpha} A_\alpha) (\bigoplus_{\alpha} u_\alpha).$$

Thus for all α , $u_\alpha A_\alpha \subset A_\alpha u_\alpha$ holds. This implies $A_\alpha \in \overline{\mathfrak{M}}_\alpha$ for all α . Hence $A \in \bigoplus_{\alpha} \overline{\mathfrak{M}}_\alpha$.

Next we consider an arbitrary element $A \in \overline{\mathfrak{M}}$. Putting $B := \text{Re}(A)$, $C := \text{Im}(A)$, then $A = \overline{B + iC}$. Since B and C are self-adjoint, by the above argument, there exist operators $B_\alpha \in \overline{\mathfrak{M}}_\alpha$ and $C_\alpha \in \overline{\mathfrak{M}}_\alpha$ such that $B = \bigoplus_{\alpha} B_\alpha$ and $C = \bigoplus_{\alpha} C_\alpha$ holds. Set $\mathcal{D} := \widehat{\bigoplus}_{\alpha} (\text{dom}(B_\alpha) \cap \text{dom}(C_\alpha))$. Since $\text{dom}(B_\alpha) \cap \text{dom}(C_\alpha)$ is a core of $\overline{B_\alpha + iC_\alpha}$, \mathcal{D} is a core of $\bigoplus_{\alpha} (\overline{B_\alpha + iC_\alpha})$ by Lemma A.2. We observe that

$$A = \overline{B + iC} \supset (\overline{B + iC}) |_{\mathcal{D}} = \left\{ \bigoplus_{\alpha} (\overline{B_\alpha + iC_\alpha}) \right\} |_{\mathcal{D}},$$

so that $A \supset \bigoplus_{\alpha} (\overline{B_\alpha + iC_\alpha})$ follows. Now we use Step 1., then we see that $\bigoplus_{\alpha} (\overline{B_\alpha + iC_\alpha}) \in \overline{\mathfrak{M}}$ because $\overline{B_\alpha + iC_\alpha} \in \overline{\mathfrak{M}}_\alpha$ for all α . Since \mathfrak{M} is a finite

von Neumann algebra, Lemma 2.14 means that $A = \oplus_{\alpha} (\overline{B_{\alpha} + iC_{\alpha}}) \in \oplus_{\alpha} \overline{\mathfrak{M}}_{\alpha}$. Hence $\overline{\mathfrak{M}} = \oplus_{\alpha} \overline{\mathfrak{M}}_{\alpha}$ follows.

Step 3. We shall show the formulae with respect to the sum, the scalar product, the product and the involution. The formulae with respect to the scalar product and the involution are trivial. We first prove the formula of the sum. Let $\oplus_{\alpha} A_{\alpha}, \oplus_{\alpha} B_{\alpha} \in \oplus_{\alpha} \overline{\mathfrak{M}}_{\alpha} = \overline{\mathfrak{M}}$. Put $\mathcal{D}_+ := \widehat{\bigoplus}_{\alpha} (\text{dom}(A_{\alpha}) \cap \text{dom}(B_{\alpha}))$. Since $\text{dom}(A_{\alpha}) \cap \text{dom}(B_{\alpha})$ is a core of $\overline{A_{\alpha} + B_{\alpha}}$, \mathcal{D}_+ is a core of $\oplus_{\alpha} (\overline{A_{\alpha} + B_{\alpha}})$. We observe that

$$\overline{(\oplus_{\alpha} A_{\alpha}) + (\oplus_{\alpha} B_{\alpha})} \supset \{ \oplus_{\alpha} (\overline{A_{\alpha} + B_{\alpha}}) \} |_{\mathcal{D}_+},$$

so that

$$\overline{(\oplus_{\alpha} A_{\alpha}) + (\oplus_{\alpha} B_{\alpha})} \supset \oplus_{\alpha} (\overline{A_{\alpha} + B_{\alpha}})$$

follows. Since both sides are elements in $\overline{\mathfrak{M}}$, we have

$$\overline{(\oplus_{\alpha} A_{\alpha}) + (\oplus_{\alpha} B_{\alpha})} = \oplus_{\alpha} (\overline{A_{\alpha} + B_{\alpha}}).$$

Next, to show the formula of the product, we put $\mathcal{D}_\times := \widehat{\bigoplus}_{\alpha} \text{dom}(A_{\alpha} B_{\alpha})$. Since $\text{dom}(A_{\alpha} B_{\alpha})$ is a core of $\overline{A_{\alpha} B_{\alpha}}$, \mathcal{D}_\times is a core of $\oplus_{\alpha} (\overline{A_{\alpha} B_{\alpha}})$. We observe that

$$\overline{(\oplus_{\alpha} A_{\alpha}) (\oplus_{\alpha} B_{\alpha})} \supset \{ \oplus_{\alpha} (\overline{A_{\alpha} B_{\alpha}}) \} |_{\mathcal{D}_\times},$$

so that

$$\overline{(\oplus_{\alpha} A_{\alpha}) (\oplus_{\alpha} B_{\alpha})} \supset \oplus_{\alpha} (\overline{A_{\alpha} B_{\alpha}})$$

follows. Since both sides are elements in $\overline{\mathfrak{M}}$, we have

$$\overline{(\oplus_{\alpha} A_{\alpha}) (\oplus_{\alpha} B_{\alpha})} = \oplus_{\alpha} (\overline{A_{\alpha} B_{\alpha}}).$$

Hence the proof of Step 3. is complete.

In the sequel, we assume that each $\overline{\mathfrak{M}}_{\alpha}$ is countably decomposable. Note that, by Lemma 3.11, each $\overline{\mathfrak{M}}_{\alpha}$ is a complete topological *-algebra with respect to the strong resolvent topology.

Step 4. Let $\{A_{\lambda}\}_{\lambda \in \Lambda}$ be a net in $\overline{\mathfrak{M}}$, A be an element of $\overline{\mathfrak{M}}$. Corresponding to $\overline{\mathfrak{M}} = \oplus_{\alpha} \overline{\mathfrak{M}}_{\alpha}$, we can write them as follows:

$$\begin{aligned} A_{\lambda} &= \oplus_{\alpha} A_{\alpha, \lambda}, & A_{\alpha, \lambda} &\in \overline{\mathfrak{M}}, \\ A &= \oplus_{\alpha} A_{\alpha}, & A_{\alpha} &\in \overline{\mathfrak{M}}. \end{aligned}$$

We shall show that A_{λ} converges to A with respect to the strong resolvent topology if and only if each $\{A_{\alpha, \lambda}\}_{\lambda \in \Lambda}$ converges to A_{α} with respect to the strong resolvent topology. From Step 3., we obtain

$$\begin{aligned} \text{Re}(A_{\lambda}) &= \oplus_{\alpha} \text{Re}(A_{\alpha, \lambda}), & \text{Im}(A_{\lambda}) &= \oplus_{\alpha} \text{Im}(A_{\alpha, \lambda}), \\ \text{Re}(A) &= \oplus_{\alpha} \text{Re}(A_{\alpha}), & \text{Im}(A) &= \oplus_{\alpha} \text{Im}(A_{\alpha}), \end{aligned}$$

so that, by Lemma 3.17, the above equivalence of convergence follows.

By Step 3. and Step 4., we see that $\overline{\mathfrak{M}}$ forms a topological *-algebra with respect to the strong resolvent topology. Next, to prove the completeness, we prepare some facts.

Step 5. Fix an arbitrary α_0 and let $V^{(\alpha_0)}$ be an arbitrary SOT-open set in \mathfrak{M}_{α_0} . Set $V^{(\alpha)} := \mathfrak{M}_{\alpha}$ ($\alpha \neq \alpha_0$) and $V := \bigoplus_{\alpha}^b V^{(\alpha)}$. Then V is a SOT-open set in \mathfrak{M} . Indeed, since for any $x = \bigoplus_{\alpha} x_{\alpha} \in V$, we have $x_{\alpha_0} \in V^{(\alpha_0)}$, there exists a positive number $\varepsilon > 0$ and finitely many vectors $\xi_1^{(\alpha_0)} \in \mathcal{H}_{\alpha_0}, \dots, \xi_n^{(\alpha_0)} \in \mathcal{H}_{\alpha_0}$ such that

$$\bigcap_{k=1}^n \left\{ y \in \mathfrak{M}_{\alpha_0} ; \|(y - x_{\alpha_0})\xi_k^{(\alpha_0)}\| < \varepsilon \right\} \subset V^{(\alpha_0)}.$$

Set $\xi_k^{(\alpha)} := 0$ ($\alpha \neq \alpha_0$), then we get $\xi_k \in \bigoplus_{\alpha} \mathcal{H}_{\alpha}$ and

$$x \in \bigcap_{k=1}^n \left\{ y \in \mathfrak{M} ; \|(y - x)\xi_k\| < \varepsilon \right\} \subset V.$$

Since $\bigcap_{k=1}^n \{y \in \mathfrak{M} ; \|(y - x)\xi_k\| < \varepsilon\}$ is a SOT-open set in \mathfrak{M} , V is a SOT-open set.

Step 6. Fix an arbitrary α_0 . Let $W^{(\alpha_0)}$ be an arbitrary SRT-neighborhood at $0 \in \overline{\mathfrak{M}}_{\alpha_0}$. Set $W^{(\alpha)} := \overline{\mathfrak{M}}_{\alpha}$ ($\alpha \neq \alpha_0$) and $W := \bigoplus_{\alpha} W^{(\alpha)}$. Then W is a neighborhood at $0 \in \overline{\mathfrak{M}}$. Indeed, $0 \in W$ is trivial. On the other hand, since $0 \in \overline{\mathfrak{M}}_{\alpha_0}$, there exists finitely many SOT-open sets $V_1^{(\alpha_0)}, \dots, V_n^{(\alpha_0)}$ in \mathfrak{M}_{α_0} such that

$$0 \in \bigcap_{k=1}^n \left\{ A \in \overline{\mathfrak{M}}_{\alpha_0} ; \{\operatorname{Re}(A) - i\}^{-1}, \{\operatorname{Im}(A) - i\}^{-1} \in V_k^{(\alpha_0)} \right\} \subset W^{(\alpha_0)}.$$

Put $V_k^{(\alpha)} := \mathfrak{M}_{\alpha}$ ($\alpha \neq \alpha_0$) and $V_k := \bigoplus_{\alpha}^b V_k^{(\alpha)}$, then, by Step 5., each V_k is a SOT-open set in \mathfrak{M} and

$$0 \in \bigcap_{k=1}^n \left\{ A \in \overline{\mathfrak{M}} ; \{\operatorname{Re}(A) - i\}^{-1}, \{\operatorname{Im}(A) - i\}^{-1} \in V_k \right\} \subset W.$$

By the definition of the strong resolvent topology,

$$\bigcap_{k=1}^n \left\{ A \in \overline{\mathfrak{M}} ; \{\operatorname{Re}(A) - i\}^{-1}, \{\operatorname{Im}(A) - i\}^{-1} \in V_k \right\}$$

is a open set, so that W is a SRT-neighborhood at $0 \in \overline{\mathfrak{M}}$.

Step 7. We shall give a proof of the completeness of $\overline{\mathfrak{M}}$. Let $\{A_{\lambda}\}_{\lambda \in \Lambda}$ be a Cauchy net in $\overline{\mathfrak{M}}$. For each $\lambda \in \Lambda$ we can write as $A_{\lambda} = \bigoplus_{\alpha} A_{\alpha, \lambda} \in \bigoplus_{\alpha} \overline{\mathfrak{M}}_{\alpha}$. Fix

an arbitrary α_0 and let $W^{(\alpha_0)}$ be an arbitrary SRT-neighborhood at $0 \in \overline{\mathfrak{M}}_{\alpha_0}$. Set $W^{(\alpha)} := \overline{\mathfrak{M}}_{\alpha}$ ($\alpha \neq \alpha_0$) and $W := \bigoplus_{\alpha} W^{(\alpha)}$, then, by Step 6., W is a SRT-neighborhood at $0 \in \overline{\mathfrak{M}}$. Therefore there exists $\lambda_0 \in \Lambda$ such that $\overline{A_{\lambda} - A_{\mu}} \in W$ for all $\lambda, \mu \geq \lambda_0$. Since $\overline{A_{\lambda} - A_{\mu}} = \bigoplus_{\alpha} (\overline{A_{\alpha, \lambda} - A_{\alpha, \mu}})$, this implies that $\overline{A_{\alpha_0, \lambda} - A_{\alpha_0, \mu}} \in W^{(\alpha_0)}$ for all $\lambda, \mu \geq \lambda_0$. Hence $\{A_{\alpha_0, \lambda}\}_{\lambda \in \Lambda}$ is a Cauchy net in $\overline{\mathfrak{M}}_{\alpha_0}$. We now use the completeness of $\overline{\mathfrak{M}}_{\alpha_0}$, then there exists an element $A_{\alpha_0} \in \overline{\mathfrak{M}}_{\alpha_0}$ such that $A_{\alpha_0, \lambda} \rightarrow A_{\alpha_0}$. Since α_0 is arbitrary, so that this means that for each α , there exists an element $A_{\alpha} \in \overline{\mathfrak{M}}_{\alpha}$ such that $A_{\alpha, \lambda} \rightarrow A_{\alpha}$. Put $A := \bigoplus_{\alpha} A_{\alpha}$, then, by Step 4., we conclude that $A_{\lambda} \rightarrow A$. Thus $\overline{\mathfrak{M}}$ is complete.

Step 8. The strong resolvent topology coincides with the strong exponential topology on $\overline{\mathfrak{M}}$. This fact follows from Lemma 3.7 and Lemma 3.17. \square

Lemma 3.19. *Let $(\mathfrak{M}, \mathcal{H})$ and $(\mathfrak{N}, \mathcal{K})$ be spatially isomorphic finite von Neumann algebras. If a unitary operator U of \mathcal{H} onto \mathcal{K} induces the spatial isomorphism, then the map*

$$\Phi : \overline{\mathfrak{M}} \rightarrow \overline{\mathfrak{N}}, X \mapsto UXU^*$$

*is a *-isomorphism. Moreover Φ is a homeomorphism with respect to the strong resolvent topology and the strong exponential topology.*

Proof. It is easy to see that $\Phi(X) \in \overline{\mathfrak{N}}$ for all $X \in \overline{\mathfrak{M}}$. Thanks to Proposition 2.14 (2), it is not difficult to show that Φ is a unital *-homomorphism:

$$\begin{aligned} U(\overline{X+Y})U^* &= \overline{UXU^* + UYU^*} \\ U(\overline{XY})U^* &= \overline{UXU^*UYU^*} \\ UX^*U^* &= (UXU^*)^*. \end{aligned}$$

Furthermore, it is straightforward to verify that Φ is invertible, the inverse of which is given by $\overline{\mathfrak{N}} \ni Y \mapsto U^*YU \in \overline{\mathfrak{M}}$. Topological property is trivial. \square

3.7 Proof of Theorem 3.9

We shall give a proof of Theorem 3.9. By Lemma 2.3, there exists a family of countably decomposable finite von Neumann algebras $\{\mathfrak{M}_{\alpha}\}_{\alpha}$ such that \mathfrak{M} is spatially isomorphic onto $\bigoplus_{\alpha}^b \mathfrak{M}_{\alpha}$. From Lemma 3.19, there exists a *-isomorphism of $\overline{\mathfrak{M}}$ onto $\bigoplus_{\alpha} \overline{\mathfrak{M}}_{\alpha}$ which is homeomorphic with respect to the strong resolvent topology and the strong exponential topology. By Lemma 3.18, $\bigoplus_{\alpha} \overline{\mathfrak{M}}_{\alpha}$ is complete topological *-algebra, so that so is $\overline{\mathfrak{M}}$. Hence the proof of Theorem 3.9 is complete.

3.8 Local Convexity

We study the local convexity of $(\overline{\mathfrak{M}}, SRT)$ here.

Proposition 3.20. *Let \mathfrak{M} be a finite von Neumann algebra. Then the following are equivalent.*

(1) $(\overline{\mathfrak{M}}, SRT)$ is locally convex.

(2) \mathfrak{M} is atomic.

We need some lemmata to prove the above proposition.

Lemma 3.21. *Let \mathfrak{M} be an atomic finite von Neumann algebra, then $(\overline{\mathfrak{M}}, SRT)$ is locally convex.*

Proof. Every atomic finite von Neumann algebra is spatially isomorphic to the direct sum of matrix algebras $\{M_{n_\lambda}(\mathbb{C})\}_{\lambda \in \Lambda}$, where each $M_{n_\lambda}(\mathbb{C})$ is the algebra of all $n_\lambda \times n_\lambda$ complex matrices. Thus we should only prove this lemma in the case that \mathfrak{M} is equal to $\bigoplus_{\lambda \in \Lambda}^b M_{n_\lambda}(\mathbb{C})$. Note that

$$\overline{\mathfrak{M}} = \overline{\bigoplus_{\lambda \in \Lambda}^b M_{n_\lambda}(\mathbb{C})} = \bigoplus_{\lambda \in \Lambda} \overline{M_{n_\lambda}(\mathbb{C})} = \bigoplus_{\lambda \in \Lambda} M_{n_\lambda}(\mathbb{C}).$$

Let p_λ be a semi-norm on $\overline{\mathfrak{M}}$ defined by

$$p_\lambda(x) := \|x_\lambda\|, \quad x = \bigoplus_{\lambda \in \Lambda} x_\lambda \in \overline{\mathfrak{M}} = \bigoplus_{\lambda \in \Lambda} M_{n_\lambda}(\mathbb{C}).$$

Then the strong resolvent topology on $\overline{\mathfrak{M}}$ coincides with the locally convex topology induced by the semi-norms $\{p_\lambda\}_{\lambda \in \Lambda}$ because there is only one Hausdorff linear topology on a finite dimensional linear space. Hence the proof is complete. \square

Lemma 3.22. *Let \mathfrak{M} be a diffuse finite von Neumann algebra, then there exists no non-zero SRT-continuous linear functional on $\overline{\mathfrak{M}}$.*

Proof. Suppose there exists a non-zero SRT-continuous linear functional f on $\overline{\mathfrak{M}}$ and we shall show a contradiction. Since SOT and SRT coincides on \mathfrak{M}_1 , the restriction of f onto \mathfrak{M} is σ -strongly continuous. This fact and the SRT-density of \mathfrak{M} in $\overline{\mathfrak{M}}$ implies that there exists a projection e_0 in \mathfrak{M} such that $f(e_0) \neq 0$.

Step 1. For any orthogonal family of non-zero projections $\{e_n\}_{n=1}^\infty$ of \mathfrak{M} , $f(e_n) = 0$ except at most finitely many $n \in \mathbb{N}$. Indeed, put

$$A := \sum_{n=1}^\infty a_n e_n \in \overline{\mathfrak{M}}, \quad a_n := \begin{cases} \frac{1}{f(e_n)} & \text{if } f(e_n) \neq 0, \\ 0 & \text{if } f(e_n) = 0, \end{cases}$$

where convergence of A is in the strong resolvent topology. Then we have

$$f(A) = \sum_{n=1}^\infty a_n f(e_n) = \sum_{f(e_n) \neq 0} 1 < \infty,$$

so that $f(e_n) = 0$ except at most finitely many $n \in \mathbb{N}$.

Step 2. For any $e \in P(\mathfrak{M})$ with $f(e) \neq 0$, there exists $e' \in P(\mathfrak{M})$ such that $0 \neq e' \leq e$ and $f(e') = 0$. Indeed, since \mathfrak{M} is diffuse, there exists an orthogonal family of non-zero projections $\{e_n\}_{n=1}^\infty$ in \mathfrak{M} such that $e = \sum_{n \geq 1} e_n$. By Step 2., $J := \{n \in \mathbb{N} ; f(e_n) \neq 0\}$ is a finite set. In particular,

$$e' := e - \sum_{n \in J} e_n \neq 0$$

satisfies $f(e') = 0$.

Step 3. We shall get a contradiction. By Step 2., we can take a maximal orthogonal family of non-zero projections $\{e_\alpha\}_{\alpha \in A}$ in \mathfrak{M} such that $e_\alpha \leq e_0$ and $f(e_\alpha) = 0$. Let $e := \sum_{\alpha \in A} e_\alpha$. The maximality of $\{e_\alpha\}_{\alpha \in A}$ and Step 2. implies $e = e_0$. Thus we have

$$0 \neq f(e_0) = \sum_{\alpha \in A} f(e_\alpha) = 0,$$

which is a contradiction. Hence there exists no non-zero SRT-continuous linear functional on $\overline{\mathfrak{M}}$. \square

Lemma 3.23. *Let \mathfrak{M}_a be an atomic finite von Neumann algebra, \mathfrak{M}_d be a diffuse finite von Neumann algebra and $\mathfrak{M} := \mathfrak{M}_a \oplus^b \mathfrak{M}_d$ be the direct sum von Neumann algebra. Denote the conjugate spaces of $(\overline{\mathfrak{M}_a}, SRT)$ and $(\overline{\mathfrak{M}}, SRT)$ by $(\overline{\mathfrak{M}_a})^*$ and $(\overline{\mathfrak{M}})^*$ respectively. For each $f \in (\overline{\mathfrak{M}_a})^*$, we define $I(f) \in (\overline{\mathfrak{M}})^*$ as*

$$I(f)(A \oplus B) := f(A), \quad A \oplus B \in \overline{\mathfrak{M}} = \overline{\mathfrak{M}_a} \oplus^b \overline{\mathfrak{M}_d},$$

then I is a bijection between $(\overline{\mathfrak{M}_a})^*$ onto $(\overline{\mathfrak{M}})^*$.

Proof. This follows immediately from Lemma 3.22. \square

Proof of Proposition 3.20. We have only to prove (1) \Rightarrow (2). Since \mathfrak{M} is spatially isomorphic to the direct sum of an atomic von Neumann algebra $\mathfrak{M}_{\text{atomic}}$ and a diffuse von Neumann algebra $\mathfrak{M}_{\text{diffuse}}$, it is enough to show that $\mathfrak{M}_{\text{diffuse}} = \{0\}$. Suppose $\mathfrak{M}_{\text{diffuse}} \neq \{0\}$ and take $y \in \mathfrak{M}_{\text{diffuse}} \setminus \{0\}$. Then, by locally convexity of $\overline{\mathfrak{M}}$, there exists a SRT-continuous linear functional f on $\overline{\mathfrak{M}}$ such that $f(0 \oplus y) \neq 0$. However this is a contradiction by Lemma 3.23. \square

Similarly one can prove the following proposition.

Proposition 3.24. *Let \mathfrak{M} be a finite von Neumann algebra. Then the following are equivalent.*

- (1) *There exists no non-zero SRT-continuous linear functional on $\overline{\mathfrak{M}}$.*
- (2) *\mathfrak{M} is diffuse.*

4 Lie Group-Lie Algebra Correspondences

In this section we state and prove the main result of this paper. As explained in the introduction, Lie theory for $U(\mathcal{H})$ is a difficult issue. What one has to resolve for discussing the Lie group-Lie algebra correspondence is a domain problem of the generators of one parameter subgroups of $G \subset U(\mathcal{H})$. The second to be discussed is a continuity of the Lie algebraic operations. However we can show that, for any strongly closed subgroup G of unitary group $U(\mathfrak{M})$ of some finite von Neumann algebra \mathfrak{M} , there exists canonically a complete topological Lie algebra. Since there are continuously many non-isomorphic finite von Neumann algebras on \mathcal{H} , there are also varieties of such groups. We hope that the ‘‘Lie Groups-Lie Algebras Correspondences’’ will play some important roles in the infinite dimensional Lie theory. We study the SRT-closed subalgebra of $\overline{\mathfrak{M}}$, too.

4.1 Existence of Lie Algebra

Let \mathfrak{M} be a finite von Neumann algebra acting on a Hilbert space \mathcal{H} . Recall that a densely defined closable operator A is called a skew-adjoint operator if $A^* = -A$, and A is called essentially skew-adjoint if \overline{A} is skew-adjoint.

Remark 4.1. In general, the strong limit of unitary operators is not necessarily unitary. It is known that $U(\mathfrak{M})$ is strongly closed if and only if \mathfrak{M} is a finite von Neumann algebra.

Definition 4.2. For a strongly closed subgroup G of $U(\mathfrak{M})$, the set

$$\mathfrak{g} = \text{Lie}(G) := \{A ; A^* = -A \text{ on } \mathcal{H}, e^{tA} \in G, \text{ for all } t \in \mathbb{R}\}$$

is called the Lie algebra of G . The complexification $\mathfrak{g}_{\mathbb{C}}$ of \mathfrak{g} is defined by

$$\mathfrak{g}_{\mathbb{C}} := \{\overline{A + iB} ; A, B \in \mathfrak{g}\}.$$

If $G = U(\mathfrak{M})$, we sometimes write \mathfrak{g} as $\mathfrak{u}(\mathfrak{M})$.

At first sight, it is not clear whether we can define algebraic operations on \mathfrak{g} . However,

Lemma 4.3. *Under the above notations, $\mathfrak{g} \subset \overline{\mathfrak{M}}$ holds.*

Proof. Let $u \in U(\mathfrak{M})$ and $A \in \mathfrak{g}$. By definition, we have $e^{tA}u = ue^{tA}$. Taking the strong derivative on each side, we have $uA \subset Au$. Since u is arbitrary we obtain $uA = Au$, which implies $A \in \overline{\mathfrak{M}}$. \square

Therefore the sum $\overline{A + B}$ and the Lie bracket $\overline{AB - BA}$ are well-defined operations in $\overline{\mathfrak{M}}$, but it is not clear whether they belong to \mathfrak{g} again. The next proposition guarantees the validity of the name ‘‘Lie algebra’’. The former part of the proof is based on the two lemmata established by Trotter-Kato and E. Nelson, which are of importance in their own.

Lemma 4.4 (Trotter-Kato, Nelson [13]). *Let A, B be skew-adjoint operators on a Hilbert space \mathcal{H} .*

(1) *If $A + B$ is essentially skew-adjoint on $\text{dom}(A) \cap \text{dom}(B)$, then it holds that*

$$e^{t(\overline{A+B})} = \text{s-}\lim_{n \rightarrow \infty} \left(e^{tA/n} e^{tB/n} \right)^n,$$

for all $t \in \mathbb{R}$.

(2) *If $(AB - BA)$ is essentially skew-adjoint on*

$$\text{dom}(A^2) \cap \text{dom}(AB) \cap \text{dom}(BA) \cap \text{dom}(B^2),$$

then it holds that

$$e^{t[A,B]} = \text{s-}\lim_{n \rightarrow \infty} \left(e^{-\sqrt{\frac{t}{n}}A} e^{-\sqrt{\frac{t}{n}}B} e^{\sqrt{\frac{t}{n}}A} e^{\sqrt{\frac{t}{n}}B} \right)^{n^2},$$

for all $t > 0$, where $[A, B] := \overline{AB - BA}$.

Lemma 4.5. *Let G be a strongly closed subgroup of $U(\mathfrak{M})$. Then \mathfrak{g} is a real Lie algebra with the Lie bracket $[X, Y] := \overline{XY - YX}$.*

Proof. Let $A, B \in \mathfrak{g}$. It suffices to prove that $\overline{A + B}$ and $\overline{AB - BA}$ belong to \mathfrak{g} . Since $\text{dom}(A) \cap \text{dom}(B)$ is completely dense, $A + B$ is essentially skew-adjoint. Therefore by Lemma 4.4 (1), we have $e^{t(\overline{A+B})} \in \overline{G^s} = G$ for all $t \in \mathbb{R}$. This implies $\overline{A + B} \in \mathfrak{g}$. It is clear that $\lambda A \in \mathfrak{g}$ for all $\lambda \in \mathbb{R}$. On the other hand, as $\overline{AB - BA}$ is essentially skew-adjoint on

$$\mathcal{D} := \text{dom}(AB) \cap \text{dom}(BA) \cap \text{dom}(A^2) \cap \text{dom}(B^2),$$

since \mathcal{D} is completely dense by Proposition 2.10 and $\overline{AB - BA} \in \overline{\mathfrak{M}}$. Therefore by Proposition 4.4 (2), we have $e^{t(\overline{AB - BA})} \in G$ for all $t > 0$. Thanks to the unitarity, this equality is also valid for $t < 0$. Thus we obtain $[A, B] \in \mathfrak{g}$. The associativity of the algebraic operations follows from the Murray-von Neumann's Theorem 2.17. \square

Now we state the main result of this paper, whose proof is almost completed in the previous arguments.

Theorem 4.6. *Let G be a strongly closed subgroup of the unitary group $U(\mathfrak{M})$ of a finite von Neumann algebra \mathfrak{M} . Then \mathfrak{g} is a complete topological real Lie algebra with respect to the strong resolvent topology. Moreover, $\mathfrak{g}_{\mathbb{C}}$ is a complete topological Lie $*$ -algebra.*

Proof. The Lie algebraic properties are proved in Lemma 4.5. By Lemma B.2, we see that \mathfrak{g} and $\mathfrak{g}_{\mathbb{C}}$ are SRT-closed Lie subalgebras of $\overline{\mathfrak{M}}$. As the algebraic operations $(X, Y) \mapsto \overline{X + Y}$, $[X, Y]$ are continuous with respect to the strong resolvent topology and, by Theorem 3.9, the topological properties follow. \square

Remark 4.7. It is easy to see that for $G = U(\mathfrak{M})$, its Lie algebra $\mathfrak{u}(\mathfrak{M})$ is equal to $\{A \in \mathfrak{M}; A^* = -A\}$ and the exponential map

$$\exp : \mathfrak{u}(\mathfrak{M}) \rightarrow U(\mathfrak{M})$$

is continuous by Lemma B.2 and surjective by the spectral theorem.

Proposition 4.8. *Let $\mathfrak{M}_1, \mathfrak{M}_2$ be finite von Neumann algebras on Hilbert spaces $\mathcal{H}_1, \mathcal{H}_2$ respectively. Let G_i be a strongly closed subgroup of $U(\mathfrak{M}_i)$ ($i = 1, 2$). For any strongly continuous group homomorphism $\varphi : G_1 \rightarrow G_2$, there exists a unique SRT-continuous Lie homomorphism $\Phi : \text{Lie}(G_1) \rightarrow \text{Lie}(G_2)$ such that $\varphi(e^A) = e^{\Phi(A)}$ for all $A \in \text{Lie}(G_1)$. In particular, if G_1 is isomorphic to G_2 , then $\text{Lie}(G_1)$ and $\text{Lie}(G_2)$ are isomorphic as a topological Lie algebra.*

Proof. Let X be an element in $\text{Lie}(G_1)$. From the strong continuity of φ , $t \mapsto \varphi(e^{tX})$ is a strongly continuous one-parameter unitary group. Therefore by Stone theorem, there exists uniquely a skew-adjoint operator $\Phi(X)$ on \mathcal{H}_2 such that $\varphi(e^{tX}) = e^{t\Phi(X)}$. This equality implies $\Phi(X) \in \text{Lie}(G_2)$. Since φ is strongly continuous, thanks to the Nelson's theorem, we see that

$$\begin{aligned} e^{t\Phi([X,Y])} &= \varphi(e^{t[X,Y]}) \\ &= \varphi\left(\text{s-}\lim_{n \rightarrow \infty} \left[e^{-\sqrt{\frac{t}{n}}X} e^{-\sqrt{\frac{t}{n}}Y} e^{\sqrt{\frac{t}{n}}X} e^{\sqrt{\frac{t}{n}}Y} \right]^n\right) \\ &= \text{s-}\lim_{n \rightarrow \infty} \left[\varphi\left(e^{-\sqrt{\frac{t}{n}}X}\right) \varphi\left(e^{-\sqrt{\frac{t}{n}}Y}\right) \varphi\left(e^{\sqrt{\frac{t}{n}}X}\right) \varphi\left(e^{\sqrt{\frac{t}{n}}Y}\right) \right]^n \\ &= \text{s-}\lim_{n \rightarrow \infty} \left[e^{-\sqrt{\frac{t}{n}}\Phi(X)} e^{-\sqrt{\frac{t}{n}}\Phi(Y)} e^{\sqrt{\frac{t}{n}}\Phi(X)} e^{\sqrt{\frac{t}{n}}\Phi(Y)} \right]^n \\ &= e^{t[\Phi(X),\Phi(Y)]}, \end{aligned}$$

for all $t > 0$. Taking the inverse of unitary operators, the equality $e^{t\Phi([X,Y])} = e^{t[\Phi(X),\Phi(Y)]}$ is also valid for all $t < 0$. Therefore from the uniqueness of a generator of one-parameter group, we have $\Phi([X, Y]) = [\Phi(X), \Phi(Y)]$. Similarly, we can prove that Φ is linear. Thus, Φ is a Lie homomorphism. The SRT-continuity of Φ follows immediately from the uniform continuity of φ . \square

As above, G has finite dimensional characters. On the other hand, it also has an infinite dimensional character.

Proposition 4.9. *Let \mathfrak{M} be a finite von Neumann algebra, then the following are equivalent.*

- (1) *The exponential map $\exp : \mathfrak{u}(\mathfrak{M}) \ni X \mapsto \exp(X) \in U(\mathfrak{M})$ is locally injective. Namely, the restriction of the map onto some SRT-neighborhood of $0 \in \mathfrak{M}$ is injective.*
- (2) *\mathfrak{M} is finite dimensional.*

Proof. (2) \Rightarrow (1) is trivial. We should only prove that (1) \Rightarrow (2).

Step 1. For each orthogonal family of non-zero projections in \mathfrak{M} , its cardinal number is finite. Indeed if there exists a orthogonal family of non-zero projections in \mathfrak{M} whose cardinal number is infinite, we can take a countably infinite subset of it and write it as $\{p_n\}_{n=1}^{\infty}$. Since p_n converges strongly to 0, it also converges to 0 in the strong resolvent topology. Define $x_n := 2\pi i p_n \neq 0$. Since the spectral set of p_n is $\{0, 1\}$, we have $e^{x_n} = 1$ for all $n \in \mathbb{N}$, while x_n converges to 0 in the strong resolvent topology. This implies that the map $\exp(\cdot)$ is not locally injective, which is a contradiction.

Step 2. \mathfrak{M} is atomic. Indeed if \mathfrak{M} is not atomic, the diffuse part of it is not $\{0\}$. Thus we can take an infinite sequence of non-zero mutually orthogonal projections in \mathfrak{M} . But this is a contradiction to Step 1..

Step 3. We shall show that \mathfrak{M} is finite dimensional. By Step 2., \mathfrak{M} is spatially isomorphic to the direct sum of a family $\{M_{n_\lambda}(\mathbb{C})\}_{\lambda \in \Lambda}$ ($n_\lambda \in \mathbb{N}$), where $M_{n_\lambda}(\mathbb{C})$ is the algebra of all $n_\lambda \times n_\lambda$ complex matrices. By Step 1., the cardinal number of Λ is finite. Hence \mathfrak{M} is finite dimensional. \square

Remark 4.10. $\text{Lie}(G)$ is not always locally convex, whereas most of infinite dimensional Lie theories, by contrast, assume locally convexity. Indeed, by Proposition 3.20, $\mathfrak{u}(\mathfrak{M})$ is locally convex if and only if \mathfrak{M} is atomic.

4.2 Closed Subalgebras of $\overline{\mathfrak{M}}$

Next, we characterize closed $*$ -subalgebras of $\overline{\mathfrak{M}}$.

Proposition 4.11. *Let \mathfrak{M} be a finite non Neumann algebra on a Hilbert space \mathcal{H} , \mathcal{R} be a SRT-closed $*$ -subalgebra of $\overline{\mathfrak{M}}$ with $1_{\mathcal{H}}$. Then there exists a unique von Neumann subalgebra \mathfrak{N} of \mathfrak{M} such that $\mathcal{R} = \overline{\mathfrak{N}}$.*

Remark 4.12. A von Neumann subalgebra of a finite von Neumann algebra is also finite.

Proof of Proposition 4.11. Put

$$\mathfrak{N} := \{x \in \mathcal{R} ; x \text{ is bounded}\}.$$

Since $0, 1 \in \mathfrak{N}$, \mathfrak{N} is not empty. We first show that \mathfrak{N} is a von Neumann algebra. It is clear that \mathfrak{N} is a subalgebra of \mathfrak{M} . Thus it is enough to check that \mathfrak{N} is closed with respect to the strong $*$ operator topology. Let $\{x_\alpha\}$ be a net in \mathfrak{N} converging to $x \in \mathfrak{N}$ with respect to the strong $*$ operator topology. So we have

$$\text{Re}(x_\alpha) \longrightarrow \text{Re}(x), \quad \text{Im}(x_\alpha) \longrightarrow \text{Im}(x)$$

with respect to the strong $*$ operator topology. By Lemma B.1,

$$\text{Re}(x_\alpha) \longrightarrow \text{Re}(x), \quad \text{Im}(x_\alpha) \longrightarrow \text{Im}(x)$$

with respect to the strong resolvent topology. As $\operatorname{Re}(x_\alpha) \in \mathcal{R}$, $\operatorname{Im}(x_\alpha) \in \mathcal{R}$ and \mathcal{R} is SRT-closed, we see that $\operatorname{Re}(x) \in \mathcal{R}$ and $\operatorname{Im}(x) \in \mathcal{R}$. Therefore

$$x = \operatorname{Re}(x) + i\operatorname{Im}(x) \in \mathcal{R}.$$

Since x is bounded, x belongs to \mathfrak{N} . Thus \mathfrak{N} is a von Neumann algebra.

Next, we show that $\mathcal{R} \subset \overline{\mathfrak{N}}$. Let A be an element of \mathcal{R} . It is enough to consider the case that A is self-adjoint. Put

$$\mathcal{C}_A := \bigcup_{n=1}^{\infty} \operatorname{ran}(E_A([-n, n])),$$

where $E_A(\cdot)$ is the spectral measure of A . \mathcal{C}_A is completely dense and all elements of \mathcal{C}_A are entire analytic vectors for A . Thus we have for all $\xi \in \mathcal{C}_A$,

$$e^{itA} \xi = \lim_{J \rightarrow \infty} \sum_{j=1}^J \frac{(itA)^j}{j!} \xi.$$

Therefore the sequence

$$\left\{ \sum_{j=1}^J \frac{(itA)^j}{j!} \right\}_{J=1}^{\infty} \subset \mathcal{R} \subset \overline{\mathfrak{N}}$$

converges almost everywhere to e^{itA} . By Proposition 3.14, it converges to e^{itA} with respect to the strong resolvent topology. Since \mathcal{R} is SRT-closed and e^{itA} is bounded, we get $e^{itA} \in \mathfrak{N}$. This implies A belongs to $\overline{\mathfrak{N}}$.

On the other hand, by the definition of \mathfrak{N} , $\mathfrak{N} \subset \mathcal{R}$. Since $\overline{\mathfrak{N}}$ is a SRT-closure of \mathfrak{N} , we see that $\overline{\mathfrak{N}} \subset \mathcal{R}$. Thus we conclude that $\overline{\mathfrak{N}} = \mathcal{R}$.

Finally, we show that the uniqueness for \mathfrak{N} . Let \mathfrak{L} be a von Neumann subalgebra of \mathfrak{M} satisfying $\overline{\mathfrak{L}} = \mathcal{R}$. Then, we have

$$\begin{aligned} \mathfrak{N} &= \{x \in \overline{\mathfrak{N}} ; x \text{ is bounded}\} \\ &= \{x \in \mathcal{R} ; x \text{ is bounded}\} \\ &= \{x \in \overline{\mathfrak{L}} ; x \text{ is bounded}\} \\ &= \mathfrak{L}. \end{aligned}$$

Thus \mathfrak{N} is unique. □

Corollary 4.13. *Let \mathfrak{M} be a finite non Neumann algebra on a Hilbert space \mathcal{H} , \mathfrak{g} be a real SRT-closed Lie subalgebra of $\mathfrak{u}(\mathfrak{M})$. Then the following are equivalent:*

- (1) *there exists a von Neumann subalgebra \mathfrak{N} of \mathfrak{M} such that $\mathfrak{g} = \mathfrak{u}(\mathfrak{N})$,*
- (2) *$1_{\mathcal{H}} \in \mathfrak{g}$ and for all $A, B \in \mathfrak{g}$, $i(\overline{AB + BA}) \in \mathfrak{g}$.*

In the above case, \mathfrak{N} is unique.

Proof. First of all, we shall show (1) \Rightarrow (2). Since $u(\mathfrak{N}) \subset \overline{\mathfrak{N}}$, we have $i(\overline{AB + BA}) \in \mathfrak{N}$. On the other hand,

$$u(\mathfrak{N}) = \{X \in \overline{\mathfrak{N}}; X^* = -X\}.$$

Thus $i(\overline{AB + BA}) \in u(\mathfrak{N}) = \mathfrak{g}$. Next we shall show (2) \Rightarrow (1). By direct computations, we see that

$$\mathcal{R} := \{\overline{X + iY} \in \overline{\mathfrak{M}}; X, Y \in \mathfrak{g}\}$$

is a SRT-closed $*$ -subalgebra of $\overline{\mathfrak{M}}$. Thus, by Proposition 4.11, there is a von Neumann subalgebra \mathfrak{N} of \mathfrak{M} such that $\mathcal{R} = \overline{\mathfrak{N}}$. Then we see that

$$\begin{aligned} \mathfrak{g} &= \{X \in \mathcal{R}; X^* = -X\} \\ &= \{X \in \overline{\mathfrak{N}}; X^* = -X\} \\ &= u(\mathfrak{N}). \end{aligned}$$

Finally, we show the uniqueness for \mathfrak{N} . Let \mathcal{L} be a von Neumann subalgebra of \mathfrak{M} satisfying $u(\mathcal{L}) = \mathfrak{g}$. Then, we have

$$\begin{aligned} \overline{\mathfrak{N}} &= \{\overline{X + iY}; X, Y \in u(\mathfrak{N})\} \\ &= \{\overline{X + iY}; X, Y \in u(\mathcal{L})\} \\ &= \overline{\mathcal{L}} \end{aligned}$$

By the uniqueness of Proposition 4.11, we get $\mathfrak{N} = \mathcal{L}$. □

5 Categorical Characterization of $\overline{\mathfrak{M}}$

5.1 Introduction

In this last section we turn the point of view and consider some categorical aspects of the $*$ -algebra $\overline{\mathfrak{M}}$. Especially, we determine when a $*$ -algebra \mathcal{R} of unbounded operators on a Hilbert space \mathcal{H} turns out to be of the form $\overline{\mathfrak{M}}$, without any reference to von Neumann algebraic structure in advance. As is well known, there are many examples of $*$ -algebra of unbounded operators that is not of the form $\overline{\mathfrak{M}}$. For example, many O^* -algebras [19] are not related to any affiliated operator algebra. Therefore, the appropriate condition to single out suitable class of $*$ -algebras of unbounded operators are necessary. For this purpose, we define the category **fRng** of unbounded operator algebras and compare this category with the category **fvN** of finite von Neumann algebras and show that both of them have natural tensor category structures (cf. Appendix C). Furthermore, we will see that they are isomorphic as a tensor category, in spite of the fact that the object in **fRng** is not locally convex in general while the one in **fvN** is a Banach space. However, the algebraic structures of $\overline{\mathfrak{M}}$ and \mathfrak{M} are very similar and in fact they constitute isomorphic categories. To begin with, let us introduce the structure of tensor category into **fRng**.

5.2 fvN and fRng as Tensor Categories

Now we turn to the question of characterizing the category **fRng** of $*$ -algebras of unbounded operators which are realized as $\overline{\mathfrak{M}}$, where \mathfrak{M} is a von Neumann algebra acting on a Hilbert space. It is well known that the usual tensor product $(\mathfrak{M}_1, \mathfrak{M}_2) \mapsto \mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2$ of von Neumann algebras and the tensor product of σ -weakly continuous homomorphisms $(\phi_1, \phi_2) \mapsto \phi_1 \otimes \phi_2$ makes the category of finite von Neumann algebras a tensor category. Therefore we define:

Definition 5.1. The category **fvN** is a category whose objects are pairs $(\mathfrak{M}, \mathcal{H})$ of a finite von Neumann algebra \mathfrak{M} acting on a Hilbert space \mathcal{H} and whose morphisms are σ -weakly continuous unital $*$ -homomorphisms. The unit object is $(\mathbb{C}1_{\mathbb{C}}, \mathbb{C})$. The tensor functor is the usual tensor product functor of von Neumann algebras. The definition of left and right unit constraint functors might be obvious.

If we are to characterize the objects in **fRng**, we must settle some subtleties due to the fact that we cannot use von Neumann algebraic structure from the outset. However, this difficulty can be overcome thanks to the the notion of the strong resolvent topology and the resolvent class whose definitions are independent of von Neumann algebras (See §3). We define **fRng** as follows.

Definition 5.2. The category **fRng** is a category whose objects $(\mathcal{R}, \mathcal{H})$ consist of a SRT-closed subset \mathcal{R} of the resolvent class $\mathcal{RC}(\mathcal{H})$ on a Hilbert space \mathcal{H} with the following properties:

- (1) $X + Y$ and XY are closable for all $X, Y \in \mathcal{R}$.
- (2) $\overline{X + Y}$, $\overline{\alpha X}$, \overline{XY} and X^* again belong to \mathcal{R} for all $X, Y \in \mathcal{R}$ and $\alpha \in \mathbb{C}$.
- (3) \mathcal{R} forms a $*$ -algebra with respect to the sum $\overline{X + Y}$, the scalar multiplication $\overline{\alpha X}$, the multiplication \overline{XY} and the involution X^* .
- (4) $1_{\mathcal{H}} \in \mathcal{R}$.

The morphism set between $(\mathcal{R}_1, \mathcal{H}_1)$ and $(\mathcal{R}_2, \mathcal{H}_2)$ consists of SRT-continuous unital $*$ -homomorphisms from \mathcal{R}_1 to \mathcal{R}_2 .

Remark 5.3. From the definition of **fRng**, it is not clear whether, for each objects in **fRng**, its algebraic operations are continuous or not. However, the next lemma shows that \mathcal{R} is a complete topological $*$ -algebra.

Lemma 5.4. *Let $(\mathcal{R}, \mathcal{H})$ be an object in **fRng**. Then there exists a unique finite von Neumann algebra \mathfrak{M} on \mathcal{H} such that $\mathcal{R} = \overline{\mathfrak{M}}$. Furthermore, $\mathfrak{M} = \mathcal{R} \cap \mathfrak{B}(\mathcal{H})$ holds.*

Proof. Define $\mathfrak{M} := \mathcal{R} \cap \mathfrak{B}(\mathcal{H})$. Then one can prove that \mathfrak{M} is von Neumann algebra by the same way as in Proposition 4.11.

We next show that $\mathcal{R} \subset \overline{\mathfrak{M}}$. Let $A \in \mathcal{R}$ be a self-adjoint operator. Define the dense subspace C_A according to the spectral decomposition of A :

$$C_A := \bigcup_{n=1}^{\infty} \text{ran}(E_A([-n, n])),$$

where

$$A = \int_{\mathbb{R}} \lambda dE_A(\lambda).$$

Since all $\xi \in C_A$ is an entire analytic vector for A , we have

$$e^{itA}\xi = \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{(itA)^k}{k!} \xi.$$

Let \mathfrak{M}_A be a von Neumann algebra generated by $\{E_A(J) ; J \in \mathcal{B}(\mathbb{R})\}$, where $\mathcal{B}(\mathbb{R})$ is the one dimensional Borel σ -field. Since \mathfrak{M}_A is abelian, it is a finite von Neumann algebra. It is also clear that

$$B_n := \sum_{k=0}^n \frac{(itA)^k}{k!} \in \overline{(\mathfrak{M}_A)} \cap \mathcal{R}$$

and $e^{itA} \in \mathfrak{M}_A$. Since C_A is completely dense for \mathfrak{M}_A , B_n converges almost everywhere to e^{itA} in $\overline{(\mathfrak{M}_A)}$. As \mathfrak{M}_A is finite, we see that B_n converges to e^{itA} in the strong resolvent topology. On the other hand, \mathcal{R} is SRT-closed and therefore $e^{itA} \in \mathcal{R} \cap \mathfrak{B}(\mathcal{H}) = \mathfrak{M}$, for all $t \in \mathbb{R}$. This implies $A \in \overline{\mathfrak{M}}$. For a general operator $B \in \mathcal{R}$, using real-imaginary part decomposition $B = \text{Re}(B) + i\text{Im}(B)$, we have $B \in \overline{\mathfrak{M}}$.

We shall show that $\overline{\mathfrak{M}} \subset \mathcal{R}$. Let $A \in \overline{\mathfrak{M}}$ and $A = U|A|$ be its polar decomposition, then $U \in \mathfrak{M} \subset \mathcal{R}$ and $|A| \in \overline{\mathfrak{M}}$. Let $|A| =: \int_0^{\infty} \lambda dE_{|A|}(\lambda)$ be the spectral decomposition of $|A|$. Put

$$x_n := \int_0^n \lambda dE_{|A|}(\lambda) \in \mathfrak{M} \subset \mathcal{R},$$

then x_n converges to $|A|$ in the strong resolvent topology. Thus $|A| \in \mathcal{R}$. Therefore $A = U|A| \in \mathcal{R}$.

The finiteness of \mathfrak{M} follows immediately from Theorem 2.21. \square

Note that for each finite von Neumann algebra \mathfrak{M} on a Hilbert space \mathcal{H} , $(\overline{\mathfrak{M}}, \mathcal{H})$ is an object in **fRng**.

The main result of this section is the next theorem.

Theorem 5.5. *The category **fRng** is a tensor category. Moreover, **fRng** and **fvN** are isomorphic as a tensor category.*

To prove this theorem, we need many lemmata. The proof is divided into several steps.

Next, we will define the tensor product $\mathcal{R}_1 \overline{\otimes} \mathcal{R}_2$ of objects \mathcal{R}_i ($i = 1, 2$) in \mathbf{fRng} (cf. Definition 5.9). For this purpose, let us review the notion of the tensor product of closed operators. Let A, B be densely defined closed operators on Hilbert spaces \mathcal{H}, \mathcal{K} , respectively. Let $A \otimes_0 B$ be an operator defined by

$$\begin{aligned} \text{dom}(A \otimes_0 B) &:= \text{dom}(A) \otimes_{\text{alg}} \text{dom}(B), \\ (A \otimes_0 B)(\xi \otimes \eta) &:= A\xi \otimes B\eta, \quad \xi \in \text{dom}(A), \eta \in \text{dom}(B). \end{aligned}$$

It is easy to see that $A \otimes_0 B$ is closable and denote its closure by $A \otimes B$.

Lemma 5.6. *Let $\mathfrak{M}_1, \mathfrak{M}_2$ be finite von Neumann algebras acting on Hilbert spaces $\mathcal{H}_1, \mathcal{H}_2$, respectively. Let $A \in \overline{\mathfrak{M}}_1$ and $B \in \overline{\mathfrak{M}}_2$. Then we have $A \otimes B \in \overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$.*

Proof. Let $x_i \in \mathfrak{M}'_i$ ($i = 1, 2$). For any $\xi \in \text{dom}(A)$ and $\eta \in \text{dom}(B)$, we have $(x_1 \otimes x_2)(\xi \otimes \eta) \in \text{dom}(A \otimes_0 B)$ and

$$\{(x_1 \otimes x_2)(A \otimes_0 B)\}(\xi \otimes \eta) = \{(A \otimes_0 B)(x_1 \otimes x_2)\}(\xi \otimes \eta).$$

Therefore, by the linearity, we have $(x_1 \otimes x_2)(A \otimes_0 B) \subset (A \otimes_0 B)(x_1 \otimes x_2)$. Since $(\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2})' = \overline{\mathfrak{M}'_1 \otimes \mathfrak{M}'_2}$ is the strong closure of $\mathfrak{M}'_1 \otimes_{\text{alg}} \mathfrak{M}'_2$, we have

$$y(A \otimes_0 B) \subset (A \otimes_0 B)y, \text{ for all } y \in (\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2})'.$$

Therefore by the limiting argument, we have $y(A \otimes B) \subset (A \otimes B)y$, which implies $A \otimes B$ is affiliated with $\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$. \square

Lemma 5.7. *Let A, B be densely defined closed operators on Hilbert spaces \mathcal{H}, \mathcal{K} with cores $\mathcal{D}_A, \mathcal{D}_B$ respectively. Then $\mathcal{D} := \mathcal{D}_A \otimes_{\text{alg}} \mathcal{D}_B$ is a core of $A \otimes B$.*

Proof. From the definition of $A \otimes B$, for any $\zeta \in \text{dom}(A \otimes B)$ and for any $\varepsilon > 0$, there exists some $\zeta_\varepsilon = \sum_{i=1}^n \xi_i \otimes \eta_i \in \text{dom}(A) \otimes_{\text{alg}} \text{dom}(B)$ such that

$$\|\zeta - \zeta_\varepsilon\| < \varepsilon, \quad \|(A \otimes B)\zeta - (A \otimes B)\zeta_\varepsilon\| < \varepsilon.$$

Put

$$C := \max_{1 \leq i \leq n} \{\|\xi_i\|, \|A\xi_i\|\} + 1 > 0.$$

Since \mathcal{D}_B is a core of B , there exists $\eta_i^\varepsilon \in \mathcal{D}_B$ such that

$$\|\eta_i - \eta_i^\varepsilon\| < \frac{\varepsilon}{nC}, \quad \|B\eta_i - B\eta_i^\varepsilon\| < \frac{\varepsilon}{nC}.$$

Put

$$C' := \max_{1 \leq i \leq n} \{\|\eta_i^\varepsilon\|, \|B\eta_i^\varepsilon\|\} + 1 > 0.$$

Similarly, since \mathcal{D}_A is a core of A , there exists $\xi_i^\varepsilon \in \mathcal{D}_A$ such that

$$\|\xi_i - \xi_i^\varepsilon\| < \frac{\varepsilon}{nC'}, \quad \|A\xi_i - A\xi_i^\varepsilon\| < \frac{\varepsilon}{nC'}.$$

Define $\zeta^\varepsilon := \sum_{i=1}^n \xi_i^\varepsilon \otimes \eta_i^\varepsilon \in \mathcal{D}$. Then we have

$$\begin{aligned}
\|\zeta - \zeta^\varepsilon\| &\leq \|\zeta - \zeta_\varepsilon\| + \|\zeta_\varepsilon - \zeta^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n \|\xi_i \otimes \eta_i - \xi_i^\varepsilon \otimes \eta_i^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n \|\xi_i \otimes \eta_i - \xi_i \otimes \eta_i^\varepsilon\| + \sum_{i=1}^n \|\xi_i \otimes \eta_i^\varepsilon - \xi_i^\varepsilon \otimes \eta_i^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n \|\xi_i\| \|\eta_i - \eta_i^\varepsilon\| + \sum_{i=1}^n \|\xi_i - \xi_i^\varepsilon\| \|\eta_i^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n C \cdot \frac{\varepsilon}{nC} + \sum_{i=1}^n \frac{\varepsilon}{nC'} \cdot C' \\
&= 3\varepsilon. \quad \dots (*)
\end{aligned}$$

Furthermore,

$$\begin{aligned}
&\|(A \otimes B)\zeta - (A \otimes B)\zeta^\varepsilon\| \\
&\leq \|(A \otimes B)\zeta - (A \otimes B)\zeta_\varepsilon\| + \|(A \otimes B)\zeta_\varepsilon - (A \otimes B)\zeta^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n \|A\xi_i \otimes B\eta_i - A\xi_i^\varepsilon \otimes B\eta_i^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n \|A\xi_i \otimes B\eta_i - A\xi_i \otimes B\eta_i^\varepsilon\| + \sum_{i=1}^n \|A\xi_i \otimes B\eta_i^\varepsilon - A\xi_i^\varepsilon \otimes B\eta_i^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n \|A\xi_i\| \|B\eta_i - B\eta_i^\varepsilon\| + \sum_{i=1}^n \|A\xi_i - A\xi_i^\varepsilon\| \|B\eta_i^\varepsilon\| \\
&\leq \varepsilon + \sum_{i=1}^n C \cdot \frac{\varepsilon}{nC} + \sum_{i=1}^n \frac{\varepsilon}{nC'} \cdot C' \\
&= 3\varepsilon \quad \dots (**).
\end{aligned}$$

(*) and (**) implies \mathcal{D} is a core of $A \otimes B$. □

Next lemma says that the tensor product of algebras of affiliated operators has a natural *-algebraic structures.

Lemma 5.8. *Let $\mathfrak{M}, \mathfrak{N}$ be finite von Neumann algebras acting on Hilbert spaces \mathcal{H}, \mathcal{K} respectively. Let $A, C \in \overline{\mathfrak{M}}, B, D \in \overline{\mathfrak{N}}$. Then we have*

- (1) $\overline{(A \otimes B)(C \otimes D)} = \overline{AC} \otimes \overline{BD}$.
- (2) $(A \otimes B)^* = A^* \otimes B^*$.
- (3) $\overline{A + C} \otimes \overline{B + D} = \overline{A \otimes B + A \otimes D + C \otimes B + C \otimes D}$.

$$(4) \quad \overline{\lambda(A \otimes B)} = \overline{\lambda A} \otimes B = A \otimes \overline{\lambda B} \quad (\lambda \in \mathbb{C}).$$

Proof. (1). From Proposition 2.13, $\mathcal{D}_1 := \{\xi \in \text{dom}(C); C\xi \in \text{dom}(A)\}$ is a core of \overline{AC} and $\mathcal{D}_2 := \{\eta \in \text{dom}(D); D\eta \in \text{dom}(B)\}$ is a core of \overline{BD} . Define $\mathcal{D} := \mathcal{D}_1 \otimes_{\text{alg}} \mathcal{D}_2$, which is a core of $\overline{AC} \otimes \overline{BD}$. Since

$$\text{dom}(\overline{(A \otimes B)(C \otimes D)}) \supset \text{dom}((A \otimes B)(C \otimes D)) \supset \mathcal{D},$$

it holds that for any $\zeta = \sum_{i=1}^n \xi_i \otimes \eta_i \in \mathcal{D}$, we have

$$\begin{aligned} \overline{(A \otimes B)(C \otimes D)}\zeta &= \sum_{i=1}^n AC\xi_i \otimes BD\eta_i = (\overline{AC} \otimes \overline{BD}) \sum_{i=1}^n \xi_i \otimes \eta_i \\ &= (\overline{AC} \otimes \overline{BD})\zeta. \end{aligned}$$

Therefore $\overline{(A \otimes B)(C \otimes D)} \supset (\overline{AC} \otimes \overline{BD})|_{\mathcal{D}}$. Since \mathcal{D} is a core of $\overline{AC} \otimes \overline{BD}$, we have (by taking the closure)

$$\overline{(A \otimes B)(C \otimes D)} \supset \overline{AC} \otimes \overline{BD}.$$

Since both operators belong to $\overline{\mathfrak{M} \otimes \mathfrak{N}}$ by Lemma 5.6, we have

$$\overline{(A \otimes B)(C \otimes D)} = \overline{AC} \otimes \overline{BD}.$$

by Proposition 2.14(2).

(2). It is easy to see that $(A \otimes B)^* \supset A^* \otimes B^*$. Since $(A \otimes B)^*$ and $A^* \otimes B^*$ are closed operators belonging to $\overline{\mathfrak{M} \otimes \mathfrak{N}}$, we have $(A \otimes B)^* = A^* \otimes B^*$ by Proposition 2.14 (2).

(3) and (4) can be easily shown in a similar manner as in (1). \square

Now we shall define the tensor product $\mathcal{R}_1 \overline{\otimes} \mathcal{R}_2$ of $(\mathcal{R}_1, \mathcal{H}_1)$ and $(\mathcal{R}_2, \mathcal{H}_2)$ in $\text{Obj}(\mathbf{fRng})$. Let \mathfrak{M}_i be finite von Neumann algebras on \mathcal{H}_i such that $\mathcal{R}_i = \overline{\mathfrak{M}_i}$ ($i = 1, 2$), respectively (cf. Lemma 5.4). From Lemma 5.8, the linear space $\mathcal{R}_1 \otimes_{\text{alg}} \mathcal{R}_2$ spanned by $\{A_1 \otimes A_2; A_i \in \mathcal{R}_i, i = 1, 2\}$ is a *-algebra. Since $\mathcal{R}_1 \otimes_{\text{alg}} \mathcal{R}_2$ is a subset of $\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$, it belongs to $\mathcal{RC}(\mathcal{H}_1 \otimes \mathcal{H}_2)$. Therefore:

Definition 5.9. Under the above notations, we define $\mathcal{R}_1 \overline{\otimes} \mathcal{R}_2$ to be the SRT-closure (for $\mathcal{H}_1 \otimes \mathcal{H}_2$) of $\mathcal{R}_1 \otimes_{\text{alg}} \mathcal{R}_2$.

Lemma 5.10. Let \mathcal{R}_i ($i = 1, 2$) be as above. Then $\mathcal{R}_1 \overline{\otimes} \mathcal{R}_2$ is also an object in \mathbf{fRng} . More precisely, if $\mathcal{R}_i = \overline{\mathfrak{M}_i}$, where \mathfrak{M}_i is a finite von Neumann algebra ($i = 1, 2$), then $\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2} = \overline{\mathfrak{M}_1} \overline{\otimes} \overline{\mathfrak{M}_2}$.

Proof. $\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2} \subset \overline{\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2}$: Let $T_i \in \overline{\mathfrak{M}_i}$ ($i = 1, 2$). Then we can show that $T_1 \otimes T_2 \in \overline{\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2}$ by Lemma 5.6. Therefore by the linearity, we obtain

$$\overline{\mathfrak{M}_1 \otimes_{\text{alg}} \mathfrak{M}_2} \subset \overline{\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2}.$$

As the left hand side is SRT-closed in $\mathcal{RC}(\mathcal{H}_1 \otimes \mathcal{H}_2)$, we have $\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2} \subset \overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$.

$\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2} \subset \overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$: It is clear that $\mathfrak{M}_1 \otimes_{\text{alg}} \mathfrak{M}_2 \subset \overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$. By the Kaplansky density theorem and Lemma 3.5, we have $\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2} \subset \overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$. By taking the SRT-closure, we obtain $\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2} \subset \overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}$. \square

The above Lemma says that $(\mathcal{R}_1 \otimes \mathcal{R}_2, \mathcal{H}_1 \otimes \mathcal{H}_2)$ is again an object in **fRng**.

Next, we discuss the extension of morphisms in **fvN** to ones in **fRng**. It requires some steps.

Lemma 5.11. *Let $(\mathfrak{M}_1, \mathcal{H}_1)$, $(\mathfrak{M}_2, \mathcal{H}_2)$ be finite von Neumann algebras. Then the mapping*

$$\begin{aligned} (\overline{\mathfrak{M}_1}, SRT) \times (\overline{\mathfrak{M}_2}, SRT) &\longrightarrow (\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}, SRT), \\ (A, B) &\longmapsto A \otimes B, \end{aligned}$$

is continuous.

Proof. Let $\{A_\alpha\}_\alpha \subset \overline{\mathfrak{M}_1}$, $\{B_\alpha\}_\alpha \subset \overline{\mathfrak{M}_2}$ be SRT-converging nets and $A \in \overline{\mathfrak{M}_1}$, $B \in \overline{\mathfrak{M}_2}$ be their limits respectively. We should only show that the net $\{A_\alpha \otimes B_\alpha\}_\alpha$ converges to $A \otimes B$ in the strong resolvent topology.

Step 1. The above claim is true if all A_α , B_α , A and B are self-adjoint. Indeed, it is easy to see that

$$e^{it(A_\alpha \otimes 1)} = e^{itA_\alpha} \otimes 1 \rightarrow e^{itA} \otimes 1 = e^{it(A \otimes 1)},$$

so that, by the limiting argument and Theorem 3.9, the SRT-convergence of $A_\alpha \otimes 1$ to $A \otimes 1$ follows. Similarly $1 \otimes B_\alpha$ converges to $1 \otimes B$ in the strong resolvent topology. Therefore, by Lemma 5.8 and the SRT-continuity of the multiplication, we have

$$A_\alpha \otimes B_\alpha = \overline{(A_\alpha \otimes 1)(1 \otimes B_\alpha)} \rightarrow \overline{(A \otimes 1)(1 \otimes B)} = A \otimes B.$$

Step 2. In a general case, by Lemma 5.8, we obtain

$$\begin{aligned} A_\alpha \otimes B_\alpha &= \overline{(\operatorname{Re}(A_\alpha) + i\operatorname{Im}(A_\alpha))} \otimes \overline{(\operatorname{Re}(B_\alpha) + i\operatorname{Im}(B_\alpha))} \\ &= \overline{\operatorname{Re}(A_\alpha) \otimes \operatorname{Re}(B_\alpha) + i\operatorname{Re}(A_\alpha) \otimes \operatorname{Im}(B_\alpha)} \\ &\quad \overline{+i\operatorname{Im}(A_\alpha) \otimes \operatorname{Re}(B_\alpha) - \operatorname{Im}(A_\alpha) \otimes \operatorname{Im}(B_\alpha)} \\ &\rightarrow \overline{\operatorname{Re}(A) \otimes \operatorname{Re}(B) + i\operatorname{Re}(A) \otimes \operatorname{Im}(B)} \\ &\quad \overline{+i\operatorname{Im}(A) \otimes \operatorname{Re}(B) - \operatorname{Im}(A) \otimes \operatorname{Im}(B)} \\ &= A \otimes B. \end{aligned}$$

Hence the proof of Lemma 5.11 is complete. \square

Lemma 5.12. *Let \mathfrak{M} be a finite von Neumann algebra on a Hilbert space \mathcal{H} and e is a projection in \mathfrak{M} , then \mathfrak{M}_e is also finite.*

Proof. Easy. □

Lemma 5.13. *Let A be a densely defined closed operator on a Hilbert space \mathcal{H} , \mathcal{K} be a closed subspace of \mathcal{K} such that $P_{\mathcal{K}}A \subset AP_{\mathcal{K}}$. Then the operator $B := A|_{\text{dom}(A) \cap \mathcal{K}}$ is a densely defined closed operator on \mathcal{K} .*

Proof. This is a straightforward verification. □

The next proposition guarantees the existence and the uniqueness of the extension of morphisms in \mathbf{fvN} to the morphisms in \mathbf{fRng} . Note that the claim is not trivial, because many σ -weakly continuous linear mappings between finite von Neumann algebras cannot be extended SRT-continuously to the algebra of affiliated operators. Indeed, we can not extend any σ -weakly continuous state on \mathfrak{M} SRT-continuously on $\overline{\mathfrak{M}}$ if \mathfrak{M} is diffuse.

Proposition 5.14. *Let $\mathfrak{M}_1, \mathfrak{M}_2$ be finite von Neumann algebras on Hilbert spaces $\mathcal{H}_1, \mathcal{H}_2$ respectively.*

- (1) *For each SRT-continuous unital $*$ -homomorphism $\Phi : \overline{\mathfrak{M}}_1 \rightarrow \overline{\mathfrak{M}}_2$, the restriction φ of Φ onto \mathfrak{M}_1 is a σ -weakly continuous unital $*$ -homomorphism from \mathfrak{M}_1 to \mathfrak{M}_2 .*
- (2) *Conversely, for each σ -weakly continuous unital $*$ -homomorphism $\varphi : \mathfrak{M}_1 \rightarrow \mathfrak{M}_2$, there exists a unique SRT-continuous unital $*$ -homomorphism $\Phi : \overline{\mathfrak{M}}_1 \rightarrow \overline{\mathfrak{M}}_2$ such that $\Phi|_{\mathfrak{M}_1} = \varphi$.*

Proof. (1). We have to prove that Φ maps all bounded operators to bounded operators. For any $u \in U(\mathfrak{M}_1)$ and $\xi \in \text{dom}(\Phi(u)^*\Phi(u))$, we have

$$\begin{aligned} \|\Phi(u)\xi\|^2 &= \langle \xi, \Phi(u)^*\Phi(u)\xi \rangle = \langle \xi, \Phi(u^*u)\xi \rangle \\ &= \langle \xi, \Phi(1)\xi \rangle = \|\xi\|^2. \end{aligned}$$

Since $\text{dom}(\Phi(u)^*\Phi(u))$ is a (completely) dense subspace, $\Phi(u) \in \mathfrak{M}_2$ and $\Phi(u)$ is an isometry. Therefore the finiteness of \mathfrak{M}_2 implies $\Phi(u) \in U(\mathfrak{M}_2)$. Thus, we see that $\Phi(U(\mathfrak{M}_1)) \subset U(\mathfrak{M}_2)$. Since any element in \mathfrak{M}_1 is a linear combination of $U(\mathfrak{M}_1)$, Φ maps \mathfrak{M}_1 into \mathfrak{M}_2 . To show that φ is σ -weakly continuous, it is sufficient to prove the (σ -) strong continuity on the unit ball, because it is a homomorphism. Since the strong resolvent topology coincides with the strong operator topology on the closed unit ball by Lemma 3.5, φ is strongly continuous on the closed unit ball. Therefore φ is a σ -weakly continuous homomorphism.

(2). Regard φ as a composition of a surjection $\varphi' : \mathfrak{M}_1 \rightarrow \varphi(\mathfrak{M}_1)$ and the inclusion map $\iota : \varphi(\mathfrak{M}_1) \hookrightarrow \mathfrak{M}_2$. Note that the σ -weak continuity of φ implies $\varphi(\mathfrak{M}_1)$ is a von Neumann algebra. Since φ' is surjective, from Theorem IV.5.5 of [21], there exists a Hilbert space \mathcal{K} , a projection $e' \in P(\mathfrak{M}'_1 \overline{\otimes} \mathfrak{B}(\mathcal{K}))$ and a unitary operator $U : e'(\mathcal{H}_1 \otimes \mathcal{K}) \xrightarrow{\sim} \mathcal{H}_2$ such that

$$\varphi'(x) = U(x \otimes 1_{\mathcal{K}})e'U^*$$

for all $x \in \mathfrak{M}_1$. Now we want to define the extension Φ' of φ' to $\overline{\mathfrak{M}_1} \rightarrow \overline{\varphi(\mathfrak{M}_1)}$. Then we define Φ' as follows:

$$\begin{array}{ccc} \overline{\mathfrak{M}_1} & \overset{\Phi'}{\dashrightarrow} & \overline{\varphi(\mathfrak{M}_1)} \\ \downarrow \cdot \mathbb{1} & \circlearrowleft & \uparrow U \cdot U^* \\ \overline{\mathfrak{M}_1} \otimes \mathbb{C}1_{\mathcal{K}} & \xrightarrow{\text{reduction by } e'} & (\overline{\mathfrak{M}_1} \otimes \mathbb{C}1_{\mathcal{K}})_{e'} \end{array}$$

More precisely, we define

$$Z = (X \otimes 1)_{e'} := e'(X \otimes 1)|_{\text{ran}(e') \cap \text{dom}(X \otimes 1)}, \quad \Phi'(X) := UZU^*.$$

We have $Z \in \overline{(\mathfrak{M}_1 \otimes \mathbb{C}1_{\mathcal{K}})_{e'}}$. Indeed, since e' commutes with $\mathfrak{M} \otimes \mathbb{C}1_{\mathcal{K}}$, it reduces the operator $X \otimes 1$ and therefore by Lemma 5.13, $(X \otimes 1)_{e'}$ is a densely defined closed operator on $\text{ran}(e')$. Since $(\mathfrak{N}_f)' = (\mathfrak{N}')_f$ for each von Neumann algebra \mathfrak{N} and $f \in P(\mathfrak{N}')$, the affiliation property is manifest. In addition, by Lemma 5.12, $(\mathfrak{M} \otimes \mathbb{C}1_{\mathcal{K}})_{e'}$ is a finite von Neumann algebra. Next, we prove the map $\overline{\mathfrak{M}} \ni X \mapsto (X \otimes 1)_{e'} \in \overline{(\mathfrak{M}_1 \otimes \mathbb{C}1_{\mathcal{K}})_{e'}}$ is a SRT-continuous unital *-homomorphism. The continuity follows from Lemma 5.11. To prove that it is a *-homomorphism, we have to show that for $X, Y \in \overline{\mathfrak{M}}$,

$$\begin{aligned} \overline{(X + Y) \otimes 1}_{e'} &= \overline{(X \otimes 1)_{e'} + (Y \otimes 1)_{e'}}, \\ \overline{(XY) \otimes 1}_{e'} &= \overline{(X \otimes 1)_{e'}(Y \otimes 1)_{e'}}, \\ \overline{(X \otimes 1)_{e'}^*} &= \overline{(X^* \otimes 1)_{e'}}. \end{aligned}$$

To prove the first equality, by Lemma 5.8, we see that

$$\begin{aligned} \overline{(X + Y) \otimes 1}_{e'} &= \overline{(X \otimes 1 + Y \otimes 1)_{e'}} \\ &\supset (X \otimes 1)_{e'} + (Y \otimes 1)_{e'}. \end{aligned}$$

Taking the closure, by Lemma 2.14, we have

$$\overline{(X + Y) \otimes 1}_{e'} = \overline{(X \otimes 1)_{e'} + (Y \otimes 1)_{e'}}.$$

The others are proved in a similar manner. Next, by Lemma 3.19, the correspondence $\overline{\mathfrak{M}_1} \ni X \mapsto U(X \otimes 1_{\mathcal{K}})_{e'}U^* \in \overline{\varphi(\mathfrak{M}_1)} \subset \overline{\mathfrak{M}_2}$ defines a SRT-continuous unital *-homomorphism Φ' which is clearly an extension of φ' . Therefore by considering $\Phi := \iota' \circ \Phi' : \overline{\mathfrak{M}_1} \rightarrow \overline{\mathfrak{M}_2}$ is the desired extension of φ , where $\iota' : \overline{\mathfrak{M}_1} \hookrightarrow \overline{\mathfrak{M}_2}$ is a mere inclusion. Finally, we prove the uniqueness of the extension. Let Ψ be another SRT-continuous unital *-homomorphism such that $\Psi|_{\mathfrak{M}_1} = \varphi$. Let $X \in \overline{\mathfrak{M}_1}$. Then from the SRT-density of \mathfrak{M}_1 in $\overline{\mathfrak{M}_1}$, there exists a sequence $\{x_n\} \subset \mathfrak{M}_1$ such that $\lim_{n \rightarrow \infty} x_n = X$ in the strong resolvent topology. Therefore we have

$$\begin{aligned} \Psi(X) &= \lim_{n \rightarrow \infty} \Psi(x_n) = \lim_{n \rightarrow \infty} \varphi(x_n) \\ &= \lim_{n \rightarrow \infty} \Phi(x_n) = \Phi(X). \end{aligned}$$

□

The next lemmata, together with Lemma 5.10, implies that \mathbf{fRng} is a tensor category.

Lemma 5.15. *Let $\mathcal{R}_i, \mathcal{S}_i$ ($i = 1, 2$) be objects in $\mathbf{Obj}(\mathbf{fRng})$. If $\Psi_1 : \mathcal{R}_1 \rightarrow \mathcal{S}_1$, $\Psi_2 : \mathcal{R}_2 \rightarrow \mathcal{S}_2$ are SRT-continuous unital *-homomorphisms, then there exists a unique SRT-continuous unital *-homomorphism $\Psi : \mathcal{R}_1 \overline{\otimes} \mathcal{R}_2 \rightarrow \mathcal{S}_1 \overline{\otimes} \mathcal{S}_2$ such that $\Psi(A \otimes B) = \Psi_1(A) \otimes \Psi_2(B)$, for all $A \in \mathcal{R}_1$ and $B \in \mathcal{R}_2$. We define $\Psi_1 \otimes \Psi_2$ to be the map Ψ .*

Proof. Let ψ_i be the restrictions of Ψ_i onto \mathfrak{M}_i ($i = 1, 2$). Then ψ_i is a σ -weakly continuous unital *-homomorphism from \mathfrak{M}_i to \mathfrak{N}_i , where $\mathfrak{N}_i = \mathcal{S}_i$. Thus there exists a σ -weakly continuous unital *-homomorphism ψ from $\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2$ to $\mathfrak{N}_1 \overline{\otimes} \mathfrak{N}_2$ such that

$$\psi(x \otimes y) = \psi_1(x) \otimes \psi_2(y), \quad x \in \mathfrak{M}_1, y \in \mathfrak{M}_2.$$

By Proposition 5.14, there exists a SRT-continuous unital *-homomorphism Ψ from $\mathcal{R}_1 \overline{\otimes} \mathcal{R}_2$ to $\mathcal{S}_1 \overline{\otimes} \mathcal{S}_2$ whose restriction to $\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2$ is equal to ψ . For all $A \in \mathcal{R}_1$, $B \in \mathcal{R}_2$, we can take sequences $\{x_k\}_{k=1}^\infty \subset \mathfrak{M}_1$, $\{y_l\}_{l=1}^\infty \subset \mathfrak{M}_2$ converging to A , B in the strong resolvent topology, respectively. Therefore, by Proposition 5.11, we have

$$\begin{aligned} \Psi(A \otimes B) &= \lim_{k \rightarrow \infty} \Psi(x_k \otimes y_k) = \lim_{k \rightarrow \infty} \psi_1(x_k) \otimes \psi_2(y_k) \\ &= \lim_{k \rightarrow \infty} \Psi_1(x_k) \otimes \Psi_2(y_k) = \Psi_1(A) \otimes \Psi_2(B). \end{aligned}$$

□

Lemma 5.16. *Let $(\mathcal{R}_i, \mathcal{H}_i)$ ($i = 1, 2, 3$) be objects in \mathbf{fRng} . Then we have a unique *-isomorphism which is homeomorphic with respect to the strong resolvent topology:*

$$\begin{aligned} (\mathcal{R}_1 \overline{\otimes} \mathcal{R}_2) \overline{\otimes} \mathcal{R}_3 &\cong \mathcal{R}_1 \overline{\otimes} (\mathcal{R}_2 \overline{\otimes} \mathcal{R}_3) \\ (X_1 \otimes X_2) \otimes X_3 &\mapsto X_1 \otimes (X_2 \otimes X_3), \text{ for all } X_i \in \mathcal{R}_i \end{aligned}$$

We denote the map as $\alpha_{\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3}$.

Proof. Let \mathfrak{M}_i be a finite von Neumann algebra such that $\mathcal{R}_i = \overline{\mathfrak{M}}_i$ ($i = 1, 2, 3$). Let α_0 be the *-isomorphism from $(\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2) \overline{\otimes} \mathfrak{M}_3$ onto $\mathfrak{M}_1 \overline{\otimes} (\mathfrak{M}_2 \overline{\otimes} \mathfrak{M}_3)$ defined by $(x_1 \otimes x_2) \otimes x_3 \mapsto x_1 \otimes (x_2 \otimes x_3)$. By Lemma 5.10, both $(\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2) \overline{\otimes} \mathfrak{M}_3$ and $\mathfrak{M}_1 \overline{\otimes} (\mathfrak{M}_2 \overline{\otimes} \mathfrak{M}_3)$ are generated by $(\mathfrak{M}_1 \overline{\otimes} \mathfrak{M}_2) \overline{\otimes} \mathfrak{M}_3$ and $\mathfrak{M}_1 \overline{\otimes} (\mathfrak{M}_2 \overline{\otimes} \mathfrak{M}_3)$, respectively. Therefore by Proposition 5.14, α_0 can be extended to the desired *-isomorphism $\alpha_{\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3}$. □

Proposition 5.17. *\mathbf{fRng} is a tensor category.*

Proof. We define the tensor product $\otimes : \mathbf{fRng} \times \mathbf{fRng} \rightarrow \mathbf{fRng}$ by

$$(\mathcal{R}_1, \mathcal{H}_1) \otimes (\mathcal{R}_2, \mathcal{H}_2) := (\mathcal{R}_1 \overline{\otimes} \mathcal{R}_2, \mathcal{H}_1 \otimes \mathcal{H}_2)$$

and for two morphisms $\Psi_i : (\mathcal{R}_i, \mathcal{H}_i) \rightarrow (\mathcal{S}_i, \mathcal{K}_i)$ ($i = 1, 2$), define $\Psi_1 \otimes \Psi_2$ according to Lemma 5.15. The unit object is $I := (\mathbb{C}1_{\mathbb{C}}, \mathbb{C})$. The associative constraint $\alpha_{\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3}$ is the map defined in Lemma 5.16. The naturality of $\alpha_{\mathcal{R}_1, \mathcal{R}_2, \mathcal{R}_3}$ follows from Proposition 5.14. The definition of left (resp. right) constraint λ . (resp. ρ .) might be clear. Now it is a routine task to verify that the data $(\mathbf{fRng}, \otimes, I, \alpha, \lambda, \rho)$ constitutes a tensor category. \square

Now we will prove that \mathbf{fvN} is isomorphic to \mathbf{fRng} as a tensor category. Define two functors $\mathcal{E} : \mathbf{fvN} \rightarrow \mathbf{fRng}$, $\mathcal{F} : \mathbf{fRng} \rightarrow \mathbf{fvN}$.

Definition 5.18. Define two correspondences \mathcal{E} , \mathcal{F} as follows:

- (1) For each object $(\mathfrak{M}, \mathcal{H})$ in \mathbf{fvN} ,

$$\mathcal{E}(\mathfrak{M}, \mathcal{H}) := (\overline{\mathfrak{M}}, \mathcal{H}),$$

which is an object in \mathbf{fRng} . For each morphism $\varphi : \mathfrak{M}_1 \rightarrow \mathfrak{M}_2$ in \mathbf{fvN} , $\mathcal{E}(\varphi) : \overline{\mathfrak{M}}_1 \rightarrow \overline{\mathfrak{M}}_2$ is the unique SRT-continuous extension of φ to $\overline{\mathfrak{M}}_1$, so that $\mathcal{E}(\varphi)$ is a morphism in \mathbf{fRng} by Proposition 5.14.

- (2) For each object $(\mathcal{R}, \mathcal{H})$ in \mathbf{fRng} ,

$$\mathcal{F}(\mathcal{R}, \mathcal{H}) := (\mathcal{R} \cap \mathfrak{B}(\mathcal{H}), \mathcal{H}).$$

For each morphism $\Phi : \mathcal{R}_1 \rightarrow \mathcal{R}_2$ in \mathbf{fRng} , $\mathcal{F}(\Phi) := \Phi|_{\mathcal{R}_1 \cap \mathfrak{B}(\mathcal{H})}$, which is a morphism in \mathbf{fvN} by Proposition 5.14.

Lemma 5.19. \mathcal{E} and \mathcal{F} are tensor functors.

Proof. We define the tensor functor (\mathcal{E}, h_1, h_2) , where

$$\begin{aligned} h_1 : (\mathbb{C}1_{\mathbb{C}}, \mathbb{C}) &\xrightarrow{\text{id}} (\mathbb{C}1_{\mathbb{C}}, \mathbb{C}) = \mathcal{E}((\mathbb{C}1_{\mathbb{C}}, \mathbb{C})), \\ h_2((\mathfrak{M}_1, \mathcal{H}_1), (\mathfrak{M}_2, \mathcal{H}_2)) &: \overline{\mathfrak{M}}_1 \otimes \overline{\mathfrak{M}}_2 \xrightarrow{\text{id}} \overline{\mathfrak{M}}_1 \otimes \overline{\mathfrak{M}}_2, \end{aligned}$$

can be taken to be identity morphisms thanks to Lemma 5.15. It is clear that $\mathcal{E}(1_{\mathfrak{M}}) = 1_{\overline{\mathfrak{M}}}$, where $1_{\mathfrak{M}}$ and $1_{\overline{\mathfrak{M}}}$ are identity map of \mathfrak{M} and $\overline{\mathfrak{M}}$, respectively. Let $\mathfrak{M}_1 \xrightarrow{\varphi_1} \mathfrak{M}_2 \xrightarrow{\varphi_2} \mathfrak{M}_3$ be a sequence of morphisms in \mathbf{fvN} . Let $x \in \mathfrak{M}_1$. It holds that

$$\begin{aligned} \mathcal{E}(\varphi_2 \circ \varphi_1)(x) &= (\varphi_2 \circ \varphi_1)(x) = \mathcal{E}(\varphi_2)(\varphi_1(x)) \\ &= \mathcal{E}(\varphi_2)(\mathcal{E}(\varphi_1)(x)) = (\mathcal{E}(\varphi_2) \circ \mathcal{E}(\varphi_1))(x). \end{aligned}$$

By Proposition 5.14 (2), we have $\mathcal{E}(\varphi_2 \circ \varphi_1) = \mathcal{E}(\varphi_2) \circ \mathcal{E}(\varphi_1)$. Therefore \mathcal{E} is a functor. The conditions for (\mathcal{E}, h_1, h_2) to be a tensor functor are described as the following three diagrams, the commutativity of which are almost obvious

by Proposition 5.14 and “ \sim ” symbols are followed from Lemma 5.16.

$$\begin{array}{ccc}
(\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}) \otimes \overline{\mathfrak{M}_3} & \xrightarrow{\sim} & \overline{\mathfrak{M}_1} \otimes (\overline{\mathfrak{M}_2 \otimes \mathfrak{M}_3}) \\
\text{id} \downarrow & & \downarrow \text{id} \\
(\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}) \otimes \overline{\mathfrak{M}_3} & \circlearrowleft & \overline{\mathfrak{M}_1} \otimes (\overline{\mathfrak{M}_2 \otimes \mathfrak{M}_3}) \\
\text{id} \downarrow & & \downarrow \text{id} \\
(\overline{\mathfrak{M}_1 \otimes \mathfrak{M}_2}) \otimes \overline{\mathfrak{M}_3} & \xrightarrow{\sim} & \overline{\mathfrak{M}_1} \otimes (\overline{\mathfrak{M}_2 \otimes \mathfrak{M}_3})
\end{array}$$

$$\begin{array}{ccc}
\mathbb{C} \otimes \overline{\mathfrak{M}} & \xrightarrow{1 \otimes X \mapsto X} & \overline{\mathfrak{M}} \\
\text{id} \downarrow & \circlearrowleft & \uparrow \\
\mathbb{C} \otimes \overline{\mathfrak{M}} & \xrightarrow{\text{id}} & \overline{\mathbb{C} \otimes \mathfrak{M}}
\end{array}
\quad
\begin{array}{ccc}
\overline{\mathfrak{M}} \otimes \mathbb{C} & \xrightarrow{X \otimes 1 \mapsto X} & \overline{\mathfrak{M}} \\
\text{id} \downarrow & \circlearrowleft & \uparrow \\
\overline{\mathfrak{M}} \otimes \mathbb{C} & \xrightarrow{\text{id}} & \overline{\mathfrak{M} \otimes \mathbb{C}}
\end{array}$$

Thus, (\mathcal{E}, h_1, h_2) is a tensor functor. The proof that $(\mathcal{F}, h'_1, h'_2)$ is a tensor functor, including the definitions of h'_1, h'_2 are easier. \square

Now we are able to prove the main theorem easily.

Proof of Theorem 5.5. We will show that \mathcal{E} and \mathcal{F} are the inverse tensor functor of each other. By Lemma 5.19, they are tensor functors. Let $(\mathfrak{M}_i, \mathcal{H}_i)$ ($i = 1, 2$) be in $\text{Obj}(\mathbf{fvN})$. Let $\varphi : \mathfrak{M}_1 \rightarrow \mathfrak{M}_2$ be a morphism in \mathbf{fvN} . Proposition 5.14 implies $\varphi = (\mathcal{F} \circ \mathcal{E})(\varphi)$. By Proposition 5.4, we have

$$(\mathfrak{M}_i, \mathcal{H}_i) = (\overline{\mathfrak{M}_i} \cap \mathfrak{B}(\mathcal{H}_i), \mathcal{H}_i) = (\mathcal{F} \circ \mathcal{E})(\mathfrak{M}_i, \mathcal{H}_i),$$

therefore $\mathcal{F} \circ \mathcal{E} = \text{id}_{\mathbf{fvN}}$.

Let $(\mathcal{R}_i, \mathcal{H}_i)$ ($i = 1, 2$) be objects in \mathbf{fRng} , $\Phi : (\mathcal{R}_1, \mathcal{H}_1) \rightarrow (\mathcal{R}_2, \mathcal{H}_2)$ be a morphism in \mathbf{fRng} . By Proposition 5.4, we have $\mathcal{R}_i = \overline{\mathfrak{M}_i}$ for a unique $(\mathfrak{M}_i, \mathcal{H}_i)$ in $\text{Obj}(\mathbf{fvN})$. Similarly, we can prove that

$$(\mathcal{R}_i, \mathcal{H}_i) = (\mathcal{E} \circ \mathcal{F})(\mathcal{R}_i, \mathcal{H}_i), \quad (\mathcal{E} \circ \mathcal{F})(\Phi) = \Phi,$$

hence $\mathcal{E} \circ \mathcal{F} = \text{id}_{\mathbf{fRng}}$. \square

Finally, we remark the correspondence of factors in \mathbf{fvN} and ones in \mathbf{fRng} . Recall that, for a *-algebra \mathcal{A} , its *center* $Z(\mathcal{A})$ is defined by

$$Z(\mathcal{A}) := \{x \in \mathcal{A} ; xy = yx, \text{ for all } y \in \mathcal{A}\}.$$

$Z(\mathcal{A})$ is also a *-algebra.

Proposition 5.20. *Let \mathfrak{M} be a finite von Neumann algebra on \mathcal{H} . The following conditions are equivalent.*

- (1) *The center $Z(\overline{\mathfrak{M}})$ of $\overline{\mathfrak{M}}$ is trivial. I.e., $Z(\overline{\mathfrak{M}}) = \mathbb{C}1_{\mathcal{H}}$.*

(2) The center $Z(\mathfrak{M})$ of \mathfrak{M} is trivial.

Proof. (1) \Rightarrow (2) is evident.

(2) \Rightarrow (1). Let $A \in \overline{\mathfrak{M}}$ be a self-adjoint element of the center $Z(\overline{\mathfrak{M}})$. For any $u \in U(\mathfrak{M}')$, we have $uAu^* = A$. Therefore from the unitary covariance of the functional calculus, it holds that $u(A-i)^{-1}u^* = (A-i)^{-1}$ and $(A-i)^{-1} \in \mathfrak{M} \cap \mathfrak{M}' = \mathbb{C}1$. Hence $(A-i)^{-1} = \alpha 1$ for some $\alpha \in \mathbb{C}$. By operating $A-i$ on both sides, we see that $A \in \mathbb{C}1$. For a general closed operator $A \in Z(\overline{\mathfrak{M}})$, we know that there is a canonical decomposition $A = \overline{\operatorname{Re}(A)} + i \overline{\operatorname{Im}(A)}$. Since A belongs to $Z(\overline{\mathfrak{M}})$, $\operatorname{Re}(A)$, $\operatorname{Im}(A)$ also belong to $Z(\mathfrak{M}) = \mathbb{C}1$. Therefore $A \in \mathbb{C}1$. \square

A Direct Sums of Operators

We recall the theory of direct sums of operators and show some facts. We do not give proofs for well-known facts. See e.g., [2].

Let $\{\mathcal{H}_\alpha\}_\alpha$ be a family of Hilbert spaces and $\mathcal{H} = \bigoplus_\alpha \mathcal{H}_\alpha$ be the direct sum Hilbert space of $\{\mathcal{H}_\alpha\}_\alpha$, i.e.,

$$\mathcal{H} := \left\{ \xi = \{\xi^{(\alpha)}\}_\alpha ; \xi^{(\alpha)} \in \mathcal{H}_\alpha, \sum_\alpha \|\xi^{(\alpha)}\|^2 < \infty. \right\}.$$

For a subspace \mathcal{D}_α of \mathcal{H}_α , we set

$$\widehat{\bigoplus_\alpha \mathcal{D}_\alpha} := \left\{ \xi = \{\xi^{(\alpha)}\}_\alpha \in \mathcal{H} ; \xi^{(\alpha)} \in \mathcal{D}_\alpha, \xi^{(\alpha)} = 0 \text{ except finitely many } \alpha. \right\}.$$

It is known that $\widehat{\bigoplus_\alpha \mathcal{D}_\alpha}$ is dense in \mathcal{H} whenever each \mathcal{D}_α is dense in \mathcal{H}_α .

Next we recall the direct sum of unbounded operators. Let A_α be a (possibly unbounded) linear operator on \mathcal{H}_α . We define the linear operator $A = \bigoplus_\alpha A_\alpha$ on \mathcal{H} as follows:

$$\begin{aligned} \operatorname{dom}(A) &:= \left\{ \xi = \{\xi^{(\alpha)}\}_\alpha \in \mathcal{H} ; \xi^{(\alpha)} \in \operatorname{dom}(A_\alpha), \sum_\alpha \|A_\alpha \xi^{(\alpha)}\|^2 < \infty. \right\}, \\ (A\xi)^{(\alpha)} &:= A_\alpha \xi^{(\alpha)}, \quad \xi \in \operatorname{dom}(A). \end{aligned}$$

A is said to be the *direct sum* of $\{A_\alpha\}_\alpha$. It is easy to see that if each A_α is a densely defined closed operator then so is A . In this case,

$$A^* = \bigoplus_\alpha A_\alpha^*$$

holds. The following lemmata are well-known.

Lemma A.1. *Assume the above notations.*

(1) $A \in \mathfrak{B}(\mathcal{H})$ if and only if each A_α is in $\mathfrak{B}(\mathcal{H}_\alpha)$ and $\sup_\alpha \|A_\alpha\| < \infty$.
In this case,

$$\|A\| = \sup_\alpha \|A_\alpha\|$$

holds.

(2) A is unitary if and only if each A_α is unitary.

(3) A is projection if and only if each A_α is projection. In this case,

$$\text{ran}(A) = \bigoplus_\alpha \text{ran}(A_\alpha)$$

holds.

Lemma A.2. Assume that each A_α is closed. Let \mathcal{D}_α be a core of A_α . Then $\widehat{\bigoplus_\alpha \mathcal{D}_\alpha}$ is a core of A .

Lemma A.3. Assume that each A_α is (possibly unbounded) self-adjoint.

(1) A is self-adjoint.

(2) For any complex valued Borel function f on \mathbb{R} ,

$$f(A) = \bigoplus_\alpha f(A_\alpha)$$

holds.

Finally, we study the direct sum of algebras of operators. Let \mathcal{S}_α be a set of densely defined closed operators on \mathcal{H}_α . Put

$$\bigoplus_\alpha \mathcal{S}_\alpha := \{ \bigoplus_\alpha A_\alpha ; A_\alpha \in \mathcal{S}_\alpha \}.$$

Note that each element in $\bigoplus_\alpha \mathcal{S}_\alpha$ is a densely defined closed operator on $\mathcal{H} = \bigoplus_\alpha \mathcal{H}_\alpha$. If each \mathcal{S}_α consists only of bounded operators, we also define

$$\bigoplus_\alpha^b \mathcal{S}_\alpha := \left\{ \bigoplus_\alpha x_\alpha ; x_\alpha \in \mathcal{S}_\alpha, \sup_\alpha \|x_\alpha\| < \infty \right\}.$$

By Lemma A.1, each element in $\bigoplus_\alpha^b \mathcal{S}_\alpha$ is bounded. The following is also well-known.

Lemma A.4. Let \mathfrak{M}_α be a von Neumann algebra acting on \mathcal{H}_α , and put

$$\mathfrak{M} := \bigoplus_\alpha^b \mathfrak{M}_\alpha.$$

Then \mathfrak{M} is von Neumann algebra acting on $\mathcal{H} = \bigoplus_{\alpha} \mathcal{H}_{\alpha}$. The sum, the scalar multiplication, the multiplication and the involution are given by

$$\begin{aligned} (\bigoplus_{\alpha} x_{\alpha}) + (\bigoplus_{\alpha} y_{\alpha}) &= \bigoplus_{\alpha} (x_{\alpha} + y_{\alpha}), \\ \lambda (\bigoplus_{\alpha} x_{\alpha}) &= \bigoplus_{\alpha} (\lambda x_{\alpha}), \quad \text{for all } \lambda \in \mathbb{C}, \\ (\bigoplus_{\alpha} x_{\alpha}) (\bigoplus_{\alpha} y_{\alpha}) &= \bigoplus_{\alpha} (x_{\alpha} y_{\alpha}), \\ (\bigoplus_{\alpha} x_{\alpha})^{*} &= \bigoplus_{\alpha} (x_{\alpha}^{*}). \end{aligned}$$

Furthermore the followings hold.

(1) $\mathfrak{M}' = \bigoplus_{\alpha}^b \mathfrak{M}'_{\alpha}$.

(2) \mathfrak{M} is a finite von Neumann algebra if and only if each \mathfrak{M}_{α} is finite von Neumann algebra.

We call $\bigoplus_{\alpha}^b \mathfrak{M}_{\alpha}$ the direct sum von Neumann algebra of $\{\mathfrak{M}_{\alpha}\}_{\alpha}$.

B Fundamental Results of SRT

Let \mathcal{H} be a Hilbert space. The following lemmata are well-known [18]:

Lemma B.1. Let $\{A_{\lambda}\}_{\lambda \in \Lambda}$ be a net of self-adjoint operators on \mathcal{H} , A be a self-adjoint operator on \mathcal{H} , and \mathcal{D} be a dense subspace of \mathcal{H} which is a core of A and $\mathcal{D} \subset \bigcap_{\lambda \in \Lambda} \text{dom}(A_{\lambda}) \cap \text{dom}(A)$. Suppose for all $\xi \in \mathcal{D}$, $\lim_{\lambda \in \Lambda} A_{\lambda} \xi = A \xi$, then A_{λ} converges to A in the strong resolvent topology.

Lemma B.2. Let $\{A_n\}_{n=1}^{\infty}$ be a sequence of self-adjoint operators on \mathcal{H} , A be a self-adjoint operator on \mathcal{H} . Then A_n converges to A in the strong resolvent topology if and only if e^{itA_n} converges strongly to e^{itA} for all $t \in \mathbb{R}$. In this case, the strong convergence of e^{itA_n} to e^{itA} is uniform on every finite interval of t .

Lemma B.3. Let $\{A_n\}_{n=1}^{\infty}$ be a sequence of self-adjoint operators on \mathcal{H} , A be a self-adjoint operator on \mathcal{H} . Suppose A_n converges to A in the strong resolvent topology, then $E_{A_n}((a, b))$ converges strongly to $E_A((a, b))$ for each $a, b \in \mathbb{R}$ with $a < b$ and $a, b \notin \sigma_p(A)$, where $\sigma_p(A)$ is the set of point spectra of A .

Lemma B.4. Let $\{A_n\}_{n=1}^{\infty}$ be a sequence of self-adjoint operators on \mathcal{H} , A be a self-adjoint operator on \mathcal{H} . Suppose A_n converges to A in the strong resolvent topology, then for all complex valued bounded continuous function f on \mathbb{R} , $f(A_n)$ converges strongly to $f(A)$.

Lemma B.5. Let $\{x_{\lambda}\}_{\lambda \in \Lambda}$ be a net of bounded self-adjoint operators on \mathcal{H} , x be a bounded self-adjoint operator on \mathcal{H} . Suppose that

$$\sup_{\lambda \in \Lambda} \|x_{\lambda}\| < \infty,$$

and x_{λ} converges to x in the strong resolvent topology, then x_{λ} converges strongly to x .

C Tensor Categories

We briefly review the definition of tensor categories. For more details about category theory, see MacLane [9] (we follow the style in Kassel [8], Chapter XI).

Definition C.1. Let $\mathcal{C}, \mathcal{C}'$ be categories, \mathcal{F}, \mathcal{G} be functors from \mathcal{C} to \mathcal{C}' . A *natural transformation* $\theta : \mathcal{F} \rightarrow \mathcal{G}$ is a function which assigns to each object A in \mathcal{C} a morphism $\theta(A) : \mathcal{F}(A) \rightarrow \mathcal{G}(A)$ of \mathcal{C}' in such a way that for every morphism $f : A \rightarrow B$ in \mathcal{C} , the following diagram commutes:

$$\begin{array}{ccc} \mathcal{F}(A) & \xrightarrow{\theta(A)} & \mathcal{G}(A) \\ \mathcal{F}(f) \downarrow & \circlearrowleft & \downarrow \mathcal{G}(f) \\ \mathcal{F}(B) & \xrightarrow{\theta(B)} & \mathcal{G}(B) \end{array}$$

If $\theta(A)$ is an invertible morphism for every A , we call θ a *natural isomorphism*.

Definition C.2. A *tensor category* $(\mathcal{C}, \otimes, I, \alpha, \lambda, \rho)$ is a category \mathcal{C} equipped with

- (1) a bifunctor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ called a *tensor product*²,
- (2) an object I in \mathcal{C} called a *unit object*,
- (3) a natural isomorphism $\alpha : \otimes(\otimes \times 1_{\mathcal{C}})^3 \rightarrow \otimes(1_{\mathcal{C}} \times \otimes)$ called an *associativity constraint*.

(3) means for any objects A, B, C in \mathcal{C} , there is an isomorphism $\alpha_{A,B,C} : (A \otimes B) \otimes C \rightarrow A \otimes (B \otimes C)$ such that the diagram

$$\begin{array}{ccc} (A \otimes B) \otimes C & \xrightarrow{\alpha_{A,B,C}} & A \otimes (B \otimes C) \\ (f \otimes g) \otimes h \downarrow & \circlearrowleft & \downarrow f \otimes (g \otimes h) \\ (A' \otimes B') \otimes C' & \xrightarrow{\alpha_{A',B',C'}} & A' \otimes (B' \otimes C') \end{array}$$

commutes for all morphisms f, g, h in \mathcal{C} .

- (4) a natural isomorphism $\lambda : \otimes(I \times 1_{\mathcal{C}})^4 \rightarrow 1_{\mathcal{C}}$ (resp. $\rho : \otimes(1_{\mathcal{C}} \times I) \rightarrow 1_{\mathcal{C}}$) called a *left* (resp. *right*) *unit constraint* with respect to I .

²This implies $(f' \otimes g') \circ (f \otimes g) = (f' \circ f) \otimes (g' \circ g)$ for all morphisms in \mathcal{C} , and $1_A \otimes 1_B = 1_{A \otimes B}$ for all objects in \mathcal{C} .

³ $\otimes(\otimes \times 1_{\mathcal{C}})$ is the composition of the functors $\otimes \times 1_{\mathcal{C}} : (\mathcal{C} \times \mathcal{C}) \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$ and $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$.

⁴ $I \times 1_{\mathcal{C}}$ is the functor from \mathcal{C} to $\mathcal{C} \times \mathcal{C}$ given by $A \mapsto (I, A)$ for all objects in \mathcal{C} and $f \mapsto (1_I, f)$ for all morphisms in \mathcal{C} .

(4) means for any object A in \mathcal{C} , there is an isomorphism $\lambda_A : I \otimes A \rightarrow A$ (resp. $\rho_A : A \otimes I \rightarrow A$) such that the following two diagrams commute:

$$\begin{array}{ccc} I \otimes A & \xrightarrow{\lambda_A} & A \\ 1_I \otimes f \downarrow & \circlearrowleft & \downarrow f \\ I \otimes A' & \xrightarrow{\lambda_{A'}} & A' \end{array} \quad \begin{array}{ccc} A \otimes I & \xrightarrow{\rho_A} & A \\ f \otimes 1_I \downarrow & \circlearrowleft & \downarrow f \\ A' \otimes I & \xrightarrow{\rho_{A'}} & A' \end{array}$$

for each morphism $f : A \rightarrow A'$ in \mathcal{C} . These functors and natural isomorphisms satisfy the *Pentagon Axiom* and the *Triangle Axiom*. Namely, for all objects A, B, C and D , the following diagrams commute:

$$\begin{array}{ccccc} & & (A \otimes (B \otimes C)) \otimes D & & \\ & \nearrow^{\alpha_{A,B,C} \otimes 1_D} & & \searrow_{\alpha_{A,B \otimes C,D}} & \\ ((A \otimes B) \otimes C) \otimes D & & \circlearrowleft & & A \otimes ((B \otimes C) \otimes D) \\ \alpha_{A \otimes B,C,D} \downarrow & & & & \downarrow 1_A \otimes \alpha_{B,C,D} \\ (A \otimes B) \otimes (C \otimes D) & \xrightarrow{\alpha_{A,B,C \otimes D}} & & \xrightarrow{\alpha_{A,B,C \otimes D}} & A \otimes (B \otimes (C \otimes D)) \end{array}$$

$$\begin{array}{ccc} (A \otimes I) \otimes B & \xrightarrow{\alpha_{A,I,B}} & A \otimes (I \otimes B) \\ \rho_A \otimes 1_B \searrow & \circlearrowleft & \swarrow 1_A \otimes \lambda_B \\ & A \otimes B & \end{array}$$

Definition C.3. Let $(\mathcal{C}, \otimes, I, \alpha, \lambda, \rho)$, $(\mathcal{C}', \otimes, I', \alpha', \lambda', \rho')$ be tensor categories.

- (1) A triple (\mathcal{F}, h_1, h_2) is called a *tensor functor* from \mathcal{C} to \mathcal{C}' if $\mathcal{F} : \mathcal{C} \rightarrow \mathcal{C}'$ is a functor, h_1 is an isomorphism $I' \xrightarrow{\sim} \mathcal{F}(I)$ and h_2 is a natural isomorphism $\otimes(\mathcal{F} \times \mathcal{F}) \xrightarrow{\sim} \mathcal{F} \otimes$, and they satisfy

$$\begin{array}{ccccc} (\mathcal{F}(A) \otimes \mathcal{F}(B)) \otimes \mathcal{F}(C) & \xrightarrow{\alpha_{\mathcal{F}(A), \mathcal{F}(B), \mathcal{F}(C)}} & & \xrightarrow{\alpha_{\mathcal{F}(A), \mathcal{F}(B), \mathcal{F}(C)}} & \mathcal{F}(A) \otimes (\mathcal{F}(B) \otimes \mathcal{F}(C)) \\ h_2(A,B) \otimes 1_{\mathcal{F}(C)} \downarrow & & & & \downarrow 1_{\mathcal{F}(A)} \otimes h_2(B,C) \\ \mathcal{F}(A \otimes B) \otimes \mathcal{F}(C) & & \circlearrowleft & & \mathcal{F}(A) \otimes \mathcal{F}(B \otimes C) \\ h_2(A \otimes B, C) \downarrow & & & & \downarrow h_2(A, B \otimes C) \\ \mathcal{F}((A \otimes B) \otimes C) & \xrightarrow{\mathcal{F}(\alpha_{A,B,C})} & & \xrightarrow{\mathcal{F}(\alpha_{A,B,C})} & \mathcal{F}(A \otimes (B \otimes C)) \end{array}$$

⁵ $\otimes(\mathcal{F} \times \mathcal{F})$ is a functor $\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ which assigns $\mathcal{F}(A) \otimes \mathcal{F}(B)$ for each object (A, B) in $\mathcal{C} \times \mathcal{C}$ and $\mathcal{F}(f) \otimes \mathcal{F}(g)$ for each morphism (f, g) in $\mathcal{C} \times \mathcal{C}$

$$\begin{array}{ccc}
I' \otimes \mathcal{F}(A) & \xrightarrow{\lambda'_{\mathcal{F}(A)}} & \mathcal{F}(A) \\
\downarrow h_1 \otimes 1_{\mathcal{F}(A)} & \circlearrowleft & \uparrow \mathcal{F}(\lambda_A) \\
\mathcal{F}(I) \otimes \mathcal{F}(A) & \xrightarrow{h_2(I,A)} & \mathcal{F}(I \otimes A)
\end{array}
\qquad
\begin{array}{ccc}
\mathcal{F}(A) \otimes I' & \xrightarrow{\rho'_{\mathcal{F}(A)}} & \mathcal{F}(A) \\
\downarrow 1_{\mathcal{F}(A)} \otimes h_1 & \circlearrowleft & \uparrow \mathcal{F}(\rho_A) \\
\mathcal{F}(A) \otimes \mathcal{F}(I) & \xrightarrow{h_2(A,I)} & \mathcal{F}(A \otimes I)
\end{array}$$

for all objects A, B, C in \mathcal{C} .

- (2) A *natural tensor transformation* $\eta : (\mathcal{F}, h_1, h_2) \rightarrow (\mathcal{F}', h'_1, h'_2)$ between tensor functors from \mathcal{C} to \mathcal{C}' is a natural transformation $\mathcal{F} \rightarrow \mathcal{F}'$ such that the following diagrams commute:

$$\begin{array}{ccccc}
& & \mathcal{F}(I) & & \mathcal{F}(A) \otimes \mathcal{F}(B) \xrightarrow{h_2(A,B)} \mathcal{F}(A \otimes B) \\
& \nearrow h_1 & \downarrow \eta(I) & \eta(A) \otimes \eta(B) & \downarrow \eta(A \otimes B) \\
I & \circlearrowleft & & \circlearrowleft & \\
& \searrow h'_1 & \mathcal{F}'(I) & & \mathcal{F}'(A) \otimes \mathcal{F}'(B) \xrightarrow{h'_2(A,B)} \mathcal{F}'(A \otimes B)
\end{array}$$

for all objects A, B in \mathcal{C} . If η is also a natural isomorphism, it is called a *natural tensor isomorphism*.

- (3) A *tensor equivalence* between tensor categories $\mathcal{C}, \mathcal{C}'$ is a tensor functor $\mathcal{F} : \mathcal{C} \rightarrow \mathcal{C}'$ such that there exists a tensor functor $\mathcal{F}' : \mathcal{C}' \rightarrow \mathcal{C}$ and natural tensor isomorphisms $\eta : 1_{\mathcal{C}'} \xrightarrow{\sim} \mathcal{F} \circ \mathcal{F}'$ and $\theta : \mathcal{F}' \circ \mathcal{F} \xrightarrow{\sim} 1_{\mathcal{C}}$. If η and θ can be taken to be identity transformations, then \mathcal{F} is called a *tensor isomorphism* and we say \mathcal{C} is isomorphic to \mathcal{C}' as a tensor category.

Acknowledgement

The authors would like to express their sincere thanks to Professor Asao Arai at Hokkaido University, Professor Izumi Ojima at Kyoto University for the fruitful discussions, insightful comments and encouragements. H.A. also thanks to his colleagues: Mr. Ryo Harada, Mr. Takahiro Hasebe, Mr. Kazuya Okamura and Mr. Hayato Saigo for their useful comments and discussions during the seminar. Y.M. also thanks to Mr. Yutaka Shikano for his professional advice about LaTeX. Finally, the authors thank to Professor Izumi Ojima for his careful proofreading and suggestions again.

References

- [1] S. Albeverio, R. Høegh-Krohn, J. Marion, D. Testard and B. Torresani, Noncommutative Distributions. Unitary Representation of Gauge Groups and Algebras, Monogr. Textbooks Pure Appl. Math., **175**, Marcel Dekker, Inc., New York, 1993.

- [2] A. Arai, Fock Spaces and Quantum Fields, Nippon-Hyouronsha, Tokyo, 2000 (in Japanese).
- [3] A. Banyaga, The Structure of Classical Diffeomorphism Groups, Math. Appl., **400**, Kluwer Academic Publishers, Dordrecht, 1997.
- [4] T. Fack and H. Kosaki, Generalized s -numbers of τ -measurable operators, *Pacific J. Math.*, **123** (1986), 260–300.
- [5] M. Gordina, Hilbert-Schmidt groups as infinite-dimensional Lie groups and their Riemannian geometry, *J. Funct. Anal.* **227** (2005), 245–272.
- [6] K. Hofmann and S. Morris, The Lie Theory of Connected Pro-Lie Groups, Europ. Math. Soc. Publ. House, 2007.
- [7] K. Hofmann and K.-H. Neeb, Pro-Lie groups which are infinite-dimensional Lie groups, *Math. Proc. Cambridge Philos. Soc.* **146** (2009), 351–378.
- [8] C. Kassel, Quantum Groups, Grad. Texts in Math., **155**, Springer-Verlag, New York, 1995.
- [9] S. MacLane, Categories for the Working Mathematician, Grad. Texts in Math., **5**, Springer-Verlag, New York, 1998.
- [10] J. Milnor, Remarks on infinite-dimensional Lie groups, In: B. DeWitt and R. Stora (Eds.), *Relativity, Groups and Topology II* (Les Houches, 1983), North-Holland, Amsterdam, 1984, 1007–1057.
- [11] F. Murray and J. von Neumann, On Rings of Operators, *Ann. Math.* **37** (1936), 116–229.
- [12] K.-H. Neeb, Towards a Lie theory of locally convex groups, *Jpn. J. Math.*, **1** (2006), 291–468.
- [13] E. Nelson, *Topics in Dynamics I*, Princeton University Press, Princeton, 1969.
- [14] E. Nelson, Notes on non-commutative integration, *J. Funct. Anal.* **15** (1974), 103–116.
- [15] H. Omori, Infinite-Dimensional Lie Groups, Transl. Math. Monogr., **158**, Amer. Math. Soc., 1997.
- [16] H. Omori, On Banach-Lie groups acting on finite dimensional manifolds, *Tôhoku Math. J.* **30** (1978), 223–250.
- [17] A. Pressley and G. Segal, *Loop Groups*, Clarendon Press, Oxford, 1986.
- [18] M. Reed and B. Simon, *Methods of Modern Mathematical Physics I*, Academic Press, New York, 1972.

- [19] K. Schmüdgen, *Unbounded Operator Algebras and Representation Theory*. Birkhäuser Verlag, Basel, 1990.
- [20] W. Stinespring, Integration theorems for gages and duality for unimodular groups. *Trans. Amer. Math. Soc.* **90** (1959), 15–56.
- [21] M. Takesaki, *Theory of Operator Algebras I*, Springer-Verlag, Berlin, 1979.