

RIMS-1703

**Affine nil-Hecke algebras and braided  
differential structure on affine Weyl groups**

By

Anatol N. KIRILLOV and Toshiaki MAENO

August 2010



**京都大学 数理解析研究所**

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

# Affine nil-Hecke algebras and braided differential structure on affine Weyl groups

Anatol N. Kirillov and Toshiaki Maeno

## Abstract

We construct a model of the affine nil-Hecke algebra as a subalgebra of the Nichols-Woronowicz algebra associated to a Yetter-Drinfeld module over the affine Weyl group. We also discuss the Peterson isomorphism between the homology of the affine Grassmannian and the small quantum cohomology ring of the flag variety in terms of the braided differential calculus.

## Introduction

The cohomology ring of the flag variety is a fundamental object of research in the study of the Schubert calculus. Fomin and the first author [4] gave a combinatorial model of the cohomology  $H^*(Fl_n)$  ring of the flag variety of type  $A$  as a commutative subalgebra of a quadratic algebra  $\mathcal{E}_n$ . It is remarkable that the algebra  $\mathcal{E}_n$  has a natural quantum deformation  $\mathcal{E}_n^q$  so that  $\mathcal{E}_n^q$  contains the quantum cohomology ring  $QH^*(Fl_n)$  as a commutative subalgebra.

It has been observed by Milinski and Schneider [11] and by Majid [10] that the defining relations of the Fomin-Kirillov quadratic algebra  $\mathcal{E}_n$  are understandable from the viewpoint of a certain kind of braided Hopf algebra called the Nichols-Woronowicz algebra. Bazlov [2] constructed the model of the coinvariant algebra of the finite Coxeter groups as a commutative subalgebra of the Nichols-Woronowicz algebra. At the same time, the nil-Coxeter algebra, which is dual to the coinvariant algebra, is also realized as a subalgebra of the Nichols-Woronowicz algebra.

The braided analogue of the symmetric or exterior algebra was introduced by Woronowicz [14] for the study of the differential forms on the quantum

groups. For a given braided vector space  $M$  over a field  $K$  of characteristic zero, the braided analogue  $\mathcal{B}(M)$  of the symmetric algebra of  $M$  is defined to be the quotient of the free tensor algebra of  $M$  by the kernel of the braided symmetrizer. It is known that the algebra  $\mathcal{B}(M)$  is a braided graded Hopf algebra characterized by the following conditions:

- (1)  $\mathcal{B}^0(M) = K$ ,
- (2)  $\mathcal{B}^1(M) = M = \{\text{primitive elements in } \mathcal{B}(M)\}$ ,
- (3)  $\mathcal{B}^1(M)$  generates  $\mathcal{B}(M)$  as an algebra.

The Hopf algebra characterized by the above conditions has been studied by Nichols [12] and named the Nichols algebra by Andruskiewitsch and Schneider [1]. The study of the algebra  $\mathcal{B}(M)$  from the viewpoint of the free braided differential calculus was developed by [9]. In this paper we will call  $\mathcal{B}(M)$  the Nichols-Woronowicz algebra simply following [2].

The aim of this paper is to construct the nil-Hecke algebra as a subalgebra of an extension of the Nichols-Woronowicz algebra  $\mathcal{B}_{\text{aff}}$  associated to a Yetter-Drinfeld module over the affine Weyl groups. Our construction is analogous to the one in [2, Section 6].

It is known that the affine Grassmannian  $\widehat{\text{Gr}} := G(\mathbb{C}((t)))/G(\mathbb{C}[[t]])$  of a semisimple Lie group  $G$  is homotopic to the loop group  $\Omega K$  of the maximal compact subgroup  $K \subset G$ . The homology  $H_*(\widehat{\text{Gr}}) \cong H_*(\Omega K)$  carries an associative algebra structure induced by the Pontryagin product. The structure of the Pontryagin ring  $H_*(\Omega K)$  has been determined by Bott [3]. The Schubert calculus for Kac-Moody flag varieties was studied by Kostant and Kumar [6] by using the nil-Hecke algebra. Peterson [13] stated that the torus-equivariant homology  $H_*^T(\widehat{\text{Gr}})$  of the affine Grassmannian is isomorphic to the so-called Peterson subalgebra of the affine nil-Hecke algebra. So our construction gives a model of  $H_*^T(\widehat{\text{Gr}})$  as a commutative subalgebra of the Nichols-Woronowicz algebra  $\mathcal{B}_{\text{aff}}(S)$ , see Theorem 3.1.

Peterson [13] also pointed out that the Pontryagin ring  $H_*^T(\widehat{\text{Gr}})$  is isomorphic to the small quantum cohomology ring  $QH_T^*(G/B)$  of the corresponding flag variety  $G/B$  as an algebra after a suitable localization. The affine Bruhat operator acting on  $H_*^T(\widehat{\text{Gr}})$  introduced by Lam and Shimozono [7] gives an explicit comparison between the multiplicative structure of  $H_*^T(\widehat{\text{Gr}})$  and that of  $QH_T^*(G/B)$ . In this paper, we will realize the affine Bruhat operator as a braided differential operator acting on our algebra  $\mathcal{B}_{\text{aff}}$ .

## Acknowledgement.

The second author is supported by Grant-in-Aid for Scientific Research.

## 1 Affine nil-Hecke algebra

Let  $G$  be a simply-connected semisimple complex Lie group and  $W$  its Weyl group. Denote by  $\Delta$  the set of the roots. We fix the set  $\Delta_+$  of the positive roots by choosing a set of simple roots  $\alpha_1, \dots, \alpha_r$ . The Weyl group  $W$  acts on the weight lattice  $P$  and the coroot lattice  $Q^\vee$  of  $G$ . The affine Weyl group  $W_{\text{aff}}$  is generated by the affine reflections  $s_{\alpha, k}$ ,  $\alpha \in \Delta$ ,  $k \in \mathbb{Z}$ , with respect to the affine hyperplanes  $H_{\alpha, k} := \{\lambda \in P \otimes \mathbb{R} \mid \langle \lambda, \alpha^\vee \rangle = k\}$ . The affine Weyl group is the semidirect product of  $W$  and  $Q^\vee$ , i.e.,  $W_{\text{aff}} = W \ltimes Q^\vee$ . The affine Weyl group  $W_{\text{aff}}$  is generated by the simple reflections  $s_1 := s_{\alpha_1, 0}, \dots, s_r := s_{\alpha_r, 0}$  and  $s_0 := s_{\theta, 1}$  where  $\theta = -\alpha_0$  is the highest root. The affine Weyl group  $W$  has the presentation as a Coxeter group as follows:

$$W_{\text{aff}} = \langle s_0, \dots, s_r \mid s_0^2 = \dots = s_r^2 = 1, (s_i s_j)^{m_{ij}} = 1 \rangle.$$

**Definition 1.1.** The affine nil-Coxeter algebra  $\mathbb{A}_0$  is the associative algebra generated by  $\tau_0, \dots, \tau_r$  subject to the relations

$$\tau_0^2 = \dots = \tau_r^2 = 0, \quad (\tau_i \tau_j)^{[m_{ij}/2]} \tau_i^{\nu_{ij}} = (\tau_j \tau_i)^{[m_{ij}/2]} \tau_j^{\nu_{ij}},$$

where  $\nu_{ij} := m_{ij} - 2[m_{ij}/2]$ .

For a reduced expression  $x = s_{i_1} \cdots s_{i_l}$  of an element  $x \in W_{\text{aff}}$ , the element  $\tau_x := \tau_{i_1} \cdots \tau_{i_l} \in \mathbb{A}_0$  is independent of the choice of the reduced expression of  $x$ . It is known that  $\{\tau_x\}_{x \in W_{\text{aff}}}$  form a linear basis of  $\mathbb{A}_0$ .

The nil-Coxeter algebra  $\mathbb{A}_0$  acts on  $S := \text{Sym} P_{\mathbb{Q}}$  via

$$\tau_0(f) := \partial_{\alpha_0}(f) = -(f - s_{\theta, 0}f)/\theta,$$

$$\tau_i(f) := \partial_{\alpha_i}(f) = (f - s_{\alpha_i, 0}f)/\alpha_i, \quad i = 1, \dots, r,$$

for  $f \in S$ .

**Definition 1.2.** ([6]) The nil-Hecke algebra  $\mathbb{A}$  is defined to be the cross product  $\mathbb{A}_0 \ltimes S$ , where the cross relation is given by

$$\tau_i f = \partial_{\alpha_i}(f) + s_i(f) \tau_i \quad f \in S, i = 1, \dots, r.$$

Here, we summarize some known results on the homology of the affine Grassmannian. The affine Grassmannian  $\widehat{\text{Gr}} := G(\mathbb{C}((t)))/G(\mathbb{C}[[t]])$  is homotopic to the loop group  $\Omega K$  of the maximal compact subgroup  $K \subset G$ . Let  $T \subset G$  be the maximal torus. An associative algebra structure on the  $T$ -equivariant homology group  $H_*^T(\widehat{\text{Gr}}) \cong H_*^T(\Omega K)$  is induced from the group multiplication

$$\Omega K \times \Omega K \rightarrow \Omega K.$$

It is known that the algebra  $H_*^T(\widehat{\text{Gr}})$  is commutative. The algebra  $H_*^T(\Omega K)$  is called the Pontryagin ring.

We regard the  $T$ -equivariant homology  $H_*^T(\widehat{\text{Gr}})$  as an  $S$ -algebra by identifying  $S = H_T^*(pt)$ . The diagonal embedding

$$\Omega K \rightarrow \Omega K \times \Omega K$$

induces a coproduct on  $H_*^T(\widehat{\text{Gr}})$ .

**Proposition 1.1.** ([13]) *The  $T$ -equivariant homology  $H_*^T(\widehat{\text{Gr}})$  is isomorphic to the centralizer  $Z_{\mathbb{A}}(S)$  of  $S$  in  $\mathbb{A}$  as Hopf algebras.*

## 2 Nichols-Woronowicz algebra for affine Weyl groups

We briefly recall the construction of the Nichols-Woronowicz algebra associated to a braided vector space. Let  $M$  be a vector space over a field of characteristic zero and  $\psi : M^{\otimes 2} \rightarrow M^{\otimes 2}$  be a fixed linear endomorphism satisfying the braid relations  $\psi_i \psi_{i+1} \psi_i = \psi_{i+1} \psi_i \psi_{i+1}$  where  $\psi_i : M^{\otimes n} \rightarrow M^{\otimes n}$  is a linear endomorphism obtained by applying  $\psi$  to the  $i$ -th and  $(i+1)$ -st components. Denote by  $s_i$  the simple transposition  $(i, i+1) \in S_n$ . For any reduced expression  $w = s_{i_1} \cdots s_{i_l} \in S_n$ , the endomorphism  $\Psi_w = \psi_{i_1} \cdots \psi_{i_l} : M^{\otimes n} \rightarrow M^{\otimes n}$  is well-defined. The Woronowicz symmetrizer [14] is given by  $\sigma_n := \sum_{w \in S_n} \Psi_w$ .

**Definition 2.1.** ([14]) The Nichols-Woronowicz algebra associated to a braided vector space  $M$  is defined by

$$\mathcal{B}(M) := \bigoplus_{n \geq 0} M^{\otimes n} / \text{Ker}(\sigma_n),$$

where  $\sigma_n : M^{\otimes n} \rightarrow M^{\otimes n}$  is the Woronowicz symmetrizer.

**Definition 2.2.** A vector space  $M$  is called a Yetter-Drinfeld module over a group  $\Gamma$ , if the following conditions are satisfied:

- (1)  $M$  is a  $\Gamma$ -module,
- (2)  $M$  is  $\Gamma$ -graded, i.e.  $M = \bigoplus_{g \in \Gamma} M_g$ , where  $M_g$  is a linear subspace of  $M$ ,
- (3) for  $h \in \Gamma$  and  $v \in M_g$ ,  $h(v) \in M_{hgh^{-1}}$ .

The Yetter-Drinfeld module  $M$  over a group  $\Gamma$  is naturally braided with the braiding  $\psi : M^{\otimes 2} \rightarrow M^{\otimes 2}$  defined by  $\psi(a \otimes b) = g(b) \otimes a$  for  $a \in M_g$  and  $b \in M$ .

In the following we are interested in the Yetter-Drinfeld module over the affine Weyl groups  $W_{\text{aff}}$ . Denote by  $t_\lambda \in W_{\text{aff}}$  the translation by  $\lambda \in Q^\vee$ . We define a Yetter-Drinfeld module  $V_{\text{aff}}$  over  $W_{\text{aff}}$  by

$$V_{\text{aff}} := \bigoplus_{\alpha \in \Delta, k \in \mathbb{Z}} \mathbb{Q} \cdot [\alpha, k] / ([\alpha, k] + [-\alpha, -k]),$$

where the  $W_{\text{aff}}$  acts on  $V_{\text{aff}}$  by

$$w[\alpha, k] := [w(\alpha), k], \quad w \in W, \quad t_\lambda[\alpha, k] := [\alpha, k + (\alpha, \lambda)], \quad \lambda \in Q^\vee.$$

The  $W_{\text{aff}}$ -grading is given by  $\deg_{W_{\text{aff}}}([\alpha, k]) := s_{\alpha, k}$ . Then it is easy to check the conditions in Definition 2.1. Now we have the Nichols-Woronowicz algebra  $\mathcal{B}_{\text{aff}} := \mathcal{B}(V_{\text{aff}})$  associated to the Yetter-Drinfeld module  $V_{\text{aff}}$ .

Let  $\mathcal{B}_W$  be the Nichols-Woronowicz algebra associated to the Yetter-Drinfeld module  $V = \bigoplus_{\alpha \in \Delta} \mathbb{Q} \cdot [\alpha] / ([\alpha] + [-\alpha])$  as in [2, Section 4].

**Lemma 2.1.** (1) We have a surjective homomorphism  $\pi : \mathcal{B}_{\text{aff}} \rightarrow \mathcal{B}_W$ ,  $\pi([\alpha, k]) := [\alpha]$ .

(2) The algebra  $\mathcal{B}_{\text{aff}}$  acts on  $S$  via  $[\alpha, k]f = \partial_\alpha(f)$  for all  $k \in \mathbb{Z}$ .

*Proof.* (1) Denote by  $\psi$  and  $\bar{\psi}$  the braidings on  $V_{\text{aff}}$  and  $V$  respectively. Let  $\tilde{\pi} : \bigoplus_n V_{\text{aff}}^{\otimes n} \rightarrow \bigoplus_n V^{\otimes n}$  be the lift of  $\pi$ . Since

$$\psi([\alpha, k] \otimes [\beta, l]) = [s_\alpha(\beta), l - \langle \alpha^\vee, \beta \rangle k] \otimes [\alpha, k]$$

and  $\bar{\psi}([\alpha] \otimes [\beta]) = [s_\alpha(\beta)] \otimes [\alpha]$ , the map  $\tilde{\pi}$  sends the kernel of the braided symmetrizer  $\sigma_n$  of  $V_{\text{aff}}^{\otimes n}$  to that of  $V^{\otimes n}$ .

(2) In [2], it is shown that the algebra  $\mathcal{B}_W$  acts on the coinvariant algebra  $S_W$  via  $[\alpha] \mapsto \partial_\alpha$ . Let  $S^W$  be the  $W$ -invariant subalgebra of  $S$ . Then we have the decomposition  $S = S^W \otimes S_W$ . The operator  $\partial_\alpha$  extends  $S^W$ -linearly to

the operator on  $S$ . Hence  $\mathcal{B}_W$  acts on  $S$ . We have seen the existence of the natural projection  $\pi$  from  $\mathcal{B}_{\text{aff}}$  to  $\mathcal{B}$ , so  $\pi$  induces the action of  $\mathcal{B}_{\text{aff}}$  on  $S$ .

Let us define the extension  $\mathcal{B}_{\text{aff}}(S) = \mathcal{B}_{\text{aff}} \ltimes S$  by the cross relation

$$[\alpha, k]f = \partial_\alpha f + s_{\alpha,0}(f)[\alpha, k], \quad [\alpha, k] \in V_{\text{aff}}, f \in S.$$

**Proposition 2.1.** *There exists a homomorphism  $\varphi : \mathbb{A} \rightarrow \mathcal{B}_{\text{aff}}(S)$  given by  $\tau_0 \mapsto [\alpha_0, -1]$ ,  $\tau_i \mapsto [\alpha_i, 0]$ ,  $i = 1, \dots, r$ , and  $f \mapsto f$ ,  $f \in S$ .*

*Proof.* It is enough to check the Coxeter relations among  $\varphi(\tau_0), \dots, \varphi(\tau_r)$  in  $\mathcal{B}_{\text{aff}}(S)$  based on the classification of the affine root systems. This is done by the direct computation of the symmetrizer for the subsystems of rank 2 in the similar manner to [2, Section 6].

**Example 2.1.** Here we list the Coxeter relations in  $\mathcal{B}_{\text{aff}}$  involving  $[\theta, 1] = -[\alpha_0, -1]$  for the root systems of rank 2. Let  $(\varepsilon_1, \dots, \varepsilon_r)$  be an orthonormal basis of the  $r$ -dimensional Euclidean space. Put  $[ij, k] := [\varepsilon_i - \varepsilon_j, k]$ ,  $[\overline{ij}, k] := [\varepsilon_i + \varepsilon_j, k]$ ,  $[i, k] := [\varepsilon_i, k]$  and  $[\alpha] := [\alpha, 0]$ .

(i) (Type  $A_2$  case)

$$[13, 1][23][13, 1] + [23][13, 1][23] = 0, \quad [13, 1][12][13, 1] + [12][13, 1][12] = 0$$

(ii) (Type  $B_2$  case)

$$[\overline{12}, 1][2][\overline{12}, 1][2] = [2][\overline{12}, 1][2][\overline{12}, 1]$$

(iii) (Type  $G_2$  case) Let  $\alpha_1, \alpha_2$  be the simple roots for  $G_2$ -system. We assume that  $\alpha_1$  is a short root and  $\alpha_2$  is a long one. Then we have  $\theta = 3\alpha_1 + 2\alpha_2$ .

$$[\theta, 1][\alpha_2][\theta, 1] + [\alpha_2][\theta, 1][\alpha_2] = 0.$$

### 3 Model of nil-Hecke algebra

The connected components of  $P \otimes \mathbb{R} \setminus \cup_{\alpha \in \Delta_+, k \in \mathbb{Z}} H_{\alpha, k}$  are called alcoves. The affine Weyl group  $W_{\text{aff}}$  acts on the set of the alcoves simply and transitively.

**Definition 3.1.** ([8]) (1) A sequence  $(A_0, \dots, A_l)$  of alcoves  $A_i$  is called an alcove path if  $A_i$  and  $A_{i+1}$  have a common wall and  $A_i \neq A_{i+1}$ .

(2) An alcove path  $(A_0, \dots, A_l)$  is called reduced if the length  $l$  of the path is minimal among all alcove paths connecting  $A_0$  and  $A_l$ .

(3) We use the symbol  $A_i \xrightarrow{\beta, k} A_{i+1}$  when  $A_i$  and  $A_{i+1}$  have a common wall of the form  $H_{\beta, k}$  and the direction of the root  $\beta$  is from  $A_i$  to  $A_{i+1}$ .

The alcove  $A^\circ$  defined by the inequalities  $\langle \lambda, \alpha_0^\vee \rangle \geq -1$  and  $\langle \lambda, \alpha_i^\vee \rangle \geq 0$ ,  $i = 1, \dots, r$ , is called the fundamental alcove. For a reduced alcove path  $\gamma : A_0 = A^\circ \xrightarrow{\beta_1, k_1} \dots \xrightarrow{\beta_l, k_l} A_l$ , we define an element  $[\gamma] \in \mathcal{B}_{\text{aff}}$  by

$$[\gamma] := [-\beta_1, -k_1] \cdots [-\beta_l, -k_l].$$

When  $A_l = x^{-1}(A^\circ)$  for  $x \in W_{\text{aff}}$ , we will also use the symbol  $[x]$  instead of  $[\gamma]$ , since  $[\gamma]$  depends only on  $x$  thanks to the Yang-Baxter relation.

For a braided vector space  $M$ , it is known that an element  $a \in M$  acts on  $\mathcal{B}(M^*)$  as a braided differential operator (see [2], [9]). Let us identify  $M^*$  with  $M$  via the  $W_{\text{aff}}$ -invariant inner product  $(\ , \ )$  given by

$$([\alpha, k], [\beta, l]) = \begin{cases} 1, & \text{if } \alpha = \beta \text{ and } k = l, \\ 0, & \text{otherwise,} \end{cases}$$

for  $\alpha, \beta \in \Delta_+$ ,  $k, l \in \mathbb{Z}$ . In our case, the differential operator  $\overleftarrow{D}_{[\alpha, k]}$ ,  $[\alpha, k] \in V_{\text{aff}}$ , acting from the right is determined by the following characterization:

- (0)  $(c) \overleftarrow{D}_{[\alpha, k]} = 0$ ,  $c \in \mathbb{Q}$ ,
- (1)  $([\alpha, k]) \overleftarrow{D}_{[\beta, l]} = ([\alpha, k], [\beta, l])$ ,
- (2)  $(FG) \overleftarrow{D}_{[\alpha, k]} = F(G \overleftarrow{D}_{[\alpha, k]}) + (F \overleftarrow{D}_{[\alpha, k]}) s_{\alpha, k}(G)$ ,

for  $\alpha, \beta \in \Delta$ ,  $k, l \in \mathbb{Z}$ ,  $F, G \in \mathcal{B}_{\text{aff}}$ . The operator  $\overleftarrow{D}_{[\alpha, k]}$  extends to the one acting on  $\mathcal{B}_{\text{aff}}(S)$  by the commutation relation  $f \cdot \overleftarrow{D}_{[\alpha, k]} = \overleftarrow{D}_{[\alpha, k]} \cdot s_{\alpha, k}(f)$ ,  $f \in S$ .

We use the abbreviation  $\overleftarrow{D}_0 := \overleftarrow{D}_{[\alpha_0, -1]}$ ,  $\overleftarrow{D}_i := \overleftarrow{D}_{[\alpha_i, 0]}$ ,  $i = 1, \dots, r$ . For  $x \in W_{\text{aff}}$ , fix a reduced decomposition  $x = s_{i_1} \cdots s_{i_l}$ . We define the corresponding braided differential operator  $\overleftarrow{D}_x$  acting on  $\mathcal{B}_{\text{aff}}$  by the formula

$$\overleftarrow{D}_x := \overleftarrow{D}_{i_l} \cdots \overleftarrow{D}_{i_1},$$

which is also independent of the choice of the reduced decomposition of  $x$  because of the braid relations.

**Lemma 3.1.** *For  $x \in W_{\text{aff}}$ , take a reduced alcove path  $\gamma$  from the fundamental alcove  $A^\circ$  to  $x^{-1}(A^\circ)$ . Then, we have  $([\gamma]) \overleftarrow{D}_x = 1$ .*

*Proof.* Let us take a reduced path

$$\gamma : A_0 = A^\circ \xrightarrow{\beta_1, k_1} A_1 \xrightarrow{\beta_2, k_2} \dots \xrightarrow{\beta_l, k_l} A_l = x^{-1}(A^\circ).$$

Define a sequence  $\sigma_1, \dots, \sigma_l \in W_{\text{aff}}$  inductively by

$$\sigma_1 := s_{\beta_1, k_1}, \quad \sigma_{j+1} := \sigma_j s_{\beta_{j+1}, k_{j+1}} \sigma_j.$$

Then it is easy to see that  $\sigma_\nu(A_j) \neq A^\circ$ ,  $1 \leq \nu \leq j-1$ ,  $\sigma_j(A_j) = A^\circ$  and the walls  $\sigma_j(H_{\beta_{j+1}, k_{j+1}})$  are corresponding to simple roots. Hence,  $\sigma_1, \dots, \sigma_l$  are simple reflections. This sequence gives a reduced expression  $x = \sigma_l \cdots \sigma_1$ . Put  $\sigma_i = s_{\alpha_{i_j}}$ . Since the direction of  $\beta_{j+1}$  is chosen to be from  $A_j$  to  $A_{j+1}$ , we have

$$[\gamma] \overleftarrow{D}_x = ([\beta_1, k_1]) \overleftarrow{D}_{i_1} \cdot (\sigma_1([\beta_2, k_2])) \overleftarrow{D}_{i_2} \cdots (\sigma_{l-1}([\beta_l, k_l])) \overleftarrow{D}_{i_l} = 1.$$

**Example 3.1.** ( $A_2$ -case) The standard realization is given by  $\alpha_1 = \varepsilon_1 - \varepsilon_2$ ,  $\alpha_2 = \varepsilon_2 - \varepsilon_3$ ,  $\alpha_0 = \varepsilon_3 - \varepsilon_1$ . Consider the translation  $t_{\alpha_1}$  by the simple root  $\alpha_1$ . If we take a reduced path

$$\gamma : A_0 = A^\circ \xrightarrow{-\alpha_2, 0} A_1 \xrightarrow{\alpha_1, 1} A_2 \xrightarrow{-\alpha_0, 1} A_3 \xrightarrow{\alpha_1, 2} A_4 = t_{\alpha_1}(A^\circ),$$

then we have  $[\gamma] = [23][21, -1][31, -1][21, -2]$ . On the other hand, the differential operator corresponding to  $t_{-\alpha_1}$  is given by  $\overleftarrow{D}_2 \overleftarrow{D}_0 \overleftarrow{D}_2 \overleftarrow{D}_1$ , where  $\overleftarrow{D}_0 = \overleftarrow{D}_{[31, -1]}$ ,  $\overleftarrow{D}_1 = \overleftarrow{D}_{[12]}$ ,  $\overleftarrow{D}_2 = \overleftarrow{D}_{[23]}$ . It is easy to check by direct computation

$$([23][21, -1][31, -1][12, 2]) \overleftarrow{D}_2 \overleftarrow{D}_0 \overleftarrow{D}_2 \overleftarrow{D}_1 = 1.$$

**Theorem 3.1.** *The algebra homomorphism  $\varphi : \mathbb{A} \rightarrow \mathcal{B}_{\text{aff}}(S)$  is injective.*

*Proof.* The nil-Hecke algebra  $\mathbb{A}$  is also  $W_{\text{aff}}$ -graded. Since the homomorphism  $\varphi : \mathbb{A} \rightarrow \mathcal{B}_{\text{aff}}(S)$  preserves the  $W_{\text{aff}}$ -grading, it is enough to check  $\varphi(\tau_x) \neq 0$ , for  $x \in W_{\text{aff}}$  in order to show the injectivity of  $\varphi$ . On the other hand,  $\mathcal{B}_{\text{aff}}^{\text{op}}$  acts on  $\mathcal{B}_{\text{aff}}$  itself via the braded differential operators. Let  $\gamma$  be a reduced alcove path from  $A^\circ$  to  $x^{-1}(A^\circ)$ . Then we have  $([\gamma]) \overleftarrow{D}_x = 1$  from Lemma 3.1. This shows  $\overleftarrow{D}_x \neq 0$ , so  $\varphi(\tau_x) \neq 0$ .

This theorem implies the following (see Proposition 1.1):

**Corollary 3.1.** *The  $T$ -equivariant Pontryagin ring  $H_*^T(\widehat{\text{Gr}})$  is a subalgebra of  $\mathcal{B}_{\text{aff}}(S)$ .*

By taking the non-equivariant limit, we also have:

**Corollary 3.2.** *The Pontryagin ring  $H_*(\widehat{\text{Gr}})$  is a subalgebra of  $\mathcal{B}_{\text{aff}}$ .*

## 4 Affine Bruhat operators

We denote by  $x \rightarrow y$  the cover relation in the Bruhat ordering of  $W_{\text{aff}}$ , i.e.  $y = xs_{\alpha,k}$  for some  $\alpha \in \Delta$  and  $k \in \mathbb{Z}$ , and  $l(y) = l(x) + 1$ .

We will use some terminology from [7]. Denote by  $\tilde{Q}$  the set of antidominant elements in  $Q^\vee$ . An element  $x \in W_{\text{aff}}$  can be expressed uniquely as a product of form  $x = wt_{v\lambda} \in W_{\text{aff}}$  with  $v, w \in W$ ,  $\lambda \in \tilde{Q}$ . We say that  $x = wt_{v\lambda}$  belongs to the " $v$ -chamber". An element  $\lambda \in \tilde{Q}$  is called superregular when  $|\langle \lambda, \alpha \rangle| > 2(\#W) + 2$  for all  $\alpha \in \Delta_+$ . If  $\lambda \in \tilde{Q}$  is superregular, then  $x = wt_{v\lambda}$  is called superregular. The subset of superregular elements in  $W_{\text{aff}}$  is denoted by  $W_{\text{aff}}^{\text{sreg}}$ . We say that a property holds for sufficiently superregular elements  $W_{\text{aff}}^{\text{ssreg}} \subset W_{\text{aff}}$  if there is a positive constant  $k \in \mathbb{Z}$  such that the property holds for all  $x \in W_{\text{aff}}^{\text{sreg}}$  satisfying the following condition:

$$y \in W_{\text{aff}}, y < x, \text{ and } l(x) - l(y) < k \Rightarrow y \in W_{\text{aff}}^{\text{sreg}}.$$

The meaning of  $W_{\text{aff}}^{\text{ssreg}}$  depends on the context, see [7, Section 4] for the details. For  $v \in W$ , consider the  $S$ -submodule  $M_v^{\text{ssreg}}$  in  $\mathcal{B}_{\text{aff}}$  generated by the sufficiently superregular elements  $[x]$  where  $x$  belongs to the  $v$ -chamber.

**Lemma 4.1.** *Let  $x \in W_{\text{aff}}$ . For  $\alpha \in \Delta$  and  $k \in \mathbb{Z}_{>0}$ , we have*

$$[x] \overleftarrow{D}_{[\alpha,k]} = \begin{cases} [xs_{\alpha,k}], & \text{if } l(x) = l(xs_{\alpha,k}) + 1, \\ 0, & \text{otherwise.} \end{cases}$$

*Proof.* The fundamental alcove  $A^\circ$  is contained in the region  $\{\lambda \in P \otimes \mathbb{R} | \langle \lambda, \alpha^\vee \rangle < k\}$  for  $\alpha \in \Delta$  and  $k \in \mathbb{Z}_{>0}$ . Let us choose any reduced path  $\gamma : A_0 \xrightarrow{\beta_1, k_1} \dots \xrightarrow{\beta_l, k_l} A_l = x^{-1}(A^\circ)$  with  $k_i \geq 0$ . If  $l(x) > l(xs_{\alpha,k})$ , then  $(\beta_i, k_i) = (\alpha, k)$  for some  $i$ . Take the largest  $i$  and consider the path

$$\gamma' : A_0 \xrightarrow{\beta_1, k_1} \dots \xrightarrow{\beta_{i-1}, k_{i-1}} A_{i-1} \xrightarrow{\beta'_{i+1}, k'_{i+1}} s_{\alpha,k}(A_{i+1}) \xrightarrow{\beta'_{i+2}, k'_{i+2}} \dots$$

$$\dots \xrightarrow{\beta'_l, k'_l} s_{\alpha,k}(A_l) = s_{\alpha,k}x^{-1}(A^\circ) = (xs_{\alpha,k})^{-1}(A^\circ),$$

where  $(\beta'_j, k'_j)$  is determined by the condition  $s_{\alpha,k}(H_{\beta_j, k_j}) = H_{\beta'_j, k'_j}$ . If  $l(x) = l(xs_{\alpha,k}) + 1$ , then the path  $\gamma'$  is a reduced path. In this case, we have  $[x] \overleftarrow{D}_{[\alpha,k]} = [xs_{\alpha,k}]$ . If  $l(x) > l(xs_{\alpha,k}) + 1$ , the above path  $\gamma'$  is not reduced and  $[x] \overleftarrow{D}_{[\alpha,k]} = 0$ . When  $l(x) < l(xs_{\alpha,k})$ , the element  $[\alpha, k]$  does not appear in the monomial  $[\gamma]$ , so we have  $[x] \overleftarrow{D}_{[\alpha,k]} = 0$ .

**Proposition 4.1.** ([7, Proposition 4.1]) *Let  $\lambda \in \tilde{Q}$  be superregular. For  $x = wt_{v\lambda}$  and  $y = xs_{v\alpha, -n}$  with  $v, w \in W$ , we have the cover relation  $y \rightarrow x$  if and only if one of the following conditions holds:*

- (1)  $l(wv) = l(wvs_\alpha) - 1$  and  $n = \langle \lambda, \alpha \rangle$ , giving  $y = ws_{v(\alpha)}t_{v(\lambda)}$ ,
- (2)  $l(wv) = l(wvs_\alpha) + \langle \alpha^\vee, 2\rho \rangle - 1$  and  $n = \langle \lambda, \alpha \rangle + 1$ , giving  $y = ws_{v(\alpha)}t_{v(\lambda + \alpha^\vee)}$ ,
- (3)  $l(v) = l(vs_\alpha) + 1$  and  $n = 0$ , giving  $y = ws_{v(\alpha)}t_{vs_\alpha(\lambda)}$ ,
- (4)  $l(v) = l(vs_\alpha) - \langle \alpha^\vee, 2\rho \rangle + 1$  and  $n = -1$ , giving  $y = ws_{v(\alpha)}t_{vs_\alpha(\lambda + \alpha^\vee)}$ .

In [7], the first kind of the conditions (1) and (2) are called the near relation because  $x$  and  $y$  belong to the same chamber. In this paper we denote the near relation by  $y \rightarrow_{\text{near}} x$ .

The affine Bruhat operator  $B^\mu : S\langle W_{\text{aff}}^{\text{ssreg}} \rangle \rightarrow S\langle W_{\text{aff}}^{\text{sreg}} \rangle$ ,  $\mu \in P$ , due to Lam and Shimozono [7, Section 5] is an  $S$ -linear map defined by the formula

$$B^\mu(x) = (\mu - wv\mu)x + \sum_{\alpha \in \Delta_+} \sum_{xs_{v(\alpha), k} \rightarrow_{\text{near}} x} \langle \alpha^\vee, \mu \rangle xs_{v(\alpha), k}$$

for  $x = wt_{v\lambda} \in W_{\text{aff}}^{\text{ssreg}}$ . We also introduce the operator  $\beta_v^\mu$ ,  $\mu \in P$ , acting on each  $M_v^{\text{ssreg}}$  by

$$\beta_v^\mu([x]) := (\mu - wv\mu)[x] + [x] \sum_{\alpha \in \Delta_+, k > 1} \langle \alpha^\vee, \mu \rangle \overleftarrow{D}_{[v(\alpha), k]},$$

where  $x = wt_{v\lambda} \in W_{\text{aff}}^{\text{ssreg}}$ . Denote by  $W_{\text{aff}}^{\text{ssreg}}(v)$  the subset of  $W_{\text{aff}}$  consisting of the superregular elements belonging to the  $v$ -chamber. Fix a left  $S$ -module isomorphism

$$\begin{array}{ccc} \iota : & S\langle W_{\text{aff}}^{\text{ssreg}}(v) \rangle & \rightarrow M_v^{\text{ssreg}} \\ & x & \mapsto [x]. \end{array}$$

**Proposition 4.2.** *For each  $v \in W$  and a sufficiently superregular element  $x \in W_{\text{aff}}^{\text{ssreg}}(v)$ ,*

$$\beta_v^\mu([x]) = \iota(B^\mu(x)).$$

*Proof.* This can be shown by using Lemma 4.1 and Proposition 4.1.

$$\beta_v^\mu([x]) = (\mu - wv\mu)[x] + [x] \sum_{\alpha \in \Delta_+, k > 1} \langle \alpha^\vee, \mu \rangle \overleftarrow{D}_{[v(\alpha), k]}$$

$$\begin{aligned}
&= (\mu - wv\mu)[x] + \sum_{\alpha \in \Delta_+} \sum_{k > 1, l(xs_{[v(\alpha), k]}) = l(x) - 1} \langle \alpha^\vee, \mu \rangle [xs_{v(\alpha), k}] \\
&= (\mu - wv\mu)[x] + \sum_{\alpha \in \Delta_+} \sum_{xs_{v(\alpha), k} \rightarrow_{near} x} \langle \alpha^\vee, \mu \rangle [xs_{v(\alpha), k}] = \iota(B^\mu(x)).
\end{aligned}$$

**Remark 4.1.** In [5] the authors introduced the quantization operators  $\eta_\alpha$  acting on the model of  $H^*(G/B) \otimes \mathbb{C}[q_1, \dots, q_r]$  realized as a subalgebra of  $\mathcal{B}_W \otimes \mathbb{C}[q_1, \dots, q_{n-1}]$ . For a superregular element  $\lambda \in \tilde{Q}$  and  $w \in W$ , consider a homomorphism  $\theta_w^\lambda$  from the  $\lambda$ -small elements (see [7, Section 5]) of  $H^*(G/B) \otimes \mathbb{C}[q]$  to  $\mathcal{B}_{\text{aff}}$  defined by

$$\theta_w^\lambda(q^\mu \sigma^v) := [vw^{-1}t_{w(\lambda+\mu)}],$$

where  $\sigma^v$  is the Schubert class of  $G/B$  corresponding to  $v \in W$  and  $q^\mu = q_1^{\mu_1} \cdots q_r^{\mu_r}$  for  $\mu = \sum_{i=1}^r \mu_i \alpha_i^\vee$ . The following is an interpretation of the formula of [7, Proposition 5.1] in our setting:

$$\theta_w^\lambda(\eta_\alpha(\sigma)) = \beta_w^{\varpi_\alpha}(\theta_w^\lambda(\sigma)).$$

## 5 Quadratic relations

For  $\alpha \in \Delta_+$  and  $v \in W$ , let us define the operator  $\mathbb{D}_v(\alpha)$  by

$$\mathbb{D}_v(\alpha) := \sum_{k > 1} \overleftarrow{D}_{[v(\alpha), k]}.$$

Then we have

$$\beta_v^\mu([x]) = (\mu - wv\mu)[x] + [x] \sum_{\alpha \in \Delta_+} \langle \alpha^\vee, \mu \rangle \mathbb{D}_v(\alpha).$$

In the following, we discuss the relations among the operators  $\mathbb{D}_v(\alpha)$ ,  $\alpha \in \Delta_+$ , for the root system of type  $A_{n-1}$ . For simplicity, we consider only non-equivariant case with  $v = id$ . Take the standard realization of the  $A_{n-1}$ -system:

$$\Delta = \{\varepsilon_i - \varepsilon_j \mid 1 \leq i, j \leq n, i \neq j\}.$$

Put  $\mathbb{D}(ij) := \mathbb{D}_{id.}(\varepsilon_i - \varepsilon_j)$  for  $1 \leq i < j \leq n$ , and  $\mathbb{D}(ij) := -\mathbb{D}(ji)$  for  $i > j$ . In this situation, we have a formula for the non-equivariant limit  $\bar{\beta}_{id.}^{\varepsilon_i}$  of the operator  $\beta_{id.}^{\varepsilon_i}$ :

$$\bar{\beta}_{id.}^{\varepsilon_i} = \sum_{j \neq i} \mathbb{D}(ij).$$

Note that this formula is analogous to the definition of the Dunkl elements in [4].

Let  $T_i$ ,  $1 \leq i \leq n-1$ , be linear operators on  $M^{\text{ssreg}}$  defined by  $T_i([x]) := [xt_{\alpha_i}]$ , where  $x \in W_{\text{aff}}$  and  $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ . It is easy to check from Proposition 4.1 that  $(T_i[x])\mathbb{D}(jk) = T_i([x]\mathbb{D}(jk))$ . Our next goal is to show that the operators  $\mathbb{D}(ij)$  satisfy the defining relations of the quantum deformation  $\mathcal{E}_n^q$  of the Fomin-Kirillov quadratic algebra [4].

**Proposition 5.1.** (i) For  $1 \leq i < j \leq n$ , we have

$$\mathbb{D}(ij)^2 = \begin{cases} T_i, & \text{if } j = i+1, \\ 0, & \text{otherwise.} \end{cases}$$

(ii) If  $\{i, j\} \cap \{k, l\} = \emptyset$ , then we have  $\mathbb{D}(ij)\mathbb{D}(kl) = \mathbb{D}(kl)\mathbb{D}(ij)$ .

(iii) For  $1 \leq i, j \leq n$ ,  $i \neq j$ , we have

$$\mathbb{D}(ij)\mathbb{D}(jk) + \mathbb{D}(jk)\mathbb{D}(kl) + \mathbb{D}(ki)\mathbb{D}(ij) = 0.$$

*Proof.* First of all, let us check the equality (i). We have

$$\mathbb{D}(ij)^2 = \sum_{k, l > 1} \overleftarrow{\mathcal{D}}_{[ij, k]} \overleftarrow{\mathcal{D}}_{[ij, l]}.$$

Let  $\lambda \in \tilde{Q}$  be sufficiently superregular. For  $x = wt_\lambda \in W_{\text{aff}}$ , assume that  $[x] \overleftarrow{\mathcal{D}}_{[ij, k]} \overleftarrow{\mathcal{D}}_{[ij, l]} \neq 0$ . Then we have the arrows  $xs_{ij, k} \rightarrow_{\text{near}} x$  and  $xs_{ij, k}s_{ij, l} \rightarrow_{\text{near}} xs_{ij, k}$  in the Bruhat ordering. From the conditions (1) and (2) in Proposition 4.1, one of the following conditions holds:

Case (1):  $k = -\langle \lambda, \varepsilon_i - \varepsilon_j \rangle$  and  $l(w) = l(ws_{ij}) - 1$ ,

Case (2):  $k = -\langle \lambda, \varepsilon_i - \varepsilon_j \rangle - 1$  and  $l(w) = l(ws_{ij}) + \langle \varepsilon_i - \varepsilon_j, 2\rho \rangle - 1$ .

In Case (1), since the arrow  $xs_{ij, k}s_{ij, l} = ws_{ij}t_\lambda s_{ij, l} \rightarrow_{\text{near}} xs_{ij, k}$  must come from the condition (2) of Proposition 4.1, we have  $\langle \varepsilon_i - \varepsilon_j, 2\rho \rangle - 1 = 1$ . This equality implies that  $\varepsilon_i - \varepsilon_j$  is a simple root  $\alpha_i$ , and we get

$$[x]\mathbb{D}(i \ i+1)^2 = [x] \overleftarrow{\mathcal{D}}_{[\alpha_i, -\langle \lambda, \alpha_i \rangle]} \overleftarrow{\mathcal{D}}_{[\alpha_i, -\langle \lambda, \alpha_i \rangle - 1]} = [xt_{\alpha_i}] = T_i[x].$$

In Case (2), since the arrow  $xs_{ij,k}s_{ij,l} = ws_{ij}t_{\lambda+\varepsilon_i-\varepsilon_j}s_{ij,l} \rightarrow_{near} xs_{ij,k}$  comes from the condition (1) of Proposition 4.1, we again obtain  $\langle \varepsilon_i - \varepsilon_j, 2\rho \rangle - 1 = 1$  and  $\varepsilon_i - \varepsilon_j = \alpha_i$ . Hence we get

$$[x]\mathbb{D}(i \ i + 1)^2 = [x]\overleftarrow{D}_{[\alpha_i, -\langle \lambda, \alpha_i \rangle - 1]}\overleftarrow{D}_{[\alpha_i, -\langle \lambda, \alpha_i \rangle - 2]} = [xt_{\alpha_i}] = T_i[x].$$

If  $j \neq i + 1$ , we have  $\mathbb{D}(ij)^2 = 0$ . The relations (ii) and (iii) follow from the identities  $[ij, a][kl, b] = [kl, b][ij, a]$  for  $\{i, j\} \cap \{k, l\} = \emptyset$ , and

$$[ij, a][jk, b] + [jk, b][ki, -a - b] + [ki, -a - b][ij, a] = 0$$

in  $\mathcal{B}_{\text{aff}}$ .

**Remark 5.1.** The operators  $\mathbb{D}_v(\alpha)$  induce the quantum Bruhat representation of  $\mathcal{E}_n^q$  via  $\theta_v^\lambda$ .

## References

- [1] N. Andruskiewitsch and H.-J. Schneider, *Finite quantum groups and Cartan matrices*, Adv. Math. **154** (2000), 1-45.
- [2] Y. Bazlov, *Nichols-Woronowicz algebra model for Schubert calculus on Coxeter groups*, J. Algebra, **297** (2006), 372-399.
- [3] R. Bott, *The space of loops on a Lie group*, Michigan Math. J., **5** (1958), 35-61.
- [4] S. Fomin and A. N. Kirillov, *Quadratic algebras, Dunkl elements and Schubert calculus*, Advances in Geometry, (J.-L. Brylinski, R. Brylinski, V. Nistor, B. Tsygan, and P. Xu, eds. ) Progress in Math., **172**, Birkhäuser, 1995, 147-182.
- [5] A. N. Kirillov and T. Maeno, *A note on quantization operators on Nichols algebra model for Schubert calculus on Weyl groups*, Lett. Math. Phys. **72** (2005), 233-241.
- [6] B. Kostant and S. Kumar, *The nil Hecke ring and cohomology of  $G/P$  for a Kac-Moody group  $G$* , Adv. in Math. **62** (1986), 187-237.
- [7] T. Lam and M. Shimozono, *Quantum cohomology of  $G/P$  and homology of affine Grassmannian*, Acta Math., **24** (2010), 49-90.

- [8] C. Lenart and A. Postnikov, *Affine Weyl groups in K-theory and representation theory*, Int. Math. Res. Notices **2007**, no. 12, Art. ID rnm038, 65pp.
- [9] S. Majid, *Free braided differential calculus, braided binomial theorem, and the braided exponential map*, J. Math. Phys., **34** (1993), 4843-4856.
- [10] S. Majid, *Noncommutative differentials and Yang-Mills on permutation groups  $S_N$* , Lecture Notes in Pure and Appl. Math., vol. 239, Dekker, 2004, 189-214.
- [11] A. Milinski and H.-J. Schneider, *Pointed indecomposable Hopf algebras over Coxeter groups*, Contemp. Math., **267** (2000), 215-236.
- [12] W. D. Nichols, *Bialgebras of type one*, Comm. Algebra, **6** (1978), 1521-1552.
- [13] D. Peterson, Lecture notes at MIT, 1997.
- [14] S. L. Woronowicz, *Differential calculus on compact matrix pseudogroups (quantum groups)*, Commun. Math. Phys., **122** (1989), 125-170.

Research Institute for Mathematical Sciences  
 Kyoto University  
 Sakyo-ku, Kyoto 606-8502, Japan  
 e-mail: kirillov@kurims.kyoto-u.ac.jp  
 URL: <http://www.kurims.kyoto-u.ac.jp/~kirillov>

Department of Electrical Engineering,  
 Kyoto University,  
 Sakyo-ku, Kyoto 606-8501, Japan  
 e-mail: maeno@kuee.kyoto-u.ac.jp