

RIMS-1758

**INTER-UNIVERSAL TEICHMÜLLER THEORY III:  
CANONICAL SPLITTINGS OF THE LOG-THETA-LATTICE**

By

Shinichi MOCHIZUKI

August 2012



京都大学 数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

# INTER-UNIVERSAL TEICHMÜLLER THEORY III: CANONICAL SPLITTINGS OF THE LOG-THETA-LATTICE

SHINICHI MOCHIZUKI

August 2012

ABSTRACT.

In the present paper, which is the third in a series of four papers, we study the theory surrounding the **log-theta-lattice**, a *highly non-commutative* two-dimensional diagram of “*miniature models of conventional scheme theory*”, called  $\Theta^{\pm\text{ell}}\text{NF-Hodge theaters}$ , that were associated, in the first paper of the series, to certain data, called *initial  $\Theta$ -data*, that includes an *elliptic curve*  $E_F$  over a *number field*  $F$ , together with a *prime number*  $l \geq 5$ . The **horizontal arrows** of the log-theta-lattice are defined as certain versions of the “ $\Theta$ -link” that was constructed, in the second paper of the series, by applying the theory of *Hodge-Arakelov-theoretic evaluation* — i.e., evaluation in the style of the **scheme-theoretic Hodge-Arakelov theory** established by the author in previous papers — of the [reciprocal of the  $l$ -th root of the] **theta function** at  **$l$ -torsion points**. In the present paper, we study the theory surrounding the **log-link** between  $\Theta^{\pm\text{ell}}\text{NF-Hodge theaters}$ . The **log-link** is obtained, roughly speaking, by applying, at each [say, for simplicity, nonarchimedean] valuation of the number field under consideration, the *local  $p$ -adic logarithm*. The significance of the **log-link** lies in the fact it allows one to construct **log-shells**, i.e., roughly speaking, slightly adjusted forms of the image of the local units at the valuation under consideration via the local  $p$ -adic logarithm. The theory of log-shells was studied extensively in a previous paper of the author. The **vertical arrows** of the log-theta-lattice are given by the **log-link**. Consideration of various properties of the log-theta-lattice leads naturally to the establishment of **multiradial algorithms** for constructing “**splitting monoids of logarithmic Gaussian procession monoids**”. Here, we recall that “multiradial algorithms” are algorithms that make sense from the point of view of an “**alien arithmetic holomorphic structure**”, i.e., the ring/scheme structure of a  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$  related to a given  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$  by means of a *non-ring/scheme-theoretic* horizontal arrow of the log-theta-lattice. These logarithmic Gaussian procession monoids, or **LGP-monoids**, for short, may be thought of as the log-shell-theoretic versions of the *Gaussian monoids* that were studied in the second paper of the series. Finally, by applying these multiradial algorithms for splitting monoids of LGP-monoids, we obtain **estimates** for the **log-volume** of these LGP-monoids. These estimates will be applied to verify various *diophantine results* in the fourth paper of the series.

## Contents:

Introduction

§0. Notations and Conventions

§1. The Log-theta-lattice

§2. Multiradial Theta Monoids

§3. Multiradial Logarithmic Gaussian Procession Monoids

## Introduction

In the following discussion, we shall continue to use the notation of the Introduction to the first paper of the present series of papers [cf. [IUTchI], §II]. In particular, we assume that are given an *elliptic curve*  $E_F$  over a *number field*  $F$ , together with a *prime number*  $l \geq 5$ . In the first paper of the series, we introduced and studied the basic properties of  $\Theta^{\pm\text{ell}}\text{NF-Hodge theaters}$ , which may be thought of as miniature models of the conventional scheme theory surrounding the given elliptic curve  $E_F$  over the number field  $F$ . In the present paper, which forms the third paper of the series, we study the theory surrounding the **log-link** between  $\Theta^{\pm\text{ell}}\text{NF-Hodge theaters}$ . The **log-link** induces an *isomorphism between the underlying  $\mathcal{D}$ - $\Theta^{\pm\text{ell}}\text{NF-Hodge theaters}$*  and, roughly speaking, is obtained by applying, at each [say, for simplicity, nonarchimedean] valuation  $\underline{v} \in \underline{\mathbb{V}}$ , the *local  $p_{\underline{v}}$ -adic logarithm* to the local units [cf. Proposition 1.3, (i)]. The significance of the **log-link** lies in the fact it allows one to construct **log-shells**, i.e., roughly speaking, slightly adjusted forms of the image of the local units at  $\underline{v} \in \underline{\mathbb{V}}$  via the local  $p_{\underline{v}}$ -adic logarithm. The theory of log-shells was studied extensively in [AbsTopIII]. The introduction of log-shells leads naturally to the construction of *new versions* — namely, the  $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -**links** [cf. Definition 3.8, (ii)] — of the  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -*links* studied in [IUTchI], [IUTchII]. The resulting [*highly non-commutative!*] diagram of iterates of the **log-** [i.e., the *vertical arrows*] and  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -links [i.e., the *horizontal arrows*] — which we refer to as the **log-theta-lattice** [cf. Definitions 1.4; 3.8, (iii), as well as Fig. I.1 below, in the case of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link] — plays a *central role* in theory of the present series of papers.




Fig. I.1: The [LGP-Gaussian] log-theta-lattice

Consideration of various properties of the log-theta-lattice leads naturally to the establishment of **multiradial algorithms** for constructing “**splitting monoids of logarithmic Gaussian procession monoids**” [cf. Theorem A below]. Here, we recall that “multiradial algorithms” [cf. the discussion of [IUTchII], Introduction] are algorithms that make sense from the point of view of an “**alien arithmetic holomorphic structure**”, i.e., the ring/scheme structure of a  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$  related to a given  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$  by means of a *non-ring/scheme-theoretic*  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -link. These logarithmic Gaussian procession

monoids, or **LGP-monoids**, for short, may be thought of as the log-shell-theoretic versions of the *Gaussian monoids* that were studied in [IUTchII]. Finally, by applying these multiradial algorithms for splitting monoids of LGP-monoids, we obtain **estimates** for the **log-volume** of these LGP-monoids [cf. Theorem B below]. These estimates will be applied to verify various *diophantine results* in [IUTchIV].

Recall [cf. [IUTchI], §I1] the notion of an  $\mathcal{F}$ -prime-strip. An  $\mathcal{F}$ -prime-strip consists of data indexed by the valuations  $\underline{v} \in \underline{\mathbb{V}}$ ; roughly speaking, the data at each  $\underline{v}$  consists of a *Frobenioid*, i.e., in essence, a system of *monoids* over a *base category*. For instance, at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , this data may be thought of as an isomorphic copy of the *monoid with Galois action*

$$\Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$$

— where we recall that  $\mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$  denotes the multiplicative monoid of nonzero integral elements of the completion of an algebraic closure  $\overline{F}$  of  $F$  at a valuation lying over  $\underline{v}$  [cf. [IUTchI], §I1, for more details]. The  $p_{\underline{v}}$ -adic logarithm  $\log_{\underline{v}} : \mathcal{O}_{\overline{F}_{\underline{v}}}^{\times} \rightarrow \overline{F}_{\underline{v}}$  at  $\underline{v}$  then defines a natural  $\Pi_{\underline{v}}$ -equivariant isomorphism of topological modules

$$(\mathcal{O}_{\overline{F}_{\underline{v}}}^{\times\mu} \otimes \mathbb{Q} \xrightarrow{\sim}) \quad \mathcal{O}_{\overline{F}_{\underline{v}}}^{\times} \otimes \mathbb{Q} \xrightarrow{\sim} \overline{F}_{\underline{v}}$$

— where we recall the notation “ $\mathcal{O}_{\overline{F}_{\underline{v}}}^{\times\mu} = \mathcal{O}_{\overline{F}_{\underline{v}}}^{\times} / \mathcal{O}_{\overline{F}_{\underline{v}}}^{\mu}$ ” from the discussion of [IUTchI], §1 — which allows one to equip  $\mathcal{O}_{\overline{F}_{\underline{v}}}^{\times} \otimes \mathbb{Q}$  with the *field structure* arising from the field structure of  $\overline{F}_{\underline{v}}$ . The portion at  $\underline{v}$  of the **log-link** associated to an  $\mathcal{F}$ -prime-strip [cf. Definition 1.1, (iii); Proposition 1.2] may be thought of as the correspondence

$$\left\{ \Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright} \right\} \xrightarrow{\log} \left\{ \Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright} \right\}$$

in which one thinks of the copy of “ $\mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$ ” on the *right* as obtained from the field structure induced by the  $p_{\underline{v}}$ -adic logarithm on the tensor product with  $\mathbb{Q}$  of the copy of the units “ $\mathcal{O}_{\overline{F}_{\underline{v}}}^{\times} \subseteq \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$ ” on the *left*. Since this correspondence induces an *isomorphism of topological groups* between the copies of  $\Pi_{\underline{v}}$  on either side, one may think of  $\Pi_{\underline{v}}$  as “*immune to*”/“*neutral with respect to*” — or, in the terminology of the present series of papers, “**coric**” with respect to — the transformation constituted by the **log-link**. This situation is studied in detail in [AbsTopIII], §3, and reviewed in Proposition 1.2 of the present paper.

By applying various results from **absolute anabelian geometry**, one may algorithmically reconstruct a copy of the data “ $\Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$ ” from  $\Pi_{\underline{v}}$ . Moreover, by applying *Kummer theory*, one obtains natural isomorphisms between this “*coric version*” of the data “ $\Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$ ” and the copies of this data that appear on either side of the **log-link**. On the other hand, one verifies immediately that these Kummer isomorphisms are **not compatible** with the **coricity** of the copy of the data “ $\Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$ ” algorithmically constructed from  $\Pi_{\underline{v}}$ . This phenomenon is, in some sense, the *central theme* of the theory of [AbsTopIII], §3, and is reviewed in Proposition 1.2, (iv), of the present paper.

The introduction of the **log-link** leads naturally to the construction of **log-shells** at each  $\underline{v} \in \underline{\mathbb{V}}$ . If, for simplicity,  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , then the log-shell at  $\underline{v}$  is given, roughly speaking, by the *compact additive module*

$$\mathcal{I}_{\underline{v}} \stackrel{\text{def}}{=} p_{\underline{v}}^{-1} \cdot \log_{\underline{v}}(\mathcal{O}_{K_{\underline{v}}}^{\times}) \subseteq K_{\underline{v}} \subseteq \overline{F}_{\underline{v}}$$

[cf. Definition 1.1, (i), (ii); Remark 1.2.2, (i), (ii)]. One has natural *functorial algorithms* for constructing various versions — i.e., **mono-analytic/holomorphic** and **étale-like/Frobenius-like** — from  $\mathcal{D}^{\dagger}$ -/ $\mathcal{D}$ -/ $\mathcal{F}^{\dagger}$ -/ $\mathcal{F}$ -prime-strips [cf. Proposition 1.2, (v), (vi), (vii), (viii), (ix)]. Although, as discussed above, the relevant Kummer isomorphisms are *not compatible* with the **log-link** “*at the level of elements*”, the log-shell  $\mathcal{I}_{\underline{v}}$  at  $\underline{v}$  satisfies the important property

$$\mathcal{O}_{K_{\underline{v}}}^{\triangleright} \subseteq \mathcal{I}_{\underline{v}}; \quad \log_{\underline{v}}(\mathcal{O}_{K_{\underline{v}}}^{\times}) \subseteq \mathcal{I}_{\underline{v}}$$

— i.e., it **contains the images** of the Kummer isomorphisms associated to both the domain and the codomain of the **log-link** [cf. Proposition 1.2, (v); Remark 1.2.2, (i), (ii)]. In light of the *compatibility* of the **log-link** with *log-volumes* [cf. Propositions 1.2, (iii); 3.9, (iv)], this property will ultimately lead to **upper bounds** — i.e., as opposed to “*precise equalities*” — in the computation of *log-volumes* in Corollary 3.12 [cf. Theorem B below]. Put another way, although iterates of the **log-link** *fail to be compatible* with the various Kummer isomorphisms that arise, one may nevertheless consider the *entire diagram* that results from considering such iterates of the **log-link** and related Kummer isomorphisms [cf. Proposition 1.2, (x)]. We shall refer to such diagrams

$$\begin{array}{ccccccc} \dots & \rightarrow & \bullet & \rightarrow & \bullet & \rightarrow & \bullet & \rightarrow & \dots \\ & & & \searrow & \downarrow & \swarrow & & & \\ & & & & \circ & & & & \end{array}$$

— i.e., where the *horizontal arrows* correspond to the **log-links** [that is to say, to the *vertical arrows* of the log-theta-lattice!]; the “ $\bullet$ ’s” correspond to the Frobenioid-theoretic data within a  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater; the “ $\circ$ ” corresponds to the *coric version* of this data [that is to say, in the terminology discussed below, *vertically coric* data of the log-theta-lattice]; the vertical/diagonal arrows correspond to the various *Kummer isomorphisms* — as **log-Kummer correspondences** [cf. Theorem 3.11, (ii); Theorem A, (ii), below]. Then the inclusions of the above display may be interpreted as a sort of “**upper semi-commutativity**” of such diagrams [cf. Remark 1.2.2, (iii)], which we shall also refer to as the “**upper semi-compatibility**” of the **log-link** with the relevant Kummer isomorphisms — cf. the discussion of the “**indeterminacy**” (Ind3) in Theorem 3.11, (ii).

By considering the **log-links** associated to the various  $\mathcal{F}$ -prime-strips that occur in a  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater, one obtains the notion of a **log-link** between  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters

$$\dagger \mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} \ddagger \mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$$

[cf. Proposition 1.3, (i)]. As discussed above, by considering the iterates of the **log-link** [i.e., the *vertical arrows*] and  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{tgp}}^{\times\mu}$ -links [i.e., the *horizontal*

arrows], one obtains a diagram which we refer to as the **log-theta-lattice** [cf. Definitions 1.4; 3.8, (iii), as well as Fig. I.1, in the case of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link]. As discussed above, this diagram is **highly noncommutative**, since the definition of the **log-link** depends, in an essential way, on both the *additive* and the *multiplicative* structures — i.e., on the *ring structure* — of the various local rings at  $v \in \underline{\mathbb{V}}$ , structures which are *not preserved* by the  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -links [cf. Remark 1.4.1, (i)]. So far, in the Introductions to [IUTchI], [IUTchII], as well as in the present Introduction, we have discussed various “*coricity*” properties — i.e., properties of *invariance* with respect to various types of “transformations” — in the context of  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -links, as well as in the context of **log-links**. In the context of the log-theta-lattice, it becomes necessary to distinguish between various types of coricity. That is to say, coricity with respect to **log-links** [i.e., the vertical arrows of the log-theta-lattice] will be referred to as **vertical coricity**, while coricity with respect to  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -links [i.e., the horizontal arrows of the log-theta-lattice] will be referred to as **horizontal coricity**. On the other hand, coricity properties that hold with respect to *all* of the arrows of the log-theta-lattice will be referred to as **bi-coricity** properties.

Relative to the analogy between the theory of the present series of papers and *p-adic Teichmüller theory* [cf. [IUTchI], §I4], we recall that a  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$ , which may be thought of as a miniature model of the *conventional scheme theory* surrounding the given elliptic curve  $E_F$  over the number field  $F$ , corresponds to the *positive characteristic scheme theory* surrounding a hyperbolic curve over a positive characteristic perfect field that is equipped with a nilpotent ordinary indigenous bundle [cf. Fig. I.2 below]. Then the **rotation**, or “**juggling**”, effected by the **log-link** of the **additive** and **multiplicative** structures of the conventional scheme theory represented by a  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$  may be thought of as corresponding to the **Frobenius morphism in positive characteristic** [cf. the discussion of [AbsTopIII], §I3, §I5]. Thus, just as the Frobenius morphism is completely well-defined in *positive characteristic*, the **log-link** may be thought of as a phenomenon that occurs within a *single arithmetic holomorphic structure*, i.e., a *vertical* line of the log-theta-lattice. By contrast, the essentially *non-ring/scheme-theoretic* relationship between  $\Theta^{\pm\text{ell}}\text{NF-Hodge theaters}$  constituted by the  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -links corresponds to the relationship between the “*mod p*” and “*mod p<sup>n+1</sup>*” portions of the ring of Witt vectors, in the context of a *canonical lifting* of the original positive characteristic data [cf. the discussion of Remark 1.4.1, (iii); Fig. I.2 below]. Thus, the **log-theta-lattice**, taken as a whole, may be thought of as corresponding to the **canonical lifting** of the original positive characteristic data, equipped with a corresponding **canonical Frobenius action/lifting** [cf. Fig. I.2 below]. Finally, the **non-commutativity** of the log-theta-lattice may be thought of as corresponding to the complicated “**intertwining**” that occurs in the theory of Witt vectors and canonical liftings between the Frobenius morphism in positive characteristic and the mixed characteristic nature of the ring of Witt vectors [cf. the discussion of Remark 1.4.1, (ii), (iii)].

One important consequence of this “*noncommutative intertwining*” of the two dimensions of the log-theta-lattice is the following. Since each *horizontal arrow* of the log-theta-lattice [i.e., the  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -link] may only be used to relate — i.e., via various *Frobenioids* — the *multiplicative* portions of the



ring structures in the domain and codomain of the arrow, one natural approach to relating the *additive* portions of these ring structures is to apply the theory of **log-shells**. That is to say, since each horizontal arrow is compatible with the **canonical splittings** [up to roots of unity] discussed in [IUTchII], Introduction, of the *theta/Gaussian monoids* in the domain of the horizontal arrow into *unit group* and *value group* portions, it is natural to attempt to relate the ring structures on either side of the horizontal arrow by applying the canonical splittings to

- relate the **multiplicative** structures on either side of the horizontal arrow by means of the **value group** portions of the theta/Gaussian monoids;
- relate the **additive** structures on either side of the horizontal arrow by means of the **unit group** portions of the theta/Gaussian monoids, **shifted once** via a *vertical arrow*, i.e., the **log-link**, so as to “*render additive*” the [*a priori*] multiplicative structure of these unit group portions.

Indeed, this is the approach that will ultimately be taken in Theorem 3.11 [cf. Theorem A below] to relating the ring structures on either side of a horizontal arrow. On the other hand, in order to actually implement this approach, it will be necessary to overcome numerous *technical obstacles*. Perhaps the most immediately obvious such obstacle lies in the observation [cf. the discussion of Remark 1.4.1, (ii)] that, precisely because of the “*noncommutative intertwining*” nature of the log-theta-lattice,

any sort of algorithmic construction concerning objects lying in the *domain* of a horizontal arrow that involves **vertical shifts** [e.g., such as the approach to relating additive structures in the fashion described above] **cannot be “translated”** in any immediate sense into an algorithm that makes sense from the point of view of the *codomain* of the horizontal arrow.

In a word, our approach to overcoming this technical obstacle consists of working with objects in the *vertical line* of the log-theta-lattice that contains the *domain* of the horizontal arrow under consideration that satisfy the crucial property of being

**invariant** with respect to **vertical shifts**

— i.e., **shifts** via iterates of the **log-link** [cf. the discussion of Remarks 1.2.2; 1.4.1, (ii)]. For instance, *étale-like* objects that are **vertically coric** satisfy this invariance property. On the other hand, as discussed in the beginning of [IUTchII], Introduction, in the theory of the present series of papers, it is of crucial importance to be able to relate *corresponding Frobenius-like* and *étale-like structures* to one another via *Kummer theory*. In particular, in order to obtain structures that are *invariant* with respect to *vertical shifts*, it is necessary to consider **log-Kummer correspondences**, as discussed above. Moreover, in the context of such **log-Kummer correspondences**, typically, one may only obtain structures that are invariant with respect to vertical shifts if one is willing to admit some sort of **indeterminacy**, e.g., such as the “**upper semi-compatibility**” [cf. the discussion of the “*indeterminacy*” (Ind3) in Theorem 3.11, (ii)] discussed above.

<i>Inter-universal Teichmüller theory</i>	<i>p-adic Teichmüller theory</i>
<b>number field</b> $F$	<b>hyperbolic curve</b> $C$ over a <i>positive characteristic perfect field</i>
<b>[once-punctured] elliptic curve</b> $X$ over $F$	<i>nilpotent ordinary</i> <b>indigenous bundle</b> $P$ over $C$
$\Theta$ - <b>link</b> arrows of the <i>log-theta-lattice</i>	<b>mixed characteristic extension</b> structure of a ring of <i>Witt vectors</i>
<b>log-link</b> arrows of the <i>log-theta-lattice</i>	the <b>Frobenius morphism</b> in <i>positive characteristic</i>
the entire <b>log-theta-lattice</b>	the resulting <b>canonical lifting</b> + <b>canonical Frobenius action</b> ; <b>canonical Frobenius lifting</b> over the ordinary locus
relatively straightforward <i>original construction of</i> $\Theta_{\text{LGP}}^{\times\mu}$ - <b>link</b>	relatively straightforward <i>original construction of</i> <b>canonical liftings</b>
highly nontrivial <i>description of alien arithmetic</i> <b>holomorphic structure</b> via <i>absolute anabelian geometry</i>	highly nontrivial <i>absolute anabelian</i> <i>reconstruction of</i> <b>canonical liftings</b>

Fig. I.2: Correspondence between inter-universal Teichmüller theory and  $p$ -adic Teichmüller theory

One important property of the **log-link**, and hence, in particular, of the construction of **log-shells**, is its **compatibility** with the  $\mathbb{F}_l^{\times\pm}$ -**symmetry** discussed in the Introductions to [IUTchI], [IUTchII] — cf. Remark 1.3.2. Here, we recall from the discussion of [IUTchII], Introduction, that the  $\mathbb{F}_l^{\times\pm}$ -symmetry allows one to relate the various  $\mathcal{F}$ -*prime-strips* — i.e., more concretely, the various copies of the



data “ $\Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  [and their analogues for  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}}$ ] — associated to the various **labels**  $\in \mathbb{F}_l$  that appear in the *Hodge-Arakelov-theoretic evaluation* of [IUTchII] in a fashion that is **compatible** with

- the **distinct nature** of distinct labels  $\in \mathbb{F}_l$ ;
- the **Kummer isomorphisms** used to relate *Frobenius-like* and *étale-like* versions of the  $\mathcal{F}$ -prime-strips appear, i.e., more concretely, the various copies of the data “ $\Pi_{\underline{v}} \curvearrowright \mathcal{O}_{\overline{F}_{\underline{v}}}^{\triangleright}$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  [and their analogues for  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}}$ ];
- the structure of the **underlying  $\mathcal{D}$ -prime-strips** that appear, i.e., more concretely, the various copies of the *[arithmetic] tempered fundamental group* “ $\Pi_{\underline{v}}$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  [and their analogues for  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}}$ ]

— cf. the discussion of [IUTchII], Introduction; Remark 1.5.1, Step (vii) of the proof of Corollary 3.12 of the present paper. This compatibility with the  $\mathbb{F}_l^{\times\pm}$ -symmetry gives rise to the construction of

- “**arithmetically holomorphic**”  $\mathcal{F}^{+\times\mu}$ -prime strips, log-shells which are **vertically coric**;
- **mono-analytic**  $\mathcal{F}^{+\times\mu}$ -prime strips, log-shells which are **bi-coric**

— cf. Theorem 1.5. These *bi-coric mono-analytic log-shells* play a central role in the theory of the present paper.

One notable aspect of the *compatibility* of the **log-link** with the  $\mathbb{F}_l^{\times\pm}$ -symmetry in the context of the theory of *Hodge-Arakelov-theoretic evaluation* developed in [IUTchII] is the following. One important property of *mono-theta environments* is the property of “**isomorphism class compatibility**”, i.e., in the terminology of [EtTh], “*compatibility with the topology of the tempered fundamental group*” [cf. the discussion of Remark 2.1.1]. This “isomorphism class compatibility” allows one to apply the Kummer theory of mono-theta environments [i.e., the theory of [EtTh]] relative to the **ring-theoretic basepoints** that occur on either side of the **log-link** [cf. Remark 2.1.1, (ii); [IUTchII], Remark 3.6.4, (i)], for instance, in the context of the **log-Kummer correspondences** discussed above. Here, we recall that the significance of working with such “ring-theoretic basepoints” lies in the fact that the full *ring structure* of the local rings involved [i.e., as opposed to, say, just the multiplicative portion of this ring structure] is necessary in order to construct the **log-link**. That is to say, it is precisely by establishing the *conjugate synchronization* arising from the  $\mathbb{F}_l^{\times\pm}$ -symmetry relative to *these basepoints* that occur on either side of the **log-link** that one is able to conclude the crucial **compatibility of this conjugate synchronization with the log-link** discussed in Remark 1.3.2. Thus, in summary, one important consequence of the “isomorphism class compatibility” of mono-theta environments is the **simultaneous compatibility** of

- the **Kummer theory of mono-theta environments**;
- the **conjugate synchronization** arising from the  $\mathbb{F}_l^{\times\pm}$ -symmetry;
- the construction of the **log-link**.

This simultaneous compatibility is necessary in order to perform the construction of the [crucial!] *splitting monoids of LGP-monoids* referred to above — cf. the discussion of Step (vi) of the proof of Corollary 3.12.

In §2 of the present paper, we continue our preparation for the *multiradial construction of splitting monoids of LGP-monoids* given in §3 [of the present paper] by presenting a **global formulation** of the essentially *local theory* at  $\underline{v} \in \mathbb{V}^{\text{bad}}$  [cf. [IUTchII], §1, §2, §3] concerning the interpretation, via the notion of **multiradiality**, of various **rigidity** properties of **mono-theta environments**. That is to say, although much of the [essentially routine!] task of formulating the local theory of [IUTchII], §1, §2, §3, in global terms was accomplished in [IUTchII], §4, the [again essentially routine!] task of formulating the portion of this local theory that concerns *multiradiality* was not addressed in the theory of [IUTchII], §4. One reason for this lies in the fact that, from the point of view of the theory to be developed in §3 of the present paper, this global formulation of multiradiality properties of the mono-theta environment may be presented most naturally in the framework developed in §1 of the present paper, involving the **log-theta-lattice** [cf. Theorem 2.2; Corollary 2.3]. Indeed, the **étale-like** versions of the mono-theta environment, as well as the various objects constructed from the mono-theta environment, may be interpreted, from the point of view of the log-theta-lattice, as **vertically coric** structures, and are **Kummer-theoretically** related to their **Frobenius-like** [i.e., Frobenioid-theoretic] counterparts, which arise from the [Frobenioid-theoretic portions of the] various  $\Theta^{\pm\text{ell}}$ NF-Hodge theaters in a vertical line of the log-theta-lattice [cf. Theorem 2.2, (ii); Corollary 2.3, (ii), (iii), (iv)]. Moreover, it is precisely the **horizontal arrows** of the log-theta-lattice that give rise to the  $\widehat{\mathbb{Z}}^\times$ -**indeterminacies** acting on copies of “ $\mathcal{O}^{\times\mu}$ ” that play a prominent role in the local multiradiality theory developed in [IUTchII] [cf. the discussion of [IUTchII], Introduction]. In this context, it is useful to recall from the discussion of [IUTchII], Introduction [cf. also Remark 2.2.1 of the present paper], that the essential content of this local multiradiality theory consists of the *observation* [cf. Fig. I.3 below] that, since *mono-theta-theoretic cyclotomic* and *constant multiple rigidity* only require the use of the portion of  $\mathcal{O}_{\underline{F}_v}^\times$ , for  $\underline{v} \in \mathbb{V}^{\text{bad}}$ , given by the *torsion subgroup*  $\mathcal{O}_{\underline{F}_v}^\mu \subseteq \mathcal{O}_{\underline{F}_v}^\times$  [i.e., the roots of unity], the *triviality* of the composite of natural morphisms

$$\mathcal{O}_{\underline{F}_v}^\mu \hookrightarrow \mathcal{O}_{\underline{F}_v}^\times \twoheadrightarrow \mathcal{O}_{\underline{F}_v}^{\times\mu}$$

has the effect of **insulating** the **Kummer theory** of the **étale theta function** — i.e., via the theory of the mono-theta environments developed in [EtTh] — from the  $\widehat{\mathbb{Z}}^\times$ -**indeterminacies** that act on the copies of “ $\mathcal{O}^{\times\mu}$ ” that arise in the  $\mathcal{F}^{\text{tr}\times\mu}$ -*prime-strips* that appear in the  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{tgp}}^{\times\mu}$ -link.

$$\begin{array}{ccc} \text{id} \curvearrowright & & \widehat{\mathbb{Z}}^\times \curvearrowright \\ \boxed{\mathcal{O}_{\underline{F}_v}^\mu} & \rightarrow & \boxed{\mathcal{O}_{\underline{F}_v}^{\times\mu}} \end{array}$$

Fig. I.3: Insulation from  $\widehat{\mathbb{Z}}^\times$ -indeterminacies in the context of mono-theta-theoretic cyclotomic, constant multiple rigidity

In §3 of the present paper, which, in some sense, constitutes the *conclusion* of the theory developed thus far in the present series of papers, we present the construction of the [splitting monoids of] **LGP-monoids**, which may be thought of as a **multiradial** version of the [splitting monoids of] **Gaussian monoids** that were constructed via the theory of *Hodge-Arakelov-theoretic evaluation* developed in [IUTchII]. In order to achieve this multiradiality, it is necessary to “multiradialize” the various components of the construction of the Gaussian monoids given in [IUTchII]. The first step in this process of “multiradialization” concerns the **labels**  $j \in \mathbb{F}_l^*$  that occur in the Hodge-Arakelov-theoretic evaluation performed in [IUTchII]. That is to say, the construction of these labels, together with the closely related theory of  $\mathbb{F}_l^*$ -**symmetry**, depend, in an essential way, on the *full arithmetic tempered fundamental groups* “ $\Pi_v$ ” at  $v \in \underline{\mathbb{V}}^{\text{bad}}$ , i.e., on the portion of the *arithmetic holomorphic structure* within a  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater which is *not shared* by an **alien arithmetic holomorphic structure** [i.e., an arithmetic holomorphic structure related to the original arithmetic holomorphic structure via a horizontal arrow of the log-theta-lattice]. One naive approach to remedying this state of affairs is to simply consider the *underlying set*, of cardinality  $l^*$ , associated to  $\mathbb{F}_l^*$ , which we regard as being equipped with the full set of *symmetries* given by arbitrary permutation automorphisms of this underlying set. The problem with this approach is that it yields a situation in which, for each label  $j \in \mathbb{F}_l^*$ , one must contend with an *indeterminacy of  $l^*$  possibilities* for the element of this underlying set that corresponds to  $j$  [cf. [IUTchI], Propositions 4.11, (i); 6.9, (i)]. From the point of view of the *log-volume computations* to be performed in [IUTchIV], this degree of indeterminacy gives rise to log-volumes which are “*too large*”, i.e., to estimates that are not sufficient for deriving the various *diophantine results* obtained in [IUTchIV]. Thus, we consider the following alternative approach, via **processions** [cf. [IUTchI], Propositions, 4.11, 6.9]. Instead of working just with the underlying set associated to  $\mathbb{F}_l^*$ , we consider the *diagram of inclusions* of finite sets

$$\mathbb{S}_1^\pm \hookrightarrow \mathbb{S}_{1+1=2}^\pm \hookrightarrow \dots \hookrightarrow \mathbb{S}_{j+1}^\pm \hookrightarrow \dots \hookrightarrow \mathbb{S}_{1+l^*=l^\pm}^\pm$$

— where we write  $\mathbb{S}_{j+1}^\pm \stackrel{\text{def}}{=} \{0, 1, \dots, j\}$ , for  $j = 0, \dots, l^*$ , and we think of each of these finite sets as being subject to arbitrary permutation automorphisms. That is to say, we think of the set  $\mathbb{S}_{j+1}$  as a **container** for the labels  $0, 1, \dots, j$ . Thus, for each  $j$ , one need only contend with an *indeterminacy of  $j + 1$  possibilities* for the element of this container that corresponds to  $j$ . In particular, if one allows  $j = 0, \dots, l^*$  to vary, then this approach allows one to *reduce* the resulting label indeterminacy from a total of  $(l^\pm)^{l^\pm}$  possibilities [where we write  $l^\pm = 1 + l^* = (l + 1)/2$ ] to a total of  $l^\pm!$  [i.e.,  $\approx \sqrt{(l^\pm)^{l^\pm}}$ ] possibilities. It turns out that this reduction will yield just the right estimates in the log-volume computations to be performed in [IUTchIV]. Moreover, this approach satisfies the important property of *insulating the “core label 0” from the various label indeterminacies* that occur.

Each element of each of the containers  $\mathbb{S}_{j+1}$  may be thought of as parametrizing an  $\mathcal{F}$ - or  $\mathcal{D}$ -*prime-strip* that occurs in the *Hodge-Arakelov-theoretic evaluation* of [IUTchII]. In order to render the construction multiradial, it is necessary to replace such *holomorphic  $\mathcal{F}$ -/ $\mathcal{D}$ -prime-strips* by *mono-analytic  $\mathcal{F}^+$ -/ $\mathcal{D}^+$ -prime-strips*. In particular, as discussed above, one may construct, for each such  $\mathcal{F}^+$ -/ $\mathcal{D}^+$ -prime-strip, a collection of **log-shells** associated to the various  $v \in \underline{\mathbb{V}}$ . Write  $\mathbb{V}_{\mathbb{Q}}$  for

the set of valuations of  $\mathbb{Q}$ . Then, in order to obtain objects that are *immune* to the various label indeterminacies discussed above, we consider, for each element  $* \in \mathbb{S}_{j+1}$ , and for each [say, for simplicity, *nonarchimedean*]  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ ,

- the *direct sum of the log-shells* associated to the prime-strip labeled by the given element  $* \in \mathbb{S}_{j+1}$  at the  $\underline{v} \in \underline{\mathbb{V}}$  that lie over  $v_{\mathbb{Q}}$ ;

we then form

- the **tensor product**, over the elements  $* \in \mathbb{S}_{j+1}$ , of these *direct sums*.

This collection of tensor products associated to  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$  will be referred to as the **tensor packet** associated to the collection of prime-strips indexed by elements of  $\mathbb{S}_{j+1}$ . One may carry out this construction of the tensor packet either for *holomorphic*  $\mathcal{F}$ -/ $\mathcal{D}$ -prime-strips [cf. Proposition 3.1] or for *mono-analytic*  $\mathcal{F}^+$ -/ $\mathcal{D}^+$ -prime-strips [cf. Proposition 3.2].

The tensor packets associated to  $\mathcal{D}^+$ -prime-strips will play a crucial role in the theory of §3, as “**multiradial mono-analytic containers**” for the principal objects of interest [cf. the discussion of Remark 3.12.2, (ii)], namely,

- the action of the **splitting monoids** of the **LGP-monoids** — i.e., the monoids generated by the **theta values**  $\{q^{\underline{v}j^2}\}_{j=1,\dots,l^*}$  — on the portion of the *tensor packets* just defined at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  [cf. Fig. I.4 below; Propositions 3.4, 3.5; the discussion of [IUTchII], Introduction];
- the action of copies “ $(F_{\text{mod}}^{\times})_j$ ” of [the multiplicative monoid of nonzero elements of] the **number field**  $F_{\text{mod}}$  labeled by  $j = 1, \dots, l^*$  on the product, over  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ , of the portion of the *tensor packets* just defined at  $v_{\mathbb{Q}}$  [cf. Fig. I.5 below; Propositions 3.3, 3.7, 3.10].

$$\begin{array}{ccccccc}
 & \underline{q}^1 \curvearrowright & & \underline{q}^{j^2} \curvearrowright & & \underline{q}^{(l^*)^2} \curvearrowright & \\
 /^{\pm} & \hookrightarrow /^{\pm}/^{\pm} & \hookrightarrow \dots & \hookrightarrow /^{\pm}/^{\pm} \dots /^{\pm} & \hookrightarrow \dots & \hookrightarrow /^{\pm}/^{\pm} \dots \dots /^{\pm} & \\
 \mathbb{S}_1^{\pm} & \mathbb{S}_{1+1=2}^{\pm} & & \mathbb{S}_{j+1}^{\pm} & & \mathbb{S}_{1+l^*=l^{\pm}}^{\pm} & 
 \end{array}$$

Fig. I.4: Splitting monoids of LGP-monoids acting on tensor packets

$$\begin{array}{ccccccc}
 & (F_{\text{mod}}^{\times})_1 \curvearrowright & & (F_{\text{mod}}^{\times})_j \curvearrowright & & (F_{\text{mod}}^{\times})_{l^*} \curvearrowright & \\
 /^{\pm} & \hookrightarrow /^{\pm}/^{\pm} & \hookrightarrow \dots & \hookrightarrow /^{\pm}/^{\pm} \dots /^{\pm} & \hookrightarrow \dots & \hookrightarrow /^{\pm}/^{\pm} \dots \dots /^{\pm} & \\
 \mathbb{S}_1^{\pm} & \mathbb{S}_{1+1=2}^{\pm} & & \mathbb{S}_{j+1}^{\pm} & & \mathbb{S}_{1+l^*=l^{\pm}}^{\pm} & 
 \end{array}$$

Fig. I.5: Copies of  $F_{\text{mod}}^{\times}$  acting on tensor packets

Indeed, these [splitting monoids of] **LGP-monoids** and copies “ $(F_{\text{mod}}^\times)_j$ ” of [the multiplicative monoid of nonzero elements of] the **number field**  $F_{\text{mod}}$  admit *natural embeddings into/actions on the various tensor packets* associated to *labeled  $\mathcal{F}$ -prime-strips* in each  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  of the log-theta-lattice. One then obtains **vertically coric** versions of these splitting monoids of LGP-monoids and labeled copies “ $(F_{\text{mod}}^\times)_j$ ” of [the multiplicative monoid of nonzero elements of] the number field  $F_{\text{mod}}$  by applying appropriate **Kummer isomorphisms** between

- *log-shells/tensor packets associated to [labeled]  $\mathcal{F}$ -prime-strips* and
- *log-shells/tensor packets associated to [labeled]  $\mathcal{D}$ -prime-strips.*

Finally, by passing to the

- *log-shells/tensor packets associated to [labeled]  $\mathcal{D}^+$ -prime-strips*

— i.e., by *forgetting the arithmetic holomorphic structure* associated to a *specific vertical line* of the log-theta-lattice — one obtains the desired **multiradial representation**, i.e., description in terms that make sense from the point of view of an **alien arithmetic holomorphic structure**, of the **splitting monoids of LGP-monoids** and *labeled copies of the number field*  $F_{\text{mod}}$  discussed above. This passage to the multiradial representation is obtained by admitting the following *three types of indeterminacy*:

- (Ind1): This is the indeterminacy that arises from the *automorphisms of processions of  $\mathcal{D}^+$ -prime-strips* that appear in the multiradial representation — i.e., more concretely, from *permutation automorphisms* of the label sets  $\mathbb{S}_{j+1}$  that appear in the processions discussed above, as well as from the *automorphisms of the  $\mathcal{D}^+$ -prime-strips* that appear in these processions.
- (Ind2): This is the indeterminacy that arises from the *automorphisms of the  $\mathcal{F}^{+\times\mu}$ -prime-strips* that appear in the  $\Theta$ -/ $\Theta^{\times\mu}$ -/ $\Theta_{\text{gau}}^{\times\mu}$ -/ $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -link — i.e., in particular, at [for simplicity]  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , the  $\widehat{\mathbb{Z}}^\times$ -**indeterminacies** acting on local copies of “ $\mathcal{O}^{\times\mu}$ ” [cf. the above discussion].
- (Ind3): This is the indeterminacy that arises from the **upper semi-compatibility of the log-Kummer correspondences** associated to the specific vertical line of the log-theta-lattice under consideration [cf. the above discussion].

A detailed description of this multiradial representation, together with the indeterminacies (Ind1), (Ind2) is given in Theorem 3.11, (i) [and summarized in Theorem A, (i), below; cf. also Fig. I.6 below].

One important property of the multiradial representation discussed above concerns the relationship between the three main components — i.e., roughly speaking, *log-shells*, *splitting monoids of LGP-monoids*, and *number fields* — of this multiradial representation and the **log-Kummer correspondence** of the specific *vertical line* of the log-theta-lattice under consideration. This property — which may be thought of as a sort of “**non-interference**”, or “**mutual compatibility**”, property — asserts that the multiplicative monoids constituted by the splitting monoids



of LGP-monoids and copies of  $F_{\text{mod}}^\times$  “do not interfere”, relative to the various arrows that occur in the **log-Kummer correspondence**, with the *local integers* at  $\underline{v} \in \underline{\mathbb{V}}$ , hence, in particular, with the *local units* at  $\underline{v} \in \underline{\mathbb{V}}$ , that give rise to the *log-shells*. In the case of splitting monoids of LGP-monoids, this *non-interference/mutual compatibility* property is, in essence, a formal consequence of the existence of the **canonical splittings** [up to roots of unity] discussed in [IUTchII], Introduction, of the *theta/Gaussian monoids* that appear into *unit group* and *value group* portions. Here, we recall that, in the case of the theta monoids, these canonical splittings are, in essence, a formal consequence of the **constant multiple rigidity** property of mono-theta environments reviewed above. In the case of copies of  $F_{\text{mod}}^\times$ , this *non-interference/mutual compatibility* property is, in essence, a formal consequence of the well-known fact in elementary algebraic number theory that any nonzero element of a number field that is **integral** at every valuation of the number field is necessarily a **root of unity**. These mutual compatibility properties are described in detail in Theorem 3.11, (ii), and summarized in Theorem A, (ii), below.

Another important property of the multiradial representation discussed above concerns the relationship between the three main components — i.e., roughly speaking, *log-shells*, *splitting monoids of LGP-monoids*, and *number fields* — of this multiradial representation and the  $\Theta_{\text{LGP}}^{\times\mu}$ -links, i.e., the *horizontal arrows* of the log-theta-lattice under consideration. This property — which may be thought of as a property of **compatibility** with the  $\Theta_{\text{LGP}}^{\times\mu}$ -link — asserts that the *cyclotomic rigidity isomorphisms* that appear in the Kummer theory concerning the splitting monoids of LGP-monoids and copies of  $F_{\text{mod}}^\times$  are *immune* to the  $\widehat{\mathbb{Z}}^\times$ -indeterminacies that act on the copies of “ $\mathcal{O}^{\times\mu}$ ” that arise in the  $\mathcal{F}^{+\times\mu}$ -prime-strips that appear in the  $\Theta_{\text{LGP}}^{\times\mu}$ -link. In the case of splitting monoids of LGP-monoids, this property amounts precisely to the *multiradiality* theory developed in §2 [cf. the above discussion], i.e., in essence, to the **mono-theta-theoretic cyclotomic rigidity** property reviewed in the above discussion. In the case of copies of  $F_{\text{mod}}^\times$ , this property follows from the theory surrounding the construction of the cyclotomic rigidity isomorphism discussed in [IUTchI], Example 5.1, (v), together with the well-known fact in elementary algebraic number theory that any nonzero element of a number field that is **integral** at every valuation of the number field is necessarily a **root of unity**. These compatibility properties are described in detail in Theorem 3.11, (iii), and summarized in Theorem A, (iii), below.




Fig. I.6: The full multiradial representation

At this point, we pause to observe that although considerable attention has been devoted so far in the present series of papers, especially in [IUTchII], to



the theory of *Gaussian monoids*, not so much attention has been devoted [i.e., outside of [IUTchI], §5; [IUTchII], Corollaries 4.7, 4.8] to [the multiplicative monoids constituted by] copies of  $F_{\text{mod}}^\times$ . These copies of  $F_{\text{mod}}^\times$  enter into the theory of the *multiradial representation* discussed above in the form of various types of *global Frobenioids* in the following way. If one starts from the *number field*  $F_{\text{mod}}$ , one natural Frobenioid that can be associated to  $F_{\text{mod}}$  is the Frobenioid  $\mathcal{F}_{\text{mod}}^\circledast$  of [stack-theoretic] *arithmetic line bundles* on [the spectrum of the ring of integers of]  $F_{\text{mod}}$  discussed in [IUTchI], Example 5.1, (iii) [cf. also Example 3.6 of the present paper]. From the point of view of the theory surrounding the *multiradial representation* discussed above, there are *two natural ways* to approach the construction of “ $\mathcal{F}_{\text{mod}}^\circledast$ ”:

- ( $\circledast_{\text{MOD}}$ ) (**Rational Function Torsor Version**): This approach consists of considering the category  $\mathcal{F}_{\text{MOD}}^\circledast$  of  $F_{\text{mod}}^\times$ -torsors equipped with *trivializations* at each  $\underline{v} \in \underline{\mathbb{V}}$  [cf. Example 3.6, (i), for more details].
- ( $\circledast_{\text{mod}}$ ) (**Local Fractional Ideal Version**): This approach consists of considering the category  $\mathcal{F}_{\text{mod}}^\circledast$  of collections of *integral structures* on the various completions  $K_{\underline{v}}$  at  $\underline{v} \in \underline{\mathbb{V}}$  and morphisms between such collections of integral structures that arise from multiplication by elements of  $F_{\text{mod}}^\times$  [cf. Example 3.6, (ii), for more details].

Then one has *natural isomorphisms of Frobenioids*

$$\mathcal{F}_{\text{mod}}^\circledast \xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^\circledast \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^\circledast$$

that induce the respective *identity morphisms*  $F_{\text{mod}}^\times \rightarrow F_{\text{mod}}^\times \rightarrow F_{\text{mod}}^\times$  on the associated *rational function monoids* [cf. [FrdI], Corollary 4.10]. In particular, at first glance,  $\mathcal{F}_{\text{MOD}}^\circledast$  and  $\mathcal{F}_{\text{mod}}^\circledast$  appear to be “essentially equivalent” objects.

On the other hand, when regarded from the point of view of the *multiradial representations* discussed above, these two constructions exhibit a number of significant differences — cf. Fig. I.7 below; the discussion of Remarks 3.6.2, 3.10.2. For instance, whereas the construction of ( $\circledast_{\text{MOD}}$ ) depends only on the **multiplicative** structure of  $F_{\text{mod}}^\times$ , the construction of ( $\circledast_{\text{mod}}$ ) involves the *module*, i.e., the **additive**, structure of the localizations  $K_{\underline{v}}$ . The global portion of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link (respectively, the  $\Theta_{\text{lgp}}^{\times\mu}$ -link) is, by definition [cf. Definition 3.8, (ii)], constructed by means of the *realification* of the Frobenioid that appears in the construction of ( $\circledast_{\text{MOD}}$ ) (respectively, ( $\circledast_{\text{mod}}$ )). This means that the construction of the global portion of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link — which is the version of the  $\Theta$ -link that is in fact ultimately used in the theory of the multiradial representation — depends only on the *multiplicative* monoid structure of a copy of  $F_{\text{mod}}^\times$ , together with the various valuation homomorphisms  $F_{\text{mod}}^\times \rightarrow \mathbb{R}$  associated to  $\underline{v} \in \underline{\mathbb{V}}$ . Thus, the *mutual compatibility* [discussed above] of copies of  $F_{\text{mod}}^\times$  with the *log-Kummer correspondence* implies that one may perform this construction of the global portion of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link in a fashion that is *immune* to the “upper semi-compatibility” indeterminacy (Ind3) [discussed above]. By contrast, the construction of ( $\circledast_{\text{mod}}$ ) involves integral structures on the underlying local *additive* modules “ $K_{\underline{v}}$ ”, i.e., from the point of view of the multiradial representation, integral structures on *log-shells* and *tensor packets* of log-shells, which *are* subject to the “upper semi-compatibility” indeterminacy

(Ind3) [discussed above]. In particular, the **log-Kummer** correspondence subjects the construction of  $(\odot_{\text{mod}})$  to “*substantial distortion*”. On the other hand, the essential role played by local integral structures in the construction of  $(\odot_{\text{mod}})$  enables one to compute the *global arithmetic degree* of the arithmetic line bundles constituted by objects of the category “ $\mathcal{F}_{\text{mod}}^{\odot}$ ” in terms of **log-volumes** on **log-shells** and **tensor packets** of log-shells [cf. Proposition 3.9, (iii)]. This property of the construction of  $(\odot_{\text{mod}})$  will play a *crucial role* in deriving the **explicit estimates** for such log-volumes that are obtained in Corollary 3.12 [cf. Theorem B below].

$\mathcal{F}_{\text{MOD}}^{\odot}$	$\mathcal{F}_{\text{mod}}^{\odot}$
biased toward <b>multiplicative</b> structures	biased toward <b>additive</b> structures
easily related to <b>value group/non-coric</b> portion “ $(-)^{\text{lf}\blacktriangleright}$ ” of $\Theta_{\text{LGP}}^{\times\mu}$ -link	easily related to <b>unit group/coric</b> portion “ $(-)^{\text{f}\times\mu}$ ” of $\Theta_{\text{LGP}}^{\times\mu}$ -/ $\Theta_{\text{lgp}}^{\times\mu}$ -link, i.e., <b>mono-analytic log-shells</b>
admits <b>precise log-Kummer</b> <b>correspondence</b>	only admits <b>“upper semi-compatible”</b> <b>log-Kummer correspondence</b>
<b>rigid</b> , but not suited to explicit computation	subject to <b>substantial distortion</b> , but suited to <b>explicit estimates</b>

Fig. I.7:  $\mathcal{F}_{\text{MOD}}^{\odot}$  versus  $\mathcal{F}_{\text{mod}}^{\odot}$

Thus, in summary, the natural isomorphism  $\mathcal{F}_{\text{MOD}}^{\odot} \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\odot}$  discussed above plays the important role, in the context of the *multiradial representation* discussed above, of *relating*

- the **multiplicative** structure of the global number field  $F_{\text{mod}}$  to the **additive** structure of  $F_{\text{mod}}$ ;
- the **unit group/coric** portion “ $(-)^{\text{f}\times\mu}$ ” of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link to the **value group/non-coric** portion “ $(-)^{\text{lf}\blacktriangleright}$ ” of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link.

Finally, in Corollary 3.12 [cf. also Theorem B below], we apply the *multiradial representation* discussed above to estimate certain *log-volumes* as follows. We begin by introducing some terminology [cf. Definition 3.8, (i)]. We shall refer to the object that arises in any of the versions [including *realifications*] of the global Frobenioid “ $\mathcal{F}_{\text{mod}}^{\odot}$ ” discussed above — such as, for instance, the realified global Frobenioid

that occurs in the *codomain* of the  $\Theta_{\text{gau}}^{\times\mu}/\Theta_{\text{LGP}}^{\times\mu}/\Theta_{\text{lgp}}^{\times\mu}$ -link — by considering the arithmetic divisor determined by the zero locus of the elements “ $q$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  as a *q-pilot object*. The **log-volume** of the **q-pilot object** will be denoted by

$$- |\log(\underline{q})| \in \mathbb{R}$$

— so  $|\log(\underline{q})| > 0$  [cf. Corollary 3.12; Theorem B]. In a similar vein, we shall refer to the object that arises in the realified global Frobenioid that occurs in the *domain* of the  $\Theta_{\text{gau}}^{\times\mu}/\Theta_{\text{LGP}}^{\times\mu}/\Theta_{\text{lgp}}^{\times\mu}$ -link by considering the arithmetic divisor determined by the zero locus of the collection of *theta values* “ $\{q^{j^2}\}_{j=1,\dots,l^*}$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  as a  $\Theta$ -*pilot object*. The **log-volume** of the **union** of the collection of **possible images** of the  $\Theta$ -**pilot object** in the **multiradial representation** — i.e., where we recall that these “possible images” are subject to the **indeterminacies** (Ind1), (Ind2), (Ind3) — will be denoted by

$$- |\log(\underline{\Theta})| \in \mathbb{R} \cup \{+\infty\}$$

[cf. Corollary 3.12; Theorem B]. Then the content of Corollary 3.12, Theorem B may be summarized, roughly speaking [cf. Remark 3.12.1, (ii)], as a result concerning the

**negativity of the  $\Theta$ -pilot log-volume**  $|\log(\underline{\Theta})|$ .

Relative to the analogy between the theory of the present series of papers and *complex/p-adic Teichmüller theory* [cf. [IUTchI], §I4], this result may be thought of as a statement to the effect that

*“the pair consisting of a number field equipped with an elliptic curve is **metrically hyperbolic**, i.e., has **negative curvature**”.*

That is to say, it may be thought of as a sort of analogue of the inequality

$$\chi_S = - \int_S d\mu_S < 0$$

arising from the classical **Gauss-Bonnet formula** on a hyperbolic Riemann surface of finite type  $S$  [where we write  $\chi_S$  for the *Euler characteristic* of  $S$  and  $d\mu_S$  for the Kähler metric on  $S$  determined by the *Poincaré metric* on the upper half-plane — cf. the discussion of Remark 3.12.3], or, alternatively, of the inequality

$$(1-p)(2g_X - 2) \leq 0$$

that arises by computing *global degrees of line bundles* in the context of the **Hasse invariant** that arises in  $p$ -adic Teichmüller theory [where  $X$  is a *smooth, proper hyperbolic curve* of genus  $g_X$  over the ring of Witt vectors of a perfect field of characteristic  $p$  which is *canonical* in the sense of  $p$ -adic Teichmüller theory — cf. the discussion of Remark 3.12.4, (v)].

The proof of Corollary 3.12 [i.e., Theorem B] is based on the following *fundamental observation*: the **multiradial representation** discussed above yields

**two tautologically equivalent ways to compute  
the  $q$ -pilot log-volume** —  $|\log(\underline{q})|$

— cf. Fig. I.8 below; Step (xi) of the proof of Corollary 3.12. That is to say, suppose that one starts with the  **$q$ -pilot object** in the  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  ${}_{1,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  at  $(1, 0)$ , which we think of as being represented, via the approach of  $(\otimes_{\text{mod}})$ , by means of the action of the various  $\underline{q}$ , for  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , on the **log-shells** that arise, via the **log-link**  ${}_{1,-1}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} {}_{1,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , from the various local “ $\mathcal{O}^{\times\mu}$ ’s” in the  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  ${}_{1,-1}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  at  $(1, -1)$ . Thus, if one considers the *value group* “ $(-)^{\text{lp}}$ ” and *unit group* “ $(-)^{\text{r}\times\mu}$ ” portions of the *codomain* of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link  ${}_{0,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\Theta_{\text{LGP}}^{\times\mu}} {}_{1,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  in the context of the *arithmetic holomorphic structure* of the vertical line  $(1, \circ)$ , this action on log-shells may be thought of as a somewhat *intricate “intertwining”* between these value group and unit group portions. On the other hand, the  $\Theta_{\text{LGP}}^{\times\mu}$ -link  ${}_{0,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\Theta_{\text{LGP}}^{\times\mu}} {}_{1,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  constitutes a sort of **gluing isomorphism** between the *arithmetic holomorphic structures* associated to the vertical lines  $(0, \circ)$  and  $(1, \circ)$  that is based on

**forgetting this intricate intertwining**, i.e., by working solely with *abstract isomorphisms of  $\mathcal{F}^{\text{lp}\times\mu}$ -prime-strips*.

Thus, in order to relate the arithmetic holomorphic structures, say, at  $(0, 0)$  and  $(1, 0)$ , one must apply the *multiradial representation* discussed above. That is to say, one starts by applying the theory of **bi-coric mono-analytic log-shells** given in Theorem 1.5. One then applies the **Kummer theory** surrounding the **splitting monoids of theta/Gaussian monoids** and copies of the **number field  $F_{\text{mod}}$** , which allows one to pass from the **Frobenius-like** versions of various objects that appear in — i.e., that are necessary in order to consider — the  $\Theta_{\text{LGP}}^{\times\mu}$ -link to the corresponding **étale-like** versions of these objects that appear in the *multiradial representation*. This passage from Frobenius-like versions to étale-like versions is referred to as the operation of **Kummer-detachment** [cf. Fig. I.8; Remark 1.5.4, (i)]. As discussed above, this operation of Kummer-detachment is possible precisely as a consequence of the **compatibility** of the multiradial representation with the  $\Theta_{\text{LGP}}^{\times\mu}$ -link, i.e., with the *indeterminacy* (Ind2). Moreover, since the log-theta-lattice is, as discussed above, *far from commutative*, in order to represent the various “**log-link-conjugates**” at  $(0, m)$  [for  $m \in \mathbb{Z}$ ] in terms that may be understood from the point of view of the arithmetic holomorphic structure at  $(1, 0)$ , one must work [not only with the Kummer isomorphisms at a *single*  $(0, m)$ , but rather with] the **entire log-Kummer correspondence**. In particular, one must take into account the *indeterminacy* (Ind3). Once one completes the operation of Kummer-detachment so as to obtain *vertically coric* versions of objects on the vertical line  $(0, \circ)$ , one then passes to *multiradial objects*, i.e., to the “final form” of the *multiradial representation*, by introducing the *indeterminacy* (Ind1), i.e., that arises from working with [*mono-analytic!*]  $\mathcal{D}^+$  - [as opposed to  $\mathcal{D}$ -!] *prime-strips*. Finally, one computes the **log-volume** of this “final form” multiradial representation of the  $\Theta$ -pilot object — i.e., subject to the *indeterminacies* (Ind1), (Ind2), (Ind3)! — and concludes the desired estimates from the *tautological observation* that

the **log-theta-lattice**, and, in particular, the “**gluing isomorphism**” constituted by the  $\Theta_{\text{LGP}}^{\times\mu}$ -link, were **constructed** precisely in such a way as to assure that this computation of  $|\log(\underline{\Theta})|$ , i.e., of possible log-volumes of the  $\Theta$ -pilot object, necessarily **contains**  $|\log(\underline{q})|$  [i.e., as a “possible log-volume of the  $\Theta$ -pilot object”]

— cf. Fig. I.8 below; Step (xi) of the proof of Corollary 3.12. That is to say, the “gluing isomorphism” constituted by the  $\Theta_{\text{LGP}}^{\times\mu}$ -link relates two distinct “*arithmetic holomorphic structures*”, i.e., two distinct *copies* of conventional ring/scheme theory, that are glued together precisely by means of a relation that identifies the  $\Theta$ -pilot object in the *domain* of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link with the  $q$ -pilot object in the *codomain* of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link. Thus, once one sets up such an *apparatus*, the computation of the log-volume of the  $\Theta$ -pilot object in the domain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link in terms of the  $q$ -pilot object in the codomain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link amounts — *tautologically!* — to the computation of the log-volume of the  $q$ -pilot object [in the codomain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link] in terms of *itself*, i.e., to a computation that reflects certain *intrinsic properties* of this  $q$ -pilot object. This is the content of Corollary 3.12 [i.e., Theorem B]. As discussed above, this sort of “*computation of intrinsic properties*” in the present context of a number field equipped with an elliptic curve may be regarded as analogous to the “*computations of intrinsic properties*” reviewed above in the classical complex and  $p$ -adic cases.




Fig. I.8: Two tautologically equivalent ways to compute the log-volume of the  $q$ -pilot object at  $(1, 0)$

We conclude the present Introduction with the following summaries of the *main results* of the present paper.

**Theorem A. (Multiradial Algorithms for Logarithmic Gaussian Procession Monoids)** *Fix a collection of initial  $\Theta$ -data  $(\overline{F}/F, X_F, l, \underline{C}_K, \underline{V}, \underline{\epsilon})$  as in [IUTchI], Definition 3.1. Let*

$$\{^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}\}_{n,m \in \mathbb{Z}}$$

*be a collection of distinct  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters [relative to the given initial  $\Theta$ -data] — which we think of as arising from a **LGP-Gaussian log-theta-lattice** [cf. Definition 3.8, (iii)]. For each  $n \in \mathbb{Z}$ , write*

$$^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$$

*for the  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater determined, up to isomorphism, by the various  $^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , where  $m \in \mathbb{Z}$ , via the **vertical coricity** of Theorem 1.5, (i) [cf. Remark 3.8.2].*

(i) **(Multiradial Representation)** *Write*

$$^{n,\circ}\mathfrak{R}^{\text{LGP}}$$

*for the collection of data consisting of*

- (a) **tensor packets of log-shells;**
- (b) **splitting monoids of LGP-monoids** acting on the tensor packets of (a);
- (c) *copies, labeled by  $j \in \mathbb{F}_l^*$ , of [the multiplicative monoid of nonzero elements of] the **number field**  $F_{\text{mod}}$  acting on the tensor packets of (a)*

*[cf. Theorem 3.11, (i), (a), (b), (c), for more details] regarded up to **indeterminacies** of the following two types:*

- (Ind1) *the indeterminacies induced by the **automorphisms** of the **procession of  $\mathcal{D}^+$ -prime-strips**  $\text{Prc}^{(n,\circ)\mathcal{D}_T^+}$  that gives rise to the tensor packets of (a);*
- (Ind2) *the indeterminacies that arise from the **automorphisms of the  $\mathcal{F}^{+\times\mu}$ -prime-strips** that appear in the  $\Theta_{\text{LGP}}^{\times\mu}$ -link, i.e., in particular, at [for simplicity]  $\underline{v} \in \underline{V}^{\text{non}}$ , the  $\widehat{\mathbb{Z}}^{\times}$ -**indeterminacies** acting on local copies of “ $\mathcal{O}^{\times\mu}$ ”*

— *cf. Theorem 3.11, (i), for more details. Then  $^{n,\circ}\mathfrak{R}^{\text{LGP}}$  may be constructed via an **algorithm** in the procession of  $\mathcal{D}^+$ -prime-strips  $\text{Prc}^{(n,\circ)\mathcal{D}_T^+}$ , which is **functorial** with respect to isomorphisms of processions of  $\mathcal{D}^+$ -prime-strips. For  $n, n' \in \mathbb{Z}$ , the **permutation symmetries** of the **étale-picture** discussed in [IUTchI], Corollary 6.10, (iii); [IUTchII], Corollary 4.11, (ii), (iii) [cf. also Corollary 2.3, (ii); Remark 3.8.2, of the present paper], induce **compatible poly-isomorphisms***

$$\text{Prc}^{(n,\circ)\mathcal{D}_T^+} \xrightarrow{\sim} \text{Prc}^{(n',\circ)\mathcal{D}_T^+}; \quad ^{n,\circ}\mathfrak{R}^{\text{LGP}} \xrightarrow{\sim} ^{n',\circ}\mathfrak{R}^{\text{LGP}}$$



which are, moreover, compatible with the **bi-coricity** poly-isomorphisms

$${}^{n,\circ}\mathcal{D}_0^+ \xrightarrow{\sim} {}^{n',\circ}\mathcal{D}_0^+$$

of Theorem 1.5, (iii) [cf. also [IUTchII], Corollaries 4.10, (iv); 4.11, (i)].

(ii) (**log-Kummer Correspondence**) For  $n, m \in \mathbb{Z}$ , the inverses of the **Kummer isomorphisms** associated to the various  **$\mathcal{F}$ -prime-strips** and **NF-bridges** that appear in the  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$   ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  induce “inverse Kummer” **isomorphisms** between the **vertically coric data** (a), (b), (c) of (i) and the corresponding **Frobenioid-theoretic data** arising from each  $\Theta^{\pm\text{ell}}\text{NF-Hodge theater}$   ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  [cf. Theorem 3.11, (ii), (a), (b), (c), for more details]. Moreover, as one varies  $m \in \mathbb{Z}$ , the corresponding **Kummer isomorphisms** [i.e., inverses of “inverse Kummer” isomorphisms] of **splitting monoids** of **LGP-monoids** [cf. (i), (b)] and labeled copies of the **number field**  $F_{\text{mod}}$  [cf. (i), (c)] are **mutually compatible**, relative to the **log-links** of the  $n$ -th column of the **LGP-Gaussian log-theta-lattice** under consideration, in the sense that the only portions of the [Frobenioid-theoretic] domains of these Kummer isomorphisms that are related to one another via the **log-links** consist of **roots of unity** [multiplication by which corresponds, via the **log-link**, to an “**addition by zero**” indeterminacy, i.e., to **no indeterminacy!**] — cf. Proposition 3.5, (ii), (c); Proposition 3.10, (ii); Theorem 3.11, (ii), for more details. On the other hand, the Kummer isomorphisms of **tensor packets** of **log-shells** [cf. (i), (a)] are subject to a certain “**indeterminacy**” as follows:

(Ind3) as one varies  $m \in \mathbb{Z}$ , these Kummer isomorphisms of tensor packets of log-shells are “**upper semi-compatible**”, relative to the **log-links** of the  $n$ -th column of the **LGP-Gaussian log-theta-lattice** under consideration, in a sense that involves certain **natural inclusions** “ $\subseteq$ ” at  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$  and certain **natural surjections** “ $\twoheadrightarrow$ ” at  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$  — cf. Proposition 3.5, (ii), (a), (b); Theorem 3.11, (ii), for more details.

Finally, as one varies  $m \in \mathbb{Z}$ , these Kummer isomorphisms of tensor packets of log-shells are [precisely!] **compatible**, relative to the **log-links** of the  $n$ -th column of the **LGP-Gaussian log-theta-lattice** under consideration, with the respective **log-volumes** [cf. Proposition 3.9, (iv)].

(iii) ( **$\Theta_{\text{LGP}}^{\times\mu}$ -Link Compatibility**) The various Kummer isomorphisms of (ii) satisfy compatibility properties with the various **horizontal arrows** — i.e.,  $\Theta_{\text{LGP}}^{\times\mu}$ -links — of the **LGP-Gaussian log-theta-lattice** under consideration as follows: The **tensor packets** of **log-shells** [cf. (i), (a)] are compatible, relative to the relevant Kummer isomorphisms, with [the unit group portion “ $(-)^{\times\mu}$ ” of] the  $\Theta_{\text{LGP}}^{\times\mu}$ -link [cf. the indeterminacy “(Ind2)” of (i)]; we refer to Theorem 3.11, (iii), (a), (b), for more details. The identity automorphism on the objects that appear in the construction of the **splitting monoids** of **LGP-monoids** via mono-theta environments [cf. (i), (b)] is compatible, relative to the relevant Kummer isomorphisms and isomorphisms of mono-theta environments, with the  $\Theta_{\text{LGP}}^{\times\mu}$ -link [cf. the indeterminacy “(Ind2)” of (i)]; we refer to Theorem 3.11, (iii), (c), for more details. The identity automorphism on the objects that appear in the construction of the

labeled copies of the **number field**  $F_{\text{mod}}$  [cf. (i), (c)] is compatible, relative to the relevant Kummer isomorphisms and cyclotomic rigidity isomorphisms [cf. the discussion of Remark 3.10.1, (ii); the construction of [IUTchI], Example 5.1, (v)], with the  $\Theta_{\text{LGP}}^{\times\mu}$ -link [cf. the indeterminacy “(Ind2)” of (i)]; we refer to Theorem 3.11, (iii), (d), for more details.

**Theorem B. (Log-volume Estimates for Multiradially Represented Splitting Monoids of Logarithmic Gaussian Procession Monoids)** *Suppose that we are in the situation of Theorem A. Write*

$$- |\log(\underline{\Theta})| \in \mathbb{R} \cup \{+\infty\}$$

for the **procession-normalized mono-analytic log-volume** [where the average is taken over  $j \in \mathbb{F}_l^*$  — cf. Remark 3.1.1, (ii); Proposition 3.9, (i), (ii); Theorem 3.11, (i), (a), for more details] of the **union** of the **possible images** of a  **$\Theta$ -pilot object** [cf. Definition 3.8, (i)], relative to the relevant **Kummer isomorphisms** [cf. Theorems A, (ii); 3.11, (ii)], in the **multiradial representation** of Theorems A, (i); 3.11, (i), which we regard as **subject to the indeterminacies** (Ind1), (Ind2), (Ind3) described in Theorems A, (i), (ii); 3.11, (i), (ii). Write

$$- |\log(\underline{q})| \in \mathbb{R}$$

for the **procession-normalized mono-analytic log-volume** of the image of a  **$q$ -pilot object** [cf. Definition 3.8, (i)], relative to the relevant **Kummer isomorphisms** [cf. Theorems A, (ii); 3.11, (ii)], in the **multiradial representation** of Theorems A, (i); 3.11, (i), which we do **not** regard as subject to the indeterminacies (Ind1), (Ind2), (Ind3) described in Theorems A, (i), (ii); 3.11, (i), (ii). Here, we recall the definition of the symbol “ $\Delta$ ” as the result of identifying the labels

$$“0” \text{ and } “\langle \mathbb{F}_l^* \rangle”$$

[cf. [IUTchII], Corollary 4.10, (i)]. In particular,  $|\log(\underline{q})| > 0$  is easily computed in terms of the various  **$q$ -parameters** of the elliptic curve  $E_F$  [cf. [IUTchI], Definition 3.1, (b)] at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}} (\neq \emptyset)$ . Then it holds that

$$C_{\Theta} \geq -1$$

for any real number  $C_{\Theta} \in \mathbb{R}$  such that

$$- |\log(\underline{\Theta})| \leq C_{\Theta} \cdot |\log(\underline{q})| \quad (*_{C_{\Theta}})$$

[i.e.,  $- |\log(\underline{\Theta})| \in \mathbb{R} \subseteq \mathbb{R} \cup \{+\infty\}$  and satisfies the inequality  $(*_{C_{\Theta}})$ ].

### Acknowledgements:

I would like to thank *Fumiharu Kato* and *Akio Tamagawa* for many helpful discussions concerning the material presented in this paper.

### Notations and Conventions:

We shall continue to use the “Notations and Conventions” of [IUTchI], §0.

## Section 1: The Log-theta-lattice

In the present §1, we discuss various enhancements to the theory of **log-shells**, as developed in [AbsTopIII]. In particular, we develop the theory of the **log-link** [cf. Definition 1.1; Propositions 1.2, 1.3], which, together with the  $\Theta^{\times\mu}$ - and  $\Theta_{\text{gau}}^{\times\mu}$ -links of [IUTchII], Corollary 4.10, (iii), leads naturally to the construction of the **log-theta-lattice**, an apparatus that is *central* to the theory of the present series of papers. We conclude the present §1 with a discussion of various **coric structures** associated to the log-theta-lattice [cf. Theorem 1.5].

In the following discussion, we assume that we have been given *initial*  $\Theta$ -data as in [IUTchI], Definition 3.1. We begin by reviewing various aspects of the theory of *log-shells* developed in [AbsTopIII].

**Definition 1.1.** Let

$$\dagger\mathfrak{F} = \{\dagger\mathcal{F}_{\underline{v}}\}_{\underline{v} \in \mathbb{V}}$$

be an  $\mathcal{F}$ -*prime-strip* [relative to the given initial  $\Theta$ -data — cf. [IUTchI], Definition 5.2, (i)]. Write

$$\dagger\mathfrak{F}^+ = \{\dagger\mathcal{F}_{\underline{v}}^+\}_{\underline{v} \in \mathbb{V}}; \quad \dagger\mathfrak{F}^{+\times\mu} = \{\dagger\mathcal{F}_{\underline{v}}^{+\times\mu}\}_{\underline{v} \in \mathbb{V}}$$

for the associated  $\mathcal{F}^+$ -,  $\mathcal{F}^{+\times\mu}$ -*prime-strips* [cf. [IUTchI], Remark 5.2.1, (ii); [IUTchII], Definition 4.9, (vi), (vii)]. Recall the *functorial algorithm* of [IUTchII], Corollary 4.6, (i), in the  $\mathcal{F}$ -prime-strip  $\dagger\mathfrak{F}$  for constructing the assignment  $\Psi_{\text{cns}}(\dagger\mathfrak{F})$  given by

$$\begin{aligned} \mathbb{V}^{\text{non}} \ni \underline{v} &\mapsto \Psi_{\text{cns}}(\dagger\mathfrak{F})_{\underline{v}} \stackrel{\text{def}}{=} \left\{ G_{\underline{v}}(\dagger\Pi_{\underline{v}}) \curvearrowright \Psi_{\dagger\mathcal{F}_{\underline{v}}} \right\} \\ \mathbb{V}^{\text{arc}} \ni \underline{v} &\mapsto \Psi_{\text{cns}}(\dagger\mathfrak{F})_{\underline{v}} \stackrel{\text{def}}{=} \Psi_{\dagger\mathcal{F}_{\underline{v}}} \end{aligned}$$

— where the data in brackets “ $\{-\}$ ” is to be regarded as being well-defined only up to a  $\dagger\Pi_{\underline{v}}$ -*conjugacy indeterminacy* [cf. [IUTchII], Corollary 4.6, (i), for more details]. In the following, we shall write

$$(-)^{\text{gp}} \stackrel{\text{def}}{=} (-)^{\text{gp}} \bigcup \{0\}$$

for the *formal union* with  $\{0\}$  of the groupification  $(-)^{\text{gp}}$  of a monoid “ $(-)$ ”. Thus, by setting the product of all elements of  $(-)^{\text{gp}}$  with 0 to be equal to 0, one obtains a natural monoid structure on  $(-)^{\text{gp}}$ .

(i) Let  $\underline{v} \in \mathbb{V}^{\text{non}}$ . Write

$$(\Psi_{\dagger\mathcal{F}_{\underline{v}}} \supseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times} \rightarrow) \quad \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim} \stackrel{\text{def}}{=} (\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times})^{\text{pf}}$$

for the *perfection*  $(\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times})^{\text{pf}}$  of the submonoid of units  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times}$  of  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}$ . Now let us recall from the theory of [AbsTopIII] [cf. [AbsTopIII], Definition 3.1, (iv); [AbsTopIII], Proposition 3.2, (iii), (v)] that the natural, algorithmically constructible *topological field structure* on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  allows one to define a  $p_{\underline{v}}$ -*adic logarithm* on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ ,

which, in turn, yields a *functorial algorithm* in the Frobenioid  $\dagger\mathcal{F}_{\underline{v}}$  for constructing a *topological field structure* on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ . Write

$$\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})} \subseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$$

for the resulting *multiplicative monoid* of nonzero integers. Here, we observe that the resulting *diagram*

$$\Psi_{\dagger\mathcal{F}_{\underline{v}}} \supseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times} \rightarrow \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim} = \Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$$

is *compatible* with the various *natural actions* of  $\dagger\Pi_{\underline{v}} \rightarrow G_{\underline{v}}(\dagger\Pi_{\underline{v}})$  on each of the [four] “ $\Psi$ ’s” appearing in this diagram. The pair  $\{\dagger\Pi_{\underline{v}} \curvearrowright \Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}\}$  now determines a *Frobenioid*

$$\log(\dagger\mathcal{F}_{\underline{v}})$$

[cf. [AbsTopIII], Remark 3.1.1; [IUTchI], Remark 3.3.2] — which is, in fact, *naturally isomorphic* to the Frobenioid  $\dagger\mathcal{F}_{\underline{v}}$ , but which we wish to think of as being related to  $\dagger\mathcal{F}_{\underline{v}}$  via the *above diagram*. We shall denote this diagram by means of the notation

$$\dagger\mathcal{F}_{\underline{v}} \xrightarrow{\log} \log(\dagger\mathcal{F}_{\underline{v}})$$

and refer to this relationship between  $\dagger\mathcal{F}_{\underline{v}}$  and  $\log(\dagger\mathcal{F}_{\underline{v}})$  as the **tautological log-link** *associated to*  $\dagger\mathcal{F}_{\underline{v}}$  [or, when  $\dagger\mathfrak{F}$  is fixed, *at*  $\underline{v}$ ]. If  $\log(\dagger\mathcal{F}_{\underline{v}}) \xrightarrow{\sim} \ddagger\mathcal{F}_{\underline{v}}$  is any [poly-]isomorphism of Frobenioids, then we shall write

$$\dagger\mathcal{F}_{\underline{v}} \xrightarrow{\log} \ddagger\mathcal{F}_{\underline{v}}$$

for the diagram obtained by post-composing the tautological **log-link** associated to  $\dagger\mathcal{F}_{\underline{v}}$  with the given [poly-]isomorphism  $\log(\dagger\mathcal{F}_{\underline{v}}) \xrightarrow{\sim} \ddagger\mathcal{F}_{\underline{v}}$  and refer to this relationship between  $\dagger\mathcal{F}_{\underline{v}}$  and  $\ddagger\mathcal{F}_{\underline{v}}$  as a **log-link** from  $\dagger\mathcal{F}_{\underline{v}}$  to  $\ddagger\mathcal{F}_{\underline{v}}$ ; when the given [poly-]isomorphism  $\log(\dagger\mathcal{F}_{\underline{v}}) \xrightarrow{\sim} \ddagger\mathcal{F}_{\underline{v}}$  is the *full poly-isomorphism*, then we shall refer to the resulting **log-link** as the *full log-link* from  $\dagger\mathcal{F}_{\underline{v}}$  to  $\ddagger\mathcal{F}_{\underline{v}}$ . Finally, we recall from [AbsTopIII], Definition 3.1, (iv), that the image in  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$  of the submonoid of  $G_{\underline{v}}(\dagger\Pi_{\underline{v}})$ -invariants of  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times}$  constitutes a *compact topological module*, which we shall refer to as the *pre-log-shell*. Write  $p_{\underline{v}}^* \stackrel{\text{def}}{=} p_{\underline{v}}$  when  $p_{\underline{v}}$  is *odd* and  $p_{\underline{v}}^* \stackrel{\text{def}}{=} p_{\underline{v}}^2$  when  $p_{\underline{v}}$  is *even*. Then we shall refer to the result of multiplying the pre-log-shell by the factor  $(p_{\underline{v}}^*)^{-1}$  as the **log-shell**

$$\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}} \subseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim} = \Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$$

[cf. [AbsTopIII], Definition 5.4, (iii)]. In particular, by applying the natural, algorithmically constructible topological field structure on  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$  [cf. [AbsTopIII], Proposition 3.2, (iii)], it thus follows that one may think of this *log-shell* as an object associated to the *codomain* of *any* [that is to say, not necessarily tautological!] **log-link**

$$\dagger\mathcal{F}_{\underline{v}} \xrightarrow{\log} \ddagger\mathcal{F}_{\underline{v}}$$

— i.e., an object that is determined by the image of a certain portion [namely, the  $G_{\underline{v}}(\dagger\Pi_{\underline{v}})$ -invariants of  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times}$ ] of the *domain* of this **log-link**.

(ii) Let  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ . For  $N \in \mathbb{N}_{\geq 1}$ , write  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\mu_N} \subseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times} \subseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  for the subgroup of  $N$ -th roots of unity and  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim} \twoheadrightarrow \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  for the [pointed] universal covering of the topological group determined by the *groupification*  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  of the topological monoid  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}$ . Then one verifies immediately that one may think of the composite covering of topological groups

$$\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim} \twoheadrightarrow \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}} \twoheadrightarrow \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}} / \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\mu_N}$$

— where the second “ $\twoheadrightarrow$ ” is the natural surjection — as a [pointed] universal covering of  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}} / \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\mu_N}$ . That is to say, one may think of  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$  as an *object constructed from*  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}} / \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\mu_N}$  [cf. also Remark 1.2.1 below]. Now let us recall from the theory of [AbsTopIII] [cf. [AbsTopIII], Definition 4.1, (iv); [AbsTopIII], Proposition 4.2, (i), (ii)] that the natural, algorithmically constructible *topological field structure* on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  allows one to define a [complex archimedean] *logarithm* on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ , which, in turn, yields a *functorial algorithm* in the collection of data  $\dagger\mathcal{F}_{\underline{v}}$  [cf. [IUTchI], Definition 5.2, (i), (b)] for constructing a *topological field structure* on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ , together with a  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ -Kummer structure on  $\dagger\mathbb{U}_{\underline{v}}$  [cf. [AbsTopIII], Definition 4.1, (iv); [IUTchII], Proposition 4.4, (i)]. Write

$$\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})} \subseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$$

for the resulting *multiplicative monoid* of nonzero integers. Here, we observe that the resulting *diagram*

$$\Psi_{\dagger\mathcal{F}_{\underline{v}}} \subseteq \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}} \leftarrow \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim} = \Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$$

is *compatible* [cf. the discussion of [AbsTopIII], Definition 4.1, (iv)] with the *co-holomorphizations* determined by the *natural*  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$ -Kummer [cf. [IUTchII], Proposition 4.4, (i)] and  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ -Kummer [cf. the above discussion] structures on  $\dagger\mathbb{U}_{\underline{v}}$ . The triple of data consisting of the topological monoid  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}$ , the Aut-holomorphic space  $\dagger\mathbb{U}_{\underline{v}}$ , and the  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ -Kummer structure on  $\dagger\mathbb{U}_{\underline{v}}$  discussed above determines a *collection of data* [i.e., as in [IUTchI], Definition 5.2, (i), (b)]

$$\log(\dagger\mathcal{F}_{\underline{v}})$$

which is, in fact, *naturally isomorphic* to the collection of data  $\dagger\mathcal{F}_{\underline{v}}$ , but which we wish to think of as being related to  $\dagger\mathcal{F}_{\underline{v}}$  via the *above diagram*. We shall denote this diagram by means of the notation

$$\dagger\mathcal{F}_{\underline{v}} \xrightarrow{\log} \log(\dagger\mathcal{F}_{\underline{v}})$$

and refer to this relationship between  $\dagger\mathcal{F}_{\underline{v}}$  and  $\log(\dagger\mathcal{F}_{\underline{v}})$  as the **tautological log-link** associated to  $\dagger\mathcal{F}_{\underline{v}}$  [or, when  $\dagger\mathfrak{F}$  is fixed, at  $\underline{v}$ ]. If  $\log(\dagger\mathcal{F}_{\underline{v}}) \xrightarrow{\sim} \dagger\mathfrak{F}_{\underline{v}}$  is any

[poly-]isomorphism of collections of data [i.e., as in [IUTchI], Definition 5.2, (i), (b)], then we shall write

$$\dagger \mathcal{F}_{\underline{v}} \xrightarrow{\text{log}} \ddagger \mathcal{F}_{\underline{v}}$$

for the diagram obtained by post-composing the tautological **log**-link associated to  $\dagger \mathcal{F}_{\underline{v}}$  with the given [poly-]isomorphism  $\text{log}(\dagger \mathcal{F}_{\underline{v}}) \xrightarrow{\sim} \ddagger \mathcal{F}_{\underline{v}}$  and refer to this relationship between  $\dagger \mathcal{F}_{\underline{v}}$  and  $\ddagger \mathcal{F}_{\underline{v}}$  as a **log-link** from  $\dagger \mathcal{F}_{\underline{v}}$  to  $\ddagger \mathcal{F}_{\underline{v}}$ ; when the given [poly-]isomorphism  $\text{log}(\dagger \mathcal{F}_{\underline{v}}) \xrightarrow{\sim} \ddagger \mathcal{F}_{\underline{v}}$  is the *full poly-isomorphism*, then we shall refer to the resulting **log-link** as the *full log-link* from  $\dagger \mathcal{F}_{\underline{v}}$  to  $\ddagger \mathcal{F}_{\underline{v}}$ . Finally, we recall from [AbsTopIII], Definition 4.1, (iv), that the submonoid of units  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\times} \subseteq \Psi_{\dagger \mathcal{F}_{\underline{v}}}$  determines a *compact topological subquotient* of  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim}$ , which we shall refer to as the *pre-log-shell*. We shall refer to the  $\Psi_{\text{log}(\dagger \mathcal{F}_{\underline{v}})}^{\times}$ -orbit of the [uniquely determined] closed line segment of  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim}$  which is *preserved by multiplication by  $\pm 1$*  and whose endpoints differ by a *generator* of the kernel of the natural surjection  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim} \rightarrow \Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\text{gp}}$  — or, equivalently, the  $\Psi_{\text{log}(\dagger \mathcal{F}_{\underline{v}})}^{\times}$ -orbit of the result of *multiplying by  $N$*  the [uniquely determined] closed line segment of  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim}$  which is *preserved by multiplication by  $\pm 1$*  and whose endpoints differ by a *generator* of the kernel of the natural surjection  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim} \rightarrow \Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\text{gp}} / \Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\mu_N}$  — as the **log-shell**

$$\mathcal{I}_{\dagger \mathcal{F}_{\underline{v}}} \subseteq \Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim} = \Psi_{\text{log}(\dagger \mathcal{F}_{\underline{v}})}^{\text{gp}}$$

[cf. [AbsTopIII], Definition 5.4, (v)]. Thus, one may think of the *log-shell* as an *object constructed from  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\text{gp}} / \Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\mu_N}$* . Moreover, by applying the natural, algorithmically constructible topological field structure on  $\Psi_{\text{log}(\dagger \mathcal{F}_{\underline{v}})}^{\text{gp}}$  ( $= \Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim}$ ), it thus follows that one may think of this *log-shell* as an object associated to the *codomain* of *any* [that is to say, not necessarily tautological!] **log-link**

$$\dagger \mathcal{F}_{\underline{v}} \xrightarrow{\text{log}} \ddagger \mathcal{F}_{\underline{v}}$$

— i.e., an object that is determined by the image of a certain portion [namely, the *subquotient*  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\times}$  of  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim}$ ] of the *domain* of this **log-link**.

(iii) Write

$$\underline{\text{log}}(\dagger \mathfrak{F}) \stackrel{\text{def}}{=} \left\{ \underline{\text{log}}(\dagger \mathcal{F}_{\underline{v}}) \stackrel{\text{def}}{=} \Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim} \right\}_{\underline{v} \in \underline{\mathbb{V}}}$$

for the collection of topological modules constructed in (i), (ii) above indexed by  $\underline{v} \in \underline{\mathbb{V}}$  — where the group structure arises from the *additive* portion of the field structures on  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim}$  discussed in (i), (ii); for  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , we regard  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\sim}$  as equipped with its natural  $G_{\underline{v}}(\dagger \Pi_{\underline{v}})$ -*action*. Write

$$\text{log}(\dagger \mathfrak{F}) \stackrel{\text{def}}{=} \{ \text{log}(\dagger \mathcal{F}_{\underline{v}}) \}_{\underline{v} \in \underline{\mathbb{V}}}$$

for the  $\mathcal{F}$ -prime-strip determined by the data  $\text{log}(\dagger \mathcal{F}_{\underline{v}})$  constructed in (i), (ii) for  $\underline{v} \in \underline{\mathbb{V}}$ . We shall denote by

$$\dagger \mathfrak{F} \xrightarrow{\text{log}} \text{log}(\dagger \mathfrak{F})$$



the collection of diagrams  $\{\dagger\mathcal{F}_v \xrightarrow{\log} \mathbf{log}(\dagger\mathcal{F}_v)\}_{v \in \underline{\mathbb{V}}}$  constructed in (i), (ii) for  $v \in \underline{\mathbb{V}}$  and refer to this relationship between  $\dagger\mathfrak{F}$  and  $\mathbf{log}(\dagger\mathfrak{F})$  as the **tautological log-link associated to  $\dagger\mathfrak{F}$** . If  $\mathbf{log}(\dagger\mathfrak{F}) \xrightarrow{\sim} \ddagger\mathfrak{F}$  is any [poly-]isomorphism of  $\mathcal{F}$ -prime-strips, then we shall write

$$\dagger\mathfrak{F} \xrightarrow{\log} \ddagger\mathfrak{F}$$

for the diagram obtained by post-composing the tautological **log-link** associated to  $\dagger\mathfrak{F}$  with the given [poly-]isomorphism  $\mathbf{log}(\dagger\mathfrak{F}) \xrightarrow{\sim} \ddagger\mathfrak{F}$  and refer to this relationship between  $\dagger\mathfrak{F}$  and  $\ddagger\mathfrak{F}$  as a **log-link** from  $\dagger\mathfrak{F}$  to  $\ddagger\mathfrak{F}$ ; when the given [poly-]isomorphism  $\mathbf{log}(\dagger\mathfrak{F}) \xrightarrow{\sim} \ddagger\mathfrak{F}$  is the *full poly-isomorphism*, then we shall refer to the resulting **log-link** as the *full log-link* from  $\dagger\mathfrak{F}$  to  $\ddagger\mathfrak{F}$ . Finally, we shall write

$$\mathcal{I}_{\dagger\mathfrak{F}} \stackrel{\text{def}}{=} \{\mathcal{I}_{\dagger\mathcal{F}_v}\}_{v \in \underline{\mathbb{V}}}$$

for the collection of log-shells constructed in (i), (ii) for  $v \in \underline{\mathbb{V}}$  and refer to this collection as the **log-shell associated to  $\dagger\mathfrak{F}$**  and [by a slight abuse of notation]

$$\mathcal{I}_{\dagger\mathfrak{F}} \subseteq \underline{\mathbf{log}}(\dagger\mathfrak{F})$$

for the collection of natural inclusions indexed by  $v \in \underline{\mathbb{V}}$ . In particular, [cf. the discussion of (i), (ii)], it thus follows that one may think of this *log-shell* as an object associated to the *codomain* of *any* [that is to say, not necessarily tautological!] **log-link**

$$\dagger\mathfrak{F} \xrightarrow{\log} \ddagger\mathfrak{F}$$

— i.e., an object that is determined by the image of a certain portion [cf. the discussion of (i), (ii)] of the *domain* of this **log-link**.

(iv) Let  $v \in \underline{\mathbb{V}}^{\text{non}}$ . Then observe that it follows immediately from the constructions of (i) that the *topological modules with  $G_v(\dagger\Pi_v)$ -action  $\mathcal{I}_{\dagger\mathcal{F}_v} \subseteq \underline{\mathbf{log}}(\dagger\mathcal{F}_v)$*  may be constructed *solely* from the collection of data  $\dagger\mathcal{F}_v^{+ \times \mu}$  [i.e., the portion of the  $\mathcal{F}^{+ \times \mu}$ -prime-strip  $\dagger\mathfrak{F}^{+ \times \mu}$  labeled by  $v$ ]. That is to say, in light of the definition of a  $\times \mu$ -Kummer structure [cf. [IUTchII], Definition 4.9, (i), (ii), (iv), (vi), (vii)], these constructions only require the *perfection* “ $(-)^{\text{pf}}$ ” of the units and are *manifestly unaffected* by the operation of forming the quotient by a torsion subgroup of the units. Write

$$\mathcal{I}_{\dagger\mathcal{F}_v^{+ \times \mu}} \subseteq \underline{\mathbf{log}}(\dagger\mathcal{F}_v^{+ \times \mu})$$

for the resulting topological modules with  $G_v(\dagger\Pi_v)$ -action, *regarded as objects constructed from  $\dagger\mathcal{F}_v^{+ \times \mu}$* .

(v) Let  $v \in \underline{\mathbb{V}}^{\text{arc}}$ . Then by applying the algorithms for constructing “ $k \sim (G)$ ”, “ $\mathcal{I}(G)$ ” given in [AbsTopIII], Proposition 5.8, (v), to the [object of the category “ $\text{TMI}^+$ ” of split topological monoids discussed in [IUTchI], Example 3.4, (ii), determined by the] *split Frobenioid* portion of the collection of data  $\dagger\mathcal{F}_v^+$ , one obtains a *functorial algorithm* in the collection of data  $\dagger\mathcal{F}_v^+$  for constructing a *topological module  $\underline{\mathbf{log}}(\dagger\mathcal{F}_v^+)$*  [i.e., corresponding to “ $k \sim (G)$ ”] and a *topological subspace  $\mathcal{I}_{\dagger\mathcal{F}_v^+}$*  [i.e., corresponding to “ $\mathcal{I}(G)$ ”]. In fact, this functorial algorithm only makes use of the *unit portion* of this split Frobenioid, together with a *pointed universal covering*

of this unit portion. Moreover, by arguing as in (ii), one may in fact regard this functorial algorithm as an algorithm that only makes use of the *quotient of this unit portion by its  $N$ -torsion subgroup*, for  $N \in \mathbb{N}_{\geq 1}$ , together with a *pointed universal covering* of this quotient. That is to say, this functorial algorithm may, in fact, be regarded as a *functorial algorithm* in the collection of data  $\dagger \mathcal{F}_v^{\dagger \times \mu}$ . Write

$$\mathcal{I}_{\dagger \mathcal{F}_v^{\dagger \times \mu}} \subseteq \underline{\log}(\dagger \mathcal{F}_v^{\dagger \times \mu})$$

for the resulting topological module equipped with a closed subspace, *regarded as objects constructed from  $\dagger \mathcal{F}_v^{\dagger \times \mu}$* .

(vi) Finally, just as in (iii), we shall write

$$\mathcal{I}_{\dagger \mathfrak{F}^{\dagger \times \mu}} \stackrel{\text{def}}{=} \{\mathcal{I}_{\dagger \mathcal{F}_v^{\dagger \times \mu}}\}_{v \in \mathbb{V}} \subseteq \underline{\log}(\dagger \mathfrak{F}^{\dagger \times \mu}) \stackrel{\text{def}}{=} \{\underline{\log}(\dagger \mathcal{F}_v^{\dagger \times \mu})\}_{v \in \mathbb{V}}$$

for the resulting collections of data constructed *solely* from the  $\mathcal{F}^{\dagger \times \mu}$ -prime-strip  $\dagger \mathfrak{F}^{\dagger \times \mu}$  [i.e., which we do *not* regard as objects constructed from  $\dagger \mathfrak{F}!$ ].

From the point of view of the present series of papers, the theory of [AbsTopIII] may be summarized as follows.

**Proposition 1.2.** (**log-links Between  $\mathcal{F}$ -prime-strips**) *Let*

$$\dagger \mathfrak{F} = \{\dagger \mathcal{F}_v\}_{v \in \mathbb{V}}; \quad \ddagger \mathfrak{F} = \{\ddagger \mathcal{F}_v\}_{v \in \mathbb{V}}$$

be  $\mathcal{F}$ -prime-strips [relative to the given initial  $\Theta$ -data — cf. [IUTchI], Definition 5.2, (i)] and

$$\dagger \mathfrak{F} \xrightarrow{\log} \ddagger \mathfrak{F}$$

a **log-link** from  $\dagger \mathfrak{F}$  to  $\ddagger \mathfrak{F}$ . Write  $\dagger \mathfrak{F}^{\dagger \times \mu}$ ,  $\ddagger \mathfrak{F}^{\dagger \times \mu}$  for the associated  $\mathcal{F}^{\dagger \times \mu}$ -prime-strips [cf. [IUTchII], Definition 4.9, (vi), (vii)];  $\dagger \mathfrak{D}$ ,  $\ddagger \mathfrak{D}$  for the associated  $\mathcal{D}$ -prime-strips [cf. [IUTchI], Remark 5.2.1, (i)];  $\dagger \mathfrak{D}^{\dagger}$ ,  $\ddagger \mathfrak{D}^{\dagger}$  for the associated  $\mathcal{D}^{\dagger}$ -prime-strips [cf. [IUTchI], Definition 4.1, (iv)]. Also, let us recall the **diagrams**

$$\Psi_{\dagger \mathcal{F}_v} \supseteq \Psi_{\dagger \mathcal{F}_v}^{\times} \rightarrow \underline{\log}(\dagger \mathcal{F}_v) = \Psi_{\log(\dagger \mathcal{F}_v)}^{\text{gp}} \xrightarrow{\sim} \Psi_{\ddagger \mathcal{F}_v}^{\text{gp}} \quad (*_{\text{non}})$$

$$\Psi_{\dagger \mathcal{F}_v} \subseteq \Psi_{\dagger \mathcal{F}_v}^{\text{gp}} \leftarrow \underline{\log}(\dagger \mathcal{F}_v) = \Psi_{\log(\dagger \mathcal{F}_v)}^{\text{gp}} \xrightarrow{\sim} \Psi_{\ddagger \mathcal{F}_v}^{\text{gp}} \quad (*_{\text{arc}})$$

— where the  $v$  of  $(*_{\text{non}})$  (respectively,  $(*_{\text{arc}})$ ) belongs to  $\mathbb{V}^{\text{non}}$  (respectively,  $\mathbb{V}^{\text{arc}}$ ), and the [poly]-isomorphisms on the right are induced by the “ $\xrightarrow{\log}$ ” — of Definition 1.1, (i), (ii).

(i) (**Coricity of Associated  $\mathcal{D}$ -Prime-Strips**) *The log-link  $\dagger \mathfrak{F} \xrightarrow{\log} \ddagger \mathfrak{F}$  induces [poly]-isomorphisms*

$$\dagger \mathfrak{D} \xrightarrow{\sim} \ddagger \mathfrak{D}; \quad \dagger \mathfrak{D}^{\dagger} \xrightarrow{\sim} \ddagger \mathfrak{D}^{\dagger}$$

between the associated  $\mathcal{D}$ - and  $\mathcal{D}^+$ -prime-strips. In particular, the [poly-]isomorphism  $\dagger\mathcal{D} \xrightarrow{\sim} \ddagger\mathcal{D}$  induced by  $\dagger\mathfrak{F} \xrightarrow{\log} \ddagger\mathfrak{F}$  induces a [poly-]isomorphism

$$\Psi_{\text{cns}}(\dagger\mathcal{D}) \xrightarrow{\sim} \Psi_{\text{cns}}(\ddagger\mathcal{D})$$

between the collections of monoids equipped with auxiliary data of [IUTchII], Corollary 4.5, (i).

(ii) **(Simultaneous Compatibility with Ring Structures)** At  $\underline{v} \in \mathbb{V}^{\text{non}}$ , the **natural  $\dagger\Pi_{\underline{v}}$ -actions** on the “ $\Psi$ ’s” appearing in the diagram  $(*\text{non})$  are compatible with the topological ring structures on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  and  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$ . At  $\underline{v} \in \mathbb{V}^{\text{arc}}$ , the **co-holomorphizations** determined by the **natural  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$ - and  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$  (=  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\sim}$ )-Kummer structures** on  $\dagger\mathbb{U}_{\underline{v}}$  — which [cf. the discussion of Definition 1.1, (ii)] are compatible with the diagram  $(*\text{arc})$  — are compatible with the topological ring structures on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  and  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$ .

(iii) **(Simultaneous Compatibility with Log-volumes)** At  $\underline{v} \in \mathbb{V}^{\text{non}}$ , the diagram  $(*\text{non})$  is compatible with the **natural  $p_{\underline{v}}$ -adic log-volumes** [cf. [AbsTopIII], Proposition 5.7, (i); [AbsTopIII], Corollary 5.10, (ii)] on the subsets of  $\dagger\Pi_{\underline{v}}$ -invariants of  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$  and  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$ . At  $\underline{v} \in \mathbb{V}^{\text{arc}}$ , the diagram  $(*\text{arc})$  is compatible with the **natural angular log-volume** [cf. Remark 1.2.1 below; [AbsTopIII], Proposition 5.7, (ii); [AbsTopIII], Corollary 5.10, (ii)] on  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times}$  and the **natural radial log-volume** [cf. [AbsTopIII], Proposition 5.7, (ii); [AbsTopIII], Corollary 5.10, (ii)] on  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}^{\text{gp}}$ .

(iv) **(Kummer theory)** The **Kummer isomorphisms**

$$\Psi_{\text{cns}}(\dagger\mathfrak{F}) \xrightarrow{\sim} \Psi_{\text{cns}}(\dagger\mathcal{D}); \quad \Psi_{\text{cns}}(\ddagger\mathfrak{F}) \xrightarrow{\sim} \Psi_{\text{cns}}(\ddagger\mathcal{D})$$

of [IUTchII], Corollary 4.6, (i), **fail to be compatible** with the [poly-]isomorphism  $\Psi_{\text{cns}}(\dagger\mathcal{D}) \xrightarrow{\sim} \Psi_{\text{cns}}(\ddagger\mathcal{D})$  of (i), relative to the diagrams  $(*\text{non})$ ,  $(*\text{arc})$  [and the notational conventions of Definition 1.1] — cf. [AbsTopIII], Corollary 5.5, (iv). [Here, we regard the diagrams  $(*\text{non})$ ,  $(*\text{arc})$  as diagrams that relate  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}$  and  $\Psi_{\ddagger\mathcal{F}_{\underline{v}}}$ , via the [poly-]isomorphism  $\log(\dagger\mathfrak{F}) \xrightarrow{\sim} \ddagger\mathfrak{F}$  that determines the **log-link**  $\dagger\mathfrak{F} \xrightarrow{\log} \ddagger\mathfrak{F}$ .]

(v) **(Holomorphic Log-shells)** At  $\underline{v} \in \mathbb{V}^{\text{non}}$ , the **log-shell**

$$\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}} \subseteq \underline{\log}(\dagger\mathcal{F}_{\underline{v}}) \quad (\xrightarrow{\sim} \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}})$$

satisfies the following properties:  $(a_{\text{non}})$   $\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}}$  is **compact**, hence of **finite log-volume** [cf. [AbsTopIII], Corollary 5.10, (i)];  $(b_{\text{non}})$   $\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}}$  contains the submonoid of  $\dagger\Pi_{\underline{v}}$ -invariants of  $\Psi_{\log(\dagger\mathcal{F}_{\underline{v}})}$  [cf. [AbsTopIII], Definition 5.4, (iii)];  $(c_{\text{non}})$   $\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}}$  contains the image of the submonoid of  $\dagger\Pi_{\underline{v}}$ -invariants of  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\times}$ . At  $\underline{v} \in \mathbb{V}^{\text{arc}}$ , the **log-shell**

$$\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}} \subseteq \underline{\log}(\dagger\mathcal{F}_{\underline{v}}) \quad (\xrightarrow{\sim} \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}})$$

satisfies the following properties: ( $a_{\text{arc}}$ )  $\mathcal{I}_{\dagger\mathcal{F}_v}$  is **compact**, hence of **finite radial log-volume** [cf. [AbsTopIII], Corollary 5.10, (i)]; ( $b_{\text{arc}}$ )  $\mathcal{I}_{\dagger\mathcal{F}_v}$  contains  $\Psi_{\log(\dagger\mathcal{F}_v)}$  [cf. [AbsTopIII], Definition 5.4, (v)]; ( $c_{\text{arc}}$ ) the image of  $\mathcal{I}_{\dagger\mathcal{F}_v}$  in  $\Psi_{\dagger\mathcal{F}_v}^{\text{gp}}$  contains  $\Psi_{\dagger\mathcal{F}_v}^{\times}$  [i.e., in essence, the pre-log-shell].

(vi) (**Nonarchimedean Mono-analytic Log-shells**) At  $v \in \mathbb{V}^{\text{non}}$ , if we write  $\dagger\mathcal{D}_v^+ = \mathcal{B}^{\text{temp}}(\dagger G_v)^0$  for the portion of  $\dagger\mathcal{D}^+$  indexed by  $v$  [cf. the notation of [IUTchII], Corollary 4.5], then the algorithms for constructing “ $k^\sim(G)$ ”, “ $\mathcal{I}(G)$ ” given in [AbsTopIII], Proposition 5.8, (ii), yield a **functorial algorithm** in the category  $\dagger\mathcal{D}_v^+$  for constructing a topological module equipped with a continuous  $\dagger G_v$ -action

$$\underline{\log}(\dagger\mathcal{D}_v^+) \stackrel{\text{def}}{=} \left\{ \dagger G_v \curvearrowright k^\sim(\dagger G_v) \right\}$$

and a topological submodule — i.e., a “**mono-analytic log-shell**” —

$$\mathcal{I}_{\dagger\mathcal{D}_v^+} \stackrel{\text{def}}{=} \mathcal{I}(\dagger G_v) \subseteq k^\sim(\dagger G_v)$$

equipped with a  $p_v$ -adic log-volume [cf. [AbsTopIII], Corollary 5.10, (iv)]. Moreover, there is a natural **functorial algorithm** [cf. the second display of [IUTchII], Corollary 4.6, (ii)] in the collection of data  $\dagger\mathcal{F}_v^{+\times\mu}$  [i.e., the portion of  $\dagger\mathcal{F}^{+\times\mu}$  labeled by  $v$ ] for constructing an **Ism-orbit of isomorphisms** [cf. [IUTchII], Example 1.8, (iv); [IUTchII], Definition 4.9, (i), (vii)]

$$\underline{\log}(\dagger\mathcal{D}_v^+) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_v^{+\times\mu})$$

of topological modules [cf. Definition 1.1, (iv)], as well as a **functorial algorithm** [cf. [AbsTopIII], Corollary 5.10, (iv), (c), (d); the fourth display of [IUTchII], Corollary 4.5, (ii); the final display of [IUTchII], Corollary 4.6, (i)] in the collection of data  $\dagger\mathcal{F}_v$  for constructing **isomorphisms**

$$\underline{\log}(\dagger\mathcal{D}_v^+) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_v^{+\times\mu}) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_v) \quad \left( \xrightarrow{\sim} \Psi_{\dagger\mathcal{F}_v}^{\text{gp}} \right)$$

of topological modules. The various isomorphisms of the last two displays are **compatible** with one another, as well as with the respective  $\dagger G_v$ - and  $G_v(\dagger\Pi_v)$ -actions [relative to the natural identification  $\dagger G_v = G_v(\dagger\Pi_v)$  that arises from regarding  $\dagger\mathcal{D}_v^+$  as an object constructed from  $\dagger\mathcal{F}_v^{+\times\mu}$ ], the respective **log-shells**, and the respective **log-volumes** on these log-shells.

(vii) (**Archimedean Mono-analytic Log-shells**) At  $v \in \mathbb{V}^{\text{arc}}$ , the algorithms for constructing “ $k^\sim(G)$ ”, “ $\mathcal{I}(G)$ ” given in [AbsTopIII], Proposition 5.8, (v), yield a **functorial algorithm** in  $\dagger\mathcal{D}_v^+$  [regarded as an object of the category “ $\text{TM}^+$ ” of split topological monoids discussed in [IUTchI], Example 3.4, (ii)] for constructing a topological module

$$\underline{\log}(\dagger\mathcal{D}_v^+) \stackrel{\text{def}}{=} k^\sim(\dagger G_v)$$

and a topological subspace — i.e., a “**mono-analytic log-shell**” —

$$\mathcal{I}_{\dagger\mathcal{D}_v^+} \stackrel{\text{def}}{=} \mathcal{I}(\dagger G_v) \subseteq k^\sim(\dagger G_v)$$

equipped with angular and radial log-volumes [cf. [AbsTopIII], Corollary 5.10, (iv)]. Moreover, there is a natural **functorial algorithm** [cf. the second display of [IUTchII], Corollary 4.6, (ii)] in the collection of data  $\dagger\mathcal{F}_{\underline{v}}^{\dagger\times\mu}$  for constructing a **poly-isomorphism** [i.e., an orbit of isomorphisms with respect to the **independent** actions of  $\{\pm 1\}$  on each of the direct factors that occur in the construction of [AbsTopIII], Proposition 5.8, (v)]

$$\underline{\log}(\dagger\mathcal{D}_{\underline{v}}^{\dagger}) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_{\underline{v}}^{\dagger\times\mu})$$

of topological modules [cf. Definition 1.1, (v)], as well as a **functorial algorithm** [cf. [AbsTopIII], Corollary 5.10, (iv), (c), (d); the fourth display of [IUTchII], Corollary 4.5, (ii); the final display of [IUTchII], Corollary 4.6, (i)] in the collection of data  $\dagger\mathcal{F}_{\underline{v}}$  for constructing **poly-isomorphisms** [i.e., orbits of isomorphisms with respect to the **independent** actions of  $\{\pm 1\}$  on each of the direct factors that occur in the construction of [AbsTopIII], Proposition 5.8, (v)]

$$\underline{\log}(\dagger\mathcal{D}_{\underline{v}}^{\dagger}) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_{\underline{v}}^{\dagger\times\mu}) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_{\underline{v}}) \quad \left( \xrightarrow{\sim} \Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}} \right)$$

of topological modules. The various isomorphisms of the last two displays are **compatible** with one another, as well as with the respective **log-shells** and the respective **angular** and **radial log-volumes** on these log-shells.

(viii) (**Mono-analytic Log-shells**) The various [poly]-isomorphisms of (vi), (vii) [cf. also Definition 1.1, (iii), (vi)] yield collections of [poly]-isomorphisms indexed by  $\underline{v} \in \mathbb{V}$

$$\underline{\log}(\dagger\mathcal{D}^{\dagger}) \stackrel{\text{def}}{=} \{\underline{\log}(\dagger\mathcal{D}_{\underline{v}}^{\dagger})\}_{\underline{v} \in \mathbb{V}} \xrightarrow{\sim} \underline{\log}(\dagger\mathfrak{F}^{\dagger\times\mu}) \stackrel{\text{def}}{=} \{\underline{\log}(\dagger\mathcal{F}_{\underline{v}}^{\dagger\times\mu})\}_{\underline{v} \in \mathbb{V}}$$

$$\mathcal{I}_{\dagger\mathcal{D}^{\dagger}} \stackrel{\text{def}}{=} \{\mathcal{I}_{\dagger\mathcal{D}_{\underline{v}}^{\dagger}}\}_{\underline{v} \in \mathbb{V}} \xrightarrow{\sim} \mathcal{I}_{\dagger\mathfrak{F}^{\dagger\times\mu}} \stackrel{\text{def}}{=} \{\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}^{\dagger\times\mu}}\}_{\underline{v} \in \mathbb{V}}$$

$$\begin{aligned} \underline{\log}(\dagger\mathcal{D}^{\dagger}) &\xrightarrow{\sim} \underline{\log}(\dagger\mathfrak{F}^{\dagger\times\mu}) \xrightarrow{\sim} \underline{\log}(\dagger\mathfrak{F}) \stackrel{\text{def}}{=} \{\underline{\log}(\dagger\mathcal{F}_{\underline{v}})\}_{\underline{v} \in \mathbb{V}} \\ &\left( \xrightarrow{\sim} \Psi_{\text{cns}}^{\text{gp}}(\dagger\mathfrak{F}) \stackrel{\text{def}}{=} \{\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}\}_{\underline{v} \in \mathbb{V}} \right) \end{aligned}$$

$$\mathcal{I}_{\dagger\mathcal{D}^{\dagger}} \xrightarrow{\sim} \mathcal{I}_{\dagger\mathfrak{F}^{\dagger\times\mu}} \xrightarrow{\sim} \mathcal{I}_{\dagger\mathfrak{F}} \stackrel{\text{def}}{=} \{\mathcal{I}_{\dagger\mathcal{F}_{\underline{v}}}\}_{\underline{v} \in \mathbb{V}}$$

— where, in the definition of “ $\Psi_{\text{cns}}^{\text{gp}}(\dagger\mathfrak{F})$ ”, we regard each  $\Psi_{\dagger\mathcal{F}_{\underline{v}}}^{\text{gp}}$ , for  $\underline{v} \in \mathbb{V}^{\text{non}}$ , as being equipped with its natural  $G_{\underline{v}}(\dagger\Pi_{\underline{v}})$ -action [cf. the discussion at the beginning of Definition 1.1].

(ix) (**Coric Holomorphic Log-shells**) Let  $*\mathcal{D}$  be a  $\mathcal{D}$ -prime-strip; write

$$\mathfrak{F}(*\mathcal{D})$$

for the  $\mathcal{F}$ -prime-strip naturally determined by  $\Psi_{\text{cns}}(*\mathcal{D})$  [cf. [IUTchII], Remark 4.5.1, (i)]. Suppose that  $\dagger\mathfrak{F} = \dagger\mathfrak{F} = \mathfrak{F}(*\mathcal{D})$ , and that the given log-link  $\mathfrak{F}(*\mathcal{D}) = \dagger\mathfrak{F} \xrightarrow{\text{log}} \dagger\mathfrak{F} = \mathfrak{F}(*\mathcal{D})$  is the full log-link. Then there exists a **functorial algorithm**

in the  $\mathcal{D}$ -prime-strip  ${}^*\mathcal{D}$  for constructing a collection of topological subspaces — i.e., a collection of “**coric holomorphic log-shells**” —

$$\mathcal{I}_{*{}^*\mathcal{D}} \stackrel{\text{def}}{=} \mathcal{I}_{\ddagger\mathfrak{F}}$$

of the collection  $\Psi_{\text{cns}}^{\text{gp}}({}^*\mathcal{D})$ , which may be naturally identified with  $\Psi_{\text{cns}}^{\text{gp}}(\ddagger\mathfrak{F})$ , together with a collection of natural isomorphisms [cf. (viii); the fourth display of [IUTchII], Corollary 4.5, (ii)]

$$\mathcal{I}_{*{}^*\mathcal{D}^\dagger} \xrightarrow{\sim} \mathcal{I}_{*{}^*\mathcal{D}}$$

— where we write  ${}^*\mathcal{D}^\dagger$  for the  $\mathcal{D}^\dagger$ -prime-strip determined by  ${}^*\mathcal{D}$ .

(x) (**Frobenius-picture**) Let  $\{{}^n\mathfrak{F}\}_{n \in \mathbb{Z}}$  be a collection of distinct  $\mathcal{F}$ -prime-strips [relative to the given initial  $\Theta$ -data — cf. [IUTchI], Definition 5.2, (i)] indexed by the integers. Write  $\{{}^n\mathcal{D}\}_{n \in \mathbb{Z}}$  for the associated  $\mathcal{D}$ -prime-strips [cf. [IUTchI], Remark 5.2.1, (i)] and  $\{{}^n\mathcal{D}^\dagger\}_{n \in \mathbb{Z}}$  for the associated  $\mathcal{D}^\dagger$ -prime-strips [cf. [IUTchI], Definition 4.1, (iv)]. Then the full log-links  ${}^n\mathfrak{F} \xrightarrow{\text{log}} ({}^{n+1})\mathfrak{F}$ , for  $n \in \mathbb{Z}$ , give rise to an infinite chain

$$\dots \xrightarrow{\text{log}} ({}^{n-1})\mathfrak{F} \xrightarrow{\text{log}} {}^n\mathfrak{F} \xrightarrow{\text{log}} ({}^{n+1})\mathfrak{F} \xrightarrow{\text{log}} \dots$$

of log-linked  $\mathcal{F}$ -prime-strips which induces chains of full poly-isomorphisms

$$\dots \xrightarrow{\sim} {}^n\mathcal{D} \xrightarrow{\sim} ({}^{n+1})\mathcal{D} \xrightarrow{\sim} \dots \quad \text{and} \quad \dots \xrightarrow{\sim} {}^n\mathcal{D}^\dagger \xrightarrow{\sim} ({}^{n+1})\mathcal{D}^\dagger \xrightarrow{\sim} \dots$$

on the associated  $\mathcal{D}$ - and  $\mathcal{D}^\dagger$ -prime-strips. These chains may be represented symbolically as an oriented graph  $\vec{\Gamma}$  [cf. [AbsTopIII], §0]

$$\begin{array}{ccccccc} \dots & \rightarrow & \bullet & \rightarrow & \bullet & \rightarrow & \bullet & \rightarrow & \dots \\ & & & \searrow & \downarrow & \swarrow & & & \\ & & & & \circ & & & & \end{array}$$

— i.e., where the horizontal arrows correspond to the “ $\xrightarrow{\text{log}}$ ’s”; the “ $\bullet$ ’s” correspond to the “ ${}^n\mathfrak{F}$ ”; the “ $\circ$ ” corresponds to the “ ${}^n\mathcal{D}$ ”, identified up to isomorphism; the vertical/diagonal arrows correspond to the **Kummer isomorphisms** of (iv). This oriented graph  $\vec{\Gamma}$  admits a natural action by  $\mathbb{Z}$  [cf. [AbsTopIII], Corollary 5.5, (v)] — i.e., a **translation symmetry** — that fixes the “**core**”  $\circ$ , but it does **not admit arbitrary permutation symmetries**. For instance,  $\vec{\Gamma}$  does not admit an automorphism that switches two adjacent vertices, but leaves the remaining vertices fixed.

*Proof.* The various assertions of Proposition 1.2 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 1.2.1.** Suppose that we are in the situation of Definition 1.1, (ii). Then at the level of *metrics* — i.e., which give rise to **angular log-volumes** as in



Proposition 1.2, (iii) — we suppose that  $\Psi_{\dagger\mathcal{F}_v}^{\text{gp}}/\Psi_{\dagger\mathcal{F}_v}^{\mu N}$  is equipped with the metric obtained by descending the metric of  $\Psi_{\dagger\mathcal{F}_v}^{\text{gp}}$ , but we regard the object

$$\Psi_{\dagger\mathcal{F}_v}^{\text{gp}}/\Psi_{\dagger\mathcal{F}_v}^{\mu N} \text{ [or } \Psi_{\dagger\mathcal{F}_v}^{\times}/\Psi_{\dagger\mathcal{F}_v}^{\mu N}] \text{ as being equipped with a “weight } N\text{”}$$

— i.e., which has the effect of ensuring that the **log-volume** of  $\Psi_{\dagger\mathcal{F}_v}^{\times}/\Psi_{\dagger\mathcal{F}_v}^{\mu N}$  is **equal** to that of  $\Psi_{\dagger\mathcal{F}_v}^{\times}$ . That is to say, this convention concerning “weights” ensures that working with  $\Psi_{\dagger\mathcal{F}_v}^{\text{gp}}/\Psi_{\dagger\mathcal{F}_v}^{\mu N}$  does not have any effect on various computations of log-volume.

**Remark 1.2.2.** Before proceeding, we pause to consider the significance of the various properties discussed in Proposition 1.2, (v). For simplicity, we suppose that “ $\mathfrak{F}$ ” is the  $\mathcal{F}$ -prime-strip that arises from the data constructed in [IUTchI], Examples 3.2, (iii); 3.3, (i); 3.4, (i) [cf. [IUTchI], Definition 5.2, (i)].

(i) Suppose that  $v \in \mathbb{V}^{\text{non}}$ . Thus,  $K_v$  [cf. the notation of [IUTchI], Definition 3.1, (e)] is a **mixed-characteristic nonarchimedean local field**. Write  $k \stackrel{\text{def}}{=} K_v$ ,  $\mathcal{O}_k \subseteq k$  for the ring of integers of  $k$ ,  $\mathcal{O}_k^{\times} \subseteq \mathcal{O}_k$  for the group of units, and  $\log_k : \mathcal{O}_k^{\times} \rightarrow k$  for the  $p_v$ -adic logarithm. Then, at a more concrete level — i.e., relative to the notation of the present discussion — the **log-shell** “ $\mathcal{I}_{\dagger\mathcal{F}_v}$ ” corresponds to the submodule

$$\mathcal{I}_k \stackrel{\text{def}}{=} (p_v^*)^{-1} \cdot \log_k(\mathcal{O}_k^{\times}) \subseteq k$$

— where  $p_v^* = p_v$  if  $p_v$  is *odd*,  $p_v^* = p_v^2$  if  $p_v$  is *even* — while the properties  $(b_{\text{non}})$ ,  $(c_{\text{non}})$  of Proposition 1.2, (v), correspond, respectively, to the evident **inclusions**

$$\mathcal{O}_k^{\triangleright} \stackrel{\text{def}}{=} \mathcal{O}_k \setminus \{0\} \subseteq \mathcal{O}_k \subseteq \mathcal{I}_k; \quad \log_k(\mathcal{O}_k^{\times}) \subseteq \mathcal{I}_k$$

of subsets of  $k$ .

(ii) Suppose that  $v \in \mathbb{V}^{\text{arc}}$ . Thus,  $K_v$  [cf. the notation of [IUTchI], Definition 3.1, (e)] is a **complex archimedean field**. Write  $k \stackrel{\text{def}}{=} K_v$ ,  $\mathcal{O}_k \subseteq k$  for the subset of elements of absolute value  $\leq 1$ ,  $\mathcal{O}_k^{\times} \subseteq \mathcal{O}_k$  for the group of elements of absolute value  $= 1$ , and  $\exp_k : k \rightarrow k^{\times}$  for the *exponential map*. Then, at a more concrete level — i.e., relative to the notation of the present discussion — the **log-shell** “ $\mathcal{I}_{\dagger\mathcal{F}_v}$ ” corresponds to the subset

$$\mathcal{I}_k \stackrel{\text{def}}{=} \{a \in k \mid |a| \leq \pi\} \subseteq k$$

of elements of absolute value  $\leq \pi$ , while the properties  $(b_{\text{arc}})$ ,  $(c_{\text{arc}})$  of Proposition 1.2, (v), correspond, respectively, to the evident **inclusions**

$$\mathcal{O}_k^{\triangleright} \stackrel{\text{def}}{=} \mathcal{O}_k \setminus \{0\} \subseteq \mathcal{O}_k \subseteq \mathcal{I}_k; \quad \mathcal{O}_k^{\times} \subseteq \exp_k(\mathcal{I}_k)$$

— where we note the slightly different roles played, in the archimedean [cf. the present (ii)] and nonarchimedean [cf. (i)] cases, by the exponential and logarithmic functions, respectively [cf. [AbsTopIII], Remark 4.5.2].

(iii) The diagram represented by the oriented graph  $\vec{\Gamma}$  of Proposition 1.2, (x), is, of course, **far from commutative** [cf. Proposition 1.2, (iv)]! Ultimately, however, [cf. the discussion of Remark 1.4.1, (ii), below] we shall be interested in

- (a) constructing **invariants** *with respect to the  $\mathbb{Z}$ -action on  $\vec{\Gamma}$*  — i.e., in effect, constructing objects via functorial algorithms in the **coric**  $\mathcal{D}$ -prime-strips “ ${}^n\mathfrak{D}$ ” —

while, at the same time,

- (b) *relating the corically constructed objects* of (a) to the non-coric “ ${}^n\mathfrak{F}$ ” via the various **Kummer isomorphisms** of Proposition 1.2, (iv).

That is to say, from the point of view of (a), (b), the content of the *inclusions* discussed in (i) and (ii) above may be interpreted, at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , as follows:

the **coric holomorphic log-shells** of Proposition 1.2, (ix), contain *not only* the images, via the Kummer isomorphisms [i.e., the vertical/diagonal arrows of  $\vec{\Gamma}$ ], of the various “ $\mathcal{O}^\triangleright$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , *but also the images*, via the composite of the Kummer isomorphisms with the *various iterates of the log-link* [i.e., the horizontal arrows of  $\vec{\Gamma}$ ], of the portions of the various “ $\mathcal{O}^\triangleright$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  on which these iterates are defined.

An analogous statement in the case of  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$  may be formulated by adjusting the wording appropriately so as accommodate the latter portion of this statement, which corresponds to a certain *surjectivity* — we leave the routine details to the reader. Thus, although the diagram [corresponding to]  $\vec{\Gamma}$  fails to be *commutative*,

the *coric holomorphic log-shells* involved exhibit a sort of “**upper semi-commutativity**” with respect to containing/surjecting onto *the various images arising from composites of arrows in  $\vec{\Gamma}$* .

(iv) Note that although the diagram  $\vec{\Gamma}$  admits a natural “upper semi-commutativity” interpretation as discussed in (iii) above, it **fails to admit a corresponding “lower semi-commutativity”** interpretation. Indeed, such a “lower semi-commutativity” interpretation would amount to the existence of some sort of *collection of portions of the various “ $\mathcal{O}^\triangleright$ ’s” involved* [cf. the discussion of (i), (ii) above] — i.e., a sort of “**core**” — that are mapped to one another *isomorphically* by the various maps “ $\log_k$ ”/“ $\exp_k$ ” [cf. the discussion of (i), (ii) above] in a fashion that is **compatible** with the various **Kummer isomorphisms** that appear in the diagram  $\vec{\Gamma}$ . On the other hand, it is difficult to see how to construct such a collection of portions of the various “ $\mathcal{O}^\triangleright$ ’s” involved.

(v) Proposition 1.2, (iii), may be interpreted in the spirit of the discussion of (iii) above. That is to say, although the diagram corresponding to  $\vec{\Gamma}$  fails to be *commutative*, it is nevertheless “**commutative with respect to log-volumes**”, in the sense discussed in Proposition 1.2, (iii). This “commutativity with respect to log-volumes” allows one to work with log-volumes in a fashion that is *consistent*

with all composites of the various arrows of  $\vec{\Gamma}$ . Log-volumes will play an important role in the theory of §3, below, as a sort of *mono-analytic version* of the notion of the *degree of a global arithmetic line bundle* [cf. the theory of [AbsTopIII], §5].

(vi) As discussed in [AbsTopIII], §I3, the **log-links** of  $\vec{\Gamma}$  may be thought of as a sort of “**juggling of  $\boxplus, \boxtimes$** ” [i.e., of the two combinatorial dimensions of the ring structure constituted by addition and multiplication]. The “**arithmetic holomorphic structure**” constituted by the coric  $\mathcal{D}$ -prime-strips is *immune to this juggling*, and hence may be thought as representing a sort of **quotient of the horizontal arrow portion of  $\vec{\Gamma}$  by the action of  $\mathbb{Z}$**  [cf. (iii), (a)] — i.e., at the level of abstract oriented graphs, as a sort of “**oriented copy of  $\mathbb{S}^1$** ”. That is to say, the horizontal arrow portion of  $\vec{\Gamma}$  may be thought of as a sort of “**unraveling**” of this “oriented copy of  $\mathbb{S}^1$ ”, which is subject to the “*juggling of  $\boxplus, \boxtimes$* ” constituted by the  $\mathbb{Z}$ -action. Here, it is useful to recall that

- (a) the *Frobenius-like structures* constituted by the monoids that appear in the horizontal arrow portion of  $\vec{\Gamma}$  play the *crucial* role in the theory of the present series of papers of allowing one to construct such “**non-ring/scheme-theoretic filters**” as the  $\Theta$ -link [cf. the discussion of [IUTchII], Remark 3.6.2, (ii)].

By contrast,

- (b) the *étale-like structures* constituted by the coric  $\mathcal{D}$ -prime-strips play the *crucial* role in the theory of the present series of papers of allowing one to construct objects that are capable of “**functorially permeating**” such non-ring/scheme-theoretic filters as the  $\Theta$ -link [cf. the discussion of [IUTchII], Remark 3.6.2, (ii)].

Finally, in order to *relate* the theory of (a) to the theory of (b), one must avail oneself of **Kummer theory** [cf. (iii), (b), above].

<b>mono-anabelian</b> coric <i>étale-like</i> structures	<i>invariant differential</i> $d\theta$ on $\mathbb{S}^1$
<b>post-anabelian</b> <i>Frobenius-like</i> structures	<i>coordinate functions</i> $\int_{\bullet} d\theta$ on $\vec{\Gamma}$

Fig. 1.1: Analogy with the differential geometry of  $\mathbb{S}^1$

(vii) From the point of view of the discussion in (vi) above of the “*oriented copy of  $\mathbb{S}^1$* ” obtained by forming the quotient of the horizontal arrow portion of  $\vec{\Gamma}$  by  $\mathbb{Z}$ , one may think of the *coric étale-like structures* of Proposition 1.2, (i) — as well as the various objects constructed from these coric étale-like structures via the various *mono-anabelian algorithms* discussed in [AbsTopIII] — as corresponding to

the “*canonical invariant differential*  $d\theta$ ” on  $\mathbb{S}^1$  [which is, in particular, *invariant* with respect to the action of  $\mathbb{Z}$ !]. On the other hand, the various *post-anabelian Frobenius-like structures* obtained by *forgetting* the mono-anabelian algorithms applied to construct these objects — cf., e.g., the “ $\Psi_{\text{cns}}(\dagger\mathfrak{F})$ ” that appear in the *Kummer isomorphisms* of Proposition 1.2, (iv) — may be thought of as *coordinate functions* on the horizontal arrow portion of  $\vec{\Gamma}$  [which are *not* invariant with respect to the action of  $\mathbb{Z}$ !] of the form “ $\int_{\bullet} d\theta$ ” obtained by integrating the invariant differential  $d\theta$  along various paths of  $\vec{\Gamma}$  that emanate from some fixed vertex “ $\bullet$ ” of  $\vec{\Gamma}$ . This point of view is summarized in Fig. 1.1 above. Finally, we observe that this point of view is reminiscent of the discussion of [AbsTopIII], §I5, concerning the analogy between the theory of [AbsTopIII] and the construction of canonical coordinates via integration of Frobenius-invariant differentials in the classical  $p$ -adic theory.

### Remark 1.2.3.

(i) Observe that, relative to the notation of Remark 1.2.2, (i), any **multiplicative indeterminacy** with respect to the action on  $\mathcal{O}_k^\times$  of some *subgroup*  $H \subseteq \mathcal{O}_k^\times$  at some “ $\bullet$ ” of the diagram  $\vec{\Gamma}$  gives rise to an **additive indeterminacy** with respect to the action of  $\log_k(H)$  on the copy of “ $\mathcal{O}_k$ ” that corresponds to the subsequent “ $\bullet$ ” of the diagram  $\vec{\Gamma}$ . In particular, if  $H$  consists of *roots of unity*, then  $\log_k(H) = \{0\}$ , so *the resulting additive indeterminacy ceases to exist*. This observation will play a *crucial role* in the theory of §3, below, when it is applied in the context of the *constant multiple rigidity* properties constituted by the *canonical splittings of theta and Gaussian monoids* discussed in [IUTchII], Proposition 3.3, (i); [IUTchII], Corollary 3.5, (iii) [cf. also [IUTchII], Corollary 1.12, (ii); the discussion of [IUTchII], Remark 1.12.2, (iv)].

(ii) In the theory of §3, below, we shall consider *global arithmetic line bundles*. This amounts, in effect, to considering **multiplicative translates** by  $f \in F_{\text{mod}}^\times$  of the product of the various “ $\mathcal{O}_k^\times$ ” of Remark 1.2.2, (i), (ii), as  $\underline{v}$  ranges over the elements of  $\underline{\mathbb{V}}$ . Such translates are *disjoint* from one another, *except* in the case where  $f$  is a *unit* at all  $\underline{v} \in \underline{\mathbb{V}}$ . By elementary algebraic number theory [cf., e.g., [Lang], p. 144, the proof of Theorem 5], this corresponds precisely to the case where  $f$  is a *root of unity*. In particular, to consider quotients by this *multiplicative* action by  $F_{\text{mod}}^\times$  at one “ $\bullet$ ” of the diagram  $\vec{\Gamma}$  [where we allow  $\underline{v}$  to range over the elements of  $\underline{\mathbb{V}}$ ] gives rise to an **additive indeterminacy** by “*logarithms of roots of unity*” at the subsequent “ $\bullet$ ” of the diagram  $\vec{\Gamma}$ . In particular, at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , *the resulting additive indeterminacy ceases to exist* [cf. the discussion of (i); Definition 1.1, (iv)]; at  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ , the resulting indeterminacy corresponds to considering certain quotients of the copies of “ $\mathcal{O}_k^\times$ ” — i.e., of “ $\mathbb{S}^1$ ” — that appear by *some finite group* [cf. the discussion of Definition 1.1, (ii)]. These observations will be of use in the development of the theory of §3, below.

### Remark 1.2.4.

(i) At this point, we pause to recall the important observation that the **log-link** is **incompatible** with the **ring structures** of  $\Psi_{\dagger\mathcal{F}_v}^{\text{gp}}$  and  $\Psi_{\text{log}(\dagger\mathcal{F}_v)}^{\text{gp}}$  [cf. the notation

of Proposition 1.2, (ii)], in the sense that it does not arise from a *ring homomorphism* between these two rings. The barrier constituted by this incompatibility between the ring structures on either side of the **log-link** is precisely what is referred to as the “**log-wall**” in the theory of [AbsTopIII] [cf. the discussion of [AbsTopIII], §I4]. This incompatibility with the respective ring structures implies that it is not possible, *a priori*, to transport objects whose structure depends on these ring structures via the **log-link** by invoking the principle of “transport of structure”. From the point of view of the theory of the present series of papers, this means, in particular, that

the **log-wall** is **incompatible** with **conventional scheme-theoretic basepoints**, which are defined by means of geometric points [i.e., *ring homomorphisms* of a certain type]

— cf. the discussion of [IUTchII], Remark 3.6.3, (i); [AbsTopIII], Remark 3.7.7, (i). In this context, it is useful to recall that étale fundamental groups — i.e., Galois groups — are defined as certain *automorphism groups of fields/rings*; in particular, the definition of such a Galois group “as a certain automorphism group of some ring structure” is *incompatible*, in a quite essential way, with the **log-wall**. In a similar vein, **Kummer theory**, which depends on the *multiplicative structure* of the ring under consideration, is also *incompatible*, in a quite essential way, with the **log-wall** [cf. Proposition 1.2, (iv)]. That is to say, in the context of the **log-link**,

the only structure of interest that is manifestly **compatible** with the **log-link** [cf. Proposition 1.2, (i), (ii)] is the associated  **$\mathcal{D}$ -prime-strip**

— i.e., the *abstract topological groups* [isomorphic to “ $\Pi_{\underline{v}}$ ” — cf. the notation of [IUTchI], Definition 3.1, (e), (f)] at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  and *abstract Aut-holomorphic spaces* [isomorphic to “ $\mathcal{U}_{\underline{v}}$ ” — cf. the notation of [IUTchII], Proposition 4.3] at  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ . Indeed, this observation is precisely the *starting point* of the theory of [AbsTopIII] [cf. the discussion of [AbsTopIII], §I1, §I4].

(ii) Other important examples of structures which are *incompatible with the log-wall* include

- (a) the *additive structure* on the image of the Kummer map [cf. the discussion of [AbsTopIII], Remark 3.7.5];
- (b) in the “*birational*” situation — i.e., where one replaces “ $\Pi_{\underline{v}}$ ” by the absolute Galois group  $\Pi_{\underline{v}}^{\text{birat}}$  of the *function field* of the *affine curve* that gave rise to  $\Pi_{\underline{v}}$  — the datum of the *collection of closed points* that determines the affine curve [cf. [AbsTopIII], Remark 3.7.7, (ii)].

Note, for instance in the case of (b), that one may think of the additional datum under consideration as consisting of the natural outer surjection  $\Pi_{\underline{v}}^{\text{birat}} \twoheadrightarrow \Pi_{\underline{v}}$  that arises from the *scheme-theoretic morphism* from the spectrum of the function field to the given affine curve. On the other hand, just as in the case of the discussion of scheme-theoretic basepoints in (i), the construction of such an object  $\Pi_{\underline{v}}^{\text{birat}} \twoheadrightarrow \Pi_{\underline{v}}$  whose structure depends, in an essential way, on the scheme [i.e., ring!] structures

involved necessarily *fails to be compatible with the log-link* [cf. the discussion of [AbsTopIII], Remark 3.7.7, (ii)].

(iii) One way to understand the *incompatibility* discussed in (ii), (b), is as follows. Write  $\Delta_{\underline{v}}^{\text{birat}}$ ,  $\Delta_{\underline{v}}$  for the respective kernels of the natural surjections  $\Pi_{\underline{v}}^{\text{birat}} \twoheadrightarrow G_{\underline{v}}$ ,  $\Pi_{\underline{v}} \twoheadrightarrow G_{\underline{v}}$ . Then if one forgets about the *scheme-theoretic* basepoints discussed in (i),  $G_{\underline{v}}$ ,  $\Delta_{\underline{v}}^{\text{birat}}$ , and  $\Delta_{\underline{v}}$  may be understood on *both* sides of the **log-wall** as “*some topological group*”, and each of the topological groups  $\Delta_{\underline{v}}^{\text{birat}}$ ,  $\Delta_{\underline{v}}$  may be understood on *both* sides of the **log-wall** as being equipped with “*some outer  $G_{\underline{v}}$ -action*” — cf. the two diagonal arrows of Fig. 1.2 below. On the other hand, the datum of a *particular outer surjection*  $\Delta_{\underline{v}}^{\text{birat}} \twoheadrightarrow \Delta_{\underline{v}}$  [cf. the dotted line in Fig. 1.2] relating these two diagonal arrows — which depends, in an essential way, on the scheme [i.e., ring] structures involved! — necessarily *fails to be compatible with the log-link* [cf. the discussion of [AbsTopIII], Remark 3.7.7, (ii)]. This issue of “*triangular compatibility between independent indeterminacies*” is formally reminiscent of the issue of compatibility of outer homomorphisms discussed in [IUTchI], Remark 4.5.1, (i) [cf. also [IUTchII], Remark 2.5.2, (ii)].




Fig. 1.2: Independent basepoint indeterminacies obstruct relationship between birational and affine geometric fundamental groups

**Proposition 1.3.** (**log-links Between  $\Theta^{\pm\text{ell}}$ NF-Hodge Theaters**) *Let*

$$\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}; \quad \ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$$

*be  $\Theta^{\pm\text{ell}}$ NF-Hodge theaters [relative to the given initial  $\Theta$ -data] — cf. [IUTchI], Definition 6.13, (i). Write  $\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$ ,  $\ddagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  for the associated  $\mathcal{D}$ - $\Theta^{\pm\text{ell}}$ NF-Hodge theaters — cf. [IUTchI], Definition 6.13, (ii). Then:*



(i) **(Construction of the log-Link)** Fix an isomorphism

$$\Xi : \dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} \ddagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$$

of  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters. Let  $\dagger\mathfrak{F}_{\square}$  be one of the  $\mathcal{F}$ -prime-strips that appear in the  $\Theta$ - and  $\Theta^{\pm}$ -bridges that constitute  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  — i.e., either one of the  **$\mathcal{F}$ -prime-strips**

$$\dagger\mathfrak{F}_{>}, \quad \dagger\mathfrak{F}_{>}$$

or one of the **constituent  $\mathcal{F}$ -prime-strips** of the capsules

$$\dagger\mathfrak{F}_J, \quad \dagger\mathfrak{F}_T$$

[cf. [IUTchI], Definition 5.5, (ii); [IUTchI], Definition 6.11, (i)]. Write  $\ddagger\mathfrak{F}_{\square}$  for the corresponding  $\mathcal{F}$ -prime-strip of  $\ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ . Then the poly-isomorphism determined by  $\Xi$  between the  $\mathcal{D}$ -prime-strips associated to  $\dagger\mathfrak{F}_{\square}$ ,  $\ddagger\mathfrak{F}_{\square}$  uniquely determines a poly-isomorphism  $\log(\dagger\mathfrak{F}_{\square}) \xrightarrow{\sim} \ddagger\mathfrak{F}_{\square}$  [cf. Definition 1.1, (iii); [IUTchI], Corollary 5.3, (ii)], hence a **log-link**  $\dagger\mathfrak{F}_{\square} \xrightarrow{\log} \ddagger\mathfrak{F}_{\square}$  [cf. Definition 1.1, (iii)]. We shall denote by

$$\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\log} \ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$$

and refer to as a **log-link** from  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  to  $\ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  the collection of data consisting of  $\Xi$ , together with the collection of **log-links**  $\dagger\mathfrak{F}_{\square} \xrightarrow{\log} \ddagger\mathfrak{F}_{\square}$ , as “ $\square$ ” ranges over all possibilities for the  $\mathcal{F}$ -prime-strips in question. When  $\Xi$  is replaced by a poly-isomorphism  $\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} \ddagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$ , we shall also refer to the resulting collection of **log-links** [i.e., corresponding to each constituent isomorphism of the poly-isomorphism  $\Xi$ ] as a **log-link** from  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  to  $\ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ . When  $\Xi$  is the full poly-isomorphism, we shall refer to the resulting **log-link** as the **full log-link**. When  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} = \ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , and  $\Xi$  is the **identity**, then we shall refer to the resulting **log-link** as the **tautological log-link**.

(ii) **(Coricity)** Any **log-link**  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\log} \ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  induces [and may be thought of as “lying over”] a **[poly-]isomorphism**

$$\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} \ddagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$$

of  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters [and indeed coincides with the **log-link** constructed in (i) from this [poly-]isomorphism of  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters].

(iii) **(Further Properties of the log-Link)** In the notation of (i), any **log-link**  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\log} \ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  satisfies, for each  $\mathcal{F}$ -prime-strip  $\dagger\mathfrak{F}_{\square}$ , properties corresponding to the properties of Proposition 1.2, (ii), (iii), (iv), (v), (vi), (vii), (viii), (ix) i.e., concerning **simultaneous compatibility with ring structures and log-volumes, Kummer theory, and log-shells**.

(iv) **(Frobenius-picture)** Let  $\{{}^n\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}\}_{n \in \mathbb{Z}}$  be a **collection of distinct  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters** [relative to the given initial  $\Theta$ -data] indexed by the integers. Write  $\{{}^n\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}\}_{n \in \mathbb{Z}}$  for the associated  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters.

Then the full **log-links**  ${}^n\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} {}^{(n+1)}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , for  $n \in \mathbb{Z}$ , give rise to an **infinite chain**

$$\dots \xrightarrow{\text{log}} {}^{(n-1)}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} {}^n\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} {}^{(n+1)}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} \dots$$

of **log-linked  $\mathcal{F}$ -prime-strips** which induces a chain of full poly-isomorphisms

$$\dots \xrightarrow{\sim} {}^n\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} {}^{(n+1)}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} \dots$$

on the associated  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters. These chains may be represented symbolically as an **oriented graph**  $\vec{\Gamma}$  [cf. [AbsTopIII], §0]

$$\begin{array}{ccccccc} \dots & \rightarrow & \bullet & \rightarrow & \bullet & \rightarrow & \bullet & \rightarrow & \dots \\ & & & \searrow & \downarrow & \swarrow & & & \\ & & & & \circ & & & & \end{array}$$

— i.e., where the horizontal arrows correspond to the “ $\xrightarrow{\text{log}}$ ’s”; the “ $\bullet$ ’s” correspond to the “ ${}^n\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ ”; the “ $\circ$ ” corresponds to the “ ${}^n\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$ ”, identified up to isomorphism; the vertical/diagonal arrows correspond to the **Kummer isomorphisms** implicit in the statement of (iii). This oriented graph  $\vec{\Gamma}$  admits a natural action by  $\mathbb{Z}$  [cf. [AbsTopIII], Corollary 5.5, (v)] — i.e., a **translation symmetry** — that fixes the “**core**”  $\circ$ , but it does **not admit arbitrary permutation symmetries**. For instance,  $\vec{\Gamma}$  does not admit an automorphism that switches two adjacent vertices, but leaves the remaining vertices fixed.

*Proof.* The various assertions of Proposition 1.3 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 1.3.1.** Note that in Proposition 1.3, (i), it was necessary to carry out the given construction of the **log-link** first for a **single**  $\Xi$  [i.e., as opposed to a poly-isomorphism  $\Xi$ ], in order to maintain *compatibility* with the crucial “ **$\pm$ -synchronization**” [cf. [IUTchI], Remark 6.12.4, (iii); [IUTchII], Remark 4.5.3, (iii)] inherent in the structure of a  $\Theta^{\pm\text{ell}}$ -Hodge theater.

**Remark 1.3.2.** In the construction of Proposition 1.3, (i), the constituent  $\mathcal{F}$ -prime-strips  ${}^\dagger\mathfrak{F}_t$ , for  $t \in T$ , of the capsule  ${}^\dagger\mathfrak{F}_T$  are considered without regard to the  $\mathbb{F}_l^{\times\pm}$ -symmetries discussed in [IUTchII], Corollary 4.6, (iii). On the other hand, one verifies immediately that the **log-links** associated, in the construction of Proposition 1.3, (i), to these  $\mathcal{F}$ -prime-strips  ${}^\dagger\mathfrak{F}_t$ , for  $t \in T$  — i.e., more precisely, associated to the **labeled collections of monoids**  $\Psi_{\text{cns}}({}^\dagger\mathfrak{F}_\succ)_t$  of [IUTchII], Corollary 4.6, (iii) — are in fact **compatible** with the  $\mathbb{F}_l^{\times\pm}$ -**symmetrizing isomorphisms** discussed in [IUTchII], Corollary 4.6, (iii), hence also with the **conjugate synchronization** determined by these  $\mathbb{F}_l^{\times\pm}$ -symmetrizing isomorphisms — cf. the discussion of Step (vi) of the proof of Corollary 3.12 of §3 below. We leave the routine details to the reader.

**Remark 1.3.3.**

(i) In the context of Proposition 1.3, it is of interest to observe that the relationship between the various **Frobenioid-theoretic** [i.e., *Frobenius-like!*] portions of the  $\Theta^{\pm\text{ell}}$ NF-Hodge theaters in the *domain* and *codomain* the **log-link** of Proposition 1.3, (i),

*does not include any data that is incompatible, relative to the relevant Kummer isomorphisms, with the log-wall*

— cf. the discussion of Remark 1.2.4, (ii), (a), (b); Remark 1.2.4, (iii). This follows immediately from the fact [cf. Remarks 1.3.1, 1.3.2; [IUTchI], Corollary 5.3, (i), (ii), (iv); [IUTchI], Corollary 5.6, (i), (ii), (iii)] that these Frobenioid-theoretic portions of the  $\Theta^{\pm\text{ell}}$ NF-Hodge theaters under consideration are completely [i.e., *fully faithfully!*] *controlled* [cf. the discussion of (ii) below for more details], via **functorial algorithms**, by the corresponding *étale-like* structures, i.e., structures that appear in the associated  $\mathcal{D}\text{-}\Theta^{\pm\text{ell}}$ NF-Hodge theaters, which satisfy the crucial **coricity** property of Proposition 1.3, (ii).

(ii) In the context of (i), it is of interest to recall that the global portion of the underlying  $\Theta^{\text{ell}}$ -bridges is defined [cf. [IUTchI], Definition 6.11, (ii)] in such a way that it does not contain any global Frobenioid-theoretic data! In particular, the issue discussed in (i) only concerns the Frobenioid-theoretic portions of the following:

- (a) the various  **$\mathcal{F}$ -prime-strips** that appear;
- (b) the underlying  **$\Theta$ -Hodge theaters** of the  $\Theta^{\pm\text{ell}}$ NF-Hodge theaters under consideration;
- (c) the global portion of the underlying **NF-bridges** of the  $\Theta^{\pm\text{ell}}$ NF-Hodge theaters under consideration.

Here, the Frobenioid-theoretic data of (c) gives rise to **independent basepoints** with respect to the  $\mathbb{F}_l^*$ -**symmetry** [cf. [IUTchI], Corollary 5.6, (iii); [IUTchI], Remark 6.12.6, (iii); [IUTchII], Remark 4.7.6]. On the other hand, the independent basepoints that arise from the Frobenioid-theoretic data of (b), as well as of the portion of (a) that lies in the underlying  $\Theta$ NF-Hodge theater, do not cause any problems since this data is only subject to relationships *defined* by means of *full poly-isomorphisms* [cf. [IUTchI], Examples 4.3, 4.4]. That is to say, the  $\mathcal{F}$ -prime-strips that lie in the underlying  $\Theta^{\pm\text{ell}}$ -Hodge theater constitute the most *delicate* [i.e., relative to the issue of independent basepoints!] portion of the Frobenioid-theoretic data of a  $\Theta^{\pm\text{ell}}$ NF-Hodge theater. This delicacy revolves around the *global synchronization of  $\pm$ -indeterminacies* in the underlying  $\Theta^{\pm\text{ell}}$ -Hodge theater [cf. [IUTchI], Remark 6.12.4, (iii); [IUTchII], Remark 4.5.3, (iii)]. On the other hand, this delicacy does not in fact cause any problems since the synchronizations of  $\pm$ -indeterminacies in the underlying  $\Theta^{\pm\text{ell}}$ -Hodge theater are *defined* [not by means of *scheme-theoretic* relationships, but rather] by means of the structure of the underlying  $\mathcal{D}\text{-}\Theta^{\pm\text{ell}}$ -Hodge theater, which satisfies the crucial **coricity** property of Proposition 1.3, (ii) [cf. the discussion of (i)].

The diagrams discussed in the following Definition 1.4 will play a *central role* in the theory of the present series of papers.

**Definition 1.4.** We maintain the notation of Proposition 1.3 [cf. also [IUTchII], Corollary 4.10, (iii)]. Let  $\{^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}\}_{n,m \in \mathbb{Z}}$  be a *collection of distinct  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters* [relative to the given initial  $\Theta$ -data] indexed by pairs of integers. Then we shall refer to either of the diagrams

$$\begin{array}{ccccccc}
 & & \vdots & & \vdots & & \\
 & & \uparrow \text{log} & & \uparrow \text{log} & & \\
 \dots & \xrightarrow{\Theta^{\times\mu}} & {}^{n,m+1}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta^{\times\mu}} & {}^{n+1,m+1}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta^{\times\mu}} & \dots \\
 & & \uparrow \text{log} & & \uparrow \text{log} & & \\
 \dots & \xrightarrow{\Theta^{\times\mu}} & {}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta^{\times\mu}} & {}^{n+1,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta^{\times\mu}} & \dots \\
 & & \uparrow \text{log} & & \uparrow \text{log} & & \\
 & & \vdots & & \vdots & & \\
 & & \vdots & & \vdots & & \\
 & & \uparrow \text{log} & & \uparrow \text{log} & & \\
 \dots & \xrightarrow{\Theta_{\text{gau}}^{\times\mu}} & {}^{n,m+1}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta_{\text{gau}}^{\times\mu}} & {}^{n+1,m+1}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta_{\text{gau}}^{\times\mu}} & \dots \\
 & & \uparrow \text{log} & & \uparrow \text{log} & & \\
 \dots & \xrightarrow{\Theta_{\text{gau}}^{\times\mu}} & {}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta_{\text{gau}}^{\times\mu}} & {}^{n+1,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} & \xrightarrow{\Theta_{\text{gau}}^{\times\mu}} & \dots \\
 & & \uparrow \text{log} & & \uparrow \text{log} & & \\
 & & \vdots & & \vdots & & 
 \end{array}$$

— where the *vertical* arrows are the *full log-links*, and the *horizontal* arrows are the  $\Theta^{\times\mu}$ - and  $\Theta_{\text{gau}}^{\times\mu}$ -links of [IUTchII], Corollary 4.10, (iii) — as the *log-theta-lattice*. We shall refer to the log-theta-lattice that involves the  $\Theta^{\times\mu}$ - (respectively,  $\Theta_{\text{gau}}^{\times\mu}$ -) links as *non-Gaussian* (respectively, *Gaussian*). Thus, either of these diagrams may be represented symbolically by an *oriented graph*

$$\begin{array}{ccccccc}
 & & \vdots & & \vdots & & \\
 & & \uparrow & & \uparrow & & \\
 \dots & \longrightarrow & \bullet & \longrightarrow & \bullet & \longrightarrow & \dots \\
 & & \uparrow & & \uparrow & & \\
 \dots & \longrightarrow & \bullet & \longrightarrow & \bullet & \longrightarrow & \dots \\
 & & \uparrow & & \uparrow & & \\
 & & \vdots & & \vdots & & 
 \end{array}$$

— where the “•’s” correspond to the “ $n, m \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$ ”.

**Remark 1.4.1.**

(i) One fundamental property of the log-theta-lattices discussed in Definition 1.4 is the following:

the various squares that appear in each of the log-theta-lattices discussed in Definition 4.1 are **far from being [1-]commutative!**

Indeed, whereas the *vertical* arrows in each log-theta-lattice are constructed by applying the various *logarithms* at  $\underline{v} \in \underline{\mathbb{V}}$  — i.e., which are defined by means of power series that depend, in an essential way, on the *local ring structures* at  $\underline{v} \in \underline{\mathbb{V}}$  — the *horizontal* arrows in each log-theta-lattice [i.e., the  $\Theta^{\times \mu}$ -,  $\Theta_{\text{gau}}^{\times \mu}$ -links] are *incompatible* with these local ring structures at  $\underline{v} \in \underline{\mathbb{V}}$  in an essential way [cf. [IUTchII], Remark 1.11.2, (i), (ii)].

(ii) Whereas the horizontal arrows in each log-theta-lattice [i.e., the  $\Theta^{\times \mu}$ -,  $\Theta_{\text{gau}}^{\times \mu}$ -links] allow one, roughly speaking, to **identify** the respective “ $\mathcal{O}^{\times \mu}$ ’s” at [for simplicity]  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  on either side of the horizontal arrow [cf. [IUTchII], Corollary 4.10, (iv)], in order to avail oneself of the theory of **log-shells** — which will play an essential role in the *multiradial representation of the Gaussian monoids* to be developed in §3 below — it is necessary for the “•” [i.e.,  $\Theta^{\pm \text{ell}} \text{NF}$ -Hodge theater] in which one operates to appear as the *codomain* of a **log-link**, i.e., of a vertical arrow of the log-theta-lattice [cf. the discussion of [AbsTopIII], Remark 5.10.2, (iii)]. That is to say, from the point of view of the goal of constructing the multiradial representation of the Gaussian monoids that is to be developed in §3 below,

each execution of a *horizontal arrow* of the log-theta-lattice necessarily obligates a subsequent execution of a *vertical arrow* of the log-theta-lattice.

On the other hand, in light of the *noncommutativity* observed in (i), this “**intertwining**” of the horizontal and vertical arrows of the log-theta-lattice means that the desired **multiradiality** — i.e., **simultaneous compatibility** with the arithmetic holomorphic structures on *both sides of a horizontal arrow* of the log-theta-lattice — can only be realized if one works with objects that are **invariant with respect to the vertical arrows** [i.e., with respect to the action of  $\mathbb{Z}$  discussed in Proposition 1.3, (iv)], that is to say, with “**vertical cores**”, of the log-theta-lattice.

(iii) From the point of view of the analogy between the theory of the present series of papers and *p-adic Teichmüller theory* [cf. [AbsTopIII], §I5], the *vertical arrows* of the log-theta-lattice correspond to the *Frobenius morphism in positive characteristic*, whereas the *horizontal arrows* of the log-theta-lattice correspond to the “*transition from  $p^n \mathbb{Z}/p^{n+1} \mathbb{Z}$  to  $p^{n-1} \mathbb{Z}/p^n \mathbb{Z}$ ”*, i.e., the *mixed characteristic extension structure of a ring of Witt vectors* [cf. [IUTchI], Remark 3.9.3, (i)]. These correspondences are summarized in Fig. 1.3 below. In particular, the “intertwining of horizontal and vertical arrows of the log-theta-lattice” discussed in (ii) above may be thought of as the analogue, in the context of the theory of the present series of papers, of the well-known “intertwining between the mixed characteristic

extension structure of a ring of Witt vectors and the Frobenius morphism in positive characteristic” that appears in the classical  $p$ -adic theory.

<b>horizontal</b> arrows of the <i>log-theta-lattice</i>	<b>mixed characteristic</b> extension structure of a ring of <i>Witt vectors</i>
<b>vertical</b> arrows of the <i>log-theta-lattice</i>	the <b>Frobenius</b> morphism in <i>positive characteristic</i>

Fig. 1.3: Analogy between the log-theta-lattice and  $p$ -adic Teichmüller theory

**Remark 1.4.2.**

(i) The horizontal and vertical arrows of the log-theta-lattices discussed in Definition 4.1 share the common property of being *incompatible with the local ring structures*, hence, in particular, with the *conventional scheme-theoretic basepoints* on either side of the arrow in question [cf. the discussion of [IUTchII], Remark 3.6.3, (i)]. On the other hand, whereas the linking data of the *vertical arrows* [i.e., the **log-link**] is **rigid** and corresponds to a **single fixed, rigid arithmetic holomorphic structure** in which *addition* and *multiplication* are subject to “*rotations*” [cf. the discussion of [AbsTopIII], §I3], the linking data of the *horizontal arrows* [i.e., the  $\Theta^{\times\mu}$ -,  $\Theta_{\text{gau}}^{\times\mu}$ -links] — i.e., more concretely, the “ $\mathcal{O}^{\times\mu}$ ’s” at [for simplicity]  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  — is subject to a  $\widehat{\mathbb{Z}}^{\times}$ -**indeterminacy**, which has the effect of *obliterating the arithmetic holomorphic structure associated to a vertical line* of the log-theta-lattice [cf. the discussion of [IUTchII], Remark 1.11.2, (i), (ii)].

(ii) If, in the spirit of the discussion of [IUTchII], Remark 1.11.2, (ii), one attempts to “*force*” the horizontal arrows of the log-theta-lattice to be compatible with the arithmetic holomorphic structures on either side of the arrow by declaring — in the style of the **log-link**! — that these horizontal arrows induce an isomorphism of the respective “ $\Pi_{\underline{v}}$ ’s” at [for simplicity]  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , then one must contend with a situation in which the “common arithmetic holomorphic structure rigidified by the isomorphic copies of  $\Pi_{\underline{v}}$ ” is obliterated each time one takes into account the action of a nontrivial element of  $\widehat{\mathbb{Z}}^{\times}$  [i.e., that arises from the  $\widehat{\mathbb{Z}}^{\times}$ -*indeterminacy* involved] on the corresponding “ $\mathcal{O}^{\times\mu}$ ”. In particular, in order to keep track of the arithmetic holomorphic structure currently under consideration, one must, in effect, consider **paths** that record the sequence of “ $\Pi_{\underline{v}}$ -rigidifying” and “ $\widehat{\mathbb{Z}}^{\times}$ -indeterminacy” operations that one invokes. On the other hand, the horizontal lines of the log-theta-lattices given in Definition 4.1 amount, in effect, to *universal covering spaces* of the loops — i.e., “*unraveling paths of the loops*” [cf. the discussion of Remark 1.2.2, (vi)] — that occur as one invokes various series of “ $\Pi_{\underline{v}}$ -rigidifying” and “ $\widehat{\mathbb{Z}}^{\times}$ -indeterminacy” operations. Thus, in summary, any attempt as described above to “*force*” the horizontal arrows of the log-theta-lattice to be compatible with the arithmetic holomorphic structures on either side of the arrow does not result in any substantive simplification of the theory of the present



series of papers. We refer the reader to [IUTchIV], Remark 3.6.3, for a discussion of a related topic.

We are now ready to state the *main result* of the present §1.

**Theorem 1.5. (Bi-cores of the Log-theta-lattice)** *Fix a collection of initial  $\Theta$ -data*

$$(\overline{F}/F, X_F, l, \underline{C}_K, \underline{V}, \underline{\epsilon})$$

*as in [IUTchI], Definition 3.1. Then any Gaussian log-theta-lattice corresponding to this collection of initial  $\Theta$ -data [cf. Definition 1.4] satisfies the following properties:*

(i) **(Vertical Coricity)** *The vertical arrows of the Gaussian log-theta-lattice induce full poly-isomorphisms between the respective associated  $\mathcal{D}$ - $\Theta^{\pm\text{ell}}$  NF-Hodge theaters*

$$\dots \xrightarrow{\sim} {}^{n,m}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} {}^{n,m+1}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} \dots$$

[cf. Proposition 1.3, (ii)]. Here,  $n \in \mathbb{Z}$  is held fixed, while  $m \in \mathbb{Z}$  is allowed to vary.

(ii) **(Horizontal Coricity)** *The horizontal arrows of the Gaussian log-theta-lattice induce full poly-isomorphisms between the respective associated  $\mathcal{F}^{\times\mu}$ -prime-strips*

$$\dots \xrightarrow{\sim} {}^{n,m}\mathfrak{F}_{\Delta}^{\times\mu} \xrightarrow{\sim} {}^{n+1,m}\mathfrak{F}_{\Delta}^{\times\mu} \xrightarrow{\sim} \dots$$

[cf. [IUTchII], Corollary 4.10, (iv)]. Here,  $m \in \mathbb{Z}$  is held fixed, while  $n \in \mathbb{Z}$  is allowed to vary.

(iii) **(Bi-coric  $\mathcal{F}^{\times\mu}$ -Prime-Strips)** *For  $n, m \in \mathbb{Z}$ , write  ${}^{n,m}\mathcal{D}_{\Delta}^{\dagger}$  for the  $\mathcal{D}^{\dagger}$ -prime-strip associated to the  $\mathcal{F}^{\dagger}$ -prime-strip  ${}^{n,m}\mathfrak{F}_{\Delta}^{\dagger}$  labeled “ $\Delta$ ” of the  $\Theta^{\pm\text{ell}}$  NF-Hodge theater  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  [cf. [IUTchII], Corollary 4.10, (i)];  ${}^{n,m}\mathcal{D}_{\succ}$  for the  $\mathcal{D}$ -prime-strip labeled “ $\succ$ ” of the  $\Theta^{\pm\text{ell}}$  NF-Hodge theater  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  [cf. [IUTchI], Definition 6.11, (i)]. Let us identify [cf. [IUTchII], Corollary 4.10, (i)] the collections of data*

$$\Psi_{\text{cns}}({}^{n,m}\mathcal{D}_{\succ})_0 \text{ and } \Psi_{\text{cns}}({}^{n,m}\mathcal{D}_{\succ})_{(\mathbb{F}_l^*)}$$

*via the isomorphism of the final display of [IUTchII], Corollary 4.5, (iii), and denote by*

$$\mathfrak{F}_{\Delta}^{\dagger}({}^{n,m}\mathcal{D}_{\succ})$$

*the resulting  $\mathcal{F}^{\dagger}$ -prime-strip. [Thus, it follows immediately from the constructions involved — cf. the discussion of [IUTchII], Corollary 4.10, (i) — that there is a natural identification isomorphism  $\mathfrak{F}_{\Delta}^{\dagger}({}^{n,m}\mathcal{D}_{\succ}) \xrightarrow{\sim} \mathfrak{F}_{\succ}^{\dagger}({}^{n,m}\mathcal{D}_{\succ})$ , where we write  $\mathfrak{F}_{\succ}^{\dagger}({}^{n,m}\mathcal{D}_{\succ})$  for the  $\mathcal{F}^{\dagger}$ -prime-strip determined by  $\Psi_{\text{cns}}({}^{n,m}\mathcal{D}_{\succ})$ .] Write*

$$\mathfrak{F}_{\Delta}^{\dagger \times}({}^{n,m}\mathcal{D}_{\succ}), \quad \mathfrak{F}_{\Delta}^{\dagger \times \mu}({}^{n,m}\mathcal{D}_{\succ})$$

for the  $\mathcal{F}^{\times -}$ ,  $\mathcal{F}^{\times \mu}$ -prime-strips determined by  $\mathfrak{F}_{\Delta}^{\dagger}(n, m \mathcal{D}_{\succ})$  [cf. [IUTchII], Definition 4.9, (vi), (vii)]. Thus, by applying the isomorphisms “ $\Psi_{\text{cns}}(n, m \mathcal{D}_{\underline{v}})^{\times} \xrightarrow{\sim} \Psi_{\text{cns}}^{\text{ss}}(n, m \mathcal{D}_{\Delta}^{\dagger})^{\times}$ ”, for  $\underline{v} \in \underline{\mathbb{V}}$ , of [IUTchII], Corollary 4.5, (ii), [it follows immediately from the definitions that] there exists a **functorial algorithm** in the  $\mathcal{D}^{\dagger}$ -prime-strip  $n, m \mathcal{D}_{\Delta}^{\dagger}$  for constructing an  $\mathcal{F}^{\times}$ -prime-strip  $\mathfrak{F}_{\Delta}^{\dagger \times}(n, m \mathcal{D}_{\Delta}^{\dagger})$ , together with a **natural isomorphism**

$$\mathfrak{F}_{\Delta}^{\dagger \times}(n, m \mathcal{D}_{\succ}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\dagger \times}(n, m \mathcal{D}_{\Delta}^{\dagger})$$

— i.e., in more intuitive terms, “ $\mathfrak{F}_{\Delta}^{\dagger \times}(n, m \mathcal{D}_{\succ})$ ”, hence also the associated  $\mathcal{F}^{\times \mu}$ -prime-strip “ $\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\succ})$ ”, may be naturally regarded as objects constructed from  $n, m \mathcal{D}_{\Delta}^{\dagger}$ . Then the poly-isomorphisms of (i) [cf. Remark 1.3.2], (ii) induce, respectively, poly-isomorphisms of  $\mathcal{F}^{\times \mu}$ -prime-strips

$$\begin{aligned} \dots &\xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\succ}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m+1 \mathcal{D}_{\succ}) \xrightarrow{\sim} \dots \\ \dots &\xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\dagger \times \mu}(n+1, m \mathcal{D}_{\Delta}^{\dagger}) \xrightarrow{\sim} \dots \end{aligned}$$

— where we note that the poly-isomorphisms of  $\mathcal{F}^{\times \mu}$ -prime-strips of the first line of the display are **not full**. In particular, by composing these isomorphisms, one obtains **full poly-isomorphisms** of  $\mathcal{F}^{\times \mu}$ -prime-strips

$$\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\dagger \times \mu}(n', m' \mathcal{D}_{\Delta}^{\dagger})$$

for arbitrary  $n', m' \in \mathbb{Z}$ . That is to say, in more intuitive terms, the  $\mathcal{F}^{\times \mu}$ -prime-strip “ $n, m \mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger})$ ”, regarded up to isomorphism, is an invariant — which we shall refer to as “**bi-coric**” — of both the horizontal and the vertical arrows of the log-theta-lattice. Finally, the Kummer isomorphisms “ $\Psi_{\text{cns}}(\dagger \mathfrak{F}) \xrightarrow{\sim} \Psi_{\text{cns}}(\dagger \mathcal{D})$ ” of [IUTchII], Corollary 4.6, (i), determine **Kummer isomorphisms**

$$n, m \mathfrak{F}_{\Delta}^{\dagger \times \mu} \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger})$$

which are **compatible** with the poly-isomorphisms of (ii), as well as with the  $\times \mu$ -Kummer structures at the  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  of the various  $\mathcal{F}^{\times \mu}$ -prime-strips involved [cf. [IUTchII], Definition 4.9, (vi), (vii)]; a similar compatibility holds for  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$  [cf. the discussion of the final portion of [IUTchII], Definition 4.9, (v)].

(iv) **(Bi-coric Mono-analytic Log-shells)** The full poly-isomorphisms that constitute the bi-coricity property discussed in (iii) induce **poly-isomorphisms**

$$\left\{ \mathcal{I}_{n, m \mathcal{D}_{\Delta}^{\dagger}} \subseteq \underline{\log}(n, m \mathcal{D}_{\Delta}^{\dagger}) \right\} \xrightarrow{\sim} \left\{ \mathcal{I}_{n', m' \mathcal{D}_{\Delta}^{\dagger}} \subseteq \underline{\log}(n', m' \mathcal{D}_{\Delta}^{\dagger}) \right\}$$

$$\left\{ \mathcal{I}_{\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger})} \subseteq \underline{\log}(\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger})) \right\} \xrightarrow{\sim} \left\{ \mathcal{I}_{\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n', m' \mathcal{D}_{\Delta}^{\dagger})} \subseteq \underline{\log}(\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n', m' \mathcal{D}_{\Delta}^{\dagger})) \right\}$$

for arbitrary  $n, m, n', m' \in \mathbb{Z}$  that are **compatible** with the **natural poly-isomorphisms**

$$\left\{ \mathcal{I}_{n, m \mathcal{D}_{\Delta}^{\dagger}} \subseteq \underline{\log}(n, m \mathcal{D}_{\Delta}^{\dagger}) \right\} \xrightarrow{\sim} \left\{ \mathcal{I}_{\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger})} \subseteq \underline{\log}(\mathfrak{F}_{\Delta}^{\dagger \times \mu}(n, m \mathcal{D}_{\Delta}^{\dagger})) \right\}$$

of Proposition 1.2, (viii). On the other hand, by applying the constructions of Definition 1.1, (i), (ii), to the collections of data “ $\Psi_{\text{cns}}(\dagger\mathfrak{F}_{\triangleright})_0$ ” and “ $\Psi_{\text{cns}}(\dagger\mathfrak{F}_{\triangleright})_{(\mathbb{R}_i^*)}$ ” used in [IUTchII], Corollary 4.10, (i), to construct  ${}^{n,m}\mathfrak{F}_{\Delta}^{\dagger}$  [cf. Remark 1.3.2], one obtains a [“**holomorphic**”] **log-shell**, together with an enveloping “ $\underline{\log}(-)$ ” [cf. the pair “ $\mathcal{I}_{\dagger\mathfrak{F}} \subseteq \underline{\log}(\dagger\mathfrak{F})$ ” of Definition 1.1, (iii)], which we denote by

$$\mathcal{I}_{n,m}\mathfrak{F}_{\Delta} \subseteq \underline{\log}({}^{n,m}\mathfrak{F}_{\Delta})$$

[by means of a slight abuse of notation, since no  $\mathcal{F}$ -prime-strip “ ${}^{n,m}\mathfrak{F}_{\Delta}$ ” has been defined!]. Then one has **natural poly-isomorphisms**

$$\begin{aligned} \left\{ \mathcal{I}_{n,m}\mathfrak{D}_{\Delta}^{\dagger} \subseteq \underline{\log}({}^{n,m}\mathfrak{D}_{\Delta}^{\dagger}) \right\} &\xrightarrow{\sim} \left\{ \mathcal{I}_{n,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu} \subseteq \underline{\log}({}^{n,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu}) \right\} \\ &\xrightarrow{\sim} \left\{ \mathcal{I}_{n,m}\mathfrak{F}_{\Delta} \subseteq \underline{\log}({}^{n,m}\mathfrak{F}_{\Delta}) \right\} \end{aligned}$$

[cf. the poly-isomorphisms obtained in Proposition 1.2, (viii)]; here, the first “ $\xrightarrow{\sim}$ ” may be regarded as being induced by the Kummer isomorphisms of (iii) and is **compatible** with the poly-isomorphisms induced by the poly-isomorphisms of (ii).

(v) (**Bi-coric Mono-analytic Realified Global Frobenioids**) Let  $n, m, n', m' \in \mathbb{Z}$ . Then the poly-isomorphisms of  $\mathcal{D}^{\dagger}$ -prime-strips  ${}^{n,m}\mathfrak{D}_{\Delta}^{\dagger} \xrightarrow{\sim} {}^{n',m'}\mathfrak{D}_{\Delta}^{\dagger}$  induced by the full poly-isomorphisms of (i), (ii) induce [cf. [IUTchII], Corollaries 4.5, (ii); 4.10, (v)] an isomorphism of collections of data

$$\begin{aligned} (\mathcal{D}^{\dagger}({}^{n,m}\mathfrak{D}_{\Delta}^{\dagger}), \text{Prime}(\mathcal{D}^{\dagger}({}^{n,m}\mathfrak{D}_{\Delta}^{\dagger}))) &\xrightarrow{\sim} \underline{\mathbb{V}}, \{ {}^{n,m}\rho_{\mathcal{D}^{\dagger}, \underline{v}} \}_{\underline{v} \in \underline{\mathbb{V}}} \\ &\xrightarrow{\sim} (\mathcal{D}^{\dagger}({}^{n',m'}\mathfrak{D}_{\Delta}^{\dagger}), \text{Prime}(\mathcal{D}^{\dagger}({}^{n',m'}\mathfrak{D}_{\Delta}^{\dagger}))) \xrightarrow{\sim} \underline{\mathbb{V}}, \{ {}^{n',m'}\rho_{\mathcal{D}^{\dagger}, \underline{v}} \}_{\underline{v} \in \underline{\mathbb{V}}} \end{aligned}$$

— i.e., consisting of a Frobenioid, a bijection, and a collection of isomorphisms of topological monoids indexed by  $\underline{\mathbb{V}}$ . Moreover, this isomorphism of collections of data is **compatible**, relative to the full poly-isomorphisms of (ii), with the  $\mathbb{R}_{>0}$ -orbits of the isomorphisms of collections of data

$$\begin{aligned} ({}^{n,m}\mathcal{C}_{\Delta}^{\dagger}, \text{Prime}({}^{n,m}\mathcal{C}_{\Delta}^{\dagger})) &\xrightarrow{\sim} \underline{\mathbb{V}}, \{ {}^{n,m}\rho_{\Delta, \underline{v}} \}_{\underline{v} \in \underline{\mathbb{V}}} \\ &\xrightarrow{\sim} (\mathcal{D}^{\dagger}({}^{n,m}\mathfrak{D}_{\Delta}^{\dagger}), \text{Prime}(\mathcal{D}^{\dagger}({}^{n,m}\mathfrak{D}_{\Delta}^{\dagger}))) \xrightarrow{\sim} \underline{\mathbb{V}}, \{ {}^{n,m}\rho_{\mathcal{D}^{\dagger}, \underline{v}} \}_{\underline{v} \in \underline{\mathbb{V}}} \end{aligned}$$

obtained by applying the functorial algorithm discussed in the final portion of [IUTchII], Corollary 4.6, (ii) [cf. also the final portion of [IUTchII], Corollary 4.10, (v)].

*Proof.* The various assertions of Theorem 1.5 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

### Remark 1.5.1.

(i) Note that the theory of **conjugate synchronization** developed in [IUTchII] [cf., especially, [IUTchII], Corollaries 4.5, (iii); 4.6, (iii)] plays an essential role in establishing the **bi-coricity** properties discussed in Theorem 1.5, (iii), (iv) — i.e., at a more technical level, in constructing the objects equipped with a subscript “ $\Delta$ ”

that appear in Theorem 1.5, (iii); [IUTchII], Corollary 4.10, (i). That is to say, the conjugate synchronization determined by the various symmetrizing isomorphisms of [IUTchII], Corollaries 4.5, (iii); 4.6, (iii), may be thought of as a sort of **descent** mechanism that allows one to descend data that, *a priori*, is **label-dependent** [i.e., depends on the labels “ $t \in \text{LabCusp}^\pm(-)$ ”] to data that is **label-independent**. Here, it is important to recall that these labels depend, in an essential way, on the “**arithmetic holomorphic structures**” involved — i.e., at a more technical level, on the *geometric fundamental groups* involved — hence only make sense within a *vertical line* of the log-theta-lattice. That is to say, the significance of this transition from label-dependence to label-independence lies in the fact that this transition is precisely what allows one to construct objects that make sense in *horizontally adjacent* “•’s” of the log-theta-lattice, i.e., to construct *horizontally coric* objects [cf. Theorem 1.5, (ii); the second line of the fifth display of Theorem 1.5, (iii)]. On the other hand, in order to construct the horizontal arrows of the log-theta-lattice, it is necessary to work with **Frobenius-like structures** [cf. the discussion of [IUTchII], Remark 3.6.2, (ii)]. In particular, in order to construct *vertically coric* objects [cf. the first line of the fifth display of Theorem 1.5, (iii)], it is necessary to pass to **étale-like structures** [cf. the discussion of Remark 1.2.4, (i)] by means of **Kummer isomorphisms** [cf. the final display of Theorem 1.5, (iii)]. Thus, in summary,

the **bi-coricity** properties discussed in Theorem 1.5, (iii), (iv) — i.e., roughly speaking, the bi-coricity of the various “ $\mathcal{O}^{\times\mu}$ ” at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  — may be thought of as a consequence of the *intricate interplay of various aspects* of the theory of **Kummer-compatible conjugate synchronization** established in [IUTchII], Corollaries 4.5, (iii); 4.6, (iii).

(ii) In light of the central role played by the theory of conjugate synchronization in the constructions that underlie Theorem 1.5 [cf. the discussion of (i)], it is of interest to examine in more detail to what extent the highly technically nontrivial theory of conjugate synchronization may be replaced by a simpler apparatus. One naive approach to this problem is the following. Let  $G$  be a *topological group* [such as one of the absolute Galois groups  $G_{\underline{v}}$  associated to  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ ]. Then one way to attempt to avoid the application of the theory of conjugate synchronization — which amounts, in essence, to the construction of a **diagonal embedding**

$$G \hookrightarrow G \times \dots \times G$$

[cf. the notation “ $\langle |\mathbb{F}_l| \rangle$ ”, “ $\langle \mathbb{F}_l^* \rangle$ ” that appears in [IUTchII], Corollaries 3.5, 3.6, 4.5, 4.6] in a product of copies of  $G$  that, *a priori*, may only be identified with one another *up to conjugacy* [i.e., up to composition with an inner automorphism] — is to try to work, instead, with the  $(G \times \dots \times G)$ -**conjugacy class of such a diagonal**. Here, to simplify the notation, let us assume that the above products of copies of  $G$  are, in fact, products of *two copies* of  $G$ . Then to *identify* the diagonal embedding  $G \hookrightarrow G \times G$  with its  $(G \times G)$ -conjugates implies that one must consider **identifications**

$$(g, g) \sim (g, hgh^{-1}) = (g, [h, g] \cdot g)$$

[where  $g, h \in G$ ] — i.e., one must identify  $(g, g)$  with the product of  $(g, g)$  with  $(1, [h, g])$ . On the other hand, the original purpose of working with distinct copies

of  $G$  lies in considering **distinct** Galois-theoretic Kummer classes — corresponding to **distinct theta values** [cf. [IUTchII], Corollaries 3.5, 3.6] — at distinct components. That is to say, to identify elements of  $(G, G)$  that differ by a factor of  $(1, [h, g])$  is **incompatible**, in an essential way, with the convention that such a factor  $(1, [h, g])$  should correspond to distinct elements [i.e., “1” and “[ $h, g$ ]”] at distinct components [cf. the discussion of Remark 1.5.3, (ii), below]. Here, we note that this incompatibility may be thought of as an essential consequence of the *highly nonabelian nature* of  $G$ , e.g., when  $G$  is taken to a copy of  $G_{\underline{v}}$ , for  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ . Thus, in summary, this naive approach to replacing the theory of conjugate synchronization by a simpler apparatus is *inadequate* from the point of view of the theory of the present series of papers.

(iii) At a purely combinatorial level, the notion of conjugate synchronization is reminiscent of the **label synchronization** discussed in [IUTchI], Remark 4.9.2, (i), (ii). Indeed, both conjugate and label synchronization may be thought of as a sort of **combinatorial representation of the arithmetic holomorphic structure** associated to a single vertical line of the log-theta-lattice [cf. the discussion of [IUTchI], Remark 4.9.2, (iv)].

### Remark 1.5.2.

(i) Recall that unlike the case with the action of the  $\mathbb{F}_l^{\times\pm}$ -symmetry on the various labeled copies of the absolute Galois group  $G_{\underline{v}}$ , for  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  [cf. [IUTchII], Corollaries 4.5, (iii); 4.6, (iii)], it is *not* possible to establish an analogous theory of **conjugate synchronization** in the case of the  $\mathbb{F}_l^*$ -symmetry for *labeled copies of  $\overline{F}$*  [cf. [IUTchII], Remark 4.7.2]. This is to say, the closest analogue of the conjugate synchronization obtained in the local case relative to the  $\mathbb{F}_l^{\times\pm}$ -symmetry is the action of the  $\mathbb{F}_l^*$ -symmetry on *labeled copies of the subfield  $F_{\text{mod}} \subseteq \overline{F}$*  discussed in [IUTchII], Corollaries 4.7, (ii); 4.8, (ii). One consequence of this incompatibility of the  $\mathbb{F}_l^*$ -symmetry with the full algebraic closure  $\overline{F}$  of  $F_{\text{mod}}$  [hence, in particular, with various roots of elements of  $F_{\text{mod}}$ ] is that the  $\mathbb{F}_l^*$ -symmetry on labeled copies of the subfield  $F_{\text{mod}} \subseteq \overline{F}$  discussed in [IUTchII], Corollaries 4.7, (ii); 4.8, (ii), **fails to be compatible with Kummer theory**. This is one quite essential reason why it is not possible to establish **bi-coricity** properties for, say, “ $F_{\text{mod}}^{\times}$ ” that are analogous to the bi-coricity properties established in Theorem 1.5, (iii), for “ $\mathcal{O}^{\times\mu}$ ” [cf. the discussion of Remark 1.5.1, (i)]. From this point of view,

the **bi-coric mono-analytic realified global Frobenioids** of Theorem 1.5, (v) — i.e., in essence, the notion of “**log-volume**” [cf. the point of view of Remark 1.2.2, (v)] — may be thought of as a sort of “**closest possible approximation**” to such a “bi-coric  $F_{\text{mod}}^{\times}$ ” [i.e., which does not exist].

Alternatively, from the point of view of the theory to be developed in §3 below,

we shall apply the **bi-coric “ $\mathcal{O}^{\times\mu}$ ’s”** of Theorem 1.5, (iii) — i.e., in the form of the **bi-coric mono-analytic log-shells** of Theorem 1.5, (iv) — to construct “**multiradial containers**” for the labeled copies of



$F_{\text{mod}}$  discussed above by applying the **localization functors** discussed in [IUTchII], Corollaries 4.7, (iii); 4.8, (iii).

That is to say, such “multiradial containers” will play the role of a **transportation mechanism** for “ $F_{\text{mod}}^{\times}$ ” — up to *certain indeterminacies!* — between *distinct arithmetic holomorphic structures* [i.e., distinct vertical lines of the log-theta-lattice].

(ii) Note that although the  $\mathbb{F}_l^*$ -symmetry fails to be compatible with labeled copies of “ $\overline{F}$ ” [cf. the discussion of (i)], if one *forgets* the  $\mathbb{F}_l^*$ -symmetry and the corresponding labels, then

the *functorial algorithms* of [IUTchII], Corollary 4.7, (i), applied to the *vertical coricity* of Theorem 1.5, (i), allow one to construct **vertically coric** versions of “ $\overline{F}$ ” which are, moreover, **compatible with Kummer theory** [cf. [IUTchII], Corollary 4.8, (i)].

This point of view will be applied in §3 below, in conjunction with the point of view of “*multiradial containers*” discussed in (i) above. Here, we recall [cf. the discussion of [IUTchII], Remark 3.6.2, (ii)] that, in general, the phenomenon of “*Kummer-incompatibility*” [i.e., incompatibility with Kummer theory — cf. the discussion of “*Kummer-blindness*” in [IUTchI], Remark 5.1.3] occurs precisely when one wishes to perform constructions — such as, for instance, the construction of the  $\Theta$ -,  $\Theta^{\times\mu}$ -, or  $\Theta_{\text{gau}}^{\times\mu}$ -links — that are “**not bound to conventional scheme theory**”. That is to say, in the case of the labeled copies of “ $F_{\text{mod}}$ ” discussed in (i), the Kummer-incompatibility that occurs may be thought of as a reflection of the **dismantling** of the **global prime-tree structure** of a number field [cf. the discussion of [IUTchII], Remark 4.11.2, (iv)] that underlies the *construction of the  $\Theta^{\pm\text{ell}}NF$ -Hodge theater* performed in [IUTchI], [IUTchII], hence, in particular, of the *conjugate synchronization* established for the  $\mathbb{F}_l^{\times\pm}$ -symmetry.

(iii) Despite the failure of labeled copies of “ $F_{\text{mod}}^{\times}$ ” to admit a natural *bi-coric structure* — a state of affairs that forces one to resort to the use of “*multiradial containers*” in order to transport such labeled copies of “ $F_{\text{mod}}^{\times}$ ” to alien arithmetic holomorphic structures [cf. the discussion of (i) above] — the global Frobenioids associated to copies of “ $F_{\text{mod}}^{\times}$ ” nevertheless possess important properties that are *not* satisfied, for instance, by the bi-coric realified global Frobenioids discussed in Theorem 1.5, (v) [cf. also [IUTchI], Definition 5.2, (iv); [IUTchII], Corollary 4.5, (ii); [IUTchII], Corollary 4.6, (ii)]. Indeed, unlike the objects contained in the *realified* global Frobenioids that appear in Theorem 1.5, (v), the objects contained in the global Frobenioids associated to copies of “ $F_{\text{mod}}^{\times}$ ” correspond to *genuine “conventional arithmetic line bundles”*. In particular, by applying the **ring structure** of the copies of “ $F_{\text{mod}}$ ” under consideration, one can *push forward* such arithmetic line bundles so as to obtain *arithmetic vector bundles* over [the ring of rational integers]  $\mathbb{Z}$  and then form *tensor products* of such arithmetic vector bundles. Such operations will play a key role in the theory of §3 below, as well as in the theory to be developed in [IUTchIV].



**Remark 1.5.3.**

(i) In [QuCnf] [cf. also [AbsTopIII], Proposition 2.6; [AbsTopIII], Corollary 2.7], a theory was developed concerning deformations of holomorphic structures on Riemann surfaces in which holomorphic structures are represented by means of **squares** or **rectangles** on the surface, while quasiconformal Teichmüller deformations of holomorphic structures are represented by **parallelograms** on the surface. That is to say, relative to appropriate choices of local coordinates, quasiconformal Teichmüller deformations may be thought of as affine linear deformations in which one of the two underlying real dimensions of the Riemann surface is *dilated* by some factor  $\in \mathbb{R}_{>0}$ , while the other underlying real dimensions is *left undeformed*. From this point of view, the theory of **conjugate synchronization** — which may be regarded as a sort of **rigidity** that represents the *arithmetic holomorphic structure* associated to a vertical line of the log-theta-lattice [cf. the discussion given in [IUTchII], Remarks 4.7.3, 4.7.4, of the *uniradiality* of the  $\mathbb{F}_l^{\times\pm}$ -symmetry that underlies the phenomenon of conjugate synchronization] — may be thought of as a sort of **nonarchimedean arithmetic analogue** of the representation of holomorphic structures by means of squares/rectangles referred to above. That is to say, the *right angles* which are characteristic of squares/rectangles may be thought of as a sort of *synchronization* between the metrics of the two underlying real dimensions of a Riemann surface [i.e., metrics which, *a priori*, may differ by some *dilating* factor] — cf. Fig. 1.4 below. Here, we mention in passing that this point of view is reminiscent of the discussion of [IUTchII], Remark 3.6.5, (ii), in which the point of view is taken that the phenomenon of conjugate synchronization may be thought of as a reflection of the **coherence** of the **arithmetic holomorphic structures** involved.




Fig. 1.4: Analogy between conjugate synchronization and the representation of complex holomorphic structures via squares/rectangles

(ii) Relative to the point of view discussed in (i), the approach described in Remark 1.5.1, (ii), to “avoiding conjugate synchronization by identifying the various

conjugates of the diagonal embedding” corresponds — in light of the *highly non-abelian* nature of the groups involved! [cf. the discussion of Remark 1.5.1, (ii)] — to thinking of a holomorphic structure on a Riemann surface as an “equivalence class of holomorphic structures in the usual sense relative to the equivalence relation of differing by a Teichmüller deformation”! That is to say, such an [unconventional!] approach to the definition of a holomorphic structure allows one to circumvent the issue of *rigidifying* the relationship between the metrics of the two underlying real dimensions of the Riemann surface — but only at the cost of rendering unfeasible any meaningful theory of “deformations of a holomorphic structure”!

(iii) The analogy discussed in (i) between conjugate synchronization [which arises from the  $\mathbb{F}_l^{\times\pm}$ -symmetry!] and the representation of a complex holomorphic structure by means of squares/rectangles may also be applied to the “**synchronization**” given in [IUTchII], Corollary 4.7, (ii); [IUTchII], Corollary 4.8, (ii), between the various labeled non-realified and realified global Frobenioids by means of the  $\mathbb{F}_l^*$ -**symmetry**. Indeed, this analogy is all the more apparent in the case of the *realified* global Frobenioids — which admit a natural  $\mathbb{R}_{>0}$ -*action*. Here, we observe in passing that, just as the theory of conjugate synchronization plays an essential role in the construction of the *local portions* of the  $\Theta^{\times\mu}$ -,  $\Theta_{\text{gau}}^{\times\mu}$ -links given in [IUTchII], Corollary 4.10, (i), (ii), (iii),

the **synchronization of realified global Frobenioids** by means of the  $\mathbb{F}_l^*$ -**symmetry** may be related — via the isomorphisms of Frobenioids of the second displays of [IUTchII], Corollary 4.7, (iii); [IUTchII], Corollary 4.8, (iii) [cf. also the discussion of [IUTchII], Remark 4.8.1] — to the construction of the *realified global Frobenioid portion* of the  $\Theta_{\text{gau}}^{\times\mu}$ -**link** given in [IUTchII], Corollary 4.10, (ii).

On the other hand, the synchronization involving the *non-realified* global Frobenioids may be thought of a sort of *further rigidification* of the realified global Frobenioids. As discussed in Remark 1.5.2, (iii), this “further rigidification” will play an important role in the theory of §3 below.

#### Remark 1.5.4.

(i) As discussed in [IUTchII], Remark 3.8.3, (iii), one of the main themes of the present series of papers is the goal of giving an **explicit description** of what **one arithmetic holomorphic structure** — i.e., one *vertical line of the log-theta-lattice* — looks like from the point of view of a **distinct arithmetic holomorphic structure** — i.e., another vertical line of the log-theta-lattice — that is only related to the original arithmetic holomorphic structure via some mono-analytic core, e.g., the various bi-coric structures discussed in Theorem 1.5, (iii), (iv), (v). Typically, the objects of interest that are constructed within the original arithmetic holomorphic structure are **Frobenius-like** structures [cf. the discussion of [IUTchII], Remark 3.6.2], which, as we recall from the discussion of Remark 1.5.2, (ii) [cf. also the discussion of [IUTchII], Remark 3.6.2, (ii)], are necessary in order to perform constructions — such as, for instance, the construction of the  $\Theta$ -,  $\Theta^{\times\mu}$ -, or  $\Theta_{\text{gau}}^{\times\mu}$ -links — that are “**not bound to conventional scheme theory**”. Indeed, the main example of such an object of interest consists precisely of the **Gaussian monoids**

discussed in [IUTchII], §3, §4. Thus, the operation of describing such an object of interest from the point of view of a *distinct arithmetic holomorphic structure* may be broken down into *two steps*:

- (a) passing from *Frobenius-like structures* to *étale-like structures* via various **Kummer isomorphisms**;
- (b) transporting the resulting *étale-like structures* from one arithmetic holomorphic structure to another by means of various **multiradiality properties**.

In particular, the computation of what the object of interest looks like from the point of view of a distinct arithmetic holomorphic structure may be broken down into the computation of the **indeterminacies** or “*departures from rigidity*” that arise — i.e., the computation of “*what sort of damage is incurred to the object of interest*” — during the execution of each of these two steps (a), (b). We shall refer to the indeterminacies that arise from (a) as **Kummer-detachment indeterminacies** and to the indeterminacies that arise from (b) as **étale-transport indeterminacies**.

(ii) *Étale-transport indeterminacies* typically amount to the indeterminacies that occur as a result of the execution of various “*anabelian*” or “*group-theoretic*” *algorithms*. One fundamental example of such indeterminacies is constituted by the indeterminacies that occur in the context of Theorem 1.5, (iii), (iv), as a result of the existence of **automorphisms** of the various [copies of] local absolute Galois groups  $G_v$ , for  $v \in \mathbb{V}^{\text{non}}$ , which are *not of scheme-theoretic origin* [cf. the discussion of [AbsTopIII], §I3].

(iii) On other hand, one important example, from the point of view of the theory of the present series of papers, of a *Kummer-detachment indeterminacy* is constituted by the **Frobenius-picture diagrams** given in Propositions 1.2, (x); 1.3, (iv) — i.e., the issue of *which path* one is to take from a particular “•” to the coric “o”. That is to say, despite the fact that these diagrams *fail to be commutative*, the “**upper semi-commutativity**” property satisfied by the **coric holomorphic log-shells** involved [cf. the discussion of Remark 1.2.2, (iii)] may be regarded as a sort of computation, in the form of an *upper estimate*, of the Kummer-detachment indeterminacy in question. Another important example, from the point of view of the theory of the present series of papers, of a *Kummer-detachment indeterminacy* is given by the  $\widehat{\mathbb{Z}}^\times$ -**indeterminacies** discussed in Remark 1.4.2 [cf. also the *Kummer isomorphisms* of the final display of Theorem 1.5, (iii)].

## Section 2: Multiradial Theta Monoids

In the present §2, we **globalize** the **multiradial** portion of the local theory of **theta monoids** developed in [IUTchII], §1, §3, at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  [cf., especially, [IUTchII], Corollary 1.12; [IUTchII], Proposition 3.4] so as to cover the theta monoids of [IUTchII], Corollaries 4.5, (iv), (v); 4.6, (iv), (v), and explain how the resulting theory may be fit into the framework of the **log-theta-lattice** developed in §1.

In the following discussion, we assume that we have been given *initial*  $\Theta$ -data as in [IUTchI], Definition 3.1. Let  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  be a  $\Theta^{\pm\text{ell}}$  *NF-Hodge theater* [relative to the given initial  $\Theta$ -data — cf. [IUTchI], Definition 6.13, (i)] and

$$\{^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}\}_{n,m \in \mathbb{Z}}$$

a *collection of distinct*  $\Theta^{\pm\text{ell}}$  *NF-Hodge theaters* [relative to the given initial  $\Theta$ -data] indexed by pairs of integers, which we think of as arising from a *Gaussian log-theta-lattice*, as in Definition 1.4. We begin by reviewing the theory of theta monoids developed in [IUTchII].

**Proposition 2.1. (Vertical Coricity and Kummer Theory of Theta Monoids)** *We maintain the notation introduced above. Also, we shall use the notation  $\text{Aut}_{\mathcal{F}^{\text{lt}}}(-)$  to denote the group of automorphisms of the  $\mathcal{F}^{\text{lt}}$ -prime-strip in parentheses. Then:*

(i) **(Vertically Coric Theta Monoids)** *In the notation of [IUTchII], Corollary 4.5, (iv), (v), there are functorial algorithms in the  $\mathcal{D}$ - and  $\mathcal{D}^+$ -prime-strips  $\dagger\mathcal{D}_{>}$ ,  $\dagger\mathcal{D}_{>}^+$  associated to the  $\Theta^{\pm\text{ell}}$  *NF-Hodge theater*  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  for constructing collections of data indexed by  $\underline{\mathbb{V}}$*

$$\underline{\mathbb{V}} \ni \underline{v} \mapsto \Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}; \quad \underline{\mathbb{V}} \ni \underline{v} \mapsto \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}$$

as well as a *global realified Frobenioid*

$$\mathcal{D}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>}^+)$$

equipped with a **bijection**  $\text{Prime}(\mathcal{D}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>}^+)) \xrightarrow{\sim} \underline{\mathbb{V}}$  and corresponding local isomorphisms, for each  $\underline{v} \in \underline{\mathbb{V}}$ , as described in detail in [IUTchII], Corollary 4.5, (v). In particular, each isomorphism of the full poly-isomorphism induced [cf. Theorem 1.5, (i)] by a **vertical arrow** of the **Gaussian log-theta-lattice** under consideration induces a compatible collection of isomorphisms

$$\Psi_{\text{env}}(^{n,m}\mathcal{D}_{>}) \xrightarrow{\sim} \Psi_{\text{env}}(^{n,m+1}\mathcal{D}_{>}); \quad \infty\Psi_{\text{env}}(^{n,m}\mathcal{D}_{>}) \xrightarrow{\sim} \infty\Psi_{\text{env}}(^{n,m+1}\mathcal{D}_{>})$$

$$\mathcal{D}_{\text{env}}^{\text{lt}}(^{n,m}\mathcal{D}_{>}^+) \xrightarrow{\sim} \mathcal{D}_{\text{env}}^{\text{lt}}(^{n,m+1}\mathcal{D}_{>}^+)$$

— where the final isomorphism of Frobenioids is compatible with the respective bijections involving “Prime(–)”, as well as with the respective local isomorphisms for each  $\underline{v} \in \underline{\mathbb{V}}$ .

(ii) (**Kummer Isomorphisms**) In the notation of [IUTchII], Corollary 4.6, (iv), (v), there are **functorial algorithms** in the  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  for constructing collections of data indexed by  $\underline{\mathbb{V}}$

$$\underline{\mathbb{V}} \ni \underline{v} \mapsto \Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}}; \quad \underline{\mathbb{V}} \ni \underline{v} \mapsto \infty\Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}}$$

as well as a global realified Frobenioid

$$\mathcal{C}_{\text{env}}^{\text{lt}}(\dagger\mathcal{HT}^{\Theta})$$

equipped with a **bijection**  $\text{Prime}(\mathcal{C}_{\text{env}}^{\text{lt}}(\dagger\mathcal{HT}^{\Theta})) \xrightarrow{\sim} \underline{\mathbb{V}}$  and corresponding local isomorphisms, for each  $\underline{v} \in \underline{\mathbb{V}}$ , as described in detail in [IUTchII], Corollary 4.6, (v). Moreover, there are **functorial algorithms** in  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  for constructing **Kummer isomorphisms**

$$\begin{aligned} \Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta}) &\xrightarrow{\sim} \Psi_{\text{env}}(\dagger\mathcal{D}_{>}); & \infty\Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta}) &\xrightarrow{\sim} \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>}) \\ \mathcal{C}_{\text{env}}^{\text{lt}}(\dagger\mathcal{HT}^{\Theta}) &\xrightarrow{\sim} \mathcal{D}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>}^+) \end{aligned}$$

— where the final isomorphism of Frobenioids is compatible with the respective bijections involving “ $\text{Prime}(-)$ ”, as well as with the respective local isomorphisms for each  $\underline{v} \in \underline{\mathbb{V}}$  — with the data discussed in (i) [cf. [IUTchII], Corollary 4.6, (iv), (v)]. Finally, the collection of data  $\Psi_{\text{env}}(\dagger\mathcal{D}_{>})$  gives rise, in a natural fashion, to an  $\mathcal{F}^+$ -prime-strip  $\mathfrak{F}_{\text{env}}^+(\dagger\mathcal{D}_{>})$  [cf. the  $\mathcal{F}^+$ -prime-strip “ $\dagger\mathfrak{F}_{\text{env}}^+$ ” of [IUTchII], Corollary 4.10, (ii)]; the global realified Frobenioid  $\mathcal{D}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>}^+)$ , equipped with the bijection  $\text{Prime}(\mathcal{D}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>}^+)) \xrightarrow{\sim} \underline{\mathbb{V}}$  and corresponding local isomorphisms, for each  $\underline{v} \in \underline{\mathbb{V}}$ , reviewed in (i), together with the  $\mathcal{F}^+$ -prime-strip  $\mathfrak{F}_{\text{env}}^+(\dagger\mathcal{D}_{>})$ , determine an  $\mathcal{F}^{\text{lt}}$ -prime-strip  $\mathfrak{F}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>})$  [cf. the  $\mathcal{F}^{\text{lt}}$ -prime-strip “ $\dagger\mathfrak{F}_{\text{env}}^{\text{lt}}$ ” of [IUTchII], Corollary 4.10, (ii)]. In particular, the first and third Kummer isomorphisms of the above display may be interpreted as [compatible] isomorphisms

$$\dagger\mathfrak{F}_{\text{env}}^+ \xrightarrow{\sim} \mathfrak{F}_{\text{env}}^+(\dagger\mathcal{D}_{>}); \quad \dagger\mathfrak{F}_{\text{env}}^{\text{lt}} \xrightarrow{\sim} \mathfrak{F}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>})$$

of  $\mathcal{F}^+$ -,  $\mathcal{F}^{\text{lt}}$ -prime-strips.

(iii) (**Kummer Theory at Bad Primes**) The portion at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  of the Kummer isomorphisms of (ii) is obtained by composing the Kummer isomorphisms of [IUTchII], Proposition 3.3, (i) — which, we recall, were defined by forming **Kummer classes** in the context of **mono-theta environments** that arise from **tempered Frobenioids** — with the isomorphisms on cohomology classes induced [cf. the upper left-hand portion of the first display of [IUTchII], Proposition 3.4, (i)] by the **full poly-isomorphism of projective systems of mono-theta environments** “ $\mathbb{M}_*^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}}) \xrightarrow{\sim} \mathbb{M}_*^{\Theta}(\dagger\underline{\mathcal{F}}_{\underline{v}})$ ” [cf. [IUTchII], Proposition 3.4; [IUTchII], Remark 4.2.1, (iv)] between projective systems of mono-theta environments that arise from tempered Frobenioids [i.e., “ $\dagger\underline{\mathcal{F}}_{\underline{v}}$ ”] and projective systems of mono-theta environments that arise from the tempered fundamental group [i.e., “ $\dagger\mathcal{D}_{>,\underline{v}}$ ”] — cf. the left-hand portion of the third display of [IUTchII], Corollary 3.6, (ii), in the context of the discussion of Remark 3.6.2, (i). Here, each “isomorphism on cohomology classes” is induced by the isomorphism on **exterior cyclotomes**

$$\Pi_{\mu}(\mathbb{M}_*^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) \xrightarrow{\sim} \Pi_{\mu}(\mathbb{M}_*^{\Theta}(\dagger\underline{\mathcal{F}}_{\underline{v}}))$$



determined by each of the isomorphisms that constitutes the full poly-isomorphism of projective systems of mono-theta environments discussed above. In particular, the **composite map**

$$\Pi_{\mu}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>, \underline{v}})) \otimes \mathbb{Q}/\mathbb{Z} \quad \rightarrow \quad (\Psi_{\dagger\mathcal{F}_{\underline{v}}^{\Theta}})^{\times\mu}$$

obtained by composing the result of applying “ $\otimes \mathbb{Q}/\mathbb{Z}$ ” to this isomorphism on exterior cyclotomes with the **natural inclusion**

$$\Pi_{\mu}(\mathbb{M}_{*}^{\Theta}(\dagger\underline{\mathcal{F}}_{\underline{v}})) \otimes \mathbb{Q}/\mathbb{Z} \quad \hookrightarrow \quad (\Psi_{\dagger\mathcal{F}_{\underline{v}}^{\Theta}})^{\times}$$

[cf. the notation of [IUTchII], Proposition 3.4, (i); the description given in [IUTchII], Proposition 1.3, (i), of the exterior cyclotome of a mono-theta environment that arises from a tempered Frobenioid] and the natural projection  $(\Psi_{\dagger\mathcal{F}_{\underline{v}}^{\Theta}})^{\times} \rightarrow (\Psi_{\dagger\mathcal{F}_{\underline{v}}^{\Theta}})^{\times\mu}$  is equal to the **zero map**.

(iv) (**Kummer Theory at Good Nonarchimedean Primes**) The **unit portion** at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}} \cap \underline{\mathbb{V}}^{\text{non}}$  of the Kummer isomorphisms of (ii) is obtained [cf. [IUTchII], Proposition 4.2, (iv)] as the unit portion of a “labeled version” of the **isomorphism of topological monoids** equipped with a **topological group action** — i.e., in the language of [AbsTopIII], Definition 3.1, (ii), the isomorphism of “**MLF-Galois TM-pairs**” — discussed in [IUTchII], Proposition 4.2, (i) [cf. also [IUTchII], Remark 1.11.1, (i), (a); [AbsTopIII], Proposition 3.2, (iv)]. In particular, the portion at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}} \cap \underline{\mathbb{V}}^{\text{non}}$  of the  $\text{Aut}_{\mathcal{F}^{\dagger}}(\dagger\mathfrak{F}_{\text{env}}^{\dagger})$ -**orbit** of the second isomorphism of the final display of (ii) may be obtained as a “labeled version” of the “**Kummer poly-isomorphism of semi-simplifications**” given in the final display of [IUTchII], Proposition 4.2, (ii).

(v) (**Kummer Theory at Good Archimedean Primes**) The **unit portion** at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}} \cap \underline{\mathbb{V}}^{\text{arc}}$  of the Kummer isomorphisms of (ii) is obtained [cf. [IUTchII], Proposition 4.4, (iv)] as the unit portion of a “labeled version” of the **isomorphism of topological monoids** discussed in [IUTchII], Proposition 4.4, (i). In particular, the portion at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}} \cap \underline{\mathbb{V}}^{\text{arc}}$  of the  $\text{Aut}_{\mathcal{F}^{\dagger}}(\dagger\mathfrak{F}_{\text{env}}^{\dagger})$ -**orbit** of the second isomorphism of the final display of (ii) may be obtained as a “labeled version” of the “**Kummer poly-isomorphism of semi-simplifications**” given in the final display of [IUTchII], Proposition 4.4, (ii) [cf. also [IUTchII], Remark 4.6.1].

(vi) (**Compatibility with Constant Monoids**) The definition of the **unit portion** of the theta monoids involved [cf. [IUTchII], Corollary 4.10, (iv)] gives rise to **natural isomorphisms**

$$\dagger\mathfrak{F}_{\Delta}^{\dagger\times} \xrightarrow{\sim} \dagger\mathfrak{F}_{\text{env}}^{\dagger\times}; \quad \mathfrak{F}_{\Delta}^{\dagger\times}(\dagger\mathcal{D}_{\Delta}^{\dagger}) \xrightarrow{\sim} \mathfrak{F}_{\text{env}}^{\dagger\times}(\dagger\mathcal{D}_{>})$$

— i.e., where the morphism induced on  $\mathcal{F}^{\dagger\times\mu}$ -prime-strips by the first displayed isomorphism is precisely the isomorphism of the first display of [IUTchII], Corollary 4.10, (iv) — of the respective associated  $\mathcal{F}^{\dagger\times}$ -prime-strips [cf. the notation of Theorem 1.5, (iii), where the label “ $n, m$ ” is replaced by the label “ $\dagger$ ”]. Moreover, these natural isomorphisms are **compatible** with the **Kummer isomorphisms** of (ii) above and Theorem 1.5, (iii).



*Proof.* The various assertions of Proposition 2.1 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 2.1.1.** The theory of **mono-theta environments** [cf. Proposition 2.1, (iii)] will play a crucial role in the theory of the present §2 [cf. Theorem 2.2, (ii); Corollary 2.3, (iv), below] in the *passage from Frobenius-like to étale-like structures* [cf. Remark 1.5.4, (i), (a)] at *bad primes*. In particular, the various **rigidity** properties of mono-theta environments established in [EtTh] play a fundamental role in ensuring that the resulting “*Kummer-detachment indeterminacies*” [cf. the discussion of Remark 1.5.4, (i)] are sufficiently mild so as to allow the establishment of the various reconstruction algorithms of interest. For this reason, we pause to review the *main properties* of mono-theta environments established in [EtTh] [cf. [EtTh], Introduction] — namely,

- (a) **cyclotomic rigidity**
- (b) **discrete rigidity**
- (c) **constant multiple rigidity**
- (d) **isomorphism class compatibility**
- (e) **Frobenioid structure compatibility**

— and the roles played by these main properties in the theory of the present series of papers. Here, we remark that “isomorphism class compatibility” [i.e., (d)] refers to compatibility with the convention that various objects of the tempered Frobenioids [and their associated base categories] under consideration are known only *up to isomorphism* [cf. [EtTh], Corollary 5.12; [EtTh], Remarks 5.12.1, 5.12.2]. In the Introduction to [EtTh], instead of referring to (d) in this form, we referred to the property of compatibility with the *topology of the tempered fundamental group*. In fact, however, this compatibility with the topology of the tempered fundamental group is a consequence of (d) [cf. [EtTh], Remarks 5.12.1, 5.12.2]. On the other hand, from the point of view of the present series of papers, the essential property of interest in this context is best understood as being the property (d).

(i) First, we recall that the significance, in the context of the theory of the present series of papers, of the *compatibility with the Frobenioid structure* of the tempered Frobenioids under consideration [i.e., (e)] — i.e., in particular, with the *monoidal portion*, equipped its natural *Galois action*, of these Frobenioids — lies in the role played by this “*Frobenius-like*” monoidal portion in performing constructions — such as, for instance, the construction of the  $\log$ -,  $\Theta$ -,  $\Theta^{\times\mu}$ -, or  $\Theta_{\text{gau}}^{\times\mu}$ -links — that are “**not bound to conventional scheme theory**”, but may be related, via **Kummer theory**, to various *étale-like structures* [cf. the discussions of Remark 1.5.4, (i); [IUTchII], Remark 3.6.2, (ii); [IUTchII], Remark 3.6.4, (ii), (v)].

(ii) Next, we consider *isomorphism class compatibility* [i.e., (d)]. As discussed above, this compatibility corresponds to regarding each of the various objects of the tempered Frobenioids [and their associated base categories] under consideration as being known only *up to isomorphism* [cf. [EtTh], Corollary 5.12; [EtTh], Remarks 5.12.1, 5.12.2]. As discussed in [IUTchII], Remark 3.6.4, (i), the significance of this property (d) in the context of the present series of papers lies in the fact that — unlike the case with the *projective systems* constituted by *Kummer towers* constructed from  $N$ -th power morphisms, which are compatible with only the

*multiplicative*, but *not the additive structures* of the  $p_v$ -adic local fields involved — *each individual object* in such a Kummer tower corresponds to a *single field* [i.e., as opposed to a projective system of multiplicative groups of fields]. This **field/ring structure** is necessary in order to apply the theory of the **log-link** developed in §1 — cf. the *vertical coricity* discussed in Proposition 2.1, (i). Note, moreover, that, unlike the **log-**,  **$\Theta$ -**,  **$\Theta^{\times\mu}$ -**, or  **$\Theta_{\text{gau}}^{\times\mu}$ -**links, the  $N$ -th power morphisms that appear in a Kummer tower are “*algebraic*”, i.e., **scheme-theoretic**, hence compatible with the conventional scheme-theoretic theory of the étale [or tempered] fundamental group. In particular, since the tempered Frobenioids under consideration may be constructed from such scheme-theoretic categories, the fundamental groups on either side of such an  $N$ -th power morphism may be related *up to an indeterminacy arising from an inner automorphism* of the tempered fundamental group [i.e., the “fundamental group” of the base category] under consideration — cf. the discussion of [IUTchII], Remark 3.6.3, (ii). On the other hand, the objects that appear in these Kummer towers necessarily arise from *nontrivial line bundles* [indeed, line bundles all of whose positive tensor powers are nontrivial!] on tempered coverings of a Tate curve — cf. the constructions underlying the Frobenioid-theoretic version of the mono-theta environment [cf. [EtTh], Proposition 1.1; [EtTh], Lemma 5.9]; the crucial role played by the *commutator* “[ $-$ ,  $-$ ]” in the theory of *cyclotomic rigidity* [i.e., (a)] reviewed in (iv) below. In particular, the extraction of various  $N$ -th roots in a Kummer tower necessarily leads to *mutually non-isomorphic line bundles*, i.e., mutually non-isomorphic objects in the Kummer tower. From the point of view of *reconstruction algorithms*, such non-isomorphic objects may be **naturally** — i.e., **algorithmically** — related to another only via **indeterminate isomorphisms** [cf., (d)!]. This point of view is precisely the starting point of the discussion of — for instance, “*constant multiple indeterminacy*” in — [EtTh], Remarks 5.12.2, 5.12.3.

(iii) Next, we recall that the significance of *constant multiple rigidity* [i.e., (c)] in the context of the present series of papers lies in the construction of the **canonical splittings of theta monoids via restriction to the zero section** discussed, for instance, in [IUTchII], Corollary 1.12, (ii); [IUTchII], Proposition 3.3, (i); [IUTchII], Remark 1.12.2, (iv) [cf. also Remark 1.2.3, (i), of the present paper].

(iv) Next, we review the significance of *cyclotomic rigidity* [i.e., (a)] in the context of the present series of papers. First, we recall that this cyclotomic rigidity is essentially a consequence of the *nondegenerate* nature of the *commutator* “[ $-$ ,  $-$ ]” of the theta groups involved [cf. the discussion of [EtTh], Introduction; [EtTh], Remark 2.19.2]. Put another way, since this commutator is quadratic in nature, one may think of this nondegenerate nature of the commutator as a statement to the effect that “*the degree of the commutator is precisely 2*”. At a more concrete level, the cyclotomic rigidity arising from a mono-theta environment consists of a certain specific isomorphism between the *interior* and *exterior cyclotomes* [cf. the discussion of [IUTchII], Definition 1.1, (ii)]. Put another way, one may think of this cyclotomic rigidity isomorphism as a sort of rigidification of a certain “*projective line of cyclotomes*”, i.e., the projectivization of the direct sum of the interior and exterior cyclotomes [cf. the computations that underlie [EtTh], Proposition 2.12]. In particular, this rigidification is fundamentally *nonlinear* in nature. Indeed, if one attempts to compose it with an  $N$ -th power morphism, then one is obliged to sacrifice constant multiple rigidity [i.e., (c)] — cf. the discussion of [EtTh],

Remark 5.12.3. That is to say, the *distinguished nature* of the “**first power**” of the cyclotomic rigidity isomorphism is an important theme in the theory of [EtTh] [cf. the discussion of [EtTh], Remark 5.12.5; [IUTchII], Remark 3.6.4, (iii), (iv)]. The **multiradiality** of mono-theta-theoretic cyclotomic rigidity [cf. [IUTchII], Corollary 1.10] — which lies in stark contrast with the indeterminacies that arise when one attempts to give a multiradial formulation [cf. [IUTchII], Corollary 1.11; the discussion of [IUTchII], Remark 1.11.3] of the more classical “*MLF-Galois pair cyclotomic rigidity*” arising from local class field theory — will play a *central role* in the theory of the present §2 [cf. Theorem 2.2, Corollary 2.3 below].

(v) Finally, we review the significance of *discrete rigidity* [i.e., (b)] in the context of the present series of papers. First, we recall that, at a technical level, whereas cyclotomic rigidity may be regarded [cf. the discussion of (iv)] as a consequence of the fact that “the degree of the commutator is precisely 2”, discrete rigidity may be regarded as a consequence of the fact that “*the degree of the commutator is  $\leq 2$* ” [cf. the statements and proofs of [EtTh], Proposition 2.14, (ii), (iii)]. At a more concrete level, discrete rigidity assures one that one may restrict one’s attentions to  **$\mathbb{Z}$ -multiples/powers** — as opposed to  **$\widehat{\mathbb{Z}}$ -multiples/powers** — of divisors, line bundles, and rational functions [such as, for instance, the *q-parameter!*] on the tempered coverings of a Tate curve that occur in the theory of [EtTh] [cf. [EtTh], Remark 2.19.4]. This prompts the following question:

Can one develop a theory of  $\widehat{\mathbb{Z}}$ -divisors/line bundles/rational functions in, for instance, a parallel fashion to the way in which one considers *perfections* and *realifications* of Frobenioids in the theory of [FrdI]?

As far as the author can see at the time of writing, the answer to this question is “*no*”. Indeed, unlike the case with  $\mathbb{Q}$  or  $\mathbb{R}$ , there is no notion of **positivity** [or negativity] in  $\widehat{\mathbb{Z}}$ . For instance,  $-1 \in \widehat{\mathbb{Z}}$  may be obtained as a limit of positive integers. In particular, if one had a theory of  $\widehat{\mathbb{Z}}$ -divisors/line bundles/rational functions, then such a theory would necessarily require one to “confuse” positive [i.e., effective] and negative divisors, hence to work *birationally*. But to work birationally means, in particular, that one must sacrifice the conventional structure of isomorphisms [e.g., automorphisms] between line bundles — which plays an indispensable role, for instance, in the constructions underlying the Frobenioid-theoretic version of the mono-theta environment [cf. [EtTh], Proposition 1.1; [EtTh], Lemma 5.9; the crucial role played by the *commutator* “[ $-$ ,  $-$ ]” in the theory of *cyclotomic rigidity* [i.e., (a)] reviewed in (iv) above].

### Remark 2.1.2.

(i) In the context of the discussion of Remark 2.1.1, (v), it is of interest to recall [cf. [IUTchII], Remark 4.5.3, (iii); [IUTchII], Remark 4.11.2, (iii)] that the essential role played, in the context of the  $\mathbb{F}_l^{\times \pm}$ -**symmetry**, by the “*global bookkeeping operations*” involving the **labels** of the evaluation points gives rise, in light of the **profinite** nature of the global étale fundamental groups involved, to a situation in which one must apply the “complements on tempered coverings” developed in [IUTchI], §2. That is to say, in the notation of the discussion given in [IUTchII],

Remark 2.1.1, (i), of the various tempered coverings that occur at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , these “complements on tempered coverings” are applied precisely so as to allow one to restrict one’s attention to the [*discrete!*]  $\underline{\mathbb{Z}}$ -conjugates — i.e., as opposed to [*profinite!*]  $\widehat{\underline{\mathbb{Z}}}$ -conjugates [where we write  $\widehat{\underline{\mathbb{Z}}}$  for the profinite completion of  $\underline{\mathbb{Z}}$ ] — of the theta functions involved. In particular, although such “evaluation-related issues”, which will become relevant in the context of the theory of §3 below, do not play a role in the theory of the present §2, the role played by the theory of [IUTchI], §2, in the theory of the present series of papers may also be thought of as a sort of “discrete rigidity” — which we shall refer to as “**evaluation discrete rigidity**” — i.e., a sort of rigidity that is concerned with similar issues to the issues discussed in the case of “*mono-theta-theoretic discrete rigidity*” in Remark 2.1.1, (v), above.

(ii) Next, let us suppose that we are in the situation discussed in [IUTchII], Proposition 2.1. Fix  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ . Write  $\Pi \stackrel{\text{def}}{=} \Pi_{\underline{v}}$ ;  $\widehat{\Pi}$  for the profinite completion of  $\Pi$ . Thus, we have *natural surjections*  $\Pi \twoheadrightarrow l \cdot \underline{\mathbb{Z}} (\subseteq \underline{\mathbb{Z}})$ ,  $\widehat{\Pi} \twoheadrightarrow l \cdot \widehat{\underline{\mathbb{Z}}} (\subseteq \widehat{\underline{\mathbb{Z}}})$ . Write  $\Pi^\dagger \stackrel{\text{def}}{=} \widehat{\Pi} \times_{\widehat{\underline{\mathbb{Z}}}} \underline{\mathbb{Z}} \subseteq \widehat{\Pi}$ . Next, we observe that from the point of view of the *evaluation points*, the evaluation discrete rigidity discussed in (i) corresponds to the issue of whether, relative to some arbitrarily chosen basepoint, the **coordinates** of the evaluation point lie  $\in \underline{\mathbb{Z}}$  or  $\in \widehat{\underline{\mathbb{Z}}}$  [cf. the discussion of the “torsor over  $\underline{\mathbb{Z}}$ ” in [IUTchII], Remark 2.1.1, (i)]. Thus, if one is only concerned with the issue of arranging for these coordinates to lie  $\in \underline{\mathbb{Z}}$ , then one is led to pose the following question:

Is it possible to simply use the “*partially tempered fundamental group*”  $\Pi^\dagger$  instead of the “full” tempered fundamental group  $\Pi$  in the theory of the present series of papers?

The answer to this question is “*no*”. One way to see this is to consider the [easily verified] natural isomorphism

$$N_{\widehat{\Pi}}(\Pi^\dagger)/\Pi^\dagger \xrightarrow{\sim} \widehat{\underline{\mathbb{Z}}}/\underline{\mathbb{Z}}$$

involving the *normalizer*  $N_{\widehat{\Pi}}(\Pi^\dagger)$  of  $\Pi^\dagger$  in  $\widehat{\Pi}$ . One consequence of this isomorphism is that — unlike the tempered fundamental group  $\Pi$  [cf., e.g., [SemiAnbd], Theorems 6.6, 6.8] — the topological group  $\Pi^\dagger$  *fails to satisfy various fundamental absolute anabelian properties* which play a *crucial role* in the theory of [EtTh], as well as in the present series of papers [cf., e.g., the theory of [IUTchII], §2]. At a more concrete level, unlike the case with the tempered fundamental group  $\Pi$ , the *profinite conjugacy indeterminacies* that act on  $\Pi^\dagger$  give rise to  $\widehat{\underline{\mathbb{Z}}}$ -translation *indeterminacies* acting on the coordinates of the evaluation points involved. That is to say, in the case of  $\Pi$ , such  $\widehat{\underline{\mathbb{Z}}}$ -translation indeterminacies are avoided precisely by applying the “complements on tempered coverings” developed in [IUTchI], §2 — i.e., in a word, as a consequence of the “*highly anabelian nature*” of the [full!] tempered fundamental group  $\Pi$ .

**Theorem 2.2.** (Kummer-compatible Multiradiality of Theta Monoids)  
Fix a collection of **initial  $\Theta$ -data**

$$(\overline{F}/F, X_F, l, \underline{C}_K, \underline{\mathbb{V}}, \underline{\epsilon})$$

as in [IUTchI], Definition 3.1. Let  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  be a  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater [relative to the given initial  $\Theta$ -data — cf. [IUTchI], Definition 6.13, (i)]. For  $\square \in \{\vdash, \vdash \blacktriangleright \times \mu, \vdash \times \mu\}$ , write  $\text{Aut}_{\mathcal{F}^\square}(-)$  for the group of automorphisms of the  $\mathcal{F}^\square$ -prime-strip in parentheses.

(i) **(Automorphisms of Prime-strips)** The natural functors determined by assigning to an  $\mathcal{F}^{\vdash}$ -prime-strip the associated  $\mathcal{F}^{\vdash \blacktriangleright \times \mu}$ - and  $\mathcal{F}^{\vdash \times \mu}$ -prime-strips [cf. [IUTchII], Definition 4.9, (vi), (vii), (viii)] and then composing with the natural isomorphisms of Proposition 2.1, (vi), determine natural homomorphisms

$$\begin{aligned} \text{Aut}_{\mathcal{F}^{\vdash}}(\mathfrak{F}_{\text{env}}^{\vdash}(\dagger\mathcal{D}_{>})) &\rightarrow \text{Aut}_{\mathcal{F}^{\vdash \blacktriangleright \times \mu}}(\mathfrak{F}_{\text{env}}^{\vdash \blacktriangleright \times \mu}(\dagger\mathcal{D}_{>})) \twoheadrightarrow \text{Aut}_{\mathcal{F}^{\vdash \times \mu}}(\mathfrak{F}_{\Delta}^{\vdash \times \mu}(\dagger\mathcal{D}_{\Delta}^{\vdash})) \\ \text{Aut}_{\mathcal{F}^{\vdash}}(\dagger\mathfrak{F}_{\text{env}}^{\vdash}) &\rightarrow \text{Aut}_{\mathcal{F}^{\vdash \blacktriangleright \times \mu}}(\dagger\mathfrak{F}_{\text{env}}^{\vdash \blacktriangleright \times \mu}) \twoheadrightarrow \text{Aut}_{\mathcal{F}^{\vdash \times \mu}}(\dagger\mathfrak{F}_{\Delta}^{\vdash \times \mu}) \end{aligned}$$

— where the second arrows in each line are surjections — that are **compatible** with the **Kummer isomorphisms** of Proposition 2.1, (ii), and Theorem 1.5, (iii) [cf. the final portions of Proposition 2.1, (iv), (v), (vi)].

(ii) **(Kummer Aspects of Multiradiality at Bad Primes)** Let  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ . Write

$$\infty\Psi_{\text{env}}^{\perp}(\dagger\mathcal{D}_{>})_{\underline{v}} \subseteq \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}; \quad \infty\Psi_{\mathcal{F}_{\text{env}}}^{\perp}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}} \subseteq \infty\Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}}$$

for the submonoids corresponding to the respective **splittings** [cf. [IUTchII], Corollaries 3.5, (iii); 3.6, (iii)], i.e., the submonoids generated by “ $\infty_{\underline{v}}^{\theta^{\perp}}(\mathbb{M}_{*}^{\Theta})$ ” [cf. the notation of [IUTchII], Proposition 3.1, (i)] and the respective **torsion subgroups**. Now consider the commutative diagram

$$\begin{array}{ccccc} \infty\Psi_{\text{env}}^{\perp}(\dagger\mathcal{D}_{>})_{\underline{v}} & \supseteq & \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}^{\mu} & \subseteq & \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}^{\times} \\ \downarrow & & \downarrow & & \downarrow \\ \infty\Psi_{\mathcal{F}_{\text{env}}}^{\perp}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}} & \supseteq & \infty\Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}}^{\mu} & \subseteq & \infty\Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}}^{\times} \\ & & \twoheadrightarrow & \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}^{\times\mu} & \xrightarrow{\sim} \Psi_{\text{cns}}^{\text{ss}}(\dagger\mathcal{D}_{\Delta}^{\vdash})_{\underline{v}}^{\times\mu} \\ & & & \downarrow & \downarrow \\ & & \twoheadrightarrow & \infty\Psi_{\mathcal{F}_{\text{env}}}(\dagger\mathcal{HT}^{\Theta})_{\underline{v}}^{\times\mu} & \xrightarrow{\sim} \Psi_{\text{cns}}^{\text{ss}}(\dagger\mathfrak{F}_{\Delta}^{\vdash})_{\underline{v}}^{\times\mu} \end{array}$$

— where the inclusions “ $\supseteq$ ”, “ $\subseteq$ ” are the natural inclusions; the surjections “ $\twoheadrightarrow$ ” are the natural surjections; the superscript “ $\mu$ ” denotes the torsion subgroup; the superscript “ $\times$ ” denotes the group of units; the superscript “ $\times\mu$ ” denotes the quotient “ $(-)^{\times}/(-)^{\mu}$ ”; the first four vertical arrows are the isomorphisms determined by the inverse of the second **Kummer isomorphism** of the third display of Proposition 2.1, (ii);  $\dagger\mathcal{D}_{\Delta}^{\vdash}$  is as discussed in Theorem 1.5, (iii);  $\dagger\mathfrak{F}_{\Delta}^{\vdash}$  is as discussed in [IUTchII], Corollary 4.10, (i); the final vertical arrow is the inverse of the **Kummer isomorphism** determined by the final displayed isomorphism of [IUTchII], Corollary 4.6, (i) [cf. also the isomorphism of the fourth display of [IUTchII], Corollary 4.5, (ii)]; the final upper horizontal arrow is the **poly-isomorphism** determined by composing the isomorphism determined by the inverse of the natural



isomorphism of Proposition 2.1, (vi), with the poly-automorphism of  $\Psi_{\text{cns}}^{\text{ss}}(\dagger\mathcal{D}_{\Delta}^{\dagger})_{\underline{v}}^{\times\mu}$  induced by the **full poly-automorphism** of the  $\mathcal{D}^{\dagger}$ -prime-strip  $\dagger\mathcal{D}_{\Delta}^{\dagger}$ ; the final lower horizontal arrow is the poly-automorphism determined by the condition that the final square be commutative. This commutative diagram is compatible with the various group actions involved relative to the following diagram

$$\begin{aligned} \Pi_{\underline{X}}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) &\twoheadrightarrow G_{\underline{v}}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) = G_{\underline{v}}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) \\ &= G_{\underline{v}}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) \xrightarrow{\sim} G_{\underline{v}}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) \end{aligned}$$

[cf. the notation of [IUTchII], Proposition 3.1; [IUTchII], Remark 4.2.1, (iv); [IUTchII], Corollary 4.5, (iv)] — where “ $\twoheadrightarrow$ ” denotes the natural surjection; “ $\xrightarrow{\sim}$ ” denotes the full poly-automorphism of  $G_{\underline{v}}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}}))$ . Finally, each of the various composite maps

$$\infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}^{\mu} \rightarrow \Psi_{\text{cns}}^{\text{ss}}(\dagger\mathfrak{F}_{\Delta}^{\dagger})_{\underline{v}}^{\times\mu}$$

is equal to the **zero map** [cf. (b<sub>v</sub>) below; the final portion of Proposition 2.1, (iii)]. In particular, the **identity** automorphism on the following objects is **compatible**, relative to the various natural morphisms involved [cf. the above commutative diagram], with the collection of automorphisms of  $\Psi_{\text{cns}}^{\text{ss}}(\dagger\mathfrak{F}_{\Delta}^{\dagger})_{\underline{v}}^{\times\mu}$  induced by **arbitrary automorphisms**  $\in \text{Aut}_{\mathcal{F}^{\dagger \times \mu}}(\dagger\mathfrak{F}_{\Delta}^{\dagger \times \mu})$  [cf. [IUTchII], Corollary 1.12, (iii); [IUTchII], Proposition 3.4, (i)]:

$$(a_{\underline{v}}) \quad \infty\Psi_{\text{env}}^{\perp}(\dagger\mathcal{D}_{>})_{\underline{v}} \supseteq \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}^{\mu};$$

(b<sub>v</sub>)  $\Pi_{\mu}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) \otimes \mathbb{Q}/\mathbb{Z}$  [cf. the discussion of Proposition 2.1, (iii)], relative to the natural isomorphism  $\Pi_{\mu}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})) \otimes \mathbb{Q}/\mathbb{Z} \xrightarrow{\sim} \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}^{\mu}$  of [IUTchII], Remark 1.5.2 [cf. (a<sub>v</sub>)];

(c<sub>v</sub>) the projective system of **mono-theta environments**  $\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}})$  [cf. (b<sub>v</sub>)];

(d<sub>v</sub>) the splittings  $\infty\Psi_{\text{env}}^{\perp}(\dagger\mathcal{D}_{>})_{\underline{v}} \twoheadrightarrow \infty\Psi_{\text{env}}(\dagger\mathcal{D}_{>})_{\underline{v}}^{\mu}$  [cf. (a<sub>v</sub>)] by means of restriction to **zero-labeled evaluation points** [cf. [IUTchII], Proposition 3.1, (i)].

*Proof.* The various assertions of Theorem 2.2 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 2.2.1.** In light of the *central importance* of Theorem 2.2, (ii), in the theory of the present §2, we pause to examine the significance of Theorem 2.2, (ii), in more conceptual terms.

(i) In the situation of Theorem 2.2, (ii), let us write [for simplicity]  $\Pi_{\underline{v}} \stackrel{\text{def}}{=} \Pi_{\underline{X}}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}}))$ ,  $\Pi_{\mu} \stackrel{\text{def}}{=} \Pi_{\mu}(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}}))$  [cf. (b<sub>v</sub>)]. Also, for simplicity, we write  $(l \cdot \Delta_{\Theta}) \stackrel{\text{def}}{=} (l \cdot \Delta_{\Theta})(\mathbb{M}_{*}^{\Theta}(\dagger\mathcal{D}_{>,\underline{v}}))$  [cf. [IUTchII], Proposition 1.5, (iii)]. Here, we recall that in fact,  $(l \cdot \Delta_{\Theta})$  may be thought of as an object *constructed from*  $\Pi_{\underline{v}}$



[cf. [IUTchII], Proposition 1.4]. Then the projective system of mono-theta environments  $\mathbb{M}_*^\Theta(\dagger\mathcal{D}_{>,v})$  [cf.  $(c_v)$ ] may be thought of as a sort of “*amalgamation of  $\Pi_v$  and  $\Pi_\mu$* ”, where the amalgamation is such that it allows the *reconstruction of the mono-theta-theoretic cyclotomic rigidity isomorphism*

$$(l \cdot \Delta_\Theta) \xrightarrow{\sim} \Pi_\mu$$

[cf. [IUTchII], Proposition 1.5, (iii)] — i.e., not just the  $\widehat{\mathbb{Z}}^\times$ -*orbit* of this isomorphism!

(ii) Now, in the notation of (i), the *Kummer classes*  $\in {}_\infty\Psi_{\text{env}}^\perp(\dagger\mathcal{D}_{>,v})$  [cf.  $(a_v)$ ] constituted by the various *étale theta functions* may be thought of, for an appropriate characteristic open subgroup  $H \subseteq \Pi_v$ , as *twisted homomorphisms*

$$(\Pi_v \supseteq) H \rightarrow \Pi_\mu$$

whose restriction to  $(l \cdot \Delta_\Theta)$  coincides with the cyclotomic rigidity isomorphism  $(l \cdot \Delta_\Theta) \xrightarrow{\sim} \Pi_\mu$  discussed in (i). Then the essential content of Theorem 2.2, (ii), lies in the observation that

since the **Kummer-theoretic link** between étale-like data and Frobenius-like data at  $v \in \underline{\mathbb{V}}^{\text{bad}}$  is established by means of projective systems of **mono-theta environments** [cf. the discussion of Proposition 2.1, (iii)] — i.e., which do *not* involve the various monoids “ $(-)^{\times\mu}$ ”! — the **mono-theta-theoretic cyclotomic rigidity isomorphism** [i.e., *not just the  $\widehat{\mathbb{Z}}^\times$ -orbit* of this isomorphism!] is **immune** to the various automorphisms of the monoids “ $(-)^{\times\mu}$ ” which, from the point of view of the **multiradial formulation** to be discussed in Corollary 2.3 below, arise from isomorphisms of *coric data*.

Put another way, this “immunity” may be thought of as a sort of **decoupling** of the “*geometric*” [i.e., in the sense of the geometric fundamental group  $\Delta_v \subseteq \Pi_v$ ] and “*base-field-theoretic*” [i.e., associated to the local absolute Galois group  $\Pi_v \twoheadrightarrow G_v$ ] data which allows one to treat the exterior cyclotome  $\Pi_\mu$  — which, *a priori*, “looks base-field-theoretic” — as being part of the “geometric” data. From the point of view of the multiradial formulation to be discussed in Corollary 2.3 below [cf. also the discussion of [IUTchII], Remark 1.12.2, (vi)], this decoupling may be thought of as a sort of **splitting** into **purely radial** and **purely coric** components — i.e., with respect to which  $\Pi_\mu$  is “*purely radial*”, while the various monoids “ $(-)^{\times\mu}$ ” are “*purely coric*”.

(iii) Note that the immunity to automorphisms of the monoids “ $(-)^{\times\mu}$ ” discussed in (ii) lies in *stark contrast* to the  $\widehat{\mathbb{Z}}^\times$ -**indeterminacies** that arise in the case of the cyclotomic rigidity isomorphisms constructed from **MLF-Galois pairs** in a fashion that makes *essential use of the monoids* “ $(-)^{\times\mu}$ ”, as discussed in [IUTchII], Corollary 1.11; [IUTchII], Remark 1.11.3. In the following discussion, let us write “ $\mathcal{O}^{\times\mu}$ ” for the various monoids “ $(-)^{\times\mu}$ ” that occur in the situation of Theorem 2.2; also, we shall use similar notation “ $\mathcal{O}^\mu$ ”, “ $\mathcal{O}^\times$ ”, “ $\mathcal{O}^\triangleright$ ”, “ $\mathcal{O}^{\text{gp}}$ ”, “ $\mathcal{O}^{\widehat{\text{gp}}}$ ” [cf. the

notational conventions of [IUTchII], Example 1.8, (iv), (vii)]. Thus, we have a diagram

$$\begin{array}{ccccccccc} \mathcal{O}^\mu & \subseteq & \mathcal{O}^\times & \subseteq & \mathcal{O}^\triangleright & \subseteq & \mathcal{O}^{\text{gp}} & \subseteq & \widehat{\mathcal{O}}^{\text{gp}} \\ & & \searrow & & \downarrow & & & & \\ & & & & \mathcal{O}^{\times\mu} & & & & \end{array}$$

of natural morphisms between monoids equipped with  $\Pi_v$ -actions. Relative to this notation, the essential *input data* for the cyclotomic rigidity isomorphism constructed from an MLF-Galois pair is given by “ $\mathcal{O}^\triangleright$ ” [cf. [IUTchII], Corollary 1.11, (a)]. On the other hand — unlike the case with  $\mathcal{O}^\mu$  — a  $\widehat{\mathbb{Z}}^\times$ -indeterminacy acting on  $\mathcal{O}^{\times\mu}$  does not lie under an *identity action* on  $\mathcal{O}^\times$ ! That is to say, a  $\widehat{\mathbb{Z}}^\times$ -indeterminacy acting on  $\mathcal{O}^{\times\mu}$  can only be *lifted naturally* to  $\widehat{\mathbb{Z}}^\times$ -indeterminacies on  $\mathcal{O}^\times$ ,  $\widehat{\mathcal{O}}^{\text{gp}}$  [cf. Fig. 2.1 below; [IUTchII], Corollary 1.11, (a), in the case where one takes “ $\Gamma$ ” to be  $\widehat{\mathbb{Z}}^\times$ ; [IUTchII], Remark 1.11.3, (ii)]. In the presence of such  $\widehat{\mathbb{Z}}^\times$ -indeterminacies, one can only recover the  $\widehat{\mathbb{Z}}^\times$ -orbit of the MLF-Galois-pair-theoretic cyclotomic rigidity isomorphism.

$$\begin{array}{ccc} \widehat{\mathbb{Z}}^\times \curvearrowright & & \widehat{\mathbb{Z}}^\times \curvearrowright & & \widehat{\mathbb{Z}}^\times \curvearrowright \\ \boxed{\begin{array}{c} \mathcal{O}^{\times\mu} \leftarrow \mathcal{O}^\times \subseteq \mathcal{O}^\triangleright \subseteq \mathcal{O}^{\text{gp}} \subseteq \widehat{\mathcal{O}}^{\text{gp}} \\ (\supseteq \mathcal{O}^\mu) \end{array}} \end{array}$$

Fig. 2.1: *Induced  $\widehat{\mathbb{Z}}^\times$ -indeterminacies* in the case of MLF-Galois pair cyclotomic rigidity

$$\begin{array}{ccc} \text{id} \curvearrowright & & \widehat{\mathbb{Z}}^\times \curvearrowright \\ \boxed{\Pi_\mu \xrightarrow{\sim} \mathcal{O}^\mu} & \rightarrow & \boxed{\mathcal{O}^{\times\mu}} \end{array}$$

Fig. 2.2: *Insulation from  $\widehat{\mathbb{Z}}^\times$ -indeterminacies* in the case of mono-theta-theoretic cyclotomic rigidity

(iv) Thus, in summary, [cf. Fig. 2.2 above]

**mono-theta-theoretic cyclotomic rigidity** plays an essential role in the theory of the present §2 — and, indeed, in the theory of the present series of papers! — in that it serves to **insulate** the **étale theta function** from the  $\widehat{\mathbb{Z}}^\times$ -**indeterminacies** which act on the **coric log-shells** [i.e., the various monoids “ $(-)^{\times\mu}$ ”].

The techniques that underlie the resulting *multiradiality of theta monoids* [cf. Corollary 2.3 below], cannot, however, be applied immediately to the case of *Gaussian*

*monoids*. That is to say, the corresponding multiradiality of Gaussian monoids, to be discussed in §3 below, requires one to apply the theory of *log-shells* developed in §1 [cf. [IUTchII], Remark 2.9.1, (iii); [IUTchII], Remark 3.4.1, (ii); [IUTchII], Remark 3.7.1]. On the other hand, as we shall see in §3 below, the multiradiality of Gaussian monoids **depends** *in an essential way* on the multiradiality of theta monoids discussed in the present §2 as a sort of “*essential first step*” constituted by the *decoupling* discussed in (ii) above. Indeed, if one tries to consider the **Kummer theory** of the **theta values** [i.e., the “ $\underline{q}^{\underline{j}^2}$ ” — cf. [IUTchII], Remark 2.5.1, (i)] just as elements of the *base field* — i.e., *without availing oneself of the theory of the étale theta function* — then it is difficult to see how to *rigidify the cyclotomes* involved by any means other than the theory of *MLF-Galois pairs* discussed in (iii) above. But, as discussed in (iii) above, this approach to cyclotomic rigidity gives rise to  $\widehat{\mathbb{Z}}^\times$ -*indeterminacies* — i.e., to *confusion* between the theta values “ $\underline{q}^{\underline{j}^2}$ ” and their  $\widehat{\mathbb{Z}}^\times$ -*powers*, which is unacceptable from the point of view of the theory of the present series of papers! For another approach to understanding the indispensability of the multiradiality of theta monoids, we refer to Remark 2.2.2 below.

**Remark 2.2.2.**

(i) One way to understand the very *special role* played by the **theta values** [i.e., the values of the theta function] in the theory of the present series of papers is to consider the following naive question:

Can one develop a similar theory to the theory of the present series of papers in which one replaces the  $\Theta_{\text{gau}}^{\times\mu}$ -link

$$\underline{q} \mapsto \underline{q}^{\begin{pmatrix} 1^2 \\ \vdots \\ (l^*)^2 \end{pmatrix}}$$

[cf. [IUTchII], Remark 4.11.1] by a correspondence of the form

$$\underline{q} \mapsto \underline{q}^\lambda$$

— where  $\lambda$  is some *arbitrary positive integer*?

The answer to this question is “*no*”. Indeed, such a correspondence does not come equipped with the extensive **multiradiality** machinery — such as **mono-theta-theoretic cyclotomic rigidity** and the **splittings** determined by **zero-labeled evaluation points** — that has been developed for the the étale theta function. For instance, the lack of mono-theta-theoretic cyclotomic rigidity means that one does not have an apparatus for **insulating** the **Kummer classes** of such a correspondence from the  $\widehat{\mathbb{Z}}^\times$ -*indeterminacies* that act on the various monoids “ $(-)^{\times\mu}$ ” [cf. the discussion of Remark 2.2.1, (iv)]. The splittings determined by zero-labeled evaluation points also play an essential role in **decoupling** these monoids “ $(-)^{\times\mu}$ ” — i.e., the **coric log-shells** — from the “**purely radial**” [or,

put another way, “value group”] portion of such a correspondence “ $\underline{q} \mapsto \underline{q}^\lambda$ ” [cf. Remark 2.2.1, (ii); [IUTchII], Remark 1.12.2, (vi)]. Note, moreover, that if one tries to realize such a multiradial splitting via *evaluation* — i.e., in accordance with the principle of “**Galois evaluation**” [cf. the discussion of [IUTchII], Remark 1.12.4] — for a correspondence “ $\underline{q} \mapsto \underline{q}^\lambda$ ” by, for instance, taking  $\lambda$  to be *one of the “ $j^2$ ”* [where  $j$  is a positive integer] that appears as a value of the étale theta function, then one must contend with issues of **symmetry** between the zero-labeled evaluation point and the evaluation point corresponding to  $\lambda$  — i.e., symmetry issues that are resolved in the theory of the present series of papers by means of the theory surrounding the  $\mathbb{F}_l^{\times\pm}$ -**symmetry** [cf. the discussion of [IUTchII], Remarks 2.6.2, 3.5.2]. As discussed in [IUTchII], Remark 2.6.3, this sort of situation leads to numerous *conditions on the collection of evaluation points* under consideration. In particular, ultimately, it is difficult to see how to construct a theory as in the present series of papers for any collection of evaluation points other than the collection that is in fact adopted in the definition of the  $\Theta_{\text{gau}}^{\times\mu}$ -link.

(ii) As discussed in Remark 2.2.1, (iv), we shall be concerned, in §3 below, with developing multiradial formulations for Gaussian monoids. These multiradial formulations will be subject to certain *indeterminacies*, which — although *sufficiently mild* to allow the execution of the *volume computations* that will be the subject of [IUTchIV] — are, nevertheless, *substantially more severe* than the indeterminacies that occur in the multiradial formulation given for theta monoids in the present §2 [cf. Corollary 2.3 below]. Indeed, the indeterminacies in the multiradial formulation given for theta monoids in the present §2 — which essentially consist of *multiplication by roots of unity* [cf. [IUTchII], Proposition 3.1, (i)] — are *essentially negligible* and may be regarded as a consequence of the highly nontrivial **Kummer theory** surrounding **mono-theta environments** [cf. Proposition 2.1, (iii); Theorem 2.2, (ii)], which, as discussed in Remark 2.2.1, (iv), cannot be mimicked for “theta values regarded just as elements of the base field”. That is to say, the quite **exact** nature of the multiradial formulation for theta monoids — i.e., which contrasts sharply with the somewhat **approximate** nature of the multiradial formulation for Gaussian monoids to be developed in §3 — constitutes another *important ingredient* of the theory of the present paper that one must sacrifice if one attempts to work with correspondences  $\underline{q} \mapsto \underline{q}^\lambda$  as discussed in (i), i.e., correspondences which do not come equipped with the extensive multiradiality machinery that arises as a consequence of the theory of the *étale theta function* developed in [EtTh].

We conclude the present §3 with the following **multiradial** interpretation [cf. [IUTchII], Remark 4.1.1, (iii); [IUTchII], Remark 4.3.1] — in the spirit of the *étale-picture of  $\mathcal{D}$ - $\Theta^{\pm\text{ell}}$  NF-Hodge theaters* of [IUTchII], Corollary 4.11, (ii) — of the theory surrounding Theorem 2.2.

**Corollary 2.3.**     **(Étale-picture of Multiradial Theta Monoids)** *In the notation of Theorem 2.2, let*

$$\{ {}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \}_{n,m \in \mathbb{Z}}$$

*be a collection of distinct  $\Theta^{\pm\text{ell}}$  NF-Hodge theaters [relative to the given initial  $\Theta$ -data] — which we think of as arising from a Gaussian log-theta-lattice [cf.*

*Definition 1.4*]. Write  ${}^{n,m}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  for the  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater associated to  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ . Consider the **radial environment** [cf. [IUTchII], Example 1.7, (ii)] defined as follows. We define a collection of **radial data**

$$\dagger\mathfrak{R} = (\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}, \mathfrak{F}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>}), \dagger\mathfrak{R}^{\text{bad}}, \mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\dagger\mathcal{D}_{\Delta}^{\text{lt}}), \mathfrak{F}_{\text{env}}^{\text{lt}\times\mu}(\dagger\mathcal{D}_{>}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\dagger\mathcal{D}_{\Delta}^{\text{lt}}))$$

to consist of

- ( $a_{\mathfrak{R}}$ ) a  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  $\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$ ;
- ( $b_{\mathfrak{R}}$ ) the  $\mathcal{F}^{\text{lt}}$ -prime-strip  $\mathfrak{F}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>})$  associated to  $\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  [cf. Proposition 2.1, (ii)];
- ( $c_{\mathfrak{R}}$ ) the data  $(a_{\underline{v}}), (b_{\underline{v}}), (c_{\underline{v}}), (d_{\underline{v}})$  of Theorem 2.2, (ii), for  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , which we denote by  $\dagger\mathfrak{R}^{\text{bad}}$ ;
- ( $d_{\mathfrak{R}}$ ) the  $\mathcal{F}^{\text{lt}\times\mu}$ -prime-strip  $\mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\dagger\mathcal{D}_{\Delta}^{\text{lt}})$  associated to  $\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  [cf. Theorem 1.5, (iii)];
- ( $e_{\mathfrak{R}}$ ) the **full poly-isomorphism** of  $\mathcal{F}^{\text{lt}\times\mu}$ -prime-strips  $\mathfrak{F}_{\text{env}}^{\text{lt}\times\mu}(\dagger\mathcal{D}_{>}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\dagger\mathcal{D}_{\Delta}^{\text{lt}})$ .

We define a morphism between two collections of radial data  $\dagger\mathfrak{R} \rightarrow \ddagger\mathfrak{R}$  [where we apply the evident notational conventions with respect to “ $\dagger$ ” and “ $\ddagger$ ”] to consist of data as follows:

- ( $a_{\text{Mor}_{\mathfrak{R}}}$ ) an isomorphism of  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters  $\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} \ddagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$ ;
- ( $b_{\text{Mor}_{\mathfrak{R}}}$ ) the isomorphism of  $\mathcal{F}^{\text{lt}}$ -prime-strips  $\mathfrak{F}_{\text{env}}^{\text{lt}}(\dagger\mathcal{D}_{>}) \xrightarrow{\sim} \mathfrak{F}_{\text{env}}^{\text{lt}}(\ddagger\mathcal{D}_{>})$  induced by the isomorphism of ( $a_{\text{Mor}_{\mathfrak{R}}}$ );
- ( $c_{\text{Mor}_{\mathfrak{R}}}$ ) the isomorphism between collections of data  $\dagger\mathfrak{R}^{\text{bad}} \xrightarrow{\sim} \ddagger\mathfrak{R}^{\text{bad}}$  induced by the isomorphism of ( $a_{\text{Mor}_{\mathfrak{R}}}$ );
- ( $d_{\text{Mor}_{\mathfrak{R}}}$ ) an isomorphism of  $\mathcal{F}^{\text{lt}\times\mu}$ -prime-strips  $\mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\dagger\mathcal{D}_{\Delta}^{\text{lt}}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\ddagger\mathcal{D}_{\Delta}^{\text{lt}})$ ;
- ( $e_{\text{Mor}_{\mathfrak{R}}}$ ) we observe that the isomorphisms of ( $b_{\text{Mor}_{\mathfrak{R}}}$ ) and ( $d_{\text{Mor}_{\mathfrak{R}}}$ ) are necessarily compatible with the poly-isomorphisms of ( $e_{\mathfrak{R}}$ ) for “ $\dagger$ ”, “ $\ddagger$ ”.

We define a collection of **coric data**

$$\dagger\mathfrak{C} = (\dagger\mathcal{D}^{\text{lt}}, \mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\dagger\mathcal{D}^{\text{lt}}))$$

to consist of

- ( $a_{\mathfrak{C}}$ ) a  $\mathcal{D}^{\text{lt}}$ -prime-strip  $\dagger\mathcal{D}^{\text{lt}}$ ;
- ( $b_{\mathfrak{C}}$ ) the  $\mathcal{F}^{\text{lt}\times\mu}$ -prime-strip  $\mathfrak{F}_{\Delta}^{\text{lt}\times\mu}(\dagger\mathcal{D}^{\text{lt}})$  associated to  $\dagger\mathcal{D}^{\text{lt}}$  [cf. Theorem 1.5, (iii)].

We define a morphism between two collections of coric data  $\dagger\mathfrak{C} \rightarrow \ddagger\mathfrak{C}$  [where we apply the evident notational conventions with respect to “ $\dagger$ ” and “ $\ddagger$ ”] to consist of data as follows:

- ( $a_{\text{Mor}_{\mathfrak{e}}}$ ) an isomorphism of  $\mathcal{D}^+$ -prime-strips  $\dagger\mathcal{D}^+ \xrightarrow{\sim} \ddagger\mathcal{D}^+$ ;
- ( $b_{\text{Mor}_{\mathfrak{e}}}$ ) an isomorphism of  $\mathcal{F}^{+\times\mu}$ -prime-strips  $\mathfrak{F}_{\Delta}^{+\times\mu}(\dagger\mathcal{D}^+) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{+\times\mu}(\ddagger\mathcal{D}^+)$  that induces the isomorphism  $\dagger\mathcal{D}^+ \xrightarrow{\sim} \ddagger\mathcal{D}^+$  on associated  $\mathcal{D}^+$ -prime-strips of ( $a_{\text{Mor}_{\mathfrak{e}}}$ ).

The **radial algorithm** is given by the assignment

$$\begin{aligned} \dagger\mathfrak{R} &= (\dagger\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}, \mathfrak{F}_{\text{env}}^{\dagger}(\dagger\mathcal{D}_{>}), \dagger\mathfrak{R}^{\text{bad}}, \mathfrak{F}_{\Delta}^{+\times\mu}(\dagger\mathcal{D}_{\Delta}^+), \mathfrak{F}_{\text{env}}^{+\times\mu}(\dagger\mathcal{D}_{>}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{+\times\mu}(\dagger\mathcal{D}_{\Delta}^+)) \\ &\mapsto \dagger\mathfrak{C} = (\dagger\mathcal{D}_{\Delta}^+, \mathfrak{F}_{\Delta}^{+\times\mu}(\dagger\mathcal{D}_{\Delta}^+)) \end{aligned}$$

— together with the assignment on morphisms determined by the data of ( $d_{\text{Mor}_{\mathfrak{R}}}$ ). Then:

(i) The functor associated to the radial algorithm defined above is **full and essentially surjective**. In particular, the radial environment defined above is **multiradial**.

(ii) Each  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  ${}^{n,m}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$ , for  $n, m \in \mathbb{Z}$ , defines, in an evident way, an associated collection of radial data  ${}^{n,m}\mathfrak{R}$ . The poly-isomorphisms induced by the **vertical** arrows of the **Gaussian log-theta-lattice** under consideration [cf. Theorem 1.5, (i)] induce poly-isomorphisms of radial data  $\dots \xrightarrow{\sim} {}^{n,m}\mathfrak{R} \xrightarrow{\sim} {}^{n,m+1}\mathfrak{R} \xrightarrow{\sim} \dots$ . Write

$${}^{n,\circ}\mathfrak{R}$$

for the collection of radial data obtained by identifying the various  ${}^{n,m}\mathfrak{R}$ , for  $m \in \mathbb{Z}$ , via these poly-isomorphisms and  ${}^{n,\circ}\mathfrak{C}$  for the collection of coric data associated, via the radial algorithm defined above, to the radial data  ${}^{n,\circ}\mathfrak{R}$ . In a similar vein, the **horizontal** arrows of the Gaussian log-theta-lattice under consideration induce full poly-isomorphisms  $\dots \xrightarrow{\sim} {}^{n,m}\mathcal{D}_{\Delta}^+ \xrightarrow{\sim} {}^{n+1,m}\mathcal{D}_{\Delta}^+ \xrightarrow{\sim} \dots$  of  $\mathcal{D}^+$ -prime-strips [cf. Theorem 1.5, (ii)]. Write

$${}^{\circ,\circ}\mathfrak{C}$$

for the collection of coric data obtained by identifying the various  ${}^{n,\circ}\mathfrak{C}$ , for  $n \in \mathbb{Z}$ , via these poly-isomorphisms. Thus, by applying the radial algorithm defined above to each  ${}^{n,\circ}\mathfrak{R}$ , for  $n \in \mathbb{Z}$ , we obtain a diagram — i.e., an **étale-picture of radial data** — as in Fig. 2.3 below. This diagram satisfies the important property of admitting **arbitrary permutation symmetries** among the spokes [i.e., the labels  $n \in \mathbb{Z}$ ] and is **compatible**, in the evident sense, with the étale-picture of  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters of [IUTchII], Corollary 4.11, (ii).

(iii) The [poly-]isomorphisms of  $\mathcal{F}^{+\times\mu}$ -prime-strips of/induced by ( $e_{\mathfrak{R}}$ ), ( $b_{\text{Mor}_{\mathfrak{R}}}$ ), ( $d_{\text{Mor}_{\mathfrak{R}}}$ ) [cf. also ( $e_{\text{Mor}_{\mathfrak{R}}}$ )] are **compatible**, relative to the **Kummer isomorphisms** of Proposition 2.1, (ii) [cf. also Proposition 2.1, (vi)], and Theorem 1.5, (iii), with the poly-isomorphisms — arising from the **horizontal arrows** of the Gaussian log-theta-lattice — of Theorem 1.5, (ii).



(iv) At  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , the isomorphism  $\dagger\mathfrak{R}^{\text{bad}} \xrightarrow{\sim} \ddagger\mathfrak{R}^{\text{bad}}$  of  $(c_{\text{Mor}_{\mathfrak{R}}})$  is **compatible** [cf. the final portion of Theorem 2.2, (ii)], relative to the **Kummer isomorphisms and poly-isomorphisms of projective systems of mono-theta environments** discussed in Proposition 2.1, (ii), (iii) [cf. also Proposition 2.1, (vi); the second display of Theorem 2.2, (ii)], and Theorem 1.5, (iii), with the poly-isomorphisms — arising from the **horizontal arrows of the Gaussian log-theta-lattice** — of Theorem 1.5, (ii).

*Proof.* The various assertions of Corollary 2.3 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circlearrowright$

$$\begin{array}{ccccc}
 & & \boxed{\begin{array}{c} \mathfrak{F}_{\text{env}}^{\text{ll-}}(n, \circ\mathcal{D}_{>}) \\ + n, \circ\mathfrak{R}^{\text{bad}} + \dots \end{array}} & & \\
 & \dots & | & & \dots \\
 & & \boxed{\begin{array}{c} \mathfrak{F}_{\Delta}^{\text{ll-} \times \mu}(\circ, \circ\mathcal{D}_{\Delta}^{\text{ll-}}) \end{array}} & & \\
 \boxed{\begin{array}{c} \mathfrak{F}_{\text{env}}^{\text{ll-}}(n', \circ\mathcal{D}_{>}) \\ + n', \circ\mathfrak{R}^{\text{bad}} + \dots \end{array}} & - & & - & \boxed{\begin{array}{c} \mathfrak{F}_{\text{env}}^{\text{ll-}}(n'', \circ\mathcal{D}_{>}) \\ + n'', \circ\mathfrak{R}^{\text{bad}} + \dots \end{array}} \\
 \dots & & | & & \dots \\
 & & \boxed{\begin{array}{c} \mathfrak{F}_{\text{env}}^{\text{ll-}}(n''', \circ\mathcal{D}_{>}) \\ + n''', \circ\mathfrak{R}^{\text{bad}} + \dots \end{array}} & & 
 \end{array}$$

Fig. 2.3: Étale-picture of radial data

**Remark 2.3.1.**

(i) In the context of the étale-picture of Fig. 2.3, it is of interest to recall the point of view of the discussion of [IUTchII], 1.12.5, (i), (ii), concerning the analogy between **étale-pictures** in the theory of the present series of papers and the **polar coordinate representation of the classical Gaussian integral**.

(ii) The étale-picture discussed in Corollary 2.3, (ii), may be thought of as a sort of **canonical splitting** of the portion of the **Gaussian log-theta-lattice** under consideration that involves **theta monoids** [cf. the discussion of [IUTchI], §I1, preceding Theorem A].

(iii) The portion of the **multiradiality** discussed in Corollary 2.3, (iv), at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  corresponds, in essence, to the multiradiality discussed in [IUTchII], Corollary 1.12, (iii); [IUTchII], Proposition 3.4, (i).

**Definition 2.4.**

(i) Let

$$\ddagger \mathcal{F}^+ = \{\ddagger \mathcal{F}_{\underline{v}}^+\}_{\underline{v} \in \mathbb{V}}$$

be an  $\mathcal{F}^+$ -*prime-strip*. Then recall from the discussion of [IUTchII], Definition 4.9, (ii), that at each  $\underline{w} \in \mathbb{V}^{\text{bad}}$ , the splittings of the split Frobenioid  $\ddagger \mathcal{F}_{\underline{w}}^+$  determine *submonoids* “ $\mathcal{O}^\perp(-) \subseteq \mathcal{O}^\triangleright(-)$ ”, as well as *quotient monoids* “ $\mathcal{O}^\perp(-) \twoheadrightarrow \mathcal{O}^\blacktriangleright(-)$ ” [i.e., by forming the quotient of “ $\mathcal{O}^\perp(-)$ ” by its torsion subgroup]. In a similar vein, for each  $\underline{w} \in \mathbb{V}^{\text{good}}$ , the splitting of the split Frobenioid determined by [indeed, “constituted by”, when  $\underline{w} \in \mathbb{V}^{\text{good}} \cap \mathbb{V}^{\text{non}}$  — cf. [IUTchI], Definition 5.2, (ii)]  $\ddagger \mathcal{F}_{\underline{w}}^+$  determines a *submonoid* “ $\mathcal{O}^\perp(-) \subseteq \mathcal{O}^\triangleright(-)$ ” whose subgroup of units is *trivial* [cf. [IUTchII], Definition 4.9, (iv), when  $\underline{w} \in \mathbb{V}^{\text{good}} \cap \mathbb{V}^{\text{non}}$ ]; in this case, we set  $\mathcal{O}^\blacktriangleright(-) \stackrel{\text{def}}{=} \mathcal{O}^\perp(-)$ . Write

$$\ddagger \mathcal{F}^{+\perp} = \{\ddagger \mathcal{F}_{\underline{v}}^{+\perp}\}_{\underline{v} \in \mathbb{V}}; \quad \ddagger \mathcal{F}^{+\blacktriangleright} = \{\ddagger \mathcal{F}_{\underline{v}}^{+\blacktriangleright}\}_{\underline{v} \in \mathbb{V}}$$

for the collections of data obtained by replacing the split Frobenioid portion of each  $\ddagger \mathcal{F}_{\underline{v}}^+$  by the *Frobenioids* determined, respectively, by the subquotient monoids “ $\mathcal{O}^\perp(-) \subseteq \mathcal{O}^\triangleright(-)$ ”, “ $\mathcal{O}^\blacktriangleright(-)$ ” just defined.

(ii) We define [in the spirit of [IUTchII], Definition 4.9, (vii)] an  $\mathcal{F}^{+\perp}$ -*prime-strip* to be a collection of data

$$* \mathcal{F}^{+\perp} = \{*\mathcal{F}_{\underline{v}}^{+\perp}\}_{\underline{v} \in \mathbb{V}}$$

that satisfies the following conditions: (a) if  $\underline{v} \in \mathbb{V}^{\text{non}}$ , then  $*\mathcal{F}_{\underline{v}}^{+\perp}$  is a *Frobenioid* that is isomorphic to  $\ddagger \mathcal{F}_{\underline{v}}^{+\perp}$  [cf. (i)]; (b) if  $\underline{v} \in \mathbb{V}^{\text{arc}}$ , then  $*\mathcal{F}_{\underline{v}}^{+\perp}$  consists of a Frobenioid and an object of  $\mathbb{T}\mathbb{M}^+$  [cf. [IUTchI], Definition 5.2, (ii)] such that  $*\mathcal{F}_{\underline{v}}^{+\perp}$  is isomorphic to  $\ddagger \mathcal{F}_{\underline{v}}^{+\perp}$ . In a similar vein, we define an  $\mathcal{F}^{+\blacktriangleright}$ -*prime-strip* to be a collection of data

$$* \mathcal{F}^{+\blacktriangleright} = \{*\mathcal{F}_{\underline{v}}^{+\blacktriangleright}\}_{\underline{v} \in \mathbb{V}}$$

that satisfies the following conditions: (a) if  $\underline{v} \in \mathbb{V}^{\text{non}}$ , then  $*\mathcal{F}_{\underline{v}}^{+\blacktriangleright}$  is a *Frobenioid* that is isomorphic to  $\ddagger \mathcal{F}_{\underline{v}}^{+\blacktriangleright}$  [cf. (i)]; (b) if  $\underline{v} \in \mathbb{V}^{\text{arc}}$ , then  $*\mathcal{F}_{\underline{v}}^{+\blacktriangleright}$  consists of a Frobenioid and an object of  $\mathbb{T}\mathbb{M}^+$  [cf. [IUTchI], Definition 5.2, (ii)] such that  $*\mathcal{F}_{\underline{v}}^{+\blacktriangleright}$  is isomorphic to  $\ddagger \mathcal{F}_{\underline{v}}^{+\blacktriangleright}$ . A *morphism of  $\mathcal{F}^{+\perp}$ - (respectively,  $\mathcal{F}^{+\blacktriangleright}$ -) prime-strips* is defined to be a collection of isomorphisms, indexed by  $\mathbb{V}$ , between the various constituent objects of the prime-strips [cf. [IUTchI], Definition 5.2, (iii)].

(iii) We define [in the spirit of [IUTchII], Definition 4.9, (viii)] an  $\mathcal{F}^{+\perp}$ -*prime-strip* to be a collection of data

$$* \mathcal{F}^{+\perp} = (*\mathcal{C}^{+\perp}, \text{Prime}(*\mathcal{C}^{+\perp}) \xrightarrow{\sim} \mathbb{V}, * \mathcal{F}^{+\perp}, \{*\rho_{\underline{v}}\}_{\underline{v} \in \mathbb{V}})$$

satisfying the conditions (a), (b), (c), (d), (e), (f) of [IUTchI], Definition 5.2, (iv), for an  $\mathcal{F}^{+\perp}$ -*prime-strip*, except that the portion of the collection of data constituted by an  $\mathcal{F}^+$ -*prime-strip* is replaced by an  $\mathcal{F}^{+\perp}$ -*prime-strip*. [We leave the routine

details to the reader.] In a similar vein, we define an  $\mathcal{F}^{\text{tr}}\blacktriangleright$ -*prime-strip* to be a collection of data

$$*\mathfrak{F}^{\text{tr}}\blacktriangleright = (*\mathcal{C}^{\text{tr}}, \text{Prime}(*\mathcal{C}^{\text{tr}}) \xrightarrow{\sim} \underline{\mathbb{V}}, *\mathfrak{F}^{\text{tr}}\blacktriangleright, \{*\rho_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}})$$

satisfying the conditions (a), (b), (c), (d), (e), (f) of [IUTchI], Definition 5.2, (iv), for an  $\mathcal{F}^{\text{tr}}$ -prime-strip, except that the portion of the collection of data constituted by an  $\mathcal{F}^{\text{tr}}$ -prime-strip is replaced by an  $\mathcal{F}^{\text{tr}}\blacktriangleright$ -prime-strip. [We leave the routine details to the reader.] A *morphism of  $\mathcal{F}^{\text{tr}\perp}$ - (respectively,  $\mathcal{F}^{\text{tr}}\blacktriangleright$ -) prime-strips* is defined to be an isomorphism between collections of data as discussed above.

**Remark 2.4.1.**

(i) Thus, by applying the constructions of Definition 2.4, (i), to the [underlying  $\mathcal{F}^{\text{tr}}$ -prime-strips associated to the]  $\mathcal{F}^{\text{tr}}$ -prime-strips “ $\mathfrak{F}_{\text{env}}^{\text{tr}}(\dagger\mathfrak{D}_{>})$ ” that appear in Corollary 2.3, one may regard the multiradiality of Corollary 2.3, (i), as implying a corresponding **multiradiality** assertion concerning the associated  $\mathcal{F}^{\text{tr}\perp}$ -prime-strips “ $\mathfrak{F}_{\text{env}}^{\text{tr}\perp}(\dagger\mathfrak{D}_{>})$ ”.

(ii) Suppose that we are in the situation discussed in (i). Then at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , the submonoids “ $\mathcal{O}^{\perp}(-) \subseteq \mathcal{O}^{\triangleright}(-)$ ” may be regarded, in a natural way [cf. Proposition 2.1, (ii); Theorem 2.2, (ii)], as *submonoids* of the monoids “ ${}_{\infty}\Psi_{\text{env}}^{\perp}(\dagger\mathfrak{D}_{>})_{\underline{v}}$ ” of Theorem 2.2, (ii), (a<sub>v</sub>). Moreover, the resulting inclusion of monoids is **compatible** with the **multiradiality** discussed in (i) and the multiradiality of the data “ $\dagger\mathfrak{R}^{\text{bad}}$ ” of Corollary 2.3, (c<sub>R</sub>), that is implied by the multiradiality of Corollary 2.3, (i).

### Section 3: Multiradial Logarithmic Gaussian Procession Monoids

In the present §3, we apply the theory developed thus far in the present series of papers to give [cf. Theorem 3.11 below] **multiradial algorithms** for a slightly modified version of the **Gaussian monoids** discussed in [IUTchII], §4. This modification revolves around the combinatorics of *processions*, as developed in [IUTchI], §4, §5, §6, and is necessary in order to establish the desired multiradiality. At a more concrete level, these combinatorics require one to apply the theory of **tensor packets** [cf. Propositions 3.1, 3.2, 3.3, 3.4, 3.7, 3.9, below]. Finally, we observe in Corollary 3.12 that these multiradial algorithms give rise to certain **estimates** concerning the **log-volumes** of the **logarithmic Gaussian procession monoids** that occur. This observation forms the starting point of the theory to be developed in [IUTchIV].

In the following discussion, we assume that we have been given *initial*  $\Theta$ -data as in [IUTchI], Definition 3.1. Also, we shall write

$$\mathbb{V}_{\mathbb{Q}} \stackrel{\text{def}}{=} \mathbb{V}(\mathbb{Q})$$

[cf. [IUTchI], §0] and apply the notation of Definition 1.1 of the present paper. We begin by discussing the theory of **tensor packets**, which may be thought of as a sort of *amalgamation* of the theory of *log-shells* developed in §1 with the theory of *processions* developed in [IUTchI], §4, §5, §6.

**Proposition 3.1. (Local Holomorphic Tensor Packets)** *Let*

$$\{\alpha \mathfrak{F}\}_{\alpha \in A} = \left\{ \{\alpha \mathcal{F}_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}} \right\}_{\alpha \in A}$$

be an  **$n$ -capsule**, with index set  $A$ , of  $\mathcal{F}$ -prime-strips [relative to the given initial  $\Theta$ -data — cf. [IUTchI], §0; [IUTchI], Definition 5.2, (i)]. Then [cf. the notation of Definition 1.1, (iii)] we shall refer to the correspondence

$$\mathbb{V}_{\mathbb{Q}} \ni v_{\mathbb{Q}} \mapsto \underline{\log}(\alpha \mathcal{F}_{v_{\mathbb{Q}}}) \stackrel{\text{def}}{=} \bigoplus_{\underline{v} \ni \underline{v} \mid v_{\mathbb{Q}}} \underline{\log}(\alpha \mathcal{F}_{\underline{v}})$$

as the **[1-]tensor packet** associated to the  $\mathcal{F}$ -prime-strip  $\alpha \mathfrak{F}$  and to the correspondence

$$\mathbb{V}_{\mathbb{Q}} \ni v_{\mathbb{Q}} \mapsto \underline{\log}({}^A \mathcal{F}_{v_{\mathbb{Q}}}) \stackrel{\text{def}}{=} \bigotimes_{\alpha \in A} \underline{\log}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$$

— where the tensor product is to be understood as a tensor product of topological modules — as the **[ $n$ -]tensor packet** associated to the collection of  $\mathcal{F}$ -prime-strips  $\{\alpha \mathfrak{F}\}_{\alpha \in A}$ . Then:

(i) **(Ring Structures)** *The topological field structures on the various  $\underline{\log}(\alpha \mathcal{F}_{\underline{v}})$  [cf. Definition 1.1, (i), (ii), (iii)], for  $\alpha \in A$ , determine a **topological ring structure** on  $\underline{\log}({}^A \mathcal{F}_{v_{\mathbb{Q}}})$  with respect to which  $\underline{\log}({}^A \mathcal{F}_{v_{\mathbb{Q}}})$  decomposes, uniquely, as a **direct sum of topological fields**, in a fashion that is compatible,*

for  $\alpha \in A$ , with the natural action of the topological group  ${}^\alpha\Pi_{\underline{v}}$  [where  $\underline{\mathbb{V}} \ni \underline{v} \mid v_{\mathbb{Q}}$ ] on the direct summand with subscript  $\underline{v}$  of the factor labeled  $\alpha$ .

(ii) **(Integral Structures)** Fix elements  $\alpha \in A$ ,  $\underline{v} \in \underline{\mathbb{V}}$ ,  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$  such that  $\underline{v} \mid v_{\mathbb{Q}}$ . Relative to the tensor product in the above definition of  $\underline{\log}({}^A\mathcal{F}_{v_{\mathbb{Q}}})$ , write

$$\underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}}) \stackrel{\text{def}}{=} \underline{\log}({}^\alpha\mathcal{F}_{\underline{v}}) \otimes \left\{ \bigotimes_{\beta \in A \setminus \{\alpha\}} \underline{\log}({}^\beta\mathcal{F}_{v_{\mathbb{Q}}}) \right\} \subseteq \underline{\log}({}^A\mathcal{F}_{v_{\mathbb{Q}}})$$

for the topological submodule determined by the tensor product of the factors labeled by  $\beta \in A \setminus \{\alpha\}$  with the tensor product of the direct summand with subscript  $\underline{v}$  of the factor labeled  $\alpha$ . Then  $\underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}})$  forms a direct summand of the topological ring  $\underline{\log}({}^A\mathcal{F}_{v_{\mathbb{Q}}})$ ; the topological ring structure on  $\underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}})$  decomposes, uniquely, as a **direct sum of topological fields**. Moreover, by forming the tensor product with “1’s” in the factors labeled by  $\beta \in A \setminus \{\alpha\}$ , one obtains a **natural injective homomorphism** of topological rings

$$\underline{\log}({}^\alpha\mathcal{F}_{\underline{v}}) \rightarrow \underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}})$$

that induces an **isomorphism** of the domain onto each of the direct summand topological fields of the codomain. In particular, the integral structure

$$\overline{\Psi}_{\underline{\log}({}^\alpha\mathcal{F}_{\underline{v}})} \stackrel{\text{def}}{=} \Psi_{\underline{\log}({}^\alpha\mathcal{F}_{\underline{v}})} \cup \{0\} \subseteq \underline{\log}({}^\alpha\mathcal{F}_{\underline{v}})$$

[cf. the notation of Definition 1.1, (i), (ii)] determines **integral structures** on each of the direct summand topological fields of  $\underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}})$ ,  $\underline{\log}({}^A\mathcal{F}_{v_{\mathbb{Q}}})$ .

*Proof.* The various assertions of Proposition 3.1 follow immediately from the definitions and the references quoted in the statements of these assertions [cf. also Remark 3.1.1, (i), below].  $\circ$

### Remark 3.1.1.

(i) Let  $\underline{v} \in \underline{\mathbb{V}}$ . In the notation of [IUTchI], Definition 3.1, write  $k \stackrel{\text{def}}{=} K_{\underline{v}}$ ; let  $\bar{k}$  be an algebraic closure of  $k$ . Then, roughly speaking, in the notation of Proposition 3.1,

$$\begin{aligned} \underline{\log}({}^\alpha\mathcal{F}_{\underline{v}}) &\xrightarrow{\sim} \bar{k}; & \overline{\Psi}_{\underline{\log}({}^\alpha\mathcal{F}_{\underline{v}})} &\xrightarrow{\sim} \mathcal{O}_{\bar{k}}; \\ \underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}}) &\xrightarrow{\sim} \bigotimes \bar{k} &\xrightarrow{\sim} \bigoplus \bar{k} &\supseteq \bigoplus \mathcal{O}_{\bar{k}} \end{aligned}$$

— i.e., one verifies immediately that each topological field  $\underline{\log}({}^\alpha\mathcal{F}_{\underline{v}})$  is isomorphic to  $\bar{k}$ ; each  $\underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}})$  is a tensor product [say, over  $\mathbb{Q}$ ] of copies of  $\bar{k}$ , hence decomposes as a direct sum of copies of  $\bar{k}$ ; each  $\overline{\Psi}_{\underline{\log}({}^\alpha\mathcal{F}_{\underline{v}})}$  is a copy of the set [i.e., a ring, when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ ] of integers  $\mathcal{O}_{\bar{k}} \subseteq \bar{k}$ . In particular, the “integral structures” discussed in the final portion of Proposition 3.1, (ii), correspond to copies of  $\mathcal{O}_{\bar{k}}$  contained in copies of  $\bar{k}$ .

(ii) Ultimately, for  $\underline{v} \in \underline{\mathbb{V}}$ , we shall be interested [cf. Proposition 3.9, (i), (ii), below] in considering **log-volumes** on the portion of  $\underline{\log}({}^\alpha\mathcal{F}_{\underline{v}})$  corresponding to

$K_{\underline{v}}$ . On the other hand, let us recall that we do not wish to consider *all* of the valuations in  $\mathbb{V}(K)$ . That is to say, we wish to restrict ourselves to considering the subset  $\underline{\mathbb{V}} \subseteq \mathbb{V}(K)$ , equipped with the *natural bijection*  $\underline{\mathbb{V}} \xrightarrow{\sim} \mathbb{V}_{\text{mod}}$  [cf. [IUTchI], Definition 3.1, (e)], which we wish to think of as a sort of “*local analytic section*” [cf. the discussion of [IUTchI], Remark 4.3.1, (i)] of the natural morphism  $\text{Spec}(K) \rightarrow \text{Spec}(F)$  [or, perhaps more precisely,  $\text{Spec}(K) \rightarrow \text{Spec}(F_{\text{mod}})$ ]. In particular, it will be necessary to consider these log-volumes on the portion of  $\underline{\log}(\alpha \mathcal{F}_{\underline{v}})$  corresponding to  $K_{\underline{v}}$  relative to the **weight**  $[K_{\underline{v}} : (F_{\text{mod}})_{\underline{v}}]^{-1}$ , where we write  $\underline{v} \in \mathbb{V}_{\text{mod}}$  for the element determined [via the natural bijection just discussed] by  $\underline{v}$  [cf. the discussion of [IUTchI], Example 3.5, (i), (ii), (iii), where similar factors appear]. When, moreover, we consider direct sums over all  $\underline{v} \in \underline{\mathbb{V}}$  lying over a given  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$  as in the case of  $\underline{\log}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$ , it will be convenient to use the **normalized weight**

$$\frac{1}{[K_{\underline{v}} : (F_{\text{mod}})_{\underline{v}}] \cdot \left( \sum_{\mathbb{V}_{\text{mod}} \ni w | v_{\mathbb{Q}}} [(F_{\text{mod}})_w : \mathbb{Q}_{v_{\mathbb{Q}}}] \right)}$$

— i.e., *normalized* so that multiplication by  $p_{v_{\mathbb{Q}}}$  affects log-volumes by addition or subtraction [that is to say, depending on whether  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$  or  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$ ] of the quantity  $\log(p_{v_{\mathbb{Q}}}) \in \mathbb{R}$ . In a similar vein, when we consider log-volumes on the portion of  $\underline{\log}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$  corresponding to the tensor product of various  $K_{\underline{v}_{\alpha}}$ , where  $\underline{\mathbb{V}} \ni \underline{v}_{\alpha} | v_{\mathbb{Q}}$ , it will be necessary to consider these log-volumes relative to the **weight**

$$\frac{1}{\prod_{\alpha \in A} [K_{\underline{v}_{\alpha}} : (F_{\text{mod}})_{\underline{v}_{\alpha}}]}$$

— where we write  $\underline{v}_{\alpha} \in \mathbb{V}_{\text{mod}}$  for the element determined by  $\underline{v}_{\alpha}$ . When, moreover, we consider direct sums over all possible choices for the data  $\{\underline{v}_{\alpha}\}_{\alpha \in A}$ , it will be convenient to use the **normalized weight**

$$\frac{1}{\left( \prod_{\alpha \in A} [K_{\underline{v}_{\alpha}} : (F_{\text{mod}})_{\underline{v}_{\alpha}}] \right) \cdot \left\{ \sum_{\{w_{\alpha}\}_{\alpha \in A}} \left( \prod_{\alpha \in A} [(F_{\text{mod}})_{w_{\alpha}} : \mathbb{Q}_{v_{\mathbb{Q}}}] \right) \right\}}$$

— where the sum is over all collections  $\{w_{\alpha}\}_{\alpha \in A}$  of [not necessarily distinct!] elements  $w_{\alpha} \in \mathbb{V}_{\text{mod}}$  lying over  $v_{\mathbb{Q}}$  and indexed by  $\alpha \in A$ . Again, these normalized weights are *normalized* so that multiplication by  $p_{v_{\mathbb{Q}}}$  affects log-volumes by addition or subtraction [that is to say, depending on whether  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$  or  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$ ] of the quantity  $\log(p_{v_{\mathbb{Q}}}) \in \mathbb{R}$ .

**Remark 3.1.2.** The constructions involving **local holomorphic tensor packets** given in Proposition 3.1 may be applied to the capsules that appear in the various  **$\mathcal{F}$ -prime-strip processions** obtained by considering the evident  $\mathcal{F}$ -prime-strip analogues [cf. [IUTchI], Remark 5.6.1; [IUTchI], Remark 6.12.1] of the **holomorphic processions** discussed in [IUTchI], Proposition 4.11, (i); [IUTchI], Proposition 6.9, (i).



**Proposition 3.2. (Local Mono-analytic Tensor Packets)** *Let*

$$\{\alpha\mathcal{D}^+\}_{\alpha\in A} = \left\{ \left\{ \alpha\mathcal{D}_{\underline{v}}^+ \right\}_{\underline{v}\in\underline{\mathbb{V}}} \right\}_{\alpha\in A}$$

be an  **$n$ -capsule**, with index set  $A$ , of  $\mathcal{D}^+$ -prime-strips [relative to the given initial  $\Theta$ -data — cf. [IUTchI], §0; [IUTchI], Definition 4.1, (iii)]. Then [cf. the notation of Proposition 1.2, (vi), (vii)] we shall refer to the correspondence

$$\mathbb{V}_{\mathbb{Q}} \ni v_{\mathbb{Q}} \mapsto \underline{\log}(\alpha\mathcal{D}_{v_{\mathbb{Q}}}^+) \stackrel{\text{def}}{=} \bigoplus_{\underline{v} \ni v \mid v_{\mathbb{Q}}} \underline{\log}(\alpha\mathcal{D}_{\underline{v}}^+)$$

as the **[1-]tensor packet** associated to the  $\mathcal{D}^+$ -prime-strip  $\alpha\mathcal{D}^+$  and to the correspondence

$$\mathbb{V}_{\mathbb{Q}} \ni v_{\mathbb{Q}} \mapsto \underline{\log}({}^A\mathcal{D}_{v_{\mathbb{Q}}}^+) \stackrel{\text{def}}{=} \bigotimes_{\alpha\in A} \underline{\log}(\alpha\mathcal{D}_{v_{\mathbb{Q}}}^+)$$

— where the tensor product is to be understood as a tensor product of topological modules — as the **[ $n$ -]tensor packet** associated to the collection of  $\mathcal{D}^+$ -prime-strips  $\{\alpha\mathcal{D}^+\}_{\alpha\in A}$ . For  $\alpha\in A$ ,  $\underline{v}\in\underline{\mathbb{V}}$ ,  $v_{\mathbb{Q}}\in\mathbb{V}_{\mathbb{Q}}$  such that  $\underline{v}\mid v_{\mathbb{Q}}$ , we shall write

$$\underline{\log}({}^{A,\alpha}\mathcal{D}_{\underline{v}}^+) \subseteq \underline{\log}({}^A\mathcal{D}_{v_{\mathbb{Q}}}^+)$$

for the topological submodule determined by the tensor product of the factors labeled by  $\beta\in A\setminus\{\alpha\}$  with the tensor product of the direct summand with subscript  $\underline{v}$  of the factor labeled  $\alpha$  [cf. Proposition 3.1, (ii)]. If the capsule of  $\mathcal{D}^+$ -prime-strips  $\{\alpha\mathcal{D}^+\}_{\alpha\in A}$  arises from a capsule of  $\mathcal{F}^{+\times\mu}$ -prime-strips

$$\{\alpha\mathfrak{F}^{+\times\mu}\}_{\alpha\in A} = \left\{ \left\{ \alpha\mathcal{F}_{\underline{v}}^{+\times\mu} \right\}_{\underline{v}\in\underline{\mathbb{V}}} \right\}_{\alpha\in A}$$

[relative to the given initial  $\Theta$ -data — cf. [IUTchI], §0; [IUTchII], Definition 4.9, (vii)], then we shall use similar notation to the notation just introduced concerning  $\{\alpha\mathcal{D}^+\}_{\alpha\in A}$  to denote objects associated to  $\{\alpha\mathfrak{F}^{+\times\mu}\}_{\alpha\in A}$ , i.e., by **replacing “ $\mathcal{D}^+$ ”** in the above notational conventions by “ $\mathcal{F}^{+\times\mu}$ ” [cf. also the notation of Proposition 1.2, (vi), (vii)]. Then:

(i) **(Mono-analytic/Holomorphic Compatibility)** Suppose that the capsule of  $\mathcal{D}^+$ -prime-strips  $\{\alpha\mathcal{D}^+\}_{\alpha\in A}$  arises from the capsule of  $\mathcal{F}$ -prime-strips  $\{\alpha\mathfrak{F}\}_{\alpha\in A}$  of Proposition 3.1; write  $\{\alpha\mathfrak{F}^{+\times\mu}\}_{\alpha\in A}$  for the capsule of  $\mathcal{F}^{+\times\mu}$ -prime-strips associated to  $\{\alpha\mathfrak{F}\}_{\alpha\in A}$ . Then the poly-isomorphisms “ $\underline{\log}(\dagger\mathcal{D}_{\underline{v}}^+) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_{\underline{v}}^{+\times\mu}) \xrightarrow{\sim} \underline{\log}(\dagger\mathcal{F}_{\underline{v}})$ ” of Proposition 1.2, (vi), (vii), induce **natural poly-isomorphisms of topological modules**

$$\underline{\log}(\alpha\mathcal{D}_{v_{\mathbb{Q}}}^+) \xrightarrow{\sim} \underline{\log}(\alpha\mathcal{F}_{v_{\mathbb{Q}}}^{+\times\mu}) \xrightarrow{\sim} \underline{\log}(\alpha\mathcal{F}_{v_{\mathbb{Q}}}); \quad \underline{\log}({}^A\mathcal{D}_{v_{\mathbb{Q}}}^+) \xrightarrow{\sim} \underline{\log}({}^A\mathcal{F}_{v_{\mathbb{Q}}}^{+\times\mu}) \xrightarrow{\sim} \underline{\log}({}^A\mathcal{F}_{v_{\mathbb{Q}}})$$

$$\underline{\log}({}^{A,\alpha}\mathcal{D}_{\underline{v}}^+) \xrightarrow{\sim} \underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}}^{+\times\mu}) \xrightarrow{\sim} \underline{\log}({}^{A,\alpha}\mathcal{F}_{\underline{v}})$$

between the various “**mono-analytic**” tensor packets of the present Proposition 3.2 and the “**holomorphic**” tensor packets of Proposition 3.1.

(ii) **(Integral Structures)** If  $\mathbb{V} \ni \underline{v} \mid v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$ , then the **mono-analytic log-shells** “ $\mathcal{I}_{\dagger \mathcal{D}_{\underline{v}}^{\dagger}}$ ” of Proposition 1.2, (vi), determine topological submodules

$$\mathcal{I}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}) \subseteq \underline{\log}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}); \quad \mathcal{I}({}^A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}) \subseteq \underline{\log}({}^A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}); \quad \mathcal{I}({}^{A,\alpha} \mathcal{D}_{\underline{v}}^{\dagger}) \subseteq \underline{\log}({}^{A,\alpha} \mathcal{D}_{\underline{v}}^{\dagger})$$

— which may be regarded as **integral structures** on the  $\mathbb{Q}$ -spans of these submodules. If  $\mathbb{V} \ni \underline{v} \mid v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$ , then by regarding the **mono-analytic log-shell** “ $\mathcal{I}_{\dagger \mathcal{D}_{\underline{v}}^{\dagger}}$ ” of Proposition 1.2, (vii), as the “closed unit ball” of a Hermitian metric on “ $\underline{\log}(\dagger \mathcal{D}_{\underline{v}}^{\dagger})$ ”, and considering the induced direct sum Hermitian metric on  $\underline{\log}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})$ , together with the induced tensor product Hermitian metric on  $\underline{\log}({}^A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})$ , one obtains Hermitian metrics on  $\underline{\log}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})$ ,  $\underline{\log}({}^A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})$ , and  $\underline{\log}({}^{A,\alpha} \mathcal{D}_{\underline{v}}^{\dagger})$ , whose associated **unit closed balls**

$$\mathcal{I}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}) \subseteq \underline{\log}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}); \quad \mathcal{I}({}^A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}) \subseteq \underline{\log}({}^A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger}); \quad \mathcal{I}({}^{A,\alpha} \mathcal{D}_{\underline{v}}^{\dagger}) \subseteq \underline{\log}({}^{A,\alpha} \mathcal{D}_{\underline{v}}^{\dagger})$$

may be regarded as **integral structures** on  $\underline{\log}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})$ ,  $\underline{\log}({}^A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})$ , and  $\underline{\log}({}^{A,\alpha} \mathcal{D}_{\underline{v}}^{\dagger})$ , respectively. For arbitrary  $\mathbb{V} \ni \underline{v} \mid v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ , we shall denote by “ $\mathcal{I}^{\mathbb{Q}}((-)$ )” the  $\mathbb{Q}$ -span of “ $\mathcal{I}((-)$ ””; also, we shall apply this notation involving “ $\mathcal{I}((-)$ ”, “ $\mathcal{I}^{\mathbb{Q}}((-)$ ” with “ $\mathcal{D}^{\dagger}$ ” replaced by “ $\mathcal{F}$ ” or “ $\mathcal{F}^{\dagger \times \mu}$ ” for the various objects obtained from the “ $\mathcal{D}^{\dagger}$ -versions” discussed above by applying the **natural poly-isomorphisms** of (i).

*Proof.* The various assertions of Proposition 3.2 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 3.2.1.** The issue of **estimating the discrepancy** between the **holomorphic integral structures** of Proposition 3.1, (ii), and the **mono-analytic integral structures** of Proposition 3.2, (ii), will form one of the main topics to be discussed in [IUTchIV] — cf. also Remark 3.9.1 below.

**Remark 3.2.2.** The constructions involving **local mono-analytic tensor packets** given in Proposition 3.2 may be applied to the capsules that appear in the various  **$\mathcal{D}^{\dagger}$ -prime-strip processions** — i.e., **mono-analytic processions** — discussed in [IUTchI], Proposition 4.11, (ii); [IUTchI], Proposition 6.9, (ii).

**Proposition 3.3. (Global Tensor Packets)** *Let*

$$\dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$$

be a  $\Theta^{\pm \text{ell}} \text{NF}$ -Hodge theater [relative to the given initial  $\Theta$ -data] — cf. [IUTchI], Definition 6.13, (i). Thus,  $\dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$  determines  $\Theta \text{NF}$ - and  $\Theta^{\pm \text{ell}}$ -Hodge theaters  $\dagger \mathcal{HT}^{\Theta \text{NF}}$ ,  $\dagger \mathcal{HT}^{\Theta^{\pm \text{ell}}}$  as in [IUTchII], Corollary 4.8. Let  $\{\alpha \mathfrak{F}\}_{\alpha \in A}$  be an  $n$ -capsule of  $\mathcal{F}$ -prime-strips as in Proposition 3.1. Suppose, further, that  $A$  is a **subset** of the index set  $J$  that appears in the  $\Theta \text{NF}$ -Hodge theater  $\dagger \mathcal{HT}^{\Theta \text{NF}}$ , and that, for each  $\alpha \in A$ , we are given a **log-link**

$$\alpha \mathfrak{F} \xrightarrow{\log} \dagger \mathfrak{F}_{\alpha}$$

— i.e., a poly-isomorphism of  $\mathcal{F}$ -prime-strips  $\log(\alpha\mathfrak{F}) \xrightarrow{\sim} \dagger\mathfrak{F}_\alpha$  [cf. Definition 1.1, (iii)]. Next, recall the field  $\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ$  discussed in [IUTchII], Corollary 4.8, (i); thus, [cf. [IUTchII], Corollary 4.8, (ii)], one also has, for  $j \in J$ , a labeled version  $(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_j$  of this field. We shall refer to

$$(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_A \stackrel{\text{def}}{=} \bigotimes_{\alpha \in A} (\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_\alpha$$

— where the tensor product is to be understood as a tensor product of modules — as the **global  $[n]$ -tensor packet** associated to the subset  $A \subseteq J$  and the  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ .

(i) **(Ring Structures)** The field structure on the various  $(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_\alpha$ , for  $\alpha \in A$ , determine a **ring structure** on  $(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_A$  with respect to which  $(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_A$  decomposes, uniquely, as a **direct sum of number fields**. Moreover, the various **localization functors** “ $(\dagger\mathcal{F}_{\text{mod}}^\circ)_j \rightarrow \dagger\mathfrak{F}_j$ ” considered in [IUTchII], Corollary 4.8, (iii), determine, by composing with the given **log-links**, a **natural injective localization ring homomorphism**

$$(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_A \rightarrow \underline{\log}({}^A\mathcal{F}_{\mathbb{V}_\mathbb{Q}}) \stackrel{\text{def}}{=} \prod_{v_\mathbb{Q} \in \mathbb{V}_\mathbb{Q}} \underline{\log}({}^A\mathcal{F}_{v_\mathbb{Q}})$$

to the product of the local holomorphic tensor packets considered in Proposition 3.1.

(ii) **(Integral Structures)** Fix an element  $\alpha \in A$ . Then by forming the tensor product with “1’s” in the factors labeled by  $\beta \in A \setminus \{\alpha\}$ , one obtains a **natural ring homomorphism**

$$(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_\alpha \rightarrow (\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_A$$

that induces an **isomorphism** of the domain onto a subfield of each of the direct summand number fields of the codomain. For each  $v_\mathbb{Q} \in \mathbb{V}_\mathbb{Q}$ , this homomorphism is **compatible**, in the evident sense, relative to the **localization** homomorphism of (i), with the natural homomorphism of topological rings considered in Proposition 3.1, (ii). Moreover, for each  $v_\mathbb{Q} \in \mathbb{V}_\mathbb{Q}^{\text{non}}$ , the composite of the above displayed homomorphism with the component at  $v_\mathbb{Q}$  of the localization homomorphism of (i) maps the ring of integers of the number field  $(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_\alpha$  into the submodule constituted by the **integral structure** on  $\underline{\log}({}^A\mathcal{F}_{v_\mathbb{Q}})$  considered in Proposition 3.1, (ii); for each  $v_\mathbb{Q} \in \mathbb{V}_\mathbb{Q}^{\text{arc}}$ , the composite of the above displayed homomorphism with the component at  $v_\mathbb{Q}$  of the localization homomorphism of (i) maps the set of archimedean integers [i.e., elements of absolute value  $\leq 1$  at all archimedean primes] of the number field  $(\dagger\overline{\mathbb{M}}_{\text{mod}}^\circ)_\alpha$  into the direct product of subsets constituted by the **integral structures** considered in Proposition 3.1, (ii), on the various direct summand topological fields of  $\underline{\log}({}^A\mathcal{F}_{v_\mathbb{Q}})$ .

*Proof.* The various assertions of Proposition 3.3 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 3.3.1.** One may perform analogous constructions to the constructions of Proposition 3.3 for the fields “ $\overline{\mathbb{M}}_{\text{mod}}^\circ(\dagger\mathcal{D}^\circ)_j$ ” of [IUTchII], Corollary 4.7, (ii) [cf.

also the localization functors of [IUTchII], Corollary 4.7, (iii)], constructed from the associated  $\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF-Hodge theater } \dagger\mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}}$ . These constructions are *compatible* with the corresponding constructions of Proposition 3.3, in the evident sense, relative to the various *labeled Kummer-theoretic isomorphisms* of [IUTchII], Corollary 4.8, (ii). We leave the routine details to the reader.

**Remark 3.3.2.**

(i) One may consider the **image** of the **localization** homomorphism of Proposition 3.3, (i), in the case of the various **local holomorphic tensor packets** arising from **processions**, as discussed in Remark 3.1.2. Indeed, at the level of the *labels* involved, this is immediate in the case of the “ $\mathbb{F}_l^*$ -processions” of [IUTchI], Proposition 4.11, (i). On the other hand, in the case of the “ $|\mathbb{F}_l|$ -processions” of [IUTchI], Proposition 6.9, (i), this may be achieved by applying the *identifying isomorphisms* between the *zero label*  $0 \in |\mathbb{F}_l|$  and the *diagonal label*  $\langle \mathbb{F}_l^* \rangle$  associated to  $\mathbb{F}_l^*$  discussed in [the final display of] [IUTchII], Corollary 4.6, (iii).

(ii) In a similar vein, one may *compose* the “ $\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF-Hodge theater version}$ ” discussed in Remark 3.3.1 of the **localization** homomorphism of Proposition 3.3, (i), with the product over  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$  of the inverses of the upper right-hand displayed isomorphisms at  $v_{\mathbb{Q}}$  of Proposition 3.2, (i), and then consider the **image** of this composite morphism in the case of the various **local mono-analytic tensor packets** arising from **processions**, as discussed in Remark 3.2.2. Just as in the holomorphic case discussed in (i), in the case of the case of the “ $|\mathbb{F}_l|$ -processions” of [IUTchI], Proposition 6.9, (ii), this obliges one to apply the *identifying isomorphisms* between the *zero label*  $0 \in |\mathbb{F}_l|$  and the *diagonal label*  $\langle \mathbb{F}_l^* \rangle$  associated to  $\mathbb{F}_l^*$  discussed in [the final display of] [IUTchII], Corollary 4.5, (iii).

(iii) The various *images of global tensor packets* discussed in (i) and (ii) above may be *identified* — i.e., in light of the *injectivity* of the homomorphisms applied to construct these images — with the *global tensor packets* themselves. These **local holomorphic/local mono-analytic global tensor packet images** will play a *central role* in the development of the theory of the present §3 [cf., e.g., Proposition 3.7, below].

**Remark 3.3.3.** The **log-shifted** nature of the **localization** homomorphism of Proposition 3.3, (i), will play a crucial role in the development of the theory of present §3 — cf. the discussion of [IUTchII], Remark 4.8.2, (i), (iii).

**Remark 3.3.4.** Relative to the notation introduced in Propositions 3.1, 3.2, 3.3, write

$$\underline{\log}_{\text{orb}}(\alpha(-)_{\underline{v}})$$

— i.e., where “ $(-)$ ” may be taken to be “ $\mathcal{F}$ ”, “ $\mathcal{D}^+$ ”, or “ $\mathcal{F}^+ \times \mu$ ” — for the set of collections of **Galois-orbits** [i.e., “ ${}^{\alpha}\Pi_{\underline{v}}$ ”- or “ ${}^{\dagger}G_{\underline{v}}$ ”-orbits when  $\underline{v} \in \mathbb{V}^{\text{non}}$  — cf. Propositions 1.2, (vi), and 3.1, (i);  $\{1\}$ -orbits, when  $\underline{v} \in \mathbb{V}^{\text{arc}}$ ] of  $\underline{\log}(\alpha(-)_{\underline{v}})$  [cf. the definition of “ $\mathbb{M}_{\text{orb}}^{\odot}(-)$ ”, “ $\overline{\mathbb{M}}_{\text{orb}}^{\odot}(-)$ ” in [IUTchII], Corollary 4.7, (i)];

$$\underline{\log}_{\text{orb}}(\alpha(-)_{v_{\mathbb{Q}}}) \stackrel{\text{def}}{=} \prod_{\mathbb{V} \ni \underline{v} \mid v_{\mathbb{Q}}} \underline{\log}_{\text{orb}}(\alpha(-)_{\underline{v}}); \quad \underline{\log}_{\text{orb}}(\alpha(-)_{\mathbb{V}_{\mathbb{Q}}}) \stackrel{\text{def}}{=} \prod_{v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \underline{\log}_{\text{orb}}(\alpha(-)_{v_{\mathbb{Q}}})$$

— where the “ $v_{\mathbb{Q}}$ ” in the first “ $\stackrel{\text{def.}}{=}$ ” is an element of  $\mathbb{V}_{\mathbb{Q}}$ . Then observe that the *natural localization homomorphism*

$$(\dagger \overline{\mathbb{M}}_{\text{mod}}^{\circ})_{\alpha} \rightarrow \underline{\log}(\alpha \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}}) \stackrel{\text{def.}}{=} \prod_{v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \underline{\log}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$$

[cf. Proposition 3.3, (i)] admits a *natural Galois-orbit-theoretic extension*

$$((\dagger \overline{\mathbb{M}}_{\text{orb}}^{\circ})_{\alpha} \subseteq) (\dagger \overline{\mathbb{M}}_{\text{orb}}^{\circ})_{\alpha} \rightarrow \underline{\log}_{\text{orb}}(\alpha \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}})$$

[cf. the notation of [IUTchII], Corollary 4.8, (ii)] — i.e., the map of sets that assigns to a collection of global Galois-orbits the resulting collection, at each  $\underline{v} \in \underline{\mathbb{V}}$ , of local Galois-orbits.

$$\begin{array}{ccccccc} \underline{q}^1 \curvearrowright & & \underline{q}^{j^2} \curvearrowright & & \underline{q}^{(l^*)^2} \curvearrowright & & \\ / \pm \hookrightarrow & / \pm / \pm \hookrightarrow & \dots \hookrightarrow & / \pm / \pm \dots / \pm \hookrightarrow & \dots \hookrightarrow & / \pm / \pm \dots \dots / \pm & \\ \mathbb{S}_1^{\pm} & \mathbb{S}_{1+1=2}^{\pm} & & \mathbb{S}_{j+1}^{\pm} & & \mathbb{S}_{1+l^*=l\pm}^{\pm} & \end{array}$$

Fig. 3.1: Splitting monoids of LGP-monoids acting on tensor packets

### Proposition 3.4. (Local Packet-theoretic Frobenioids)

(i) **(Single Packet Monoids)** *In the situation of Proposition 3.1, fix elements  $\alpha \in A$ ,  $\underline{v} \in \underline{\mathbb{V}}$ ,  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$  such that  $\underline{v} \mid v_{\mathbb{Q}}$ . Then by forming the **image** via the natural homomorphism  $\underline{\log}(\alpha \mathcal{F}_{\underline{v}}) \rightarrow \underline{\log}(A, \alpha \mathcal{F}_{\underline{v}})$  of Proposition 3.1, (ii), the monoid  $\Psi_{\underline{\log}(\alpha \mathcal{F}_{\underline{v}})}$  [cf. the notation of Definition 1.1, (i), (ii)], together with its submonoid of units  $\Psi_{\underline{\log}(\alpha \mathcal{F}_{\underline{v}})}^{\times}$  and realification  $\Psi_{\underline{\log}(\alpha \mathcal{F}_{\underline{v}})}^{\mathbb{R}}$ , determine monoids*

$$\Psi_{\underline{\log}(A, \alpha \mathcal{F}_{\underline{v}})}, \quad \Psi_{\underline{\log}(A, \alpha \mathcal{F}_{\underline{v}})}^{\times}, \quad \Psi_{\underline{\log}(A, \alpha \mathcal{F}_{\underline{v}})}^{\mathbb{R}}$$

— which are equipped with  $G_{\underline{v}}(\alpha \Pi_{\underline{v}})$ -**actions** when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  and, in the case of the first displayed monoid, with a pair consisting of an **Aut-holomorphic orbispace** and a **Kummer structure** when  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ . We shall think of these monoids as [possibly realified] **subquotients** of

$$\underline{\log}(A, \alpha \mathcal{F}_{\underline{v}})$$

that **act** [multiplicatively] on appropriate [possibly realified] subquotients of  $\underline{\log}(A, \alpha \mathcal{F}_{\underline{v}})$ . In particular, when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , the first displayed monoid, together with its  $\alpha \Pi_{\underline{v}}$ -action, determine a **Frobenioid** equipped with a natural isomorphism to  $\alpha \mathcal{F}_{\underline{v}}$ ; when  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ , the first displayed monoid, together with its Aut-holomorphic orbispace and Kummer structure, determine a collection of data equipped with a natural isomorphism to  $\alpha \mathcal{F}_{\underline{v}}$ .

(ii) **(Local Logarithmic Gaussian Procession Monoids)** *Let*

$$\dagger \mathcal{HT}^{\ominus \pm \text{ell} \text{NF}} \xrightarrow{\underline{\log}} \dagger \mathcal{HT}^{\ominus \pm \text{ell} \text{NF}}$$



be a **log-link** of  $\Theta^{\pm\text{ell}}\mathbf{NF}$ -Hodge theaters as in Proposition 1.3, (i) [cf. also the situation of Proposition 3.3]. Consider the  $\mathcal{F}$ -prime-strip processions that arise as the  $\mathcal{F}$ -prime-strip analogues [cf. Remark 3.1.2; [IUTchI], Remark 6.12.1] of the **holomorphic processions** discussed in [IUTchI], Proposition 6.9, (i), when the functor of [IUTchI], Proposition 6.9, (i), is applied to the  $\Theta^{\pm}$ -bridges associated to  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}}$ ,  $\ddagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}}$ ; we shall refer to such processions as “ $\dagger$ -” or “ $\ddagger$ -” processions. Here, we recall that for  $j \in \{1, \dots, l^*\}$ , the index set of the  $(j+1)$ -capsule that appear in such a procession is denoted  $\mathbb{S}_{j+1}^{\pm}$ . Then by applying the various constructions of “**single packet monoids**” given in (i) in the case of the various capsules of  $\mathcal{F}$ -prime-strips that appear in a holomorphic  $\ddagger$ -procession — i.e., more precisely, in the case of the **label**  $j \in \{1, \dots, l^*\}$  [which shall occasionally identify with its image in  $\mathbb{F}_l^* \subseteq |\mathbb{F}_l|$ ] that appears in the  $(j+1)$ -capsule of the  $\ddagger$ -procession — to the pull-backs, via the given **log-link**, of the [collections of] monoids  $\Psi_{\mathcal{F}_{\text{gau}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}})_{\underline{v}}$ ,  $\infty\Psi_{\mathcal{F}_{\text{gau}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}})_{\underline{v}}$  of [IUTchII], Corollary 4.6, (iv), for  $\underline{v} \in \underline{\mathbb{V}}$ , one obtains a **functorial algorithm** in the **log-link** of  $\Theta^{\pm\text{ell}}\mathbf{NF}$ -Hodge theaters  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}} \xrightarrow{\text{log}} \dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}}$  for constructing [collections of] monoids

$$\underline{v} \ni \underline{\mathbb{V}} \mapsto \Psi_{\mathcal{F}_{\text{LGP}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}})_{\underline{v}}; \quad \underline{v} \ni \underline{\mathbb{V}} \mapsto \infty\Psi_{\mathcal{F}_{\text{LGP}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}})_{\underline{v}}$$

equipped with **splittings** [up to torsion, when  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ ] — which we refer to as “**[local] LGP-monoids**”, or “**logarithmic Gaussian procession monoids**” [cf. Fig. 3.1 above]. Here, we note that the notation “ $(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}})$ ” constitutes a slight abuse of notation. Also, we note that this functorial algorithm requires one to apply the **compatibility** of the given **log-link** with the  $\mathbb{F}_l^{\times\pm}$ -**symmetrizing isomorphisms** involved [cf. Remark 1.3.2]. For  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , the component labeled  $j \in \{1, \dots, l^*\}$  of the submodule of **Galois invariants** [cf. (i)] of the **entire LGP-monoid**  $\Psi_{\mathcal{F}_{\text{LGP}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}})_{\underline{v}}$  is a subset of

$$\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm, j; \ddagger} \mathcal{F}_{\underline{v}})$$

[i.e., where the notation “ $\ddagger$ ” denotes the result of applying the discussion of (i) to the case of  $\mathcal{F}$ -prime-strips labeled “ $\ddagger$ ”; cf. also the notational conventions of Proposition 3.2, (ii)] that acts multiplicatively on  $\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm, j; \ddagger} \mathcal{F}_{\underline{v}})$  [cf. the constructions of [IUTchII], Corollary 3.6, (ii)]. For any  $\underline{v} \in \underline{\mathbb{V}}$ , the component labeled  $j \in \{1, \dots, l^*\}$  of the submodule of **Galois invariants** [cf. (i) when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ ; this Galois action is trivial when  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ ] of the **unit portion**  $\Psi_{\mathcal{F}_{\text{LGP}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\mathbf{NF}})_{\underline{v}}^{\times}$  of such an LGP-monoid is a subset of

$$\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm, j; \ddagger} \mathcal{F}_{\underline{v}})$$

[cf. the discussion of (i); the notational conventions of Proposition 3.2, (ii)] that acts multiplicatively on  $\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm, j; \ddagger} \mathcal{F}_{\underline{v}})$  [cf. the constructions of [IUTchII], Corollary 3.6, (ii); [IUTchII], Proposition 4.2, (iv); [IUTchII], Proposition 4.4, (iv)].

*Proof.* The various assertions of Proposition 3.4 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$



**Proposition 3.5. (Kummer Theory and Upper Semi-compatibility for Vertically Coric Local LGP-Monoids)** *Let  $\{^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}\}_{n,m \in \mathbb{Z}}$  be a collection of distinct  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters [relative to the given initial  $\Theta$ -data] — which we think of as arising from a Gaussian log-theta-lattice [cf. Definition 1.4]. For each  $n \in \mathbb{Z}$ , write*

$$^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$$

for the  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater determined, up to isomorphism, by the various  $^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , where  $m \in \mathbb{Z}$ , via the vertical coricity of Theorem 1.5, (i).

(i) **(Vertically Coric Local LGP-Monoids and Associated Kummer Theory)** *Write*

$$\mathfrak{F}^{(n,\circ)\mathcal{D}_{\succ}})_t$$

for the  $\mathcal{F}$ -prime-strip associated [cf. [IUTchII], Remark 4.5.1, (i)] to the labeled collection of monoids “ $\Psi_{\text{cns}}^{(n,\circ)\mathcal{D}_{\succ}})_t$ ” of [IUTchII], Corollary 4.5, (iii) [i.e., where we take “ $\natural$ ” to be “ $n,\circ$ ”]. Recall the constructions of Proposition 3.4, (ii), involving  $\mathcal{F}$ -prime-strip processions. Then by applying these constructions to the  $\mathcal{F}$ -prime-strips “ $\mathfrak{F}^{(n,\circ)\mathcal{D}_{\succ}})_t$ ” and the various full log-links associated [cf. the discussion of Proposition 1.2, (ix)] to these  $\mathcal{F}$ -prime-strips — which we consider in a fashion **compatible** with the  $\mathbb{F}_l^{\times\pm}$ -symmetries involved [cf. Remark 1.3.2; Proposition 3.4, (ii)] — we obtain a **functorial algorithm** in the  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  $^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  for constructing [collections of] monoids

$$\underline{v} \ni \underline{\mathbb{V}} \mapsto \Psi_{\text{LGP}}(^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}}; \quad \underline{v} \ni \underline{\mathbb{V}} \mapsto \infty\Psi_{\text{LGP}}(^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}}$$

equipped with **splittings** [up to torsion, when  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ ] — which we refer to as “**vertically coric [local] LGP-monoids**”. For each  $n, m \in \mathbb{Z}$ , this functorial algorithm is **compatible** [in the evident sense] with the functorial algorithm of Proposition 3.4, (ii) — i.e., where we take “ $\natural$ ” to be “ $n, m$ ” and “ $\natural$ ” to be “ $n, m-1$ ” — relative to the **Kummer isomorphisms** of labeled data

$$\Psi_{\text{cns}}(^{n,m'}\mathfrak{F}_{\succ})_t \xrightarrow{\sim} \Psi_{\text{cns}}(^{n,\circ}\mathfrak{F}_{\succ})_t$$

of [IUTchII], Corollary 4.6, (iii), and the evident identification, for  $m' = m, m-1$ , of  $^{n,m'}\mathfrak{F}_t$  [i.e., the  $\mathcal{F}$ -prime-strip that appears in the associated  $\Theta^{\pm}$ -bridge] with the  $\mathcal{F}$ -prime-strip associated to  $\Psi_{\text{cns}}(^{n,m'}\mathfrak{F}_{\succ})_t$ . In particular, for each  $n, m \in \mathbb{Z}$ , we obtain **Kummer isomorphisms** of [collections of] monoids

$$\begin{aligned} \Psi_{\mathcal{F}\text{LGP}}(^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}} &\xrightarrow{\sim} \Psi_{\text{LGP}}(^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}} \\ \infty\Psi_{\mathcal{F}\text{LGP}}(^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}} &\xrightarrow{\sim} \infty\Psi_{\text{LGP}}(^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}} \end{aligned}$$

for  $\underline{v} \in \underline{\mathbb{V}}$ .

(ii) **(Upper Semi-compatibility)** *The Kummer isomorphisms of the final two displays of (i) are “upper semi-compatible” — cf. the discussion of “upper semi-commutativity” in Remark 1.2.2, (iii) — with the various log-links of  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters  $^{n,m-1}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} ^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  [where  $m \in \mathbb{Z}$ ]*

of the Gaussian log-theta-lattice under consideration in the following sense. Let  $j \in \{0, 1, \dots, l^*\}$ . Then:

(a) **(Nonarchimedean Primes)** For  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$ , the topological module

$$\mathcal{I}(\mathbb{S}_{j+1}^{\pm} \mathcal{F}^{(n, \circ)} \mathcal{D}_{\succ})_{v_{\mathbb{Q}}}$$

— i.e., that arises from applying the constructions of Proposition 3.4, (ii), in the **vertically coric** context of (i) above [cf. also the notational conventions of Proposition 3.2, (ii)] — **contains** the image of the submodules of **Galois invariants** [where we recall the Galois actions that appear in the data of [IUTchII], Corollary 4.6, (i), (iii)] of the **groups of units**  $(\Psi_{\text{cns}}^{(n, m)} \mathfrak{F}_{\succ})_{|t|}^{\times}_{\underline{v}}$ , for  $\underline{V} \ni \underline{v} \mid v_{\mathbb{Q}}$  and  $|t| \in \{0, \dots, j\}$ , via **both**

- (1) the tensor product, over such  $|t|$ , of the [relevant] **Kummer isomorphisms** of (i), and
- (2) the tensor product, over such  $|t|$ , of the pre-composite of these Kummer isomorphisms with the  $m'$ -th **iterates** of the **log-links**, for  $m' \geq 1$ , of the  $n$ -th column of the Gaussian log-theta-lattice under consideration [cf. the discussion of Remark 1.2.2, (i), (iii)].

(b) **(Archimedean Primes)** For  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$ , the closed unit ball

$$\mathcal{I}(\mathbb{S}_{j+1}^{\pm} \mathcal{F}^{(n, \circ)} \mathcal{D}_{\succ})_{v_{\mathbb{Q}}}$$

— i.e., that arises from applying the constructions of Proposition 3.4, (ii), in the **vertically coric** context of (i) above [cf. also the notational conventions of Proposition 3.2, (ii)] — **contains** the image, via the tensor product, over  $|t| \in \{0, \dots, j\}$ , of the [relevant] **Kummer isomorphisms** of (i), of **both**

- (1) the **groups of units**  $(\Psi_{\text{cns}}^{(n, m)} \mathfrak{F}_{\succ})_{|t|}^{\times}_{\underline{v}}$ , for  $\underline{V} \ni \underline{v} \mid v_{\mathbb{Q}}$ , and
- (2) the **closed balls of radius  $\pi$**  inside  $(\Psi_{\text{cns}}^{(n, m)} \mathfrak{F}_{\succ})_{|t|}^{\text{gp}}_{\underline{v}}$  [cf. the notational conventions of Definition 1.1], for  $\underline{V} \ni \underline{v} \mid v_{\mathbb{Q}}$ .

Here, we recall from the discussion of Remark 1.2.2, (ii), (iii), that a closed ball as in (2) contains, for each  $m' \geq 1$ , a subset that **subjects**, via the  $m'$ -th **iterate** of the **log-link** of the  $n$ -th column of the Gaussian log-theta-lattice under consideration, onto the subset of the group of units  $(\Psi_{\text{cns}}^{(n, m-m')} \mathfrak{F}_{\succ})_{|t|}^{\times}_{\underline{v}}$  given by the image, via this iterate, of the entire domain on which this iterate is defined.

(c) **(Bad Primes)** Let  $\underline{v} \in \underline{V}^{\text{bad}}$ ; suppose that  $j \neq 0$ . Recall that the various monoids “ $\Psi_{\mathcal{F}_{\text{LGP}}}(-)_{\underline{v}}$ ”, “ ${}_{\infty} \Psi_{\mathcal{F}_{\text{LGP}}}(-)_{\underline{v}}$ ” constructed in Proposition 3.4, (ii), as well as the monoids “ $\Psi_{\text{LGP}}(-)_{\underline{v}}$ ”, “ ${}_{\infty} \Psi_{\text{LGP}}(-)_{\underline{v}}$ ” constructed in (i) above, are equipped with natural **splittings up to torsion**. Write

$$\Psi_{\mathcal{F}_{\text{LGP}}}^{\perp}(-)_{\underline{v}} \subseteq \Psi_{\mathcal{F}_{\text{LGP}}}(-)_{\underline{v}}; \quad {}_{\infty} \Psi_{\mathcal{F}_{\text{LGP}}}^{\perp}(-)_{\underline{v}} \subseteq {}_{\infty} \Psi_{\mathcal{F}_{\text{LGP}}}(-)_{\underline{v}}$$

$$\Psi_{\text{LGP}}^{\perp}(-)_{\underline{v}} \subseteq \Psi_{\text{LGP}}(-)_{\underline{v}}; \quad \infty\Psi_{\text{LGP}}^{\perp}(-)_{\underline{v}} \subseteq \infty\Psi_{\text{LGP}}(-)_{\underline{v}}$$

for the submonoids corresponding to these splittings [cf. the submonoids “ $\mathcal{O}^{\perp}(-) \subseteq \mathcal{O}^{\triangleright}(-)$ ” discussed in Definition 2.4, (i), in the case of “ $\Psi^{\perp}$ ”; the notational conventions of Theorem 2.2, (ii), in the case of “ $\infty\Psi^{\perp}$ ”]. [Thus, the subgroup of units of “ $\Psi^{\perp}$ ” consists of the  $2l$ -torsion subgroup of “ $\Psi$ ”, while the subgroup of units of “ $\infty\Psi^{\perp}$ ” contains of the entire torsion subgroup of “ $\infty\Psi$ ”.] Then, as  $m$  ranges over the elements of  $\mathbb{Z}$ , the actions, via the [relevant] **Kummer isomorphisms** of (i), of the various monoids  $\Psi_{\text{LGP}}^{\perp}(^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}} \subseteq \infty\Psi_{\text{LGP}}^{\perp}(^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}}$  on the topological modules

$$\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm,j}\mathcal{F}(^{n,\circ}\mathcal{D}_{\succ})_{\underline{v}}) \subseteq \underline{\log}(\mathbb{S}_{j+1}^{\pm,j}\mathcal{F}(^{n,\circ}\mathcal{D}_{\succ})_{\underline{v}})$$

[where  $j = 1, \dots, l^*$ ] — i.e., that arise from applying the constructions of Proposition 3.4, (ii), in the **vertically coric** context of (i) above [cf. also the notational conventions of Proposition 3.2, (ii)] — are **mutually compatible**, relative to the **log-links** of the  $n$ -th column of the Gaussian log-theta-lattice under consideration, in the sense that the only portions of these actions that are related to one another via these **log-links** are the **indeterminacies** with respect to **multiplication by roots of unity**, that is to say, indeterminacies at  $m$  that correspond, via the **log-link**, to “**addition by zero**” — i.e., to **no indeterminacy!** — at  $m + 1$ .

Now let us think of the various **groups of units**, for  $\underline{v} \in \mathbb{V}$ ,

$$(\Psi_{\text{cns}}(^{n,m}\mathfrak{F}_{\succ})_{|t|})_{\underline{v}}^{\times}, \quad \Psi_{\text{LGP}}(^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}}^{\times}$$

and **splitting monoids**, for  $\underline{v} \in \mathbb{V}^{\text{bad}}$ ,

$$\Psi_{\text{LGP}}^{\perp}(^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}}$$

as acting on various portions of the modules, for  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ ,

$$\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}\mathcal{F}(^{n,\circ}\mathcal{D}_{\succ})_{v_{\mathbb{Q}}})$$

not via a **single Kummer isomorphism** as in (i) — which fails to be **compatible** with the **log-links** of the Gaussian log-theta-lattice! — but rather via the **totality** of the various pre-composites of the relevant Kummer isomorphism with the various **iterates** of the **log-links** of the Gaussian log-theta-lattice — i.e., precisely as was described in detail in (a), (b), (c) above! Then one obtains a sort of “**log-Kummer correspondence**” between the **totality**, as  $m$  ranges over the elements of  $\mathbb{Z}$ , of the various groups of units and splitting monoids just discussed [i.e., which are labeled by “ $n, m$ ”] and their **actions** on the “ $\mathcal{I}^{\mathbb{Q}}$ ” labeled by “ $n, \circ$ ” which is **invariant** with respect to the **translation symmetries** [cf. Proposition 1.3, (iv)] of the  $n$ -th column of the Gaussian log-theta-lattice [cf. the discussion of Remark 1.2.2, (iii)].

*Proof.* The various assertions of Proposition 3.5 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Example 3.6. Concrete Representations of Global Frobenioids.** Before proceeding, we pause to take a closer look at the Frobenioid “ $\dagger \mathcal{F}_{\text{mod}}^\circ$ ” of [IUTchI], Example 5.1, (iii), i.e., more concretely speaking, the Frobenioid of *arithmetic line bundles on the stack* “ $S_{\text{mod}}$ ” of [IUTchI], Remark 3.1.5. Let us write

$$\mathcal{F}_{\text{mod}}^\circ$$

for the Frobenioid “ $\dagger \mathcal{F}_{\text{mod}}^\circ$ ” of [IUTchI], Example 5.1, (iii), in the case where the data denoted by the label “ $\dagger$ ” arises [in the evident sense] from data as discussed in [IUTchI], Definition 3.1. In the following discussion, we shall use the notation of [IUTchI], Definition 3.1.

(i) **(Rational Function Torsor Version)** For each  $\underline{v} \in \mathbb{V}$ , the valuation on  $K_{\underline{v}}$  determined by  $\underline{v}$  determines a *group homomorphism*  $\beta_{\underline{v}} : F_{\text{mod}}^\times \rightarrow K_{\underline{v}}^\times / \mathcal{O}_{K_{\underline{v}}}^\times$  [cf. Remark 3.6.1 below]. Then let us define a *category*  $\mathcal{F}_{\text{MOD}}^\circ$  as follows. An *object*  $\mathcal{T} = (T, \{t_{\underline{v}}\}_{\underline{v} \in \mathbb{V}})$  of  $\mathcal{F}_{\text{MOD}}^\circ$  consists of

- (a) an  $F_{\text{mod}}^\times$ -torsor  $T$ ;
- (b) a *trivialization*  $t_{\underline{v}}$  of the torsor  $T_{\underline{v}}$  obtained from  $T$  by executing the “change of structure group” operation determined by the homomorphism  $\beta_{\underline{v}}$ .

An *elementary morphism*  $\mathcal{T}_1 = (T_1, \{t_{1,\underline{v}}\}_{\underline{v} \in \mathbb{V}}) \rightarrow \mathcal{T}_2 = (T_2, \{t_{2,\underline{v}}\}_{\underline{v} \in \mathbb{V}})$  between objects of  $\mathcal{F}_{\text{MOD}}^\circ$  is defined to be an isomorphism  $T_1 \xrightarrow{\sim} T_2$  of  $F_{\text{mod}}^\times$ -torsors which is *integral* at each  $\underline{v} \in \mathbb{V}$ , i.e., maps the trivialization  $t_{1,\underline{v}}$  to an element of the  $\mathcal{O}_{K_{\underline{v}}}^\times$ -orbit of  $t_{2,\underline{v}}$ . There is an evident notion of *composition of elementary morphisms*, as well as an evident notion of *tensor powers*  $\mathcal{T}^{\otimes n}$ , for  $n \in \mathbb{Z}$ , of an object  $\mathcal{T}$  of  $\mathcal{F}_{\text{MOD}}^\circ$ . A *morphism*  $\mathcal{T}_1 = (T_1, \{t_{1,\underline{v}}\}_{\underline{v} \in \mathbb{V}}) \rightarrow \mathcal{T}_2 = (T_2, \{t_{2,\underline{v}}\}_{\underline{v} \in \mathbb{V}})$  between objects of  $\mathcal{F}_{\text{MOD}}^\circ$  is defined to be an elementary morphism  $(\mathcal{T}_1)^{\otimes n} \rightarrow \mathcal{T}_2$  for some positive integer  $n$ . There is an evident notion of *composition of morphisms*. Thus,  $\mathcal{F}_{\text{MOD}}^\circ$  forms a *category*. In fact, one verifies immediately that, from the point of view of the theory of Frobenioids developed in [FrdI], [FrdII],  $\mathcal{F}_{\text{MOD}}^\circ$  admits a natural *Frobenioid* structure [cf. [FrdI], Definition 1.3], for which the *base category* is the category with precisely one arrow. Relative to this Frobenioid structure, the elementary morphisms are precisely the *linear morphisms*, and the positive integer “ $n$ ” that appears in the definition of a morphism of  $\mathcal{F}_{\text{MOD}}^\circ$  is the *Frobenius degree* of the morphism. Moreover, by associating to an arithmetic line bundle on  $S_{\text{mod}}$  the  $F_{\text{mod}}^\times$ -torsor determined by restricting the line bundle to the generic point of  $S_{\text{mod}}$  and the local trivializations at  $\underline{v} \in \mathbb{V}$  determined by the various local integral structures, one verifies immediately that there exists a *natural isomorphism of Frobenioids*

$$\mathcal{F}_{\text{mod}}^\circ \xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^\circ$$

that induces the *identity morphism*  $F_{\text{mod}}^\times \rightarrow F_{\text{mod}}^\times$  on the associated *rational function monoids* [cf. [FrdI], Corollary 4.10].

(ii) **(Local Fractional Ideal Version)** Let us define a *category*  $\mathcal{F}_{\text{mod}}^\circ$  as follows. An *object*

$$\mathcal{J} = \{J_{\underline{v}}\}_{\underline{v} \in \mathbb{V}}$$

of  $\mathcal{F}_{\text{mod}}^{\circledast}$  consists of a collection of “fractional ideals”  $J_{\underline{v}} \subseteq K_{\underline{v}}$  for each  $\underline{v} \in \underline{\mathbb{V}}$  — i.e., a finitely generated nonzero  $\mathcal{O}_{K_{\underline{v}}}$ -submodule of  $K_{\underline{v}}$  when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ ; a positive real multiple of  $\mathcal{O}_{K_{\underline{v}}} \stackrel{\text{def}}{=} \{\lambda \in K_{\underline{v}} \mid |\lambda| \leq 1\} \subseteq K_{\underline{v}}$  when  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$  — such that  $J_{\underline{v}} = \mathcal{O}_{K_{\underline{v}}}$  for all but finitely many  $\underline{v}$ . If  $\mathcal{J} = \{J_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$  is an object of  $\mathcal{F}_{\text{mod}}^{\circledast}$ , then for any element  $f \in F_{\text{mod}}^{\times}$ , one obtains an object  $f \cdot \mathcal{J} = \{f \cdot J_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$  of  $\mathcal{F}_{\text{mod}}^{\circledast}$  by multiplying each of the fractional ideals  $J_{\underline{v}}$  by  $f$ . Moreover, if  $\mathcal{J} = \{J_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$  is an object of  $\mathcal{F}_{\text{mod}}^{\circledast}$ , then for any  $n \in \mathbb{Z}$ , there is an evident notion of the  $n$ -th tensor power  $\mathcal{J}^{\otimes n}$  of  $\mathcal{J}$ . An elementary morphism  $\mathcal{J}_1 = \{J_{1,\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}} \rightarrow \mathcal{J}_2 = \{J_{2,\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$  between objects of  $\mathcal{F}_{\text{mod}}^{\circledast}$  is defined to be an element  $f \in F_{\text{mod}}^{\times}$  that is integral with respect to  $\mathcal{J}_1$  and  $\mathcal{J}_2$  in the sense that  $f \cdot J_{1,\underline{v}} \subseteq J_{2,\underline{v}}$  for each  $\underline{v} \in \underline{\mathbb{V}}$ . There is an evident notion of composition of elementary morphisms. A morphism  $\mathcal{J}_1 = \{J_{1,\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}} \rightarrow \mathcal{J}_2 = \{J_{2,\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$  between objects of  $\mathcal{F}_{\text{mod}}^{\circledast}$  is defined to be an elementary morphism  $(\mathcal{J}_1)^{\otimes n} \rightarrow \mathcal{J}_2$  for some positive integer  $n$ . There is an evident notion of composition of morphisms. Thus,  $\mathcal{F}_{\text{mod}}^{\circledast}$  forms a category. In fact, one verifies immediately that, from the point of view of the theory of Frobenioids developed in [FrdI], [FrdII],  $\mathcal{F}_{\text{mod}}^{\circledast}$  admits a natural Frobenioid structure [cf. [FrdI], Definition 1.3], for which the base category is the category with precisely one arrow. Relative to this Frobenioid structure, the elementary morphisms are precisely the linear morphisms, and the positive integer “ $n$ ” that appears in the definition of a morphism of  $\mathcal{F}_{\text{mod}}^{\circledast}$  is the Frobenius degree of the morphism. Moreover, by associating to an object  $\mathcal{J} = \{J_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$  of  $\mathcal{F}_{\text{mod}}^{\circledast}$  the arithmetic line bundle on  $S_{\text{mod}}$  obtained from the trivial arithmetic line bundle on  $S_{\text{mod}}$  by modifying the integral structure of the trivial line bundle at  $\underline{v} \in \underline{\mathbb{V}}$  in the fashion prescribed by  $J_{\underline{v}}$ , one verifies immediately that there exists a natural isomorphism of Frobenioids

$$\mathcal{F}_{\text{mod}}^{\circledast} \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circledast}$$

that induces the identity morphism  $F_{\text{mod}}^{\times} \rightarrow F_{\text{mod}}^{\times}$  on the associated rational function monoids [cf. [FrdI], Corollary 4.10].

(iii) By composing the isomorphisms of Frobenioids of (i) and (ii), one thus obtains a natural isomorphism of Frobenioids

$$\mathcal{F}_{\text{mod}}^{\circledast} \xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circledast}$$

that induces the identity morphism  $F_{\text{mod}}^{\times} \rightarrow F_{\text{mod}}^{\times}$  on the associated rational function monoids [cf. [FrdI], Corollary 4.10]. One verifies immediately that although the above isomorphism of Frobenioids is not necessarily determined by the condition that it induce the identity morphism on  $F_{\text{mod}}^{\times}$ , the induced isomorphism between the respective *perfections* [hence also on *realifications*] of  $\mathcal{F}_{\text{mod}}^{\circledast}$ ,  $\mathcal{F}_{\text{MOD}}^{\circledast}$  is completely determined by this condition.

**Remark 3.6.1.** Note that, as far the various constructions of Example 3.6, (i), are concerned, the various homomorphisms  $\beta_{\underline{v}}$ , for  $\underline{v} \in \underline{\mathbb{V}}$ , may be thought of, alternatively, as a collection of

$$\text{subquotients of the perfection } (F_{\text{mod}}^{\times})^{\text{pf}} \text{ of } F_{\text{mod}}^{\times}$$

— each of which is equipped with a submonoid of “nonnegative elements” — that are completely determined by the ring structure of the field  $F_{\text{mod}}$  [i.e., equipped with its structure as the field of moduli of  $X_F$ ].



**Remark 3.6.2.**

(i) In the theory to be developed below, we shall be interested in relating the *realifications* of various Frobenioids isomorphic to  $\mathcal{F}_{\text{mod}}^{\circledast}$  that lie on opposite sides of the  $\Theta_{\text{gau}}^{\times\mu}$ -**link** to one another. In particular, at the level of objects of the Frobenioids involved, it only makes sense to work with **isomorphism classes** of objects that are preserved by the isomorphisms of Frobenioids that appear. Here, we note that the isomorphism classes of the sort of Frobenioids that appear in this context are determined, in an essential way, by the **rational function monoid** of the Frobenioid in question [cf. the constructions given in [FrdI], Theorem 5.2, (i)]. In this context, we observe that the rational function monoid  $F_{\text{mod}}^{\times}$  of  $\mathcal{F}_{\text{mod}}^{\circledast}$  satisfies the following *fundamental property*:

[the union with  $\{0\}$  of]  $F_{\text{mod}}^{\times}$  admits a natural **additive structure**.

In this context, we note that this property is *not* satisfied by

- (a) the rational function monoids of the *perfection* or *realification* of  $\mathcal{F}_{\text{mod}}^{\circledast}$
- (b) subgroups  $\Gamma \subseteq F_{\text{mod}}^{\times}$  — such as, for instance, the trivial subgroup  $\{1\}$  or the subgroup of *S-units*, for  $S \subseteq \mathbb{V}_{\text{mod}}$  a nonempty finite subset — that do not arise as the multiplicative group of some subfield of  $F_{\text{mod}}$ .

The significance of this fundamental property is that it allows one to represent the objects of  $\mathcal{F}_{\text{mod}}^{\circledast}$  **additively**, i.e., as *modules* — cf. the point of view of Example 3.6, (ii). At a more concrete level, if, in the notation of (b), one considers the result of “*adding*” two elements of a  $\Gamma$ -*torsor* [cf. the point of view of Example 3.6, (i)!], then the resulting “*sum*” can only be rendered meaningful, relative to the given  $\Gamma$ -torsor, if  $\Gamma$  is *additively closed*. The **additive representation** of objects of  $\mathcal{F}_{\text{mod}}^{\circledast}$  will be of *crucial importance* in the theory of the present series of papers since it will allow us to relate objects of  $\mathcal{F}_{\text{mod}}^{\circledast}$  on opposite sides of the  $\Theta_{\text{gau}}^{\times\mu}$ -link to one another — which, a priori, are only related to one another at the level of *realifications* in a **multiplicative** fashion — by means of **mono-analytic log-shells** [cf. the discussion of [IUTchII], Remark 4.7.2].

(ii) One way to understand the content of the discussion of (i) is as follows: whereas

the construction of  $\mathcal{F}_{\text{mod}}^{\circledast}$  depends on the **additive** structure of  $F_{\text{mod}}^{\times}$

in an essential way,

the construction of  $\mathcal{F}_{\text{MOD}}^{\circledast}$  is strictly **multiplicative** in nature.

Indeed, the construction of  $\mathcal{F}_{\text{MOD}}^{\circledast}$  given in Example 3.6, (i), is essentially the same as the construction of  $\mathcal{F}_{\text{mod}}^{\circledast}$  given in [FrdI], Example 6.3 [i.e., in effect, in [FrdI], Theorem 5.2, (i)]. From this point of view, it is natural to **identify**  $\mathcal{F}_{\text{MOD}}^{\circledast}$  with  $\mathcal{F}_{\text{mod}}^{\circledast}$  via the natural isomorphism of Frobenioids of Example 3.6, (i). We shall often do this in the theory to be developed below.



**Proposition 3.7. (Global Packet-theoretic Frobenioids)**

(i) **(Single Packet Rational Function Torsor Version)** *In the notation of Proposition 3.3: For each  $\alpha \in A$ , there is an **algorithm** for constructing, as discussed in Example 3.6, (i) [cf. also Remark 3.6.1], from the [number] field given by the **image***

$$(\dagger \overline{\mathbb{M}}_{\text{MOD}}^{\circ})_{\alpha}$$

of the composite

$$(\dagger \overline{\mathbb{M}}_{\text{mod}}^{\circ})_{\alpha} \rightarrow (\dagger \overline{\mathbb{M}}_{\text{mod}}^{\circ})_A \rightarrow \underline{\log}({}^A \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}})$$

of the homomorphisms of Proposition 3.3, (i), (ii), a **Frobenioid**  $(\dagger \mathcal{F}_{\text{MOD}}^{\circ})_{\alpha}$ , together with a **natural isomorphism of Frobenioids**

$$(\dagger \mathcal{F}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} (\dagger \mathcal{F}_{\text{MOD}}^{\circ})_{\alpha}$$

[cf. the notation of [IUTchII], Corollary 4.8, (ii)] that induces the **tautological isomorphism**  $(\dagger \mathbb{M}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} (\dagger \mathbb{M}_{\text{MOD}}^{\circ})_{\alpha}$  on the associated rational function monoids [cf. Example 3.6, (i)]. We shall often use this isomorphism of Frobenioids to **identify**  $(\dagger \mathcal{F}_{\text{mod}}^{\circ})_{\alpha}$  with  $(\dagger \mathcal{F}_{\text{MOD}}^{\circ})_{\alpha}$  [cf. Remark 3.6.2, (ii)]. Write  $(\dagger \mathcal{F}_{\text{MOD}}^{\circ \mathbb{R}})_{\alpha}$  for the **realification** of  $(\dagger \mathcal{F}_{\text{MOD}}^{\circ})_{\alpha}$ .

(ii) **(Single Packet Local Fractional Ideal Version)** *In the notation of Propositions 3.3, 3.4: For each  $\alpha \in A$ , there is an **algorithm** for constructing, as discussed in Example 3.6, (ii), from the [number] field  $(\dagger \overline{\mathbb{M}}_{\text{mod}}^{\circ})_{\alpha} \stackrel{\text{def}}{=} (\dagger \overline{\mathbb{M}}_{\text{MOD}}^{\circ})_{\alpha}$  [cf. (i)] and the **Galois invariants** of the local monoids*

$$\Psi_{\log({}^{A,\alpha} \mathcal{F}_{\underline{v}})} \subseteq \log({}^{A,\alpha} \mathcal{F}_{\underline{v}})$$

for  $\underline{v} \in \underline{\mathbb{V}}$  of Proposition 3.4, (i) — i.e., so the corresponding local “fractional ideal  $J_{\underline{v}}$ ” of Example 3.6, (ii), is a **subset** [indeed a submodule when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ ] of  $\mathcal{I}^{\mathbb{Q}}({}^{A,\alpha} \mathcal{F}_{\underline{v}})$  whose  $\mathbb{Q}$ -span is equal to  $\mathcal{I}^{\mathbb{Q}}({}^{A,\alpha} \mathcal{F}_{\underline{v}})$  [cf. the notational conventions of Proposition 3.2, (ii)] — a **Frobenioid**  $(\dagger \mathcal{F}_{\text{mod}}^{\circ})_{\alpha}$ , together with **natural isomorphisms of Frobenioids**

$$(\dagger \mathcal{F}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} (\dagger \mathcal{F}_{\text{mod}}^{\circ})_{\alpha}; \quad (\dagger \mathcal{F}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} (\dagger \mathcal{F}_{\text{MOD}}^{\circ})_{\alpha}$$

that induce the **tautological isomorphisms**  $(\dagger \mathbb{M}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} (\dagger \mathbb{M}_{\text{mod}}^{\circ})_{\alpha}$ ,  $(\dagger \mathbb{M}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} (\dagger \mathbb{M}_{\text{MOD}}^{\circ})_{\alpha}$  on the associated rational function monoids [cf. the natural isomorphism of Frobenioids of (i); Example 3.6, (ii), (iii)]. Write  $(\dagger \mathcal{F}_{\text{mod}}^{\circ \mathbb{R}})_{\alpha}$  for the **realification** of  $(\dagger \mathcal{F}_{\text{mod}}^{\circ})_{\alpha}$ .

(iii) **(Realified Global LGP-Frobenioids)** *In the notation of Proposition 3.4: By applying the composites of the isomorphisms of Frobenioids “ $\dagger \mathcal{C}_j^{\text{lf}} \xrightarrow{\sim} (\dagger \mathcal{F}_{\text{mod}}^{\circ \mathbb{R}})_j$ ” of [IUTchII], Corollary 4.8, (iii), with the realifications “ $(\dagger \mathcal{F}_{\text{mod}}^{\circ \mathbb{R}})_{\alpha} \xrightarrow{\sim} (\dagger \mathcal{F}_{\text{MOD}}^{\circ \mathbb{R}})_{\alpha}$ ” of the isomorphisms of Frobenioids of (i) above to the realified global Frobenioid portion  $\dagger \mathcal{C}_{\text{gau}}^{\text{lf}}$  of the  $\mathcal{F}^{\text{lf}}$ -prime-strip  $\dagger \mathfrak{F}_{\text{gau}}^{\text{lf}}$  of [IUTchII], Corollary 4.10, (ii) [cf. Remarks 1.5.3, (iii); 3.3.2, (i)], one obtains a **functorial algorithm***

in the **log-link of  $\Theta^{\pm\text{ell}}\text{NF-Hodge}$  theaters**  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} \dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  of Proposition 3.4, (ii), for constructing a Frobenioid

$$\mathcal{C}_{\text{LGP}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$$

— which we refer to as a “**global realified LGP-Frobenioid**”. Here, we note that the notation “ $(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$ ” constitutes a slight abuse of notation. In particular, the global realified Frobenioid  $\dagger\mathcal{C}_{\text{LGP}}^{\text{ll-}} \stackrel{\text{def}}{=} \mathcal{C}_{\text{LGP}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$ , together with the collection of data  $\Psi_{\mathcal{F}_{\text{LGP}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$  constructed in Proposition 3.4, (ii), give rise, in a natural fashion, to an  $\mathcal{F}^{\text{ll-}}$ -prime-strip

$$\dagger\mathfrak{F}_{\text{LGP}}^{\text{ll-}} = (\dagger\mathcal{C}_{\text{LGP}}^{\text{ll-}}, \text{Prime}(\dagger\mathcal{C}_{\text{LGP}}^{\text{ll-}}) \xrightarrow{\sim} \underline{\mathbb{V}}, \dagger\mathfrak{F}_{\text{LGP}}^{\text{ll-}}, \{\dagger\rho_{\text{LGP}, \underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}})$$

— cf. the construction of the  $\mathcal{F}^{\text{ll-}}$ -prime-strip  $\dagger\mathfrak{F}_{\text{gau}}^{\text{ll-}}$  in [IUTchII], Corollary 4.10, (ii) — together with a **natural isomorphism**

$$\dagger\mathfrak{F}_{\text{gau}}^{\text{ll-}} \xrightarrow{\sim} \dagger\mathfrak{F}_{\text{LGP}}^{\text{ll-}}$$

of  $\mathcal{F}^{\text{ll-}}$ -prime-strips [i.e., that arises **tautologically** from the construction of  $\dagger\mathfrak{F}_{\text{LGP}}^{\text{ll-}}$ !].

(iv) (**Realified Global lgp-Frobenioids**) In the situation of (iii) above, write  $\Psi_{\mathcal{F}_{\text{lgp}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}) \stackrel{\text{def}}{=} \Psi_{\mathcal{F}_{\text{LGP}}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$ ,  $\dagger\mathfrak{F}_{\text{lgp}}^{\text{ll-}} \stackrel{\text{def}}{=} \dagger\mathfrak{F}_{\text{LGP}}^{\text{ll-}}$ . Then by replacing, in the construction of (iii), the isomorphisms “ $(\dagger\mathcal{F}_{\text{mod}}^{\circledast\mathbb{R}})_{\alpha} \xrightarrow{\sim} (\dagger\mathcal{F}_{\text{MOD}}^{\circledast\mathbb{R}})_{\alpha}$ ” by the natural isomorphisms “ $(\dagger\mathcal{F}_{\text{mod}}^{\circledast\mathbb{R}})_{\alpha} \xrightarrow{\sim} (\dagger\mathcal{F}_{\text{mod}}^{\circledast\mathbb{R}})_{\alpha}$ ” [cf. (ii)], one obtains a **functorial algorithm** in the **log-link of  $\Theta^{\pm\text{ell}}\text{NF-Hodge}$  theaters**  $\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\text{log}} \dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  of Proposition 3.4, (ii), for constructing a Frobenioid

$$\mathcal{C}_{\text{lgp}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$$

— which we refer to as a “**global realified lgp-Frobenioid**” — as well as an  $\mathcal{F}^{\text{ll-}}$ -prime-strip

$$\dagger\mathfrak{F}_{\text{lgp}}^{\text{ll-}} = (\dagger\mathcal{C}_{\text{lgp}}^{\text{ll-}}, \text{Prime}(\dagger\mathcal{C}_{\text{lgp}}^{\text{ll-}}) \xrightarrow{\sim} \underline{\mathbb{V}}, \dagger\mathfrak{F}_{\text{lgp}}^{\text{ll-}}, \{\dagger\rho_{\text{lgp}, \underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}})$$

— where we write  $\dagger\mathcal{C}_{\text{lgp}}^{\text{ll-}} \stackrel{\text{def}}{=} \mathcal{C}_{\text{lgp}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$  — together with **tautological isomorphisms**

$$\dagger\mathfrak{F}_{\text{gau}}^{\text{ll-}} \xrightarrow{\sim} \dagger\mathfrak{F}_{\text{LGP}}^{\text{ll-}} \xrightarrow{\sim} \dagger\mathfrak{F}_{\text{lgp}}^{\text{ll-}}$$

of  $\mathcal{F}^{\text{ll-}}$ -prime-strips [cf. (iii)].

(v) (**Realified Product Embeddings and Non-realified Global Frobenioids**) The constructions of  $\mathcal{C}_{\text{LGP}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$ ,  $\mathcal{C}_{\text{lgp}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})$  given in (iii) and (iv) above give rise to a commutative diagram of categories

$$\begin{array}{ccc} \mathcal{C}_{\text{LGP}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}) & \hookrightarrow & \prod_{j \in \mathbb{F}_l^*} (\dagger\mathcal{F}_{\text{MOD}}^{\circledast\mathbb{R}})_j \\ \downarrow & & \downarrow \\ \mathcal{C}_{\text{lgp}}^{\text{ll-}}(\dagger\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}) & \hookrightarrow & \prod_{j \in \mathbb{F}_l^*} (\dagger\mathcal{F}_{\text{mod}}^{\circledast\mathbb{R}})_j \end{array}$$

— where the horizontal arrows are **embeddings** that arise tautologically from the constructions of (iii) and (iv) [cf. [IUTchII], Remark 4.8.1, (i)]; the vertical arrows are **isomorphisms**; the left-hand vertical arrow arises from the second isomorphism that appears in the final display of (iv); the right-hand vertical arrow is the product of the **realifications** of copies of the inverse of the second isomorphism that appears in the final display of (ii). In particular, by applying the definition of  $(\dagger \mathcal{F}_{\text{mod}}^{\odot})_j$  — i.e., in terms of **local fractional ideals** [cf. (ii)] — together with the products of **realification functors**

$$\prod_{j \in \mathbb{F}_l^*} (\dagger \mathcal{F}_{\text{mod}}^{\odot})_j \rightarrow \prod_{j \in \mathbb{F}_l^*} (\dagger \mathcal{F}_{\text{mod}}^{\odot \mathbb{R}})_j$$

[cf. [FrdI], Proposition 5.3], one obtains an **algorithm** for constructing, in a fashion compatible [in the evident sense] with the **local isomorphisms**  $\{\dagger \rho_{\text{Igp}, \underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$ ,  $\{\dagger \rho_{\text{LGP}, \underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$  of (iii) and (iv), objects of the [global!] categories  $\mathcal{C}_{\text{Igp}}^{\text{lf}}(\dagger \mathcal{HT}^{\ominus \pm \text{ell}} \text{NF})$ ,  $\mathcal{C}_{\text{LGP}}^{\text{lf}}(\dagger \mathcal{HT}^{\ominus \pm \text{ell}} \text{NF})$  from the local fractional ideals generated by elements of the **monoids** [cf. (iv); Proposition 3.4, (ii)]

$$\Psi_{\mathcal{F}_{\text{Igp}}}(\dagger \mathcal{HT}^{\ominus \pm \text{ell}} \text{NF})_{\underline{v}}$$

for  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ .

*Proof.* The various assertions of Proposition 3.7 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 3.7.1.** By analogy to the notation introduced in Proposition 3.7, (i), (ii), we shall write

$$(\dagger \mathbb{M}_{\text{ORB}}^{\odot})_{\alpha} = (\dagger \mathbb{M}_{\text{orb}}^{\odot})_{\alpha} \subseteq (\dagger \overline{\mathbb{M}}_{\text{ORB}}^{\odot})_{\alpha} = (\dagger \overline{\mathbb{M}}_{\text{orb}}^{\odot})_{\alpha}$$

for the respective *images* of the maps  $(\dagger \mathbb{M}_{\text{orb}}^{\odot})_{\alpha} \rightarrow \underline{\log}_{\text{orb}}(\alpha \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}})$ ,  $(\dagger \overline{\mathbb{M}}_{\text{orb}}^{\odot})_{\alpha} \rightarrow \underline{\log}_{\text{orb}}(\alpha \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}})$  defined in Remark 3.3.4.

**Definition 3.8.**

(i) In the situation of Proposition 3.7, (iv), (v), write  $\Psi_{\mathcal{F}_{\text{Igp}}}^{\perp}(-)_{\underline{v}} \stackrel{\text{def}}{=} \Psi_{\mathcal{F}_{\text{LGP}}}^{\perp}(-)_{\underline{v}}$ , for  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  [cf. the notation of Proposition 3.5, (ii), (c)]. Then we shall refer to the object of

$$\prod_{j \in \mathbb{F}_l^*} (\dagger \mathcal{F}_{\text{mod}}^{\odot})_j$$

— as well as its image in  $\dagger \mathcal{C}_{\text{LGP}}^{\text{lf}} = \mathcal{C}_{\text{LGP}}^{\text{lf}}(\dagger \mathcal{HT}^{\ominus \pm \text{ell}} \text{NF})$  or  $\dagger \mathcal{C}_{\text{Igp}}^{\text{lf}} = \mathcal{C}_{\text{Igp}}^{\text{lf}}(\dagger \mathcal{HT}^{\ominus \pm \text{ell}} \text{NF})$  [cf. Proposition 3.7, (iii), (iv), (v)] — determined by any collection, indexed by  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , of generators up to torsion of the monoids  $\Psi_{\mathcal{F}_{\text{Igp}}}^{\perp}(\dagger \mathcal{HT}^{\ominus \pm \text{ell}} \text{NF})_{\underline{v}}$  as a  **$\Theta$ -pilot object**. We shall refer to the object of the [global realified] Frobenioid

$$\dagger \mathcal{C}_{\Delta}^{\text{lf}}$$

of [IUTchII], Corollary 4.10, (i), determined by any collection, indexed by  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ , of generators up to torsion of the splitting monoid associated to the split Frobenioid  $(\dagger \mathfrak{F}_{\Delta}^{\pm})_{\underline{v}}$  [i.e., the data indexed by  $\underline{v}$  of the  $\mathcal{F}^{\pm}$ -prime-strip  $\dagger \mathfrak{F}_{\Delta}^{\pm}$  of [IUTchII], Corollary 4.10, (i)] — that is to say, at a more concrete level, determined by the “ $q$ ”, for  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  [cf. the notation of [IUTchI], Example 3.2, (iv)] — as a  **$q$ -pilot object**.

(ii) Let

$$\ddagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} \xrightarrow{\text{log}} \dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$$

be a **log-link** of  $\Theta^{\pm \text{ell}} \text{NF}$ -Hodge theaters and

$$* \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$$

a  $\Theta^{\pm \text{ell}} \text{NF}$ -Hodge theater [all relative to the given initial  $\Theta$ -data]. Recall the  $\mathcal{F}^{\pm}$ -prime-strip

$$* \mathfrak{F}_{\Delta}^{\pm}$$

constructed from  $* \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$  in [IUTchII], Corollary 4.10, (i). Following the notational conventions of [IUTchII], Corollary 4.10, (iii), let us write  $* \mathfrak{F}_{\Delta}^{\pm \blacktriangleright \times \mu}$  (respectively,  $\dagger \mathfrak{F}_{\text{LGP}}^{\pm \blacktriangleright \times \mu}$ ;  $\dagger \mathfrak{F}_{\text{lgp}}^{\pm \blacktriangleright \times \mu}$ ) for the  $\mathcal{F}^{\pm \blacktriangleright \times \mu}$ -prime-strip associated to the  $\mathcal{F}^{\pm}$ -prime-strip  $* \mathfrak{F}_{\Delta}^{\pm}$  (respectively,  $\dagger \mathfrak{F}_{\text{LGP}}^{\pm}$ ;  $\dagger \mathfrak{F}_{\text{lgp}}^{\pm}$ ) [cf. Proposition 3.7, (iii), (iv); [IUTchII], Definition 4.9, (viii); the functorial algorithm described in [IUTchII], Definition 4.9, (vi)]. Then — in the style of [IUTchII], Corollary 4.10, (iii) — we shall refer to the full poly-isomorphism of  $\mathcal{F}^{\pm \blacktriangleright \times \mu}$ -prime-strips  $\dagger \mathfrak{F}_{\text{LGP}}^{\pm \blacktriangleright \times \mu} \xrightarrow{\sim} * \mathfrak{F}_{\Delta}^{\pm \blacktriangleright \times \mu}$  as the  **$\Theta_{\text{LGP}}^{\times \mu}$ -link**

$$\dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} \xrightarrow{\Theta_{\text{LGP}}^{\times \mu}} * \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$$

from  $\dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$  to  $\ddagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$ , relative to the **log-link**  $\ddagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} \xrightarrow{\text{log}} \dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$ , and to the full poly-isomorphism of  $\mathcal{F}^{\pm \blacktriangleright \times \mu}$ -prime-strips  $\dagger \mathfrak{F}_{\text{lgp}}^{\pm \blacktriangleright \times \mu} \xrightarrow{\sim} * \mathfrak{F}_{\Delta}^{\pm \blacktriangleright \times \mu}$  as the  **$\Theta_{\text{lgp}}^{\times \mu}$ -link**

$$\dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} \xrightarrow{\Theta_{\text{lgp}}^{\times \mu}} * \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$$

from  $\dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$  to  $\ddagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$ , relative to the **log-link**  $\ddagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} \xrightarrow{\text{log}} \dagger \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}$ .

(iii) Let  $\{n, m \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}\}_{n, m \in \mathbb{Z}}$  be a *collection of distinct*  $\Theta^{\pm \text{ell}} \text{NF}$ -Hodge theaters [relative to the given initial  $\Theta$ -data] indexed by pairs of integers. Then we shall refer to the first (respectively, second) diagram

$$\begin{array}{ccccccc} & & \vdots & & \vdots & & \\ & & \uparrow \text{log} & & \uparrow \text{log} & & \\ \dots & \xrightarrow{\Theta_{\text{LGP}}^{\times \mu}} & n, m+1 \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} & \xrightarrow{\Theta_{\text{LGP}}^{\times \mu}} & n+1, m+1 \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} & \xrightarrow{\Theta_{\text{LGP}}^{\times \mu}} & \dots \\ & & \uparrow \text{log} & & \uparrow \text{log} & & \\ \dots & \xrightarrow{\Theta_{\text{LGP}}^{\times \mu}} & n, m \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} & \xrightarrow{\Theta_{\text{LGP}}^{\times \mu}} & n+1, m \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}} & \xrightarrow{\Theta_{\text{LGP}}^{\times \mu}} & \dots \\ & & \uparrow \text{log} & & \uparrow \text{log} & & \\ & & \vdots & & \vdots & & \end{array}$$

$$\begin{array}{ccccccc}
& & \vdots & & \vdots & & \\
& & \uparrow \text{log} & & \uparrow \text{log} & & \\
\dots & \xrightarrow{\Theta_{\text{lgp}}^{\times\mu}} & n, m+1 \mathcal{HT}^{\pm\text{ell}} \text{NF} & \xrightarrow{\Theta_{\text{lgp}}^{\times\mu}} & n+1, m+1 \mathcal{HT}^{\pm\text{ell}} \text{NF} & \xrightarrow{\Theta_{\text{lgp}}^{\times\mu}} & \dots \\
& & \uparrow \text{log} & & \uparrow \text{log} & & \\
\dots & \xrightarrow{\Theta_{\text{lgp}}^{\times\mu}} & n, m \mathcal{HT}^{\pm\text{ell}} \text{NF} & \xrightarrow{\Theta_{\text{lgp}}^{\times\mu}} & n+1, m \mathcal{HT}^{\pm\text{ell}} \text{NF} & \xrightarrow{\Theta_{\text{lgp}}^{\times\mu}} & \dots \\
& & \uparrow \text{log} & & \uparrow \text{log} & & \\
& & \vdots & & \vdots & & 
\end{array}$$

— where the *vertical* arrows are the *full log-links*, and the *horizontal* arrow of the first (respectively, second) diagram from  $n, m \mathcal{HT}^{\pm\text{ell}} \text{NF}$  to  $n+1, m \mathcal{HT}^{\pm\text{ell}} \text{NF}$  is the  $\Theta_{\text{LGP}}^{\times\mu}$ - (respectively,  $\Theta_{\text{lgp}}^{\times\mu}$ -) link from  $n, m \mathcal{HT}^{\pm\text{ell}} \text{NF}$  to  $n+1, m \mathcal{HT}^{\pm\text{ell}} \text{NF}$ , relative to the full *log*-link  $n, m-1 \mathcal{HT}^{\pm\text{ell}} \text{NF} \xrightarrow{\text{log}} n, m \mathcal{HT}^{\pm\text{ell}} \text{NF}$  [cf. (ii)] — as the [LGP-Gaussian] (respectively, [lgp-Gaussian]) **log-theta-lattice**. Thus, [cf. Definition 1.4] either of these diagrams may be represented symbolically by an *oriented graph*

$$\begin{array}{ccccccc}
& & \vdots & & \vdots & & \\
& & \uparrow & & \uparrow & & \\
\dots & \longrightarrow & \bullet & \longrightarrow & \bullet & \longrightarrow & \dots \\
& & \uparrow & & \uparrow & & \\
\dots & \longrightarrow & \bullet & \longrightarrow & \bullet & \longrightarrow & \dots \\
& & \uparrow & & \uparrow & & \\
& & \vdots & & \vdots & & 
\end{array}$$

— where the “•’s” correspond to the “ $n, m \mathcal{HT}^{\pm\text{ell}} \text{NF}$ ”.

**Remark 3.8.1.** The LGP-Gaussian and lgp-Gaussian log-theta-lattices are, of course, closely related, but, in the theory to be developed below, we shall mainly be interested in the **LGP-Gaussian log-theta-lattice** [for reasons to be explained in Remark 3.10.2, (ii), below]. On the other hand, our computation of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link will involve the  $\Theta_{\text{lgp}}^{\times\mu}$ -link, as well as related  **$\Theta$ -pilot objects**, in an essential way. Here, we note, for future reference, that both the  $\Theta_{\text{LGP}}^{\times\mu}$ - and the  $\Theta_{\text{lgp}}^{\times\mu}$ -link map  **$\Theta$ -pilot objects** to  **$q$ -pilot objects**.

**Remark 3.8.2.** One verifies immediately that the *main results* obtained so far concerning Gaussian log-theta-lattices — namely, Theorem 1.5, Proposition 2.1, Corollary 2.3, and Proposition 3.5 — generalize immediately [indeed, “formally”]

to the case of LGP- or **lgp-Gaussian log-theta-lattices**. Indeed, the substantive content of these results concerns portions of the log-theta-lattices involved that are *substantively unaffected* by the transition from “Gaussian” to “LGP- or **lgp-Gaussian**”.

**Remark 3.8.3.** In the definition of the various **horizontal arrows** of the **log-theta-lattices** discussed in Definition 3.8, (iii), it may appear to the reader, at first glance, that, instead of working with  $\mathcal{F}^{\text{tr}} \times^{\mu}$ -prime-strips, it might in fact be sufficient to replace the unit [i.e.,  $\mathcal{F}^{\text{tr}} \times^{\mu}$ -prime-strip] portions of these prime-strips by the associated **log-shells** [cf. Proposition 1.2, (vi), (vii)], on which, at nonarchimedean  $\underline{v} \in \underline{\mathbb{V}}$ , the associated local Galois groups act *trivially*. In fact, however, this is *not* the case. That is to say, the *nontrivial Galois action* on the local unit portions of the  $\mathcal{F}^{\text{tr}} \times^{\mu}$ -prime-strips involved is necessary in order to consider the **Kummer theory** [cf. Proposition 3.5, (i), (ii), as well as Proposition 3.10, (i), (iii); Remark 3.10.1, (ii), below] of the various local and global objects for which the log-shells serve as “**multiradial containers**” [cf. the discussion of Remark 1.5.2]. Here, we recall that this Kummer theory plays a crucial role in the theory of the present series of papers in relating corresponding *Frobenius-like* and *étale-like* objects [cf. the discussion of Remark 1.5.4, (i)].

**Proposition 3.9. (Log-volume for Packets and Processions)**

(i) **(Local Holomorphic Packets)** *In the situation of Proposition 3.2, (i), (ii): Suppose that  $\underline{\mathbb{V}} \ni \underline{v} \mid v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$ ,  $\alpha \in A$ . Then the  $p_{v_{\mathbb{Q}}}$ -adic log-volume on each of the direct summand  $p_{v_{\mathbb{Q}}}$ -adic fields of  $\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$ ,  $\mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})$ , and  $\mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}})$  — cf. the direct sum decompositions of Proposition 3.1, (i), together with the discussion of **normalized weights** in Remark 3.1.1, (ii) — determines [cf. [AbsTopIII], Proposition 5.7, (i)] **log-volumes***

$$\mu_{\alpha, v_{\mathbb{Q}}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})) \rightarrow \mathbb{R}; \quad \mu_{A, v_{\mathbb{Q}}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})) \rightarrow \mathbb{R}$$

$$\mu_{A, \alpha, \underline{v}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}})) \rightarrow \mathbb{R}$$

— where we write “ $\mathfrak{M}(-)$ ” for the set of **compact open subsets** of “ $(-)$ ” — such that the log-volume of each of the “**local holomorphic**” **integral structures** of Proposition 3.1, (ii) — i.e., the elements

$$\mathcal{O}_{\alpha \mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}}); \quad \mathcal{O}_{A \mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}}); \quad \mathcal{O}_{A, \alpha \mathcal{F}_{\underline{v}}} \subseteq \mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}})$$

of “ $\mathfrak{M}(-)$ ” given by the integral structures discussed in Proposition 3.1, (ii), on each of the direct summand  $p_{v_{\mathbb{Q}}}$ -adic fields — is equal to **zero**. Here, we assume that these log-volumes are normalized so that multiplication of an element of “ $\mathfrak{M}(-)$ ” by  $p_{\underline{v}}$  corresponds to adding the quantity  $-\log(p_{\underline{v}}) \in \mathbb{R}$ ; we shall refer to this normalization as the **packet-normalization**. Suppose that  $\underline{\mathbb{V}} \ni \underline{v} \mid v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$ ,  $\alpha \in A$ . Then the sum of the radial log-volumes on each of the direct summand complex archimedean fields of  $\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$ ,  $\mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})$ , and  $\mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}})$  — cf. the direct sum decompositions of Proposition 3.1, (i), together with the discussion of **normalized**



**weights** in Remark 3.1.1, (ii) — determines [cf. [AbsTopIII], Proposition 5.7, (ii)] **log-volumes**

$$\mu_{\alpha, v_{\mathbb{Q}}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})) \rightarrow \mathbb{R}; \quad \mu_{A, v_{\mathbb{Q}}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})) \rightarrow \mathbb{R}$$

$$\mu_{A, \alpha, \underline{v}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}})) \rightarrow \mathbb{R}$$

— where we write “ $\mathfrak{M}(-)$ ” for the set of **compact closures of open subsets** of “ $(-)$ ” — such that the log-volume of each of the “**local holomorphic**” **integral structures** of Proposition 3.1, (ii) — i.e., the elements

$$\mathcal{O}_{\alpha \mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}}); \quad \mathcal{O}_{A \mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}}); \quad \mathcal{O}_{A, \alpha \mathcal{F}_{\underline{v}}} \subseteq \mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}})$$

of “ $\mathfrak{M}(-)$ ” given by the products of the integral structures discussed in Proposition 3.1, (ii), on each of the direct summand complex archimedean fields — is equal to **zero**. Here, we assume that these log-volumes are normalized so that multiplication of an element of “ $\mathfrak{M}(-)$ ” by  $e = 2.71828\dots$  corresponds to adding the quantity  $1 = \log(e) \in \mathbb{R}$ ; we shall refer to this normalization as the **packet-normalization**. In both the nonarchimedean and archimedean cases, “ $\mu_{A, v_{\mathbb{Q}}}^{\log}$ ” is **invariant** with respect to **permutations** of  $A$ . Finally, when working with collections of capsules in a procession, as in Proposition 3.4, (ii), we obtain, in both the nonarchimedean and archimedean cases, log-volumes on the products of the “ $\mathfrak{M}(-)$ ” associated to the various capsules under consideration, which we normalize by taking the **average**, over the various capsules under consideration; we shall refer to this normalization as the **procession-normalization** [cf. Remark 3.9.3 below].

(ii) (**Mono-analytic Compatibility**) In the situation of Proposition 3.2, (i), (ii): Suppose that  $\mathbb{V} \ni \underline{v} \mid v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ . Then by applying the  $p_{v_{\mathbb{Q}}}$ -adic log-volume, when  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$ , or the radial log-volume, when  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$ , on the **mono-analytic log-shells** “ $\mathcal{I}_{\dagger \mathcal{D}_{\underline{v}}^{\dagger}}$ ” of Proposition 1.2, (vi), (vii), (viii), and adjusting appropriately [cf. Remark 3.9.1 below for more details] to account for the **discrepancy** between the “**local holomorphic**” **integral structures** of Proposition 3.1, (ii), and the “**mono-analytic**” **integral structures** of Proposition 3.2, (ii), one obtains [by a slight abuse of notation] **log-volumes**

$$\mu_{\alpha, v_{\mathbb{Q}}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})) \rightarrow \mathbb{R}; \quad \mu_{A, v_{\mathbb{Q}}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(A \mathcal{D}_{v_{\mathbb{Q}}}^{\dagger})) \rightarrow \mathbb{R}$$

$$\mu_{A, \alpha, \underline{v}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{D}_{\underline{v}}^{\dagger})) \rightarrow \mathbb{R}$$

— where “ $\mathfrak{M}(-)$ ” is as in (i) above — which are **compatible** with the log-volumes obtained in (i), relative to the **natural poly-isomorphisms** of Proposition 3.2, (i). In particular, these log-volumes may be constructed via a **functorial algorithm** from the  $\mathcal{D}^{\dagger}$ -prime-strips under consideration. If one considers the **mono-analyticization** [cf. [IUTchI], Proposition 6.9, (ii)] of a holomorphic procession as in Proposition 3.4, (ii), then taking the average, as in (i) above, of the **packet-normalized log-volumes** of the above display gives rise to **procession-normalized log-volumes**, which are compatible, relative to the natural poly-isomorphisms of Proposition 3.2, (i), with the procession-normalized log-volumes of (i). Finally, by **replacing** “ $\mathcal{D}^{\dagger}$ ” by “ $\mathcal{F}^{\dagger \times \mu}$ ” [cf. also the discussion of Proposition 1.2, (vi),

(vii), (viii)], one obtains a similar theory of log-volumes for the various objects associated to the mono-analytic log-shells “ $\mathcal{I}_{\dagger \mathcal{F}_{\underline{v}}^+ \times \mu}$ ”, which is **compatible** with the theory obtained for “ $\mathcal{D}^+$ ” relative to the various **natural poly-isomorphisms** of Proposition 3.2, (i).

(iii) (**Global Compatibility**) In the situation of Proposition 3.7, (i), (ii): Write

$$\mathcal{I}^{\mathbb{Q}}({}^A \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}}) \stackrel{\text{def}}{=} \prod_{v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \mathcal{I}^{\mathbb{Q}}({}^A \mathcal{F}_{v_{\mathbb{Q}}}) \subseteq \underline{\log}({}^A \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}}) = \prod_{v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \underline{\log}({}^A \mathcal{F}_{v_{\mathbb{Q}}})$$

and

$$\mathfrak{M}(\mathcal{I}^{\mathbb{Q}}({}^A \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}})) \subseteq \prod_{v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}({}^A \mathcal{F}_{v_{\mathbb{Q}}}))$$

for the subset of elements whose components, indexed by  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ , have **zero log-volume** [cf. (i)] for all but finitely many  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ . Then, by adding the log-volumes of (i) [all but finitely many of which are zero!] at the various  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}$ , one obtains a **global log-volume**

$$\mu_{A, \mathbb{V}_{\mathbb{Q}}}^{\log} : \mathfrak{M}(\mathcal{I}^{\mathbb{Q}}({}^A \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}})) \rightarrow \mathbb{R}$$

which is **invariant** with respect to **multiplication** by elements of

$$({}^{\dagger} \overline{\mathbb{M}}_{\text{mod}}^{\odot})_{\alpha} = ({}^{\dagger} \overline{\mathbb{M}}_{\text{MOD}}^{\odot})_{\alpha} \subseteq \mathcal{I}^{\mathbb{Q}}({}^A \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}})$$

as well as with respect to **permutations** of  $A$ , and, moreover, satisfies the following property concerning [the elements of “ $\mathfrak{M}(-)$ ” determined by] objects “ $\mathcal{J} = \{J_{\underline{v}}\}_{\underline{v} \in \underline{\mathbb{V}}}$ ” of  $({}^{\dagger} \mathcal{F}_{\text{mod}}^{\odot})_{\alpha}$  [cf. Example 3.6, (ii); Proposition 3.7, (ii)]: the **global log-volume**  $\mu_{A, \mathbb{V}_{\mathbb{Q}}}^{\log}(\mathcal{J})$  is equal to the **degree** of the **arithmetic line bundle** determined by  $\mathcal{J}$  [cf. the discussion of Example 3.6, (ii); the natural isomorphism  $({}^{\dagger} \mathcal{F}_{\text{mod}}^{\odot})_{\alpha} \xrightarrow{\sim} ({}^{\dagger} \mathcal{F}_{\text{mod}}^{\odot})_{\alpha}$  of Proposition 3.7, (ii)], relative to a **suitable normalization**.

(iv) (**log-link Compatibility**) Let  $\{n, m \mathcal{H}\mathcal{T}^{\Theta^{\pm \text{ell}} \text{NF}}\}_{n, m \in \mathbb{Z}}$  be a **collection of distinct  $\Theta^{\pm \text{ell}} \text{NF}$ -Hodge theaters** [relative to the given initial  $\Theta$ -data] — which we think of as arising from an **LGP-Gaussian log-theta-lattice** [cf. Definition 3.8, (iii)]. Then [cf. also the discussion of Remark 3.9.4 below]:

(a) For  $n, m \in \mathbb{Z}$ , the **log-volumes** constructed in (i), (ii), (iii) above determine log-volumes on the various “ $\mathcal{I}^{\mathbb{Q}}((-))$ ” that appear in the construction of the **local/global LGP-/lgp-monoids/Frobenioids** that appear in the  $\mathcal{F}^{\text{lt}}$ -prime-strips  ${}^{n, m} \mathfrak{S}_{\text{LGP}}^{\text{lt}}$ ,  ${}^{n, m} \mathfrak{S}_{\text{lgp}}^{\text{lt}}$  constructed in Proposition 3.7, (iii), (iv), relative to the **log-link**  ${}^{n, m-1} \mathcal{H}\mathcal{T}^{\Theta^{\pm \text{ell}} \text{NF}} \xrightarrow{\log} {}^{n, m} \mathcal{H}\mathcal{T}^{\Theta^{\pm \text{ell}} \text{NF}}$ .

(b) At the level of the  $\mathbb{Q}$ -spans of **log-shells** “ $\mathcal{I}^{\mathbb{Q}}((-))$ ” that arise from the various  **$\mathcal{F}$ -prime-strips** involved, the log-volumes of (a) indexed by  $(n, m)$  are **compatible** — in the sense discussed in Propositions 1.2, (iii); 1.3, (iii) — with the corresponding log-volumes indexed by  $(n, m-1)$ , relative to the **log-link**  ${}^{n, m-1} \mathcal{H}\mathcal{T}^{\Theta^{\pm \text{ell}} \text{NF}} \xrightarrow{\log} {}^{n, m} \mathcal{H}\mathcal{T}^{\Theta^{\pm \text{ell}} \text{NF}}$ .

*Proof.* The various assertions of Proposition 3.9 follow immediately from the definitions and the references quoted in the statements of these assertions.  $\circ$

**Remark 3.9.1.** In the spirit of the *explicit descriptions* of Remark 3.1.1, (i) [cf. also Remark 1.2.2, (i), (ii)], we make the following observations.

(i) Suppose that  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$ . Write  $\{\underline{v}_1, \dots, \underline{v}_n\}$  for the [distinct!] elements of  $\mathbb{V}$  that lie over  $v_{\mathbb{Q}}$ . For each  $i = 1, \dots, n$ , set  $k_i \stackrel{\text{def}}{=} K_{\underline{v}_i}$ ; write  $\mathcal{O}_{k_i} \subseteq k_i$  for the ring of integers of  $k_i$ ,

$$\mathcal{I}_i \stackrel{\text{def}}{=} (p_{v_{\mathbb{Q}}}^*)^{-1} \cdot \log_{k_i}(\mathcal{O}_{k_i}^{\times}) \subseteq k_i$$

— where  $p_{v_{\mathbb{Q}}}^* = p_{\underline{v}}$  if  $p_{v_{\mathbb{Q}}}$  is *odd*,  $p_{v_{\mathbb{Q}}}^* = p_{v_{\mathbb{Q}}}^2$  if  $p_{v_{\mathbb{Q}}}$  is *even* — cf. Remark 1.2.2, (i). Then, roughly speaking, in the notation of Proposition 3.9, (i), the **mono-analytic integral structures** of Proposition 3.2, (ii), in

$$\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}}) \xrightarrow{\sim} \bigoplus_{i=1}^n k_i; \quad \mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}}) \xrightarrow{\sim} \bigotimes_{\alpha \in A} \mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$$

are given by

$$\mathcal{I}(\alpha \mathcal{F}_{v_{\mathbb{Q}}}) \xrightarrow{\sim} \bigoplus_{i=1}^n \mathcal{I}_i; \quad \mathcal{I}(A \mathcal{F}_{v_{\mathbb{Q}}}) \xrightarrow{\sim} \bigotimes_{\alpha \in A} \mathcal{I}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$$

while the **local holomorphic integral structures**

$$\mathcal{O}_{\alpha \mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}}); \quad \mathcal{O}_{A \mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})$$

of Proposition 3.9, (i), in the topological rings  $\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$ ,  $\mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})$  — both of which are *direct sums* of finite extensions of  $\mathbb{Q}_{p_{v_{\mathbb{Q}}}}$  — are given by the *subrings of integers* in  $\mathcal{I}^{\mathbb{Q}}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})$ ,  $\mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})$ . Thus, by applying the formula of the final display of [AbsTopIII], Proposition 5.8, (iii), for the log-volume of  $\mathcal{I}_i$ , [one verifies easily that] one may compute the *log-volumes*

$$\mu_{\alpha, v_{\mathbb{Q}}}^{\log}(\mathcal{I}(\alpha \mathcal{F}_{v_{\mathbb{Q}}})) , \quad \mu_{A, v_{\mathbb{Q}}}^{\log}(\mathcal{I}(A \mathcal{F}_{v_{\mathbb{Q}}}))$$

entirely in terms of the given **initial  $\Theta$ -data**. We leave the routine details to the reader.

(ii) Suppose that  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$ . Write  $\{\underline{v}_1, \dots, \underline{v}_n\}$  for the [distinct!] elements of  $\mathbb{V}$  that lie over  $v_{\mathbb{Q}}$ . For each  $i = 1, \dots, n$ , set  $k_i \stackrel{\text{def}}{=} K_{\underline{v}_i}$ ; write  $\mathcal{O}_{k_i} \subseteq \{\lambda \in k_i \mid |\lambda| \leq 1\} \subseteq k_i$  for the “set of integers” of  $k_i$ ,

$$\mathcal{I}_i \stackrel{\text{def}}{=} \pi \cdot \mathcal{O}_{k_i} \subseteq k_i$$

— cf. Remark 1.2.2, (ii). Then, roughly speaking, in the notation of Proposition 3.9, (i), the **discrepancy** between the **mono-analytic integral structures** of

Proposition 3.2, (ii), determined by the  $\mathcal{I}(\dagger\mathcal{F}_{v_i}) \xrightarrow{\sim} \mathcal{I}_i \subseteq k_i$  and the **local holomorphic integral structures**

$$\mathcal{O}_{\alpha\mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(\alpha\mathcal{F}_{v_{\mathbb{Q}}}) \xrightarrow{\sim} \bigoplus_{i=1}^n k_i$$

$$\mathcal{O}_{A\mathcal{F}_{v_{\mathbb{Q}}}} \subseteq \mathcal{I}^{\mathbb{Q}}(A\mathcal{F}_{v_{\mathbb{Q}}}) \xrightarrow{\sim} \bigotimes_{\alpha \in A} \mathcal{I}^{\mathbb{Q}}(\alpha\mathcal{F}_{v_{\mathbb{Q}}})$$

of Proposition 3.9, (i), in the topological rings  $\mathcal{I}^{\mathbb{Q}}(\alpha\mathcal{F}_{v_{\mathbb{Q}}})$ ,  $\mathcal{I}^{\mathbb{Q}}(A\mathcal{F}_{v_{\mathbb{Q}}})$  — both of which are direct sums of complex archimedean fields — determined by taking the product [relative to this direct sum decomposition] of the respective “*subsets of integers*” may be computed entirely in terms of the given **initial  $\Theta$ -data**, by applying the following two [easily verified] observations:

- (a) Equip  $\mathbb{C}$  with its standard Hermitian metric, i.e., the metric determined by the complex norm. This metric on  $\mathbb{C}$  determines a tensor product metric on  $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$ , as well as a direct sum metric on  $\mathbb{C} \oplus \mathbb{C}$ . Then, relative to these metrics, any *isomorphism of topological rings* [i.e., arising from the Chinese remainder theorem]

$$\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} \xrightarrow{\sim} \mathbb{C} \oplus \mathbb{C}$$

is **compatible** with these **metrics**, up a factor of 2, i.e., the metric on the right-hand side corresponds to 2 times the metric on the left-hand side.

- (b) Relative to the notation of (a), the **direct sum decomposition**  $\mathbb{C} \oplus \mathbb{C}$ , together with its Hermitian metric, is **preserved**, relative to the displayed isomorphism of (a), by the operation of conjugation on either of the two copies of “ $\mathbb{C}$ ” that appear in  $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$ , as well as by the operations of multiplying by  $\pm 1$  or  $\pm\sqrt{-1}$  via either of the two copies of “ $\mathbb{C}$ ” that appear in  $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$ .

We leave the routine details to the reader.

(iii) The computation of the discrepancy between local holomorphic and mono-analytic integral structures will be discussed in more detail in [IUTchIV].

**Remark 3.9.2.** In the situation of Proposition 3.9, (iii), one may construct [“**mono-analytic**”] **algorithms** for recovering the **subquotient** of the *perfection* of  $(\dagger\mathbb{M}_{\text{mod}}^{\circ})_{\alpha} = (\dagger\mathbb{M}_{\text{MOD}}^{\circ})_{\alpha}$  associated to  $\underline{w} \in \underline{\mathbb{V}}$  [cf. Remark 3.6.1], together with the *submonoid of “nonnegative elements”* of such a subquotient, by considering the effect of multiplication by elements of  $(\dagger\mathbb{M}_{\text{mod}}^{\circ})_{\alpha} = (\dagger\mathbb{M}_{\text{MOD}}^{\circ})_{\alpha}$  on the *log-volumes* defined on the various  $\mathcal{I}^{\mathbb{Q}}(A, \alpha\mathcal{F}_{\underline{v}}) \xrightarrow{\sim} \mathcal{I}^{\mathbb{Q}}(A, \alpha\mathcal{D}_{\underline{v}}^{\dagger})$  [cf. Proposition 3.9, (ii)].

**Remark 3.9.3.** With regard to the **procession-normalizations** discussed in Proposition 3.9, (i), (ii), the reader might wonder the following: Is it possible to work with

more general **weighted averages**, i.e., as opposed to just *averages*, in the usual sense, over the capsules that appear in the procession?

The answer to this question is “no”. Indeed, in the situation of Proposition 3.4, (ii), for  $j \in \{1, \dots, l^*\}$ , the *packet-normalized* log-volume corresponding to the capsule with index set  $\mathbb{S}_{j+1}^\pm$  may be thought of as a log-volume that arises from “any one of the log-shells whose label  $\in \{0, 1, \dots, j\}$ ”. In particular, if  $j', j_1, j_2 \in \{1, \dots, l^*\}$ , and  $j' \leq j_1, j_2$ , then log-volumes corresponding to the **same log-shell** labeled  $j'$  might give rise to packet-normalized log-volumes corresponding to *either* of [the capsules with index sets]  $\mathbb{S}_{j_1+1}^\pm, \mathbb{S}_{j_2+1}^\pm$ . That is to say, in order for the resulting notion of a *procession-normalized log-volume* to be **compatible** with the appearance of the component labeled  $j'$  in various distinct capsules of the procession — i.e., compatible with the various **inclusion morphisms** of the procession! — one has no choice but to assign the *same weights* to [the capsules with index sets]  $\mathbb{S}_{j_1+1}^\pm, \mathbb{S}_{j_2+1}^\pm$ .

**Remark 3.9.4.** One way to understand the significance of the **log-link compatibility** of **log-volumes** discussed in Proposition 3.9, (iv), is as follows. Suppose that instead of knowing this property, one only knows that

each application of the **log-link** has the effect of **dilating volumes** by a factor  $\lambda \in \mathbb{R}_{>0}$ , i.e., which is not necessarily equal to 1.

Then in order to *compute log-volumes* in a fashion that is *consistent* with the various arrows [i.e., both Kummer isomorphisms and **log-links**!] of the “systems” constituted by the **log-Kummer correspondences** discussed in Proposition 3.5, (ii), it would be necessary, whenever  $\lambda \neq 1$ , to regard the various “log-volumes” computed as only giving rise to well-defined elements [not  $\in \mathbb{R}$ , but rather]

$$\in \mathbb{R}/\mathbb{Z} \cdot \log(\lambda) \quad (\cong \mathbb{S}^1)$$

— a situation which is *not acceptable*, relative to the goal of obtaining estimates [i.e., as in Corollary 3.12 below] for the various objects for which log-shells serve as “*multiradial containers*” [cf. the discussion of Remark 1.5.2; the content of Theorem 3.11 below].

**Proposition 3.10.** (**Global Kummer Theory and Non-interference with Local Integers**) *Let  $\{{}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}\}_{n,m \in \mathbb{Z}}$  be a collection of distinct  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theaters [relative to the given initial  $\Theta$ -data] — which we think of as arising from an **LGP-Gaussian log-theta-lattice** [cf. Definition 3.8, (iii); Proposition 3.5; Remark 3.8.2]. For each  $n \in \mathbb{Z}$ , write*

$${}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$$

for the  $\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater determined, up to isomorphism, by the various  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , where  $m \in \mathbb{Z}$ , via the **vertical coricity** of Theorem 1.5, (i) [cf. Remark 3.8.2].

(i) (**Vertically Coric Global LGP-, lgp-Frobenioids and Associated Kummer Theory**) Recall the constructions of various global Frobenioids in Proposition 3.7, (i), (ii), (iii), (iv), in the context of  $\mathcal{F}$ -prime-strip processions. Then by applying these constructions to the  $\mathcal{F}$ -prime-strips “ $\mathfrak{F}^{(n, \circ \mathcal{D}_{\succ})_t}$ ” [cf. the notation of Proposition 3.5, (i)] and the various full **log-links** associated [cf. the discussion of Proposition 1.2, (ix)] to these  $\mathcal{F}$ -prime-strips — which we consider in a fashion **compatible** with the  $\mathbb{F}_l^{\times \pm}$ -**symmetries** involved [cf. Remark 1.3.2; Proposition 3.4, (ii)] — we obtain **functorial algorithms** in the  $\mathcal{D}$ - $\Theta^{\pm \text{ell}} \mathbf{NF}$ -**Hodge theater**  ${}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}}$  for constructing [number] **fields, monoids, and Frobenioids** equipped with natural isomorphisms

$$\begin{aligned} \overline{\mathbb{M}}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha} &= \overline{\mathbb{M}}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha} \\ &\supseteq \mathbb{M}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha} = \mathbb{M}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha} \end{aligned}$$

$$\overline{\mathbb{M}}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha} \supseteq \mathbb{M}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}$$

$$\mathcal{F}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha} \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha} \xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}$$

[cf. the number fields, monoids, and Frobenioids “ $\overline{\mathbb{M}}_{\text{mod}}^{\circ}(\dagger \mathcal{D}^{\circ})_j \supseteq \mathbb{M}_{\text{mod}}^{\circ}(\dagger \mathcal{D}^{\circ})_j$ ”, “ $\mathcal{F}_{\text{mod}}^{\circ}(\dagger \mathcal{D}^{\circ})_j$ ” of [IUTchII], Corollary 4.7, (ii)] for  $\alpha \in A$ , where  $A$  is a subset of  $J$  [cf. Proposition 3.3], as well as  $\mathcal{F}^{\text{lt}}$ -**prime-strips** equipped with natural isomorphisms

$$\mathfrak{F}^{\text{lt}}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\text{gau}} \xrightarrow{\sim} \mathfrak{F}^{\text{lt}}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\text{LGP}} \xrightarrow{\sim} \mathfrak{F}^{\text{lt}}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\text{lgp}}$$

— [all of] which we shall refer to as being “**vertically coric**”. For each  $n, m \in \mathbb{Z}$ , these functorial algorithms are **compatible** [in the evident sense] with the [“non-vertically coric”] functorial algorithms of Proposition 3.7, (i), (ii), (iii), (iv) — i.e., where [in Proposition 3.7, (iii), (iv)] we take “ $\dagger$ ” to be “ $n, m$ ” and “ $\ddagger$ ” to be “ $n, m - 1$ ” — relative to the **Kummer isomorphisms** of labeled data

$$\Psi_{\text{cns}}({}^{n, m'} \mathfrak{F}_{\succ})_t \xrightarrow{\sim} \Psi_{\text{cns}}({}^{n, \circ} \mathcal{D}_{\succ})_t$$

$$({}^{n, m'} \mathbb{M}_{\text{mod}}^{\circ})_j \xrightarrow{\sim} \mathbb{M}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{D}^{\circ})_j; \quad ({}^{n, m'} \overline{\mathbb{M}}_{\text{mod}}^{\circ})_j \xrightarrow{\sim} \overline{\mathbb{M}}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{D}^{\circ})_j$$

[cf. [IUTchII], Corollary 4.6, (iii); [IUTchII], Corollary 4.8, (ii)] and the evident identification, for  $m' = m, m - 1$ , of  ${}^{n, m'} \mathfrak{F}_t$  [i.e., the  $\mathcal{F}$ -prime-strip that appears in the associated  $\Theta^{\pm}$ -bridge] with the  $\mathcal{F}$ -prime-strip associated to  $\Psi_{\text{cns}}({}^{n, m'} \mathfrak{F}_{\succ})_t$  [cf. Proposition 3.5, (i)]. In particular, for each  $n, m \in \mathbb{Z}$ , we obtain “**Kummer isomorphisms**” of fields, monoids, Frobenioids, and  $\mathcal{F}^{\text{lt}}$ -prime-strips

$$({}^{n, m} \overline{\mathbb{M}}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} \overline{\mathbb{M}}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}; \quad ({}^{n, m} \overline{\mathbb{M}}_{\text{MOD}}^{\circ})_{\alpha} \xrightarrow{\sim} \overline{\mathbb{M}}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}$$

$$({}^{n, m} \mathbb{M}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} \mathbb{M}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}; \quad ({}^{n, m} \mathbb{M}_{\text{MOD}}^{\circ})_{\alpha} \xrightarrow{\sim} \mathbb{M}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}$$

$$({}^{n, m} \mathcal{F}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}; \quad ({}^{n, m} \mathcal{F}_{\text{MOD}}^{\circ})_{\alpha} \xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}$$

$$({}^{n, m} \overline{\mathbb{M}}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} \overline{\mathbb{M}}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}; \quad ({}^{n, m} \mathbb{M}_{\text{mod}}^{\circ})_{\alpha} \xrightarrow{\sim} \mathbb{M}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \mathbf{NF}})_{\alpha}$$



$$\begin{aligned} ({}^{n,m}\mathcal{F}_{\text{mod}}^{\circ})_{\alpha} &\xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circ}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\alpha}; & {}^{n,m}\mathfrak{F}_{\text{gau}}^{\text{ll}} &\xrightarrow{\sim} \mathfrak{F}^{\text{ll}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{gau}} \\ {}^{n,m}\mathfrak{F}_{\text{LGP}}^{\text{ll}} &\xrightarrow{\sim} \mathfrak{F}^{\text{ll}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}}; & {}^{n,m}\mathfrak{F}_{\text{lgp}}^{\text{ll}} &\xrightarrow{\sim} \mathfrak{F}^{\text{ll}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{lgp}} \end{aligned}$$

that are **compatible** with the various equalities, natural inclusions, and natural isomorphisms discussed above.

(ii) **(Non-interference with Local Integers)** In the notation of Propositions 3.2, (ii); 3.4, (i); 3.7, (i), (ii); 3.9, (iii), we have

$$\begin{aligned} (\dagger\mathbb{M}_{\text{MOD}}^{\circ})_{\alpha} \cap \prod_{\underline{v} \in \underline{\mathbb{V}}} \Psi_{\text{log}(A, \alpha \mathcal{F}_{\underline{v}})} &= (\dagger\mathbb{M}_{\text{MOD}}^{\circ\mu})_{\alpha} \\ \left( \subseteq \prod_{\underline{v} \in \underline{\mathbb{V}}} \mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}}) = \prod_{v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}}) = \mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}}) \right) \end{aligned}$$

— where we write  $(\dagger\mathbb{M}_{\text{MOD}}^{\circ\mu})_{\alpha} \subseteq (\dagger\mathbb{M}_{\text{MOD}}^{\circ})_{\alpha}$  for the [finite] subgroup of torsion elements, i.e., **roots of unity**; we identify the product  $\prod_{\underline{v} \ni \underline{v} | v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \mathcal{I}^{\mathbb{Q}}(A, \alpha \mathcal{F}_{\underline{v}})$  with  $\mathcal{I}^{\mathbb{Q}}(A \mathcal{F}_{v_{\mathbb{Q}}})$ . Now let us think of the various **groups**

$$({}^{n,m}\mathbb{M}_{\text{MOD}}^{\circ})_j$$

[of nonzero elements of a number field] as acting on various portions of the modules

$$\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm} \mathcal{F}({}^{n,\circ}\mathcal{D}_{\succ})_{\mathbb{V}_{\mathbb{Q}}})$$

[cf. the notation of Proposition 3.5, (ii)] **not** via a **single Kummer isomorphism** as in (i), but rather via the **totality** of the various pre-composites of the relevant Kummer isomorphism with the various **iterates** of the **log-links** of the LGP-Gaussian log-theta-lattice — i.e., which is possible, up to [harmless!] “**identity indeterminacies**” at an adjacent “ $m$ ”, precisely as a consequence of the equality of the first display of the present (ii) [cf. the discussion of Remark 1.2.3, (ii); the discussion of Definition 1.1, (ii), concerning quotients by  $\Psi_{\dagger \mathcal{F}_{\underline{v}}}^{\mu_N}$  at  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ ; the discussion of Definition 1.1, (iv), at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ ]. Then one obtains a sort of “**log-Kummer correspondence**” between the **totality**, as  $m$  ranges over the elements of  $\mathbb{Z}$ , of the various groups [of nonzero elements of a number field] just discussed [i.e., which are labeled by “ $n, m$ ”] and their actions on the “ $\mathcal{I}^{\mathbb{Q}}$ ” labeled by “ $n, \circ$ ” which is **invariant** with respect to the **translation symmetries** [cf. Proposition 1.3, (iv)] of the  $n$ -th column of the LGP-Gaussian log-theta-lattice [cf. the discussion of Remark 1.2.2, (iii)].

(iii) **(Frobenioid-theoretic log-Kummer Correspondences)** The relevant Kummer isomorphisms of (i) induce, via the “**log-Kummer correspondence**” of (ii) [cf. also Proposition 3.7, (i); Remarks 3.6.1, 3.9.2], **isomorphisms of Frobenioids**

$$\begin{aligned} ({}^{n,m}\mathcal{F}_{\text{MOD}}^{\circ})_{\alpha} &\xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circ}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\alpha} \\ ({}^{n,m}\mathcal{F}_{\text{MOD}}^{\circ\mathbb{R}})_{\alpha} &\xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circ\mathbb{R}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\alpha} \end{aligned}$$

that are **mutually compatible**, as  $m$  varies over the elements of  $\mathbb{Z}$ , with the **log-links** of the LGP-Gaussian log-theta-lattice. Moreover, these compatible isomorphisms of Frobenioids, together with the relevant Kummer isomorphisms of (i), induce, via the **global “log-Kummer correspondence”** of (ii) and the **splitting monoid portion** of the “log-Kummer correspondence” of Proposition 3.5, (ii), **isomorphisms of associated  $\mathcal{F}^{\dagger\perp}$ -prime-strips** [cf. Definition 2.4, (iii)]

$${}^{n,m}\mathfrak{F}_{\text{LGP}}^{\dagger\perp} \xrightarrow{\sim} \mathfrak{F}^{\dagger\perp}(n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}}$$

that are **mutually compatible**, as  $m$  varies over the elements of  $\mathbb{Z}$ , with the **log-links** of the LGP-Gaussian log-theta-lattice.

*Proof.* The various assertions of Proposition 3.10 follow immediately from the definitions and the references quoted in the statements of these assertions. Here, we observe that the computation of the *intersection* of the first display of (ii) is an immediate consequence of the well-known fact that the set of nonzero elements of a number field that are *integral* at all of the places of the number field consists of the set of *roots of unity* contained in the number field [cf. the discussion of Remark 1.2.3, (ii); [Lang], p. 144, the proof of Theorem 5].  $\circ$

### Remark 3.10.1.

(i) Observe that it follows immediately from the well-known fact from *elementary algebraic number theory* applied in the proof of Proposition 3.10, (ii), that the first display of Proposition 3.10, (ii), admits a **Galois-orbit-theoretic analogue**:

$$(\dagger\mathbb{M}_{\text{ORB}}^{\circ})_{\alpha} \bigcap \prod_{\underline{v} \in \underline{\mathbb{V}}} (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})})_{\text{orb}} = (\dagger\mathbb{M}_{\text{ORB}}^{\circ\mu})_{\alpha} \left( \subseteq \underline{\log}_{\text{orb}}(\alpha \mathcal{F}_{\underline{\mathbb{V}}}) \right)$$

— where we write

$$(\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})})_{\text{orb}}$$

for the set of collections of *Galois-orbits* [i.e., “ $\alpha\Pi_{\underline{v}}$ ”- or “ $\dagger G_{\underline{v}}$ ”-orbits when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  — cf. Propositions 1.2, (vi), and 3.1, (i);  $\{1\}$ -orbits, when  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ ] of  $\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}$  [cf. the discussion of Remark 3.3.4] and, in the notation of Remark 3.7.1,

$$(\dagger\mathbb{M}_{\text{ORB}}^{\circ\mu})_{\alpha} = (\dagger\mathbb{M}_{\text{orb}}^{\circ\mu})_{\alpha} \subseteq (\dagger\mathbb{M}_{\text{ORB}}^{\circ})_{\alpha} = (\dagger\mathbb{M}_{\text{orb}}^{\circ})_{\alpha}$$

for the subset of collections of Galois-orbits all of whose members are *roots of unity*.

(ii) Write

$$\left( (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})})_{\text{orb}} \supseteq \right) (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}^{\times})_{\text{orb}} \twoheadrightarrow (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}^{\times\mu})_{\text{orb}}$$

for the respective sets of collections of *Galois-orbits* [i.e., “ $\alpha\Pi_{\underline{v}}$ ”- or “ $\dagger G_{\underline{v}}$ ”-orbits when  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  — cf. Propositions 1.2, (vi), and 3.1, (i);  $\{1\}$ -orbits, when  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$ ] of  $\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}^{\times}$ ,  $\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}^{\times\mu}$  [cf. the discussion of (i) above] and

$$\log(\alpha \mathfrak{F})^{\dagger \times \mu} \stackrel{\text{def}}{=} \{ \log(\alpha \mathcal{F}_{\underline{v}})^{\dagger \times \mu} \}_{\underline{v} \in \underline{\mathbb{V}}}$$

for the  $\mathcal{F}^{\dagger \times \mu}$ -*prime-strip* determined [cf. [IUTchII], Definition 4.9, (vi), (vii)] by the  $\mathcal{F}$ -*prime-strip*  $\log(\alpha \mathfrak{F})$  [cf. Definition 1.1, (iii)]. Then one way to interpret the computation of the **intersection** of the first display of (i) is as an analogue for “ $\mathbb{M}^{\circledast}$ ” of the *multiradiality for mono-theta environments* discussed in Theorem 2.2, (ii); Remark 2.2.1 [cf., especially, Fig. 2.2]; Corollary 2.3, (iv) — i.e., as a sort of **multiradiality for number fields** to the effect that

the **cyclotomic rigidity isomorphism** and **related Kummer theory** of [IUTchI], Example 5.1, (v), are **insulated** from the **indeterminacies** that arise from automorphisms  $\in \text{Aut}_{\mathcal{F}^{\dagger \times \mu}}(\log(\alpha \mathfrak{F})^{\dagger \times \mu})$ .

That is to say, it follows immediately from (i) that

the **identity automorphism** of  $(\dagger \mathbb{M}_{\text{ORB}}^{\circledast})_{\alpha}$  is **compatible** with the **collection** of automorphisms of  $(\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}^{\times \mu})_{\text{orb}}$  induced by **arbitrary automorphisms**  $\in \text{Aut}_{\mathcal{F}^{\dagger \times \mu}}(\log(\alpha \mathfrak{F})^{\dagger \times \mu})$ , **relative to the diagram**

$$\begin{array}{ccccc}
 & & (\dagger \mathbb{M}_{\text{ORB}}^{\circledast})_{\alpha} \cap \prod_{\underline{v} \in \underline{\mathbb{V}}} (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})})_{\text{orb}} & \longrightarrow & (\dagger \mathbb{M}_{\text{ORB}}^{\circledast})_{\alpha} \\
 & & \downarrow & & \downarrow \\
 \prod_{\underline{v} \in \underline{\mathbb{V}}} (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}^{\times})_{\text{orb}} & \hookrightarrow & \prod_{\underline{v} \in \underline{\mathbb{V}}} (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})})_{\text{orb}} & \longrightarrow & \underline{\log}_{\text{orb}}(\alpha \mathcal{F}_{\mathbb{V}_{\mathbb{Q}}}) \\
 \downarrow & & & & \\
 \prod_{\underline{v} \in \underline{\mathbb{V}}} (\Psi_{\log(\alpha \mathcal{F}_{\underline{v}})}^{\times \mu})_{\text{orb}} & & & & 
 \end{array}$$

— in which the arrows are the natural morphisms.

Here, we note that the significance of considering various versions of “orb” lies in the fact that *collections of Galois-orbits* arise naturally when one restricts the *Kummer theory* — i.e., the theory of Galois-stable collections of  $N$ -th roots of elements [for  $N \geq 1$  an integer] — of copies of the number field  $F_{\text{mod}}$  [i.e., such as  $(\dagger \mathbb{M}_{\text{MOD}}^{\circledast})_{\alpha}$ ] to the *Kummer theory* of the various localizations of the number field at  $\underline{v} \in \underline{\mathbb{V}}$ .

### Remark 3.10.2.

(i) Note that the **log-Kummer correspondence** of Proposition 3.10, (ii), induces isomorphisms of Frobenioids as in the first display of Proposition 3.10, (iii), precisely because the construction of “ $(\dagger \mathcal{F}_{\text{MOD}}^{\circledast})_{\alpha}$ ” only involves the group “ $(\dagger \mathbb{M}_{\text{MOD}}^{\circledast})_{\alpha}$ ”, together with the collection of *subquotients* of its perfection indexed by  $\underline{\mathbb{V}}$  [cf. Proposition 3.7, (i); Remarks 3.6.1, 3.9.2]. By contrast, the construction of “ $(\dagger \mathcal{F}_{\text{mod}}^{\circledast})_{\alpha}$ ” also involves the *local monoids* “ $\Psi_{\log(A, \alpha \mathcal{F}_{\underline{v}})} \subseteq \log(A, \alpha \mathcal{F}_{\underline{v}})$ ” in an essential way [cf. Proposition 3.7, (ii)]. These local monoids are subject to a somewhat more complicated “**log-Kummer correspondence**” [cf. Proposition 3.5, (ii)] that revolves around “*upper semi-compatibility*”, i.e., in a word, *one-sided inclusions*, as opposed to precise equalities. The imprecise nature of such one-sided inclusions is *incompatible* with the construction of “ $(\dagger \mathcal{F}_{\text{mod}}^{\circledast})_{\alpha}$ ”. In particular, one cannot construct **log-link-compatible** isomorphisms of Frobenioids for “ $(\dagger \mathcal{F}_{\text{mod}}^{\circledast})_{\alpha}$ ” as in the first display of Proposition 3.10, (iii).

(ii) The **precise compatibility** of “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ” with the **log**-links of the LGP-Gaussian log-theta-lattice [cf. the discussion of (i); the first “mutual compatibility” of Proposition 3.10, (iii)] makes it more suited [i.e., by comparison to “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ”] to the task of computing the **Kummer-detachment indeterminacies** [cf. Remark 1.5.4, (i), (iii)] that arise when one attempts to pass from the *Frobenius-like structures* constituted by the global portion of the domain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -links of the LGP-Gaussian log-theta-lattice to corresponding *étale-like structures*. That is to say, the mutual compatibility of the isomorphisms

$${}^{n,m}\mathfrak{F}_{\text{LGP}}^{\text{ll}\perp} \xrightarrow{\sim} \mathfrak{F}^{\text{ll}\perp}(n, \circ\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}}$$

of the second display of Proposition 3.10, (iii), asserts, in effect, that such *Kummer-detachment indeterminacies do not arise*. This is precisely the reason why we wish to work with the LGP-, as opposed to the **lgp**-, Gaussian log-theta lattice [cf. Remark 3.8.1]. On the other hand, the essentially **multiplicative** nature of “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ” [cf. Remark 3.6.2, (ii)] makes it *ill-suited* to the task of computing the **étale-transport indeterminacies** [cf. Remark 1.5.4, (i), (ii)] that occur as one passes between distinct arithmetic holomorphic structures on opposite sides of a  $\Theta_{\text{LGP}}^{\times\mu}$ -link.

(iii) By contrast, whereas the **additive nature** of the local modules [i.e., local fractional ideals] that occur in the construction of “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ” renders “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ” ill-suited to the computation of Kummer-detachment indeterminacies [cf. the discussion of (i), (ii)], the *close relationship* [cf. Proposition 3.9, (i), (ii), (iii)] of these local modules to the **mono-analytic log-shells** that are **coric** with respect to the  $\Theta_{\text{LGP}}^{\times\mu}$ -link [cf. Theorem 1.5, (iv); Remark 3.8.2] renders “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ” *well-suited* to the computation of the **étale-transport indeterminacies** that occur as one passes between distinct arithmetic holomorphic structures on opposite sides of a  $\Theta_{\text{LGP}}^{\times\mu}$ -link. That is to say, although various *distortions* of these local modules arise as a result of both [the Kummer-detachment indeterminacies constituted by] the local “**upper semi-compatibility**” of Proposition 3.5, (ii), and [the étale-transport indeterminacies constituted by] the **discrepancy between local holomorphic and mono-analytic integral structures** [cf. Remark 3.9.1, (i), (ii)], one may nevertheless compute — i.e., if one takes into account the various distortions that occur, “**estimate**” — the **global arithmetic degrees** of objects of “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ” by computing **log-volumes** [cf. Proposition 3.9, (iii)], which are **bi-coric**, i.e., coric with respect to both the  $\Theta_{\text{LGP}}^{\times\mu}$ -links [cf. Proposition 3.9, (ii)] and the **log**-links [cf. Proposition 3.9, (iv)] of the LGP-Gaussian log-theta-lattice. This **computability** is precisely the topic of Corollary 3.12 below. On the other hand, the issue of obtaining *concrete estimates* will be treated in [IUTchIV].

(iv) The various properties of “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ” and “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ” discussed in (i), (ii), (iii) above are summarized in Fig. 3.2 below. In this context, it is of interest to observe that the natural isomorphisms of Frobenioids

$$\mathcal{F}_{\text{mod}}^{\circledast}(n, \circ\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\alpha} \xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circledast}(n, \circ\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\alpha}$$

as well as the resulting isomorphisms of  $\mathcal{F}^{\text{ll}\perp}$ -prime-strips

$$\mathfrak{F}^{\text{ll}\perp}(n, \circ\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}} \xrightarrow{\sim} \mathfrak{F}^{\text{ll}\perp}(n, \circ\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{lgp}}$$

of Proposition 3.10, (i), play the *highly nontrivial* role of relating [cf. the discussion of [IUTchII], Remark 4.8.2, (i)] the “*multiplicatively biased*  $\mathcal{F}_{\text{MOD}}^{\circledast}$ ” to the “*additively biased*  $\mathcal{F}_{\text{mod}}^{\circledast}$ ” by means of the **global ring structure** of the number field  $\overline{\mathbb{M}}_{\text{mod}}^{\circledast}(n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \text{NF}})_{\alpha} = \overline{\mathbb{M}}_{\text{MOD}}^{\circledast}(n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \text{NF}})_{\alpha}$ . A similar statement holds concerning the tautological isomorphism of  $\mathcal{F}^{\text{!}}$ -prime-strips  $\dagger \mathfrak{F}_{\text{LGP}}^{\text{!}} \xrightarrow{\sim} \dagger \mathfrak{F}_{\text{lgp}}^{\text{!}}$  of Proposition 3.7, (iv).

$\mathcal{F}_{\text{MOD}}^{\circledast}/\underline{\text{LGP-structures}}$	$\mathcal{F}_{\text{mod}}^{\circledast}/\underline{\text{lgp-structures}}$
biased toward <b>multiplicative</b> structures	biased toward <b>additive</b> structures
easily related to <b>value group/non-coric</b> portion “ $(-)^{\text{!}} \blacktriangleright$ ” of $\Theta_{\text{LGP}}^{\times \mu}$ -link	easily related to <b>unit group/coric</b> portion “ $(-)^{\text{!}} \times \mu$ ” of $\Theta_{\text{LGP}}^{\times \mu}/\Theta_{\text{lgp}}^{\times \mu}$ -link, i.e., <b>mono-analytic log-shells</b>
admits <b>precise log-Kummer</b> <b>correspondence</b>	only admits <b>“upper semi-compatible”</b> <b>log-Kummer correspondence</b>
<b>rigid</b> , but not suited to explicit computation	subject to <b>substantial distortion</b> , but suited to <b>explicit estimates</b>

Fig. 3.2:  $\mathcal{F}_{\text{MOD}}^{\circledast}/\text{LGP-structures}$  versus  $\mathcal{F}_{\text{mod}}^{\circledast}/\text{lgp-structures}$

We are now ready to discuss the *main theorem* of the present series of papers.

**Theorem 3.11.** (Multiradial Algorithms via LGP-Monoids/Frobenioids)  
Fix a collection of **initial  $\Theta$ -data**

$$(\overline{F}/F, X_F, l, \underline{C}_K, \underline{V}, \underline{\epsilon})$$

as in [IUTchI], Definition 3.1. Let

$$\{n, m \mathcal{HT}^{\Theta^{\pm \text{ell}} \text{NF}}\}_{n, m \in \mathbb{Z}}$$

be a **collection of distinct  $\Theta^{\pm \text{ell}} \text{NF}$ -Hodge theaters** [relative to the given initial  $\Theta$ -data] — which we think of as arising from an **LGP-Gaussian log-theta-lattice** [cf. Definition 3.8, (iii)]. For each  $n \in \mathbb{Z}$ , write

$$n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm \text{ell}} \text{NF}}$$

for the  $\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater determined, up to isomorphism, by the various  $n, m \mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , where  $m \in \mathbb{Z}$ , via the **vertical coricity** of Theorem 1.5, (i) [cf. Remark 3.8.2].

(i) (**Multiradial Representation**) Consider the procession of  $\mathcal{D}^+$ -prime-strips  $\text{PrC}^{(n, \circ) \mathcal{D}_T^+}$

$$\{^{n, \circ} \mathcal{D}_0^+\} \hookrightarrow \{^{n, \circ} \mathcal{D}_0^+, ^{n, \circ} \mathcal{D}_1^+\} \hookrightarrow \dots \hookrightarrow \{^{n, \circ} \mathcal{D}_0^+, ^{n, \circ} \mathcal{D}_1^+, \dots, ^{n, \circ} \mathcal{D}_{l\pm}^+\}$$

obtained by applying the natural functor of [IUTchI], Proposition 6.9, (ii), to [the  $\mathcal{D}\text{-}\Theta^{\pm}$ -bridge associated to]  $^{n, \circ} \mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}}$ . Consider also the following data:

(a) for  $\underline{\mathbb{V}} \ni \underline{v} \mid v_{\mathbb{Q}}, j \in |\mathbb{F}_l|$ , the **topological modules and mono-analytic integral structures**

$$\mathcal{I}(\mathbb{S}_{j+1}^{\pm}; ^{n, \circ} \mathcal{D}_{v_{\mathbb{Q}}}^+) \subseteq \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; ^{n, \circ} \mathcal{D}_{v_{\mathbb{Q}}}^+); \quad \mathcal{I}(\mathbb{S}_{j+1}^{\pm}; j; ^{n, \circ} \mathcal{D}_{\underline{v}}^+) \subseteq \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; ^{n, \circ} \mathcal{D}_{\underline{v}}^+)$$

— where the notation “ $; n, \circ$ ” denotes the result of applying the construction in question to the case of  $\mathcal{D}^+$ -prime-strips labeled “ $n, \circ$ ” — of Proposition 3.2, (ii) [cf. also the notational conventions of Proposition 3.4, (ii)], which we regard as equipped with the **procession-normalized mono-analytic log-volumes** of Proposition 3.9, (ii);

(b) for  $\underline{\mathbb{V}}^{\text{bad}} \ni \underline{v}$ , the **splitting monoid**

$$\Psi_{\text{LGP}}^{\perp}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}}$$

of Proposition 3.5, (ii), (c) [cf. also the notation of Proposition 3.5, (i)], which we regard — via the natural poly-isomorphisms

$$\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; ^{n, \circ} \mathcal{D}_{\underline{v}}^+) \xrightarrow{\sim} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; \mathcal{F}^{\perp \times \mu}({}^{n, \circ} \mathcal{D}_{\prec})_{\underline{v}}) \xrightarrow{\sim} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; \mathcal{F}({}^{n, \circ} \mathcal{D}_{\prec})_{\underline{v}})$$

for  $j \in \mathbb{F}_l^*$  [cf. Proposition 3.2, (i), (ii)] — as a **subset** of

$$\prod_{j \in \mathbb{F}_l^*} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; ^{n, \circ} \mathcal{D}_{\underline{v}}^+)$$

equipped with  $a(n)$  [multiplicative] **action** on  $\prod_{j \in \mathbb{F}_l^*} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; ^{n, \circ} \mathcal{D}_{\underline{v}}^+)$ ;

(c) for  $j \in \mathbb{F}_l^*$ , the **number field**

$$\begin{aligned} \overline{\mathbb{M}}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}})_j &= \overline{\mathbb{M}}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}})_j \\ &\subseteq \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; ^{n, \circ} \mathcal{D}_{\mathbb{V}_{\mathbb{Q}}}^+) \stackrel{\text{def}}{=} \prod_{v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; ^{n, \circ} \mathcal{D}_{v_{\mathbb{Q}}}^+) \end{aligned}$$

[cf. the natural isomorphisms discussed in (b); Proposition 3.9, (iii); Proposition 3.10, (i)], together with **natural isomorphisms** between the associated **global non-realified/ realified Frobenioids**

$$\mathcal{F}_{\text{MOD}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}})_j \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circ}({}^{n, \circ} \mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}})_j$$



$$\mathcal{F}_{\text{MOD}}^{\circledast\mathbb{R}}(n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j \xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circledast\mathbb{R}}(n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j$$

[cf. Proposition 3.10, (i)], whose associated “**global degrees**” may be computed by means of the **log-volumes** of (a) [cf. Proposition 3.9, (iii)].

Write

$$n, \circ \mathfrak{R}^{\text{LGP}}$$

for the **collection of data** (a), (b), (c) regarded up to **indeterminacies** of the following two types:

- (Ind1) the indeterminacies induced by the **automorphisms** of the **procession of  $\mathcal{D}^+$ -prime-strips**  $\text{Prc}(n, \circ \mathcal{D}_T^+)$ ;
- (Ind2) for each  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$  (respectively,  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$ ), the indeterminacies induced by the action of **independent** copies of **Ism** [cf. Proposition 1.2, (vi)] (respectively, copies of each of the automorphisms of order 2 whose orbit constitutes the poly-automorphism discussed in Proposition 1.2, (vii)) on each of the **direct summands** of the  $j+1$  **factors** appearing in the tensor product used to define  $\mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; n, \circ \mathcal{D}_{v_{\mathbb{Q}}}^+)$  [cf. (a) above; Proposition 3.2, (ii)] — where we recall that the cardinality of the collection of direct summands is equal to the cardinality of the set of  $\underline{v} \in \underline{\mathbb{V}}$  that lie over  $v_{\mathbb{Q}}$ .

Then  $n, \circ \mathfrak{R}^{\text{LGP}}$  may be constructed via an **algorithm** in the procession of  $\mathcal{D}^+$ -prime-strips  $\text{Prc}(n, \circ \mathcal{D}_T^+)$  that is **functorial** with respect to isomorphisms of processions of  $\mathcal{D}^+$ -prime-strips. For  $n, n' \in \mathbb{Z}$ , the **permutation symmetries** of the **étale-picture** discussed in [IUTchI], Corollary 6.10, (iii); [IUTchII], Corollary 4.11, (ii), (iii) [cf. also Corollary 2.3, (ii); Remark 3.8.2, of the present paper], induce **compatible poly-isomorphisms**

$$\text{Prc}(n, \circ \mathcal{D}_T^+) \xrightarrow{\sim} \text{Prc}(n', \circ \mathcal{D}_T^+); \quad n, \circ \mathfrak{R}^{\text{LGP}} \xrightarrow{\sim} n', \circ \mathfrak{R}^{\text{LGP}}$$

which are, moreover, compatible with the poly-isomorphisms

$$n, \circ \mathcal{D}_0^+ \xrightarrow{\sim} n', \circ \mathcal{D}_0^+$$

induced by the **bi-coricity** poly-isomorphisms of Theorem 1.5, (iii) [cf. also [IUTchII], Corollaries 4.10, (iv); 4.11, (i)].

(ii) (**log-Kummer Correspondence**) For  $n, m \in \mathbb{Z}$ , the **Kummer isomorphisms** of labeled data

$$\Psi_{\text{cns}}(n, m \mathfrak{F}_{\succ})_t \xrightarrow{\sim} \Psi_{\text{cns}}(n, \circ \mathcal{D}_{\succ})_t$$

$$\left\{ \pi_1(n, \circ \mathcal{D}^{\circledast}) \curvearrowright n, m \overline{\mathbb{M}}^{\circledast} \right\} \xrightarrow{\sim} \left\{ \pi_1(n, \circ \mathcal{D}^{\circledast}) \curvearrowright \overline{\mathbb{M}}^{\circledast}(n, \circ \mathcal{D}^{\circledast}) \right\}$$

$$(n, m \overline{\mathbb{M}}_{\text{mod}}^{\circledast})_j \xrightarrow{\sim} \overline{\mathbb{M}}_{\text{mod}}^{\circledast}(n, \circ \mathcal{D}^{\circledast})_j; \quad (n, m \overline{\mathbb{M}}_{\text{orb}}^{\circledast})_j \xrightarrow{\sim} \overline{\mathbb{M}}_{\text{orb}}^{\circledast}(n, \circ \mathcal{D}^{\circledast})_j$$

— where  $t \in \text{LabCusp}^{\pm}(n, \circ \mathcal{D}_{\succ})$  — of [IUTchII], Corollary 4.6, (iii); [IUTchII], Corollary 4.8, (i), (ii) [cf. also Propositions 3.5, (i); 3.10, (i), of the present paper] induce **isomorphisms** between the **vertically coric** data (a), (b), (c) of (i) and

the corresponding data arising from each  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$ , i.e.:

(a) for  $\mathbb{V} \ni \underline{v} \mid v_{\mathbb{Q}}$ ,  $j \in |\mathbb{F}_l|$ , isomorphisms with **local mono-analytic tensor packets** and their  $\mathbb{Q}$ -spans

$$\begin{aligned} \mathcal{I}(\mathbb{S}_{j+1}^{\pm}; n, m \mathcal{F}_{v_{\mathbb{Q}}}) &\xrightarrow{\sim} \mathcal{I}(\mathbb{S}_{j+1}^{\pm}; n, m \mathcal{F}_{v_{\mathbb{Q}}}^+ \times \mu) \xrightarrow{\sim} \mathcal{I}(\mathbb{S}_{j+1}^{\pm}; n, \circ \mathcal{D}_{v_{\mathbb{Q}}}^+) \\ \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; n, m \mathcal{F}_{v_{\mathbb{Q}}}) &\xrightarrow{\sim} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; n, m \mathcal{F}_{v_{\mathbb{Q}}}^+ \times \mu) \xrightarrow{\sim} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; n, \circ \mathcal{D}_{v_{\mathbb{Q}}}^+) \\ \mathcal{I}(\mathbb{S}_{j+1}^{\pm}; j; n, m \mathcal{F}_{\underline{v}}) &\xrightarrow{\sim} \mathcal{I}(\mathbb{S}_{j+1}^{\pm}; j; n, m \mathcal{F}_{\underline{v}}^+ \times \mu) \xrightarrow{\sim} \mathcal{I}(\mathbb{S}_{j+1}^{\pm}; j; n, \circ \mathcal{D}_{\underline{v}}^+) \\ \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; n, m \mathcal{F}_{\underline{v}}) &\xrightarrow{\sim} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; n, m \mathcal{F}_{\underline{v}}^+ \times \mu) \xrightarrow{\sim} \mathcal{I}^{\mathbb{Q}}(\mathbb{S}_{j+1}^{\pm}; j; n, \circ \mathcal{D}_{\underline{v}}^+) \end{aligned}$$

[cf. Propositions 3.2, (i), (ii); 3.4, (ii); 3.5, (i)], all of which are **compatible** with the respective **log-volumes** [cf. Proposition 3.9, (ii)];

(b) for  $\mathbb{V}^{\text{bad}} \ni \underline{v}$ , isomorphisms of **splitting monoids**

$$\Psi_{\mathcal{F}_{\text{LGP}}}^{\perp}({}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}} \xrightarrow{\sim} \Psi_{\text{LGP}}^{\perp}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\underline{v}}$$

[cf. Proposition 3.5, (i); Proposition 3.5, (ii), (c)];

(c) for  $j \in \mathbb{F}_l^*$ , isomorphisms of **number fields** and **non-realified/realified global Frobenioids**

$$\begin{aligned} ({}^{n,m}\overline{\mathbb{M}}_{\text{MOD}}^{\circ})_j &\xrightarrow{\sim} \overline{\mathbb{M}}_{\text{MOD}}^{\circ}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j; & ({}^{n,m}\overline{\mathbb{M}}_{\text{mod}}^{\circ})_j &\xrightarrow{\sim} \overline{\mathbb{M}}_{\text{mod}}^{\circ}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j \\ ({}^{n,m}\mathcal{F}_{\text{MOD}}^{\circ})_j &\xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circ}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j; & ({}^{n,m}\mathcal{F}_{\text{mod}}^{\circ})_j &\xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circ}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j \\ ({}^{n,m}\mathcal{F}_{\text{MOD}}^{\circ\mathbb{R}})_j &\xrightarrow{\sim} \mathcal{F}_{\text{MOD}}^{\circ\mathbb{R}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j; & ({}^{n,m}\mathcal{F}_{\text{mod}}^{\circ\mathbb{R}})_j &\xrightarrow{\sim} \mathcal{F}_{\text{mod}}^{\circ\mathbb{R}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_j \end{aligned}$$

which are compatible with the respective natural isomorphisms between “MOD”- and “mod”-subscripted versions [cf. Proposition 3.10, (i)]; here, the isomorphisms of the third line of the display induce isomorphisms of the **global realified Frobenioid** portions

$${}^{n,m}\mathcal{C}_{\text{LGP}}^{\text{lt}} \xrightarrow{\sim} \mathcal{C}_{\text{LGP}}^{\text{lt}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}); \quad {}^{n,m}\mathcal{C}_{\text{lgp}}^{\text{lt}} \xrightarrow{\sim} \mathcal{C}_{\text{lgp}}^{\text{lt}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})$$

of the  $\mathcal{F}^{\text{lt}}$ -prime-strips  ${}^{n,m}\mathfrak{F}_{\text{LGP}}^{\text{lt}}$ ,  $\mathfrak{F}^{\text{lt}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}}$ ,  ${}^{n,m}\mathfrak{F}_{\text{lgp}}^{\text{lt}}$ , and  $\mathfrak{F}^{\text{lt}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{lgp}}$  [cf. Propositions 3.7, (iii), (iv), (v); 3.10, (i)].

Moreover, as one varies  $m \in \mathbb{Z}$ , the isomorphisms of (b), as well as the isomorphisms of the first line in the first display of (c), are **mutually compatible**, relative to the **log-links** of the  $n$ -th column of the LGP-Gaussian log-theta-lattice under consideration, in the sense that the only portions of the domains of these isomorphisms that are related to one another via the **log-links** consist of **roots of unity** [multiplication by which corresponds, via the **log-link**, to an “**addition by zero**” indeterminacy, i.e., to **no indeterminacy!**] — cf. Proposition 3.5, (ii), (c); Proposition 3.10, (ii). This mutual compatibility of the isomorphisms of the

first line in the first display of (c) implies a corresponding **mutual compatibility** between the isomorphisms of the second and third lines in the first display of (c) that **involve the subscript “MOD”** [but **not** between the isomorphisms that involve the subscript “**mod**”! — cf. Proposition 3.10, (iii); Remark 3.10.2]. On the other hand, the isomorphisms of (a) are subject to a certain **“indeterminacy”** as follows:

- (Ind3) as one varies  $m \in \mathbb{Z}$ , the isomorphisms of (a) are **“upper semi-compatible”**, relative to the **log-links** of the  $n$ -th column of the LGP-Gaussian log-theta-lattice under consideration, in a sense that involves certain **natural inclusions “ $\subseteq$ ”** at  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{non}}$  and certain **natural surjections “ $\rightarrow$ ”** at  $v_{\mathbb{Q}} \in \mathbb{V}_{\mathbb{Q}}^{\text{arc}}$  — cf. Proposition 3.5, (ii), (a), (b), for more details.

Finally, as one varies  $m \in \mathbb{Z}$ , the isomorphisms of (a) are [precisely!] **compatible**, relative to the **log-links** of the  $n$ -th column of the LGP-Gaussian log-theta-lattice under consideration, with the respective **log-volumes** [cf. Proposition 3.9, (iv)].

(iii) ( **$\Theta_{\text{LGP}}^{\times\mu}$ -Link Compatibility**) The various Kummer isomorphisms of (ii) satisfy compatibility properties with the various **horizontal arrows** — i.e.,  $\Theta_{\text{LGP}}^{\times\mu}$ -links — of the LGP-Gaussian log-theta-lattice under consideration as follows:

- (a) The first Kummer isomorphism of the first display of (ii) induces — by applying the  $\mathbb{F}_l^{\times\pm}$ -**symmetry** of the  $\Theta^{\pm\text{ell}}$ NF-Hodge theater  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  — a **Kummer isomorphism**  ${}^{n,m}\mathfrak{F}_{\Delta}^{\times\mu} \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\times\mu}(n, \circ \mathcal{D}_{\Delta}^{\dagger})$  [cf. Theorem 1.5, (iii)]. Relative to this Kummer isomorphism, the full poly-isomorphism of  $\mathcal{F}^{\times\mu}$ -prime-strips

$$\mathfrak{F}_{\Delta}^{\times\mu}(n, \circ \mathcal{D}_{\Delta}^{\dagger}) \xrightarrow{\sim} \mathfrak{F}_{\Delta}^{\times\mu}(n+1, \circ \mathcal{D}_{\Delta}^{\dagger})$$

is **compatible** with the full poly-isomorphism of  $\mathcal{F}^{\times\mu}$ -prime-strips

$${}^{n,m}\mathfrak{F}_{\Delta}^{\times\mu} \xrightarrow{\sim} {}^{n+1,m}\mathfrak{F}_{\Delta}^{\times\mu}$$

induced [cf. Theorem 1.5, (ii)] by the **horizontal arrows** of the LGP-Gaussian log-theta-lattice under consideration [cf. Theorem 1.5, (iii)].

- (b) The  $\mathcal{F}^{\dagger}$ -prime-strips  ${}^{n,m}\mathfrak{F}_{\text{env}}^{\dagger}, \mathfrak{F}_{\text{env}}^{\dagger}(n, \circ \mathcal{D}_{>})$  [cf. Proposition 2.1, (ii)] that appear **implicitly** in the construction of the local portion of the  $\mathcal{F}^{\dagger}$ -prime-strips  ${}^{n,m}\mathfrak{F}_{\text{LGP}}^{\dagger}, \mathfrak{F}^{\dagger}(n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}}, {}^{n,m}\mathfrak{F}_{\text{Igp}}^{\dagger}, \mathfrak{F}^{\dagger}(n, \circ \mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{Igp}}$  [cf. (ii), (b), above; Proposition 3.4, (ii); Proposition 3.7, (iii), (iv); [IUTchII], Corollary 4.6, (iv)] admit **natural isomorphisms** of associated  $\mathcal{F}^{\times\mu}$ -prime-strips  ${}^{n,m}\mathfrak{F}_{\Delta}^{\times\mu} \xrightarrow{\sim} {}^{n,m}\mathfrak{F}_{\text{env}}^{\times\mu}, \mathfrak{F}_{\Delta}^{\times\mu}(n, \circ \mathcal{D}_{\Delta}^{\dagger}) \xrightarrow{\sim} \mathfrak{F}_{\text{env}}^{\times\mu}(n, \circ \mathcal{D}_{>})$  [cf. Proposition 2.1, (vi)]. Relative to these natural isomorphisms and to the Kummer isomorphism discussed in (a) above, the full poly-isomorphism of  $\mathcal{F}^{\times\mu}$ -prime-strips

$$\mathfrak{F}_{\text{env}}^{\times\mu}(n, \circ \mathcal{D}_{\Delta}^{\dagger}) \xrightarrow{\sim} \mathfrak{F}_{\text{env}}^{\times\mu}(n+1, \circ \mathcal{D}_{\Delta}^{\dagger})$$

is **compatible** with the full poly-isomorphism of  $\mathcal{F}^{\dagger \times \mu}$ -prime-strips

$${}^{n,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu} \xrightarrow{\sim} {}^{n+1,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu}$$

induced [cf. Theorem 1.5, (ii)] by the **horizontal arrows** of the LGP-Gaussian log-theta-lattice under consideration [cf. Corollary 2.3, (iii)].

- (c) Recall the data “ ${}^{n,\circ}\mathfrak{R}^{\text{bad}}$ ” [cf. Corollary 2.3, (c $\mathfrak{R}$ )] associated to the  $\mathcal{D}$ - $\Theta^{\pm\text{ell}}$  NF-Hodge theater  ${}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  — data which appears **implicitly** in the construction of the local portion of the  $\mathcal{F}^{\text{lt}}$ -prime-strips  ${}^{n,m}\mathfrak{F}_{\text{LGP}}^{\text{lt}}$ ,  $\mathfrak{F}^{\text{lt}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}}$ ,  ${}^{n,m}\mathfrak{F}_{\text{tgp}}^{\text{lt}}$ ,  $\mathfrak{F}^{\text{lt}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{tgp}}$  [cf. (ii), (b), above; Proposition 3.4, (ii); Proposition 3.7, (iii), (iv); [IUTchII], Corollary 4.6, (iv)]. This data that arises from  ${}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  is related to corresponding data that arises from the projective system of mono-theta environments associated to the tempered Frobenioids of the  $\Theta^{\pm\text{ell}}$  NF-Hodge theater  ${}^{n,m}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  via the **Kummer isomorphisms and poly-isomorphisms of projective systems of mono-theta environments** discussed in Proposition 2.1, (ii), (iii) [cf. also the second display of Theorem 2.2, (ii)] and Theorem 1.5, (iii) [cf. also (a), (b) above]. Relative to these Kummer isomorphisms and poly-isomorphisms of projective systems of mono-theta environments, the poly-isomorphism

$${}^{n,\circ}\mathfrak{R}^{\text{bad}} \xrightarrow{\sim} {}^{n+1,\circ}\mathfrak{R}^{\text{bad}}$$

induced by any **permutation symmetry** of the étale-picture [cf. the final portion of (i) above; Corollary 2.3, (ii); Remark 3.8.2]  ${}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} {}^{n+1,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  is **compatible** with the full poly-isomorphism of  $\mathcal{F}^{\dagger \times \mu}$ -prime-strips

$${}^{n,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu} \xrightarrow{\sim} {}^{n+1,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu}$$

induced [cf. Theorem 1.5, (ii)] by the **horizontal arrows** of the LGP-Gaussian log-theta-lattice under consideration [cf. Corollary 2.3, (iv)].

- (d) Relative to the Kummer isomorphisms of the first display of (ii) [cf. also (a), (b) above], the poly-isomorphisms

$$\left\{ \pi_1({}^{n,\circ}\mathcal{D}^{\otimes}) \curvearrowright \overline{\mathbb{M}}^{\otimes}({}^{n,\circ}\mathcal{D}^{\otimes}) \right\} \xrightarrow{\sim} \left\{ \pi_1({}^{n+1,\circ}\mathcal{D}^{\otimes}) \curvearrowright \overline{\mathbb{M}}^{\otimes}({}^{n+1,\circ}\mathcal{D}^{\otimes}) \right\}$$

$$\overline{\mathbb{M}}_{\text{orb}}^{\otimes}({}^{n,\circ}\mathcal{D}^{\otimes})_j \xrightarrow{\sim} \overline{\mathbb{M}}_{\text{orb}}^{\otimes}({}^{n+1,\circ}\mathcal{D}^{\otimes})_j$$

induced by any **permutation symmetry** of the étale-picture [cf. the final portion of (i) above; Corollary 2.3, (ii); Remark 3.8.2]  ${}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\sim} {}^{n+1,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}}$  are **compatible** [cf. the discussion of Remark 3.10.1, (ii), for more details] with the full poly-isomorphism of  $\mathcal{F}^{\dagger \times \mu}$ -prime-strips

$${}^{n,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu} \xrightarrow{\sim} {}^{n+1,m}\mathfrak{F}_{\Delta}^{\dagger \times \mu}$$

induced [cf. Theorem 1.5, (ii)] by the **horizontal arrows** of the LGP-Gaussian log-theta-lattice under consideration.

*Proof.* The various assertions of Theorem 3.11 follow immediately from the definitions and the references quoted in the statements of these assertions — cf. also the various related observations of Remarks 3.11.1, 3.11.2 below.  $\circ$

**Remark 3.11.1.** One way to summarize the content of Theorem 3.11 is as follows:

Theorem 3.11 gives an **algorithm** for describing, up to certain relatively **mild indeterminacies**, the **LGP-monoids** [cf. Fig. 3.1] — i.e., in essence, the **theta values**

$$\left\{ \underset{\equiv}{q^{j^2}} \right\}_{j=1, \dots, l^*}$$

— which are constructed relative to the **scheme/ring structure**, i.e., **“arithmetic holomorphic structure”**, associated to *one* vertical line [i.e., “ $(n, \circ)$ ” for *fixed* some  $n \in \mathbb{Z}$ ] in the LGP-Gaussian log-theta-lattice under consideration in terms of the *a priori* **alien** arithmetic holomorphic structure of *another* vertical line [i.e., “ $(n + 1, \circ)$ ”] in the LGP-Gaussian log-theta-lattice under consideration — cf., especially, the final portion of Theorem 3.11, (i), concerning **functoriality** and *compatibility* with the **permutation symmetries** of the **étale-picture**.

This point of view is consistent with the point of view of the discussion of Remark 1.5.4; [IUTchII], Remark 3.8.3, (iii).

**Remark 3.11.2.**

(i) In Theorem 3.11, (i), we do not apply the *formalism* or *language* developed in [IUTchII], §1, for discussing multiradiality. Nevertheless, the approach taken in Theorem 3.11, (i) — i.e., by regarding the collection of data (a), (b), (c) up to the indeterminacies given by (Ind1), (Ind2) — to constructing **“multiradial representations”** amounts, in essence, to a special case of the **tautological** approach to constructing multiradial environments discussed in [IUTchII], Example 1.9, (ii). That is to say, this tautological approach is applied to the **vertically coric** constructions of Proposition 3.5, (i); 3.10, (i), which, *a priori*, are *uniradial* in the sense that they depend, in an essential way, on the **arithmetic holomorphic structure** constituted by a *particular* vertical line — i.e., “ $(n, \circ)$ ” for *fixed* some  $n \in \mathbb{Z}$  — in the LGP-Gaussian log-theta-lattice under consideration.

(ii) One important underlying aspect of the *tautological approach to multiradiality* discussed in (i) is the treatment of the various **labels** that occur in the **multiplicative** and **additive combinatorial Teichmüller theory** associated to the  $\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater  ${}^{n, \circ}\mathcal{HT}^{\mathcal{D}\text{-}\Theta^{\pm\text{ell}}\text{NF}}$  under consideration [cf. the theory of [IUTchI], §4, §6]. The various transitions between types of labels is illustrated in Fig. 3.3 below. Here, we recall that:

- (a) the passage from the  $\mathbb{F}_l^{\times\pm}$ -**symmetry** to **labels**  $\in \mathbb{F}_l$  forms the content of the associated  $\mathcal{D}\text{-}\Theta^{\pm\text{ell}}$ -*Hodge theater* [cf. [IUTchI], Remark 6.6.1];

- (b) the passage from **labels**  $\in \mathbb{F}_l$  to **labels**  $\in |\mathbb{F}_l|$  forms the content of the *functorial algorithm* of [IUTchI], Proposition 6.7;
- (c) the passage from **labels**  $\in |\mathbb{F}_l|$  to  $\pm$ -**processions** forms the content of [IUTchI], Proposition 6.9, (ii);
- (d) the passage from the  $\mathbb{F}_l^*$ -**symmetry** to **labels**  $\in \mathbb{F}_l^*$  forms the content of the associated  $\mathcal{D}$ - $\Theta NF$ -*Hodge theater* [cf. [IUTchI], Remark 4.7.2, (i)];
- (e) the passage from **labels**  $\in \mathbb{F}_l^*$  to  $*$ -**processions** forms the content of [IUTchI], Proposition 4.11, (ii);
- (f) the compatibility between  $*$ -**processions** and  $\pm$ -**processions**, relative to the natural inclusion of **labels**  $\mathbb{F}_l^* \hookrightarrow |\mathbb{F}_l|$ , forms the content of [IUTchI], Proposition 6.9, (iii).

Here, we observe in passing that, in order to perform these various *transitions*, it is absolutely necessary to work with *all of the labels in*  $\mathbb{F}_l$  or  $|\mathbb{F}_l|$ , i.e., one does not have the option of “*arbitrarily omitting certain of the labels*” [cf. the discussion of [IUTchII], Remark 2.6.3; [IUTchII], Remark 3.5.2]. Also, in this context, it is important to note that there is a fundamental difference between the **labels**  $\in \mathbb{F}_l, |\mathbb{F}_l|, \mathbb{F}_l^*$  — which are essentially **arithmetic holomorphic** in the sense that they depend, in an essential way, on the various local and global *arithmetic fundamental groups* involved — and the **index sets of the mono-analytic  $\pm$ -processions** that appear in the multiradial representation of Theorem 3.11, (i). Indeed, these index sets are just “*naked sets*” which are determined, up to isomorphism, by their *cardinality*. In particular,

*the construction of these index sets is independent of the various arithmetic holomorphic structures involved.*

Indeed, it is precisely this property of these index sets that renders them suitable for use in the construction of the *multiradial representations* of Theorem 3.11, (i). As discussed in [IUTchI], Proposition 6.9, (i), for  $j \in \{0, \dots, l^*\}$ , there are precisely  $j + 1$  *possibilities* for the “*element labeled*  $j$ ” in the index set of cardinality  $j + 1$ ; this leads to a total of  $(l^* + 1)! = l^{\pm!}$  *possibilities* for the “*label identification*” of elements of index sets of capsules appearing in the mono-analytic  $\pm$ -processions of Theorem 3.11, (i). Finally, in this context, it is of interest to recall that the “*rougher approach to symmetrization*” that arises when one works with *mono-analytic processions* is [“*downward*”] *compatible* with the *finer* arithmetically holomorphic approach to symmetrization that arises from the  $\mathbb{F}_l^{\times \pm}$ -**symmetry** [cf. [IUTchII], Remark 3.5.3; [IUTchII], Remark 4.5.2, (ii); [IUTchII], Remark 4.5.3, (ii)].

(iii) Observe that the “**Kummer isomorphism of global realified Frobenioids**” that appears in the theory of [IUTchII], §4 — i.e., more precisely, the various versions of the isomorphism of Frobenioids “ $\mathbb{F}_l^{\times \pm} \xrightarrow{\sim} \mathcal{D}^{\text{tr}}(\mathbb{F}_l^{\times \pm})$ ” discussed in [IUTchII], Corollary 4.6, (ii), (v) — is constructed by considering isomorphisms between **local value groups** obtained by forming the **quotient** of the multiplicative groups associated to the various local fields that appear by the subgroups of



**local units** [cf. [IUTchII], Propositions 4.2, (ii); 4.4, (ii)]. In particular, such “Kummer isomorphisms” *fail* to give rise to a “**log-Kummer correspondence**”, i.e., they fail to satisfy **mutual compatibility** properties of the sort discussed in the final portion of Theorem 3.11, (ii). Indeed, as discussed in Remark 1.2.3, (i) [cf. also [IUTchII], Remark 1.12.2, (iv)], at  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$ , the operation of forming a *multiplicative* quotient by *local units* corresponds, on the opposite side of the **log-link**, to forming an *additive* quotient by the submodule obtained as the  $p_{\underline{v}}$ -*adic logarithm* of these *local units*. This is precisely why, in the context of Theorem 3.11, (ii), we wish to work with the global non-realified/realified Frobenioids “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ”, “ $\mathcal{F}_{\text{MOD}}^{\circledast\mathbb{R}}$ ” that arise from copies of “ $F_{\text{mod}}$ ” which satisfy a “**log-Kummer correspondence**”, as described in the final portion of Theorem 3.11, (ii) [cf. the discussion of Remark 3.10.2]. On the other hand, the *pathologies/indeterminacies* that arise from working with global arithmetic line bundles by means of various *local data* at  $\underline{v} \in \underline{\mathbb{V}}$  in the context of the **log-link** are *formalized* via the theory of the global Frobenioids “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ”, together with the “**upper semi-compatibility**” of local units discussed in the final portion of Theorem 3.11, (ii) [cf. also the discussion of Remark 3.10.2].




Fig. 3.3: Transitions from symmetries to labels to processions in a  $\Theta^{\pm\text{ell}}\text{NF}$ -Hodge theater

(iv) In the context of the discussion of *global realified Frobenioids* given in (iii), we observe that, in the case of the global realified Frobenioids [constructed by means of “ $\mathcal{F}_{\text{MOD}}^{\circledast\mathbb{R}}$ ”!] that appear in the  $\mathcal{F}^{\text{ll}}$ -prime-strips  ${}^{n,m}\mathfrak{F}_{\text{LGP}}^{\text{ll}}, \mathfrak{F}^{\text{ll}}({}^{n,\circ}\mathcal{HT}^{\mathcal{D}-\Theta^{\pm\text{ell}}\text{NF}})_{\text{LGP}}$  [cf. Theorem 3.11, (ii), (c)], the various **localization functors** that appear [i.e., the various “ $\dagger\rho_{\underline{v}}$ ” of [IUTchI], Definition 5.2, (iv); cf. also the isomorphisms of the second display of [IUTchII], Corollary 4.6, (v)] may be **reconstructed**, in the spirit of the discussion of Remark 3.9.2, “by considering the *effect of multiplication* by elements of the [non-realified] *global monoids* under consideration on the *log-volumes* of the various local mono-analytic tensor packets that appear”. [We leave the routine details to the reader.] This reconstructibility, together with the *mutual incompatibilities* observed in (iii) above that arise when one attempts to work *simultaneously* with *log-shells* and with the *splitting monoids* of the  $\mathcal{F}^{\text{ll}}$ -prime-strip

$n, m \mathfrak{F}_{\text{LGP}}^{\text{lt}}$  at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}}$ , are the primary reasons for our omission of the the splitting monoids at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}}$  from the statement of Theorem 3.11 [cf. Theorem 3.11, (i), (b); Theorem 3.11, (ii), (b); Theorem 3.11, (iii), (c), in the case of  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$ ].

**Remark 3.11.3.** Before proceeding, we pause to discuss the relationship between the **log-Kummer correspondence** of Theorem 3.11, (ii), and the  $\Theta_{\text{LGP}}^{\times\mu}$ -**link compatibility** of Theorem 3.11, (iii).

(i) First, we recall [cf. Remarks 1.4.1, (i); 3.8.2] that the various squares that appear in the [LGP-Gaussian] *log-theta-lattice* are *far from being [1-]commutative!* On the other hand, the **bi-coricity** of  $\mathcal{F}^{+\times\mu}$ -*prime-strips* and *mono-analytic log-shells* discussed in Theorem 1.5, (iii), (iv), may be interpreted as the statement that

the various squares that appear in the [LGP-Gaussian] *log-theta-lattice* **are** in fact **[1-]commutative** with respect to [the portion of the data associated to each “•” in the log-theta-lattice that is constituted by] these bi-coric  $\mathcal{F}^{+\times\mu}$ -*prime-strips* and *mono-analytic log-shells*.

(ii) Next, let us observe that in order to relate both the *unit* and *value group* portions of the domain and codomain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link corresponding to *adjacent vertical lines* — i.e.,  $(n-1, *)$  and  $(n, *)$  — of the [LGP-Gaussian] log-theta-lattice to one another,

it is necessary to relate these **unit** and **value group** portions to one another by means of a **single**  $\Theta_{\text{LGP}}^{\times\mu}$ -**link**, i.e., from  $(n-1, m)$  to  $(n, m)$ .

That is to say, from the point of view of constructing the various LGP-*monoids* that appear in the multiradial representation of Theorem 3.11, (i), one is tempted to work with correspondences between *value groups* on adjacent vertical lines that lie in a **vertically once-shifted** position — i.e., say, at  $(n-1, m)$  and  $(n, m)$  — relative to the correspondence between *unit groups* on adjacent vertical lines, i.e., say, at  $(n-1, m-1)$  and  $(n, m-1)$ . On the other hand, such an approach *fails*, at least from an *a priori* point of view, precisely on account of the *noncommutativity* discussed in (i). Finally, we observe that in order to relate both unit and value groups by means of a *single*  $\Theta_{\text{LGP}}^{\times\mu}$ -*link*,

it is necessary to avail oneself of the  $\Theta_{\text{LGP}}^{\times\mu}$ -*link compatibility* properties discussed in Theorem 3.11, (iii) — i.e., of the theory of §2 and [IUTchI], Example 5.1, (v) — so as to **insulate** the **cyclotomes** that appear in the construction of the **étale theta function** via **mono-theta environments** and the construction of **number fields** via global Kummer theory from the **Aut $_{\mathcal{F}^{+\times\mu}}(-)$ -indeterminacies** that act on the  $\mathcal{F}^{+\times\mu}$ -*prime-strips* involved as a result of the application of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link

— cf. the discussion of Remarks 2.2.1; 3.10.1, (ii).

(iii) As discussed in (ii) above, a “vertically once-shifted” approach to relating units on adjacent vertical lines *fails* on account of the *noncommutativity* discussed in

(i). Thus, one natural approach to treating the units in a “vertically once-shifted” fashion — which, we recall, is necessary in order to relate the LGP-*monoids* on adjacent vertical lines to one another! — is to apply the **bi-coricity of mono-analytic log-shells** discussed in (i). On the other hand, to take this approach means that one must work in a *framework* that allows one to *relate* [cf. the discussion of Remark 1.5.4, (i)] the “Frobenius-like” structure constituted by the *Frobenioid-theoretic* units [i.e., which occur in the domain and codomain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link] to corresponding étale-like structures **simultaneously** via both

- (a) the usual **Kummer isomorphisms** — i.e., so as to be compatible with the application of the compatibility properties of Theorem 3.11, (iii), as discussed in (ii) — and
- (b) the **composite** of the usual Kummer isomorphisms with [a single iterate of] the **log-link** — i.e., so as to be compatible with the bi-coric treatment of mono-analytic log-shells [as well as the closely related construction of LGP-monoids] proposed above.

Such a framework may only be realized if one relates Frobenius-like structures to étale-like structures in a fashion that is **invariant** with respect to pre-composition of various iterates of the **log-link** [cf. the final portions of Propositions 3.5, (ii); 3.10, (ii)]. This is precisely what is achieved by the **log-Kummer correspondences** of the final portion of Theorem 3.11, (ii).

(iv) The discussion of (i), (ii), (iii) above may be summarized as follows: The **log-Kummer correspondences** of the final portion of Theorem 3.11, (ii), allow one to

- (a) relate both the **unit** and the **value group** portions of the domain and codomain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link corresponding to *adjacent vertical lines* of the [LGP-Gaussian] log-theta-lattice to one another, in a fashion that
- (b) **insulates** the **cyclotomes/Kummer theory** surrounding the **étale theta function** and **number fields** involved from the  $\text{Aut}_{\mathcal{F}^+ \times \mu}(-)$ -**indeterminacies** that act on  $\mathcal{F}^+ \times \mu$ -prime-strips involved as a result of the application of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link [cf. Theorem 3.11, (iii)], and, moreover,
- (c) is **compatible** with the **bi-coricity** of the **mono-analytic log-shells** [cf. Theorem 1.5, (iv)], hence also with the operation of relating the **LGP-monoids** that appear in the multiradial representation of Theorem 3.11, (i), corresponding to *adjacent vertical lines* of the [LGP-Gaussian] log-theta-lattice to one another.

These observations will play a *key role* in the proof of Corollary 3.12 below.

The following result may be thought of as a relatively *concrete consequence* of the somewhat abstract content of Theorem 3.11.

**Corollary 3.12.** (**Log-volume Estimates for  $\Theta$ -Pilot Objects**) *Suppose that we are in the situation of Theorem 3.11. Write*

$$- |\log(\underline{\Theta})| \in \mathbb{R} \cup \{+\infty\}$$

for the **procession-normalized mono-analytic log-volume** [i.e., where the average is taken over  $j \in \mathbb{F}_l^*$  — cf. Remark 3.1.1, (ii); Proposition 3.9, (i), (ii); Theorem 3.11, (i), (a)] of the **union of the possible images of a  $\Theta$ -pilot object** [cf. Definition 3.8, (i)], relative to the relevant **Kummer isomorphisms** [cf. Theorem 3.11, (ii)], in the **multiradial representation** of Theorem 3.11, (i), which we regard as **subject to the indeterminacies** (Ind1), (Ind2), (Ind3) described in Theorem 3.11, (i), (ii). Write

$$- |\log(\underline{q})| \in \mathbb{R}$$

for the **procession-normalized mono-analytic log-volume** of the image of a  **$q$ -pilot object** [cf. Definition 3.8, (i)], relative to the relevant **Kummer isomorphisms** [cf. Theorem 3.11, (ii)], in the **multiradial representation** of Theorem 3.11, (i), which we do **not** regard as subject to the indeterminacies (Ind1), (Ind2), (Ind3) described in Theorem 3.11, (i), (ii). Here, we recall the definition of the symbol “ $\triangle$ ” as the result of identifying the labels

$$“0” \text{ and } “\langle \mathbb{F}_l^* \rangle”$$

[cf. [IUTchII], Corollary 4.10, (i)]. In particular,  $|\log(\underline{q})| > 0$  is easily computed in terms of the various  **$q$ -parameters** of the elliptic curve  $E_F$  [cf. [IUTchI], Definition 3.1, (b)] at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}} (\neq \emptyset)$ . Then it holds that

$$C_\Theta \geq -1$$

for any real number  $C_\Theta \in \mathbb{R}$  such that

$$- |\log(\underline{\Theta})| \leq C_\Theta \cdot |\log(\underline{q})| \quad (*_{C_\Theta})$$

[i.e.,  $- |\log(\underline{\Theta})| \in \mathbb{R} \subseteq \mathbb{R} \cup \{+\infty\}$  and satisfies the inequality  $(*_{C_\Theta})$ ].

*Proof.* Suppose that we are in the situation of Theorem 3.11. We begin by reviewing precisely what is achieved by the various portions of Theorem 3.11 and, indeed, by the theory developed thus far in the present series of papers. This review leads naturally to an interpretation of the theory that gives rise to the *inequality* asserted in the statement of Corollary 3.12. For ease of reference, we divide our discussion into *steps*, as follows.

(i) In the following discussion, we concentrate on a *single arrow* — i.e., a *single  $\Theta_{\text{LGP}}^{\times\mu}$ -link*

$${}_{0,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}} \xrightarrow{\Theta_{\text{LGP}}^{\times\mu}} {}_{1,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$$

— of the [LGP-Gaussian] log-theta-lattice under consideration. This arrow consists of the *full poly-isomorphism* of  $\mathcal{F}^{\text{H}\blacktriangleright\times\mu}$ -prime-strips

$${}_{0,0}\mathfrak{F}_{\text{LGP}}^{\text{H}\blacktriangleright\times\mu} \xrightarrow{\sim} {}_{1,0}\mathfrak{F}_{\triangle}^{\text{H}\blacktriangleright\times\mu}$$

[cf. Definition 3.8, (ii)]. This poly-isomorphism may be thought of as consisting of a “**unit portion**” constituted by the associated [full] poly-isomorphism of  $\mathcal{F}^{\text{+}\times\mu}$ -prime-strips

$${}_{0,0}\mathfrak{F}_{\text{LGP}}^{\text{+}\times\mu} \xrightarrow{\sim} {}_{1,0}\mathfrak{F}_{\triangle}^{\text{+}\times\mu}$$

and a “**value group portion**” constituted by the associated [full] poly-isomorphism of  $\mathcal{F}^{\text{tr}}\blacktriangleright$ -prime-strips

$${}^{0,0}\mathfrak{F}_{\text{LGP}}^{\text{tr}} \xrightarrow{\sim} {}^{1,0}\mathfrak{F}_{\Delta}^{\text{tr}}$$

[cf. Definition 2.4, (iii)]. This value group portion of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link maps  $\Theta$ -*pilot objects* of  ${}^{0,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  to  $q$ -*pilot objects* of  ${}^{1,0}\mathcal{HT}^{\Theta^{\pm\text{ell}}\text{NF}}$  [cf. Remark 3.8.1].

(ii) Whereas the units of the Frobenioids that appear in the  $\mathcal{F}^{\text{tr}}\times\mu$ -prime-strip  ${}^{0,0}\mathfrak{F}_{\text{LGP}}^{\text{tr}\times\mu}$  are subject to  $\text{Aut}_{\mathcal{F}^{\text{tr}}\times\mu}(-)$ -indeterminacies [i.e., “(Ind2)” — cf. Theorem 3.11, (iii), (a), (b)], the **cyclotomes** that appear in the Kummer theory related to the **étale theta function** and the **number fields** involved, i.e., which give rise to the “value group portion”  ${}^{0,0}\mathfrak{F}_{\text{LGP}}^{\text{tr}}$ , are **insulated** from these  $\text{Aut}_{\mathcal{F}^{\text{tr}}\times\mu}(-)$ -**indeterminacies** — cf. Theorem 3.11, (iii), (c), (d); the discussion of Remark 3.11.3, (iv); Fig. 3.4 below. Here, we recall that in the case of the étale theta function, this follows from the theory of §2, i.e., in essence, from the **cyclotomic rigidity of mono-theta environments**, as discussed in [EtTh]. On the other hand, in the case of number fields, this follows, in essence, from the elementary algebraic number theory fact applied in the proof of Proposition 3.10 to the effect that the nonzero **global integers** of a number field are necessarily **roots of unity**, together with the algorithms discussed in [IUTchI], Example 5.1, (v).

	<i><math>\Theta</math>-related objects</i>	<i>NF-related objects</i>
<p><b>require</b> <i>mono-analytic</i> <i>containers,</i> <b>Kummer-incompatible</b></p>	<p>local LGP-monoids  [cf. Proposition 3.4, (ii)]</p>	<p><math>F_{\text{mod}}</math>  [cf. Proposition 3.7, (i)]</p>
<p><b>independent of</b> <i>mono-analytic</i> <i>containers,</i> <b>Kummer-compatible</b></p>	<p>étale theta function, mono-theta environments [cf. Corollary 2.3]</p>	<p><math>\overline{F}</math> [cf. [IUTchI], Example 5.1, (v); [IUTchII], Corollary 4.8, (i)]</p>

Fig. 3.4: Relationship of theta- and number field-related objects to mono-analytic containers

(iii) In the following discussion, it will be of crucial importance to relate **simultaneously** both the **unit** and the **value group** portions of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link(s) involved on the *0-column* [i.e., the vertical line indexed by 0] of the log-theta-lattice under consideration to the corresponding unit and value group portions on the *1-column* [i.e., the vertical line indexed by 1] of the log-theta-lattice under consideration. On the other hand, if one attempts to relate the *unit* portions via

one  $\Theta_{\text{LGP}}^{\times\mu}$ -link [say, from  $(0, m)$  to  $(1, m)$ ] and the *value group* portions via another  $\Theta_{\text{LGP}}^{\times\mu}$ -link [say, from  $(0, m')$  to  $(1, m')$ , for  $m' \neq m$ ], then the **non-commutativity** of the log-theta-lattice renders it practically impossible to obtain conclusions that require one to relate both the unit and the value group portions *simultaneously* [cf. the discussion of Remark 3.11.3, (i), (ii)]. This is precisely why we concentrate on a **single  $\Theta_{\text{LGP}}^{\times\mu}$ -link** [cf. (i)].

(iv) The issue discussed in (iii) is relevant in the context of the present discussion for the following reason. Ultimately, we wish to apply the **bi-coricity** of the **units** [cf. Theorem 1.5, (iii), (iv)] in order to compute the 0-*column*  $\Theta$ -*pilot object* in terms of the arithmetic holomorphic structure of the 1-*column*. In order to do this, one must work with *units* that are **vertically once-shifted** [i.e., lie at  $(n, m - 1)$ ] relative to the value group structures involved [i.e., which lie at  $(n, m)$ ] — cf. the discussion of Remark 3.11.3, (ii). The solution to the problem of simultaneous accommodating these apparently *contradictory requirements* — i.e., “vertical shift” vs. “impossibility of vertical shift” [cf. (iii)] — is given precisely by working, on the 0-column, with structures that are **invariant** with respect to **vertical shifts** [i.e., “ $(0, m) \mapsto (0, m + 1)$ ”] of the log-theta-lattice [cf. the discussion surrounding Remark 1.2.2, (iii), (a)] such as **vertically coric structures** [i.e., indexed by “ $(n, \circ)$ ”] that are related to the “*Frobenius-like*” structures which are *not* vertically coric by means of the **log-Kummer correspondences** of Theorem 3.11, (ii). Here, we note that this “solution” may be implemented only at the cost of admitting the “*indeterminacy*” constituted by the **upper semi-compatibility** of (Ind3).

(v) Thus, we begin our computation of the 0-*column*  $\Theta$ -*pilot object* in terms of the arithmetic holomorphic structure of the 1-*column* by relating the units on the 0- and 1-columns by means of the **unit portion**

$${}_{0,0}\mathfrak{F}_{\text{LGP}}^{\dagger \times \mu} \xrightarrow{\simeq} {}_{1,0}\mathfrak{F}_{\Delta}^{\dagger \times \mu}$$

of the  $\Theta_{\text{LGP}}^{\times\mu}$ -link from  $(0, 0)$  to  $(1, 0)$  [cf. (i)] and then applying the **bi-coricity** of the **units** of Theorem 1.5, (iii), (iv). In particular, the **mono-analytic log-shell** interpretation of this bi-coricity given in Theorem 1.5, (iv), will be applied to regard these mono-analytic log-shells as “**multiradial mono-analytic containers**” [cf. the discussion of Remark 1.5.2, (i), (ii), (iii)] for the various [local and global] value group structures that constitute the  $\Theta$ -pilot object on the 0-column — cf. Fig. 3.4 above. [Here, we observe that the parallel treatment of “*theta-related*” and “*number field-related*” objects is reminiscent of the discussion of [IUTchII], Remark 4.11.2, (iv).] That is to say, we will relate the various Frobenioid-theoretic [i.e., “Frobenius-like” — cf. Remark 1.5.4, (i)]

- *local units* at  $\underline{v} \in \mathbb{V}$ ,
- *splitting monoids* at  $\underline{v} \in \mathbb{V}^{\text{bad}}$ , and
- *global Frobenioids*

indexed by  $(0, m)$ , for  $m \in \mathbb{Z}$ , to the *vertically coric* [i.e., indexed by “ $(0, \circ)$ ”] versions of these bi-coric mono-analytic containers by means of the **log-Kummer**



**correspondences** of Theorem 3.11, (ii), (a), (b), (c) — i.e., by *varying* the “*Kummer input index*”  $(0, m)$  along the 0-column.

(vi) In the context of (v), it is useful to recall that the **log-Kummer** correspondences of Theorem 3.11, (ii), (b), (c), are obtained precisely as a consequence of the **splittings**, up to roots of unity, of the relevant monoids into **unit** and **value group** portions constructed by applying the **constant multiple rigidity** established in [EtTh] [in the case of Theorem 3.11, (ii), (b)] and the elementary algebraic number theory characterization of the **global integers** of a number field that was applied in the proof of Proposition 3.10 [in the case of Theorem 3.11, (ii), (c)]. Moreover, we recall that the Kummer theory surrounding the local LGP-monoids of Proposition 3.4, (ii), depends, in an essential way, on the theory of [IUTchII], §3 [cf., especially, [IUTchII], Corollaries 3.5, 3.6], which, in turn, depends, in an essential way, on the Kummer theory surrounding **mono-theta environments** established in [EtTh]. Thus, for instance, we recall that the **discrete rigidity** established in [EtTh] is applied so as to avoid working, in the tempered Frobenioids that occur, with “ **$\widehat{\mathbb{Z}}$ -divisors/line bundles**” [i.e., “ $\widehat{\mathbb{Z}}$ -completions” of  $\mathbb{Z}$ -modules of divisors/line bundles], which are *fundamentally incompatible* with conventional notions of divisors/line bundles, hence, in particular, with *mono-theta-theoretic cyclotomic rigidity* [cf. Remark 2.1.1, (v)]. Also, we recall that “**isomorphism class compatibility**” — i.e., in the terminology of [EtTh], “*compatibility with the topology of the tempered fundamental group*” [cf. the discussion at the beginning of Remark 2.1.1] — allows one to apply the Kummer theory of mono-theta environments [i.e., the theory of [EtTh]] relative to the **ring-theoretic basepoints** that occur on either side of the **log-link** [cf. Remark 2.1.1, (ii); [IUTchII], Remark 3.6.4, (i)], for instance, in the context of the **log-Kummer correspondence for the splitting monoids of local LGP-monoids**, whose construction depends, in an essential way [cf. the theory of [IUTchII], §3, especially, [IUTchII], Corollaries 3.5, 3.6], on the **conjugate synchronization** arising from the  $\mathbb{F}_l^{\times\pm}$ -**symmetry**. That is to say,

it is precisely by establishing this conjugate synchronization arising from the  $\mathbb{F}_l^{\times\pm}$ -symmetry relative to *these basepoints* that occur on either side of the **log-link** that one is able to conclude the crucial **compatibility of this conjugate synchronization with the log-link** discussed in Remark 1.3.2.

A similar observation may be made concerning the *MLF-Galois pair* approach to the *cyclotomic rigidity isomorphism* that is applied at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}} \cap \underline{\mathbb{V}}^{\text{non}}$  [cf. [IUTchII], Corollary 1.11, (a); [IUTchII], Remark 1.11.1, (i), (a); [IUTchII], Proposition 4.2, (i); [AbsTopIII], Proposition 3.2, (iv)], which amounts, in essence, to

computations involving the Galois cohomology groups of *various subquotients* — *such as torsion subgroups [i.e., roots of unity] and associated value groups* — of the [multiplicative] module of nonzero elements of an algebraic closure of the mixed characteristic local field involved

[cf. the proof of [AbsAnab], Proposition 1.2.1, (vii)] — i.e., algorithms that are *compatible with the topology of the profinite groups involved*, in the sense that they do not require one to pass to *Kummer towers* [cf. the discussion of [IUTchII],

Remark 3.6.4, (i)], which have are *fundamentally incompatible* with the *ring structure* of the fields involved. Here, we note in passing that the corresponding property for  $\underline{v} \in \underline{\mathbb{V}}^{\text{arc}}$  [cf. [IUTchII], Proposition 4.4, (i)] holds for the simple reason that the complex archimedean local fields that occur at such  $\underline{v}$  are already *algebraically closed* [so that there is no issue of possibly having to pass to Kummer towers]! On the other hand, the approaches to cyclotomic rigidity just discussed for  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  and  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}} \cap \underline{\mathbb{V}}^{\text{non}}$  differ quite *fundamentally* from the approach to cyclotomic rigidity taken in the case of [global] number fields in the algorithms described in [IUTchI], Example 5.1, (v), which depend, in an essential way, on the property

$$\mathbb{Q}_{>0} \cap \widehat{\mathbb{Z}}^\times = \{1\}$$

— i.e., which is **fundamentally incompatible** with the **topology** of the profinite groups involved in the sense that it clearly cannot be obtained as some sort of limit of corresponding properties of  $(\mathbb{Z}/N\mathbb{Z})^\times$ ! Nevertheless, with regard to uni-/multi-radiality issues, this approach to cyclotomic rigidity in the case of the number fields resembles the theory of mono-theta-theoretic cyclotomic rigidity at  $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$  in that it admits a *natural multiradial formulation* [cf. Theorem 3.11, (iii), (c), (d); the discussion of Remark 3.11.3], in sharp contrast to the essentially **uniradial** nature of the approach to cyclotomic rigidity via MLF-Galois pairs at  $\underline{v} \in \underline{\mathbb{V}}^{\text{good}} \cap \underline{\mathbb{V}}^{\text{non}}$  [cf. the discussion of [IUTchII], Remark 1.11.3]. These observations are summarized in Fig. 3.5 below. Finally, we recall that [one verifies immediately that] the various approaches to cyclotomic rigidity just discussed are *mutually compatible* in the sense that they yield the same cyclotomic rigidity isomorphism in any setting in which more than one of these approaches may be applied.

<u>Approach to cyclotomic rigidity</u>	<u>Uni-/multi-radiality</u>	<u>Compatibility with ring structure, log-link in the context of the <math>\mathbb{F}_l^{\times \pm}</math>-symmetry</u>
<i>mono-theta environments</i>	<b>multiradial</b>	<b>compatible</b>
<i>MLF-Galois pairs, via Brauer groups</i>	<b>uniradial</b>	<b>compatible</b>
<i>number fields, via <math>\mathbb{Q}_{&gt;0} \cap \widehat{\mathbb{Z}}^\times = \{1\}</math></i>	<b>multiradial</b> ( $\Leftarrow \exists$ splitting)	<b>incompatible</b>

Fig. 3.5: Three approaches to cyclotomic rigidity

(vii) In the context of the discussion in the final portion of (vi), it is of interest to recall that the constructions underlying the crucial **bi-coricity** theory of

Theorem 1.5, (iii), (iv), depend, in an essential way, on the **conjugate synchronization** arising from the  $\mathbb{F}_l^{\times\pm}$ -**symmetry**, which allows one to relate the local monoids and Galois groups at distinct labels  $\in |\mathbb{F}_l|$  to one another in a fashion that is *simultaneously compatible* both with

- the **vertically coric** structures and **Kummer theory** that give rise to the **log-Kummer correspondences** of Theorem 3.11, (ii),

and with

- the property of **distinguishing** [i.e., not identifying] data indexed by **distinct labels**  $\in |\mathbb{F}_l|$

— cf. the discussion of Remark 1.5.1, (i), (ii). Since, moreover, this crucial conjugate synchronization is *fundamentally incompatible* with the  $\mathbb{F}_l^*$ -**symmetry**, it is necessary to work with these two symmetries *separately*, as was done in [IUTchI], §4, §5, §6 [cf. [IUTchII], Remark 4.7.6]. Here, it is useful to recall that the  $\mathbb{F}_l^*$ -symmetry also plays a crucial role, in that it allows one to “*descend to  $F_{\text{mod}}$ ” at the level of **absolute Galois groups** [cf. [IUTchII], Remark 4.7.6]. On the other hand, *both* the  $\mathbb{F}_l^{\times\pm}$ - and  $\mathbb{F}_l^*$ -symmetries share the property of being compatible with the **vertical coricity** and relevant **Kummer isomorphisms** of the 0-column — cf. the **log-Kummer correspondences** of Theorem 3.11, (ii), (b) [in the case of the  $\mathbb{F}_l^{\times\pm}$ -symmetry], (c) [in the case of the  $\mathbb{F}_l^*$ -symmetry]. Indeed, the vertically coric versions of both the  $\mathbb{F}_l^{\times\pm}$ - and the  $\mathbb{F}_l^*$ -symmetries depend, in an essential way, on the **arithmetic holomorphic structure** of the 0-column, hence give rise to **multiradial** structures via the **tautological** approach to constructing such structures discussed in Remark 3.11.2, (i), (ii).*

(viii) In the context of (vii), it is useful to recall that in order to construct the  $\mathbb{F}_l^{\times\pm}$ -symmetry, it is necessary to make use of **global  $\pm$ -synchronizations** of various local  $\pm$ -indeterminacies. Since the local tempered fundamental groups at  $\underline{v} \in \mathbb{V}^{\text{bad}}$  do not extend to a “global tempered fundamental group”, these global  $\pm$ -synchronizations give rise to **profinite conjugacy indeterminacies** in the vertically coric construction of the LGP-monoids [i.e., the theta values at torsion points] given in [IUTchII], §2, which are *resolved* by applying the theory of [IUTchI], §2 — cf. the discussion of [IUTchI], Remark 6.12.4, (iii); [IUTchII], Remark 4.5.3, (iii); [IUTchII], Remark 4.11.2, (iii).

(ix) In the context of (vii), it is also useful to recall the important role played, in the theory of the present series of papers, by the various “copies of  $F_{\text{mod}}$ ”, i.e., more concretely, in the form of the various copies of the **global Frobenioids** “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ”, “ $\mathcal{F}_{\text{mod}}^{\circledast}$ ” and their realifications. That is to say, the **ring structure** of the global field  $F_{\text{mod}}$  allows one to bridge the gap — i.e., furnishes a **translation apparatus** — between the **multiplicative** structures constituted by the global realified Frobenioids related via the  $\Theta_{\text{LGP}}^{\times\mu}$ -link and the **additive** representations of these global Frobenioids that arise from the “mono-analytic containers” furnished by the *mono-analytic log-shells* [cf. (v)]. Here, the **precise compatibility** of the ingredients for “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ” with the **log-Kummer** correspondence renders “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ” better suited to describing the relation to the  $\Theta_{\text{LGP}}^{\times\mu}$ -link [cf. Remark 3.10.2, (ii)].

On the other hand, the local portion of “ $\mathcal{F}_{\text{mod}}^{\odot}$ ” — i.e., which is subject to “**upper semi-compatibility**” [cf. (Ind3)], hence only “*approximately compatible*” with the **log-Kummer** correspondence — renders it better suited to **explicit estimates** of global arithmetic degrees, by means of **log-volumes** [cf. Remark 3.10.2, (iii)].

(x) Thus, one may summarize the discussion thus far as follows. The theory of “**Kummer-detachment**” — cf. Remarks 1.5.4, (i); 2.1.1; 3.10.2, (ii), (iii) — furnished by Theorem 3.11, (ii), (iii), allows one to relate the **Frobenoid-theoretic** [i.e., “Frobenius-like”] structures that appear in the domain [i.e., at  $(0, 0)$ ] of the  $\Theta_{\text{LGP}}^{\times\mu}$ -**link** [cf. (i)] to the **multiradial representation** described in Theorem 3.11, (i), (a), (b), (c), but only at the cost of introducing the **indeterminacies**

- (Ind1) — which may be thought of as arising from the requirement of *compatibility* with the **permutation symmetries** of the **étale-picture** [cf. Theorem 3.11, (i)];
- (Ind2) — which may be thought of as arising from the requirement of *compatibility* with the  $\text{Aut}_{\mathcal{F}^{\times\mu}}(-)$ -**indeterminacies** that act on the domain/codomain of the  $\Theta_{\text{LGP}}^{\times\mu}$ -**link** [cf. (ii); Theorem 3.11, (i), (iii)], i.e., with the **horizontal arrows** of the log-theta-lattice;
- (Ind3) — which may be thought of as arising from the requirement of *compatibility* with the **log-Kummer correspondences** of Theorem 3.11, (ii), i.e., with the **vertical arrows** of the log-theta-lattice.

The various indeterminacies (Ind1), (Ind2), (Ind3) to which the multiradial representation is subject may be thought of as data that describes some sort of “**formal quotient**”, like the “**fine moduli spaces**” that appear in algebraic geometry. In this context, the **procession-normalized mono-analytic log-volumes** [i.e., where the average is taken over  $j \in \mathbb{F}_l^*$ ] of Theorem 3.11, (i), (a), (c), furnish a means of constructing a sort of associated “**coarse space**” or “**inductive limit**” [of the “inductive system” constituted by this “formal quotient”] — i.e., in the sense that [one verifies immediately — cf. Proposition 3.9, (ii) — that] the resulting log-volumes  $\in \mathbb{R}$  are **invariant** with respect to the indeterminacies (Ind1), (Ind2), and have the effect of converting the indeterminacy (Ind3) into an **inequality** (from above). Moreover, the **log-link compatibility** of the various log-volumes that appear [cf. Proposition 3.9, (iv); the final portion of Theorem 3.11, (ii)] ensures that these log-volumes are *compatible* with [the portion of the “formal quotient”/“inductive system” constituted by] the various arrows [i.e., *Kummer isomorphisms* and **log-links**] of the **log-Kummer correspondence** of Theorem 3.11, (ii). Here, we note that the *averages over*  $j \in \mathbb{F}_l^*$  that appear in the definition of the procession-normalized volumes involved may be thought of as a consequence of the  $\mathbb{F}_l^*$ -**symmetry** acting on the labels of the theta values that give rise to the LGP-monoids — cf. the also the definition of the symbol “ $\Delta$ ” in [IUTchII], Corollary 4.10, (i), via the *identification of the symbols* “0” and “ $\langle \mathbb{F}_l^* \rangle$ ”; the discussion of Remark 3.9.3. Also, in this context, it is of interest to observe that the various **tensor products** that appear in the various local mono-analytic tensor packets that arise in the multiradial representation of Theorem 3.11, (i), (a), have the effect of **identifying** the operation of “multiplication by elements of  $\mathbb{Z}$ ” — and hence also

the effect on **log-volumes** of such multiplication operations! — at *different labels*  $\in \mathbb{F}_l^*$ .

(xi) Now let us consider a ***q*-pilot object** at  $(1, 0)$ , which we think of — relative to the relevant copy of “ $\mathcal{F}_{\text{mod}}^\circ$ ” — in terms of the **mono-analytic log-shells** constructed at  $(1, 0)$  [cf. (v)]. Then the  $\Theta_{\text{LGP}}^{\times\mu}$ -**link** from  $(0, 0)$  to  $(1, 0)$  may be interpreted as a sort of **gluing isomorphism** that relates the **arithmetic holomorphic structure** — i.e., the “*conventional ring/scheme-theory*” — at  $(1, 0)$  to the arithmetic holomorphic structure at  $(0, 0)$  in such a way that the  **$\Theta$ -pilot object** at  $(0, 0)$  [thought of as an object of the relevant global realified Frobenioid] corresponds to the ***q*-pilot object** at  $(1, 0)$  [cf. (i)]. On the other hand, the discussion of (x) furnishes **another way of computing** the *global arithmetic degree* — i.e., the *log-volume* [cf. Theorem 3.11, (i), (c)] — of this *q*-pilot object at  $(1, 0)$ , namely, by computing the log-volume of the  $\Theta$ -pilot object  $(0, 0)$ , constructed relative to the **alien** [i.e., from the point of view of the arithmetic holomorphic structure at  $(1, 0)$ ] arithmetic holomorphic structure at  $(0, 0)$ , in **terms** of the arithmetic holomorphic structure at  $(1, 0)$  [cf. the final portion of Theorem 3.11, (i); Remark 3.11.1]. That is to say,

*the theory of the present series of papers yields two tautologically equivalent ways to compute the log-volume of the *q*-pilot object at  $(1, 0)$*




Fig. 3.6: Two tautologically equivalent ways to compute the log-volume of the *q*-pilot object at  $(1, 0)$

— cf. Fig. 3.6. If one interprets this observation in terms of the notation introduced in the statement of Corollary 3.12, then one concludes that

$$- |\log(\underline{q})| \leq - |\log(\underline{\Theta})| \in \mathbb{R}$$



— hence, in particular, that

$$- |\log(\underline{\underline{\Theta}})| \leq C_{\Theta} \cdot |\log(\underline{\underline{q}})|$$

for any  $C_{\Theta} \in \mathbb{R}$  such that  $- |\log(\underline{\underline{\Theta}})| \leq C_{\Theta} \cdot |\log(\underline{\underline{q}})|$ . Since [one verifies immediately that]  $|\log(\underline{\underline{q}})| \in \mathbb{R}$  is *positive*, we thus conclude that  $C_{\Theta} \geq -1$ , as desired. In this context, it is useful to recall that the above argument depends, in an essential way [cf. the discussion of (ii), (vi)], on the theory of [EtTh], which *does not admit any evident generalization* to the case of  $N$ -th tensor powers of  $\Theta$ -pilot objects, for  $N \geq 2$ . That is to say, the *log-volume* of such an  $N$ -th tensor power of a  $\Theta$ -pilot object must always be computed as the result of *multiplying the log-volume of the original  $\Theta$ -pilot object by  $N$*  — cf. Remark 2.1.1, (iv); [IUTchII], Remark 3.6.4, (iii), (iv). In particular, although the analogue of the above argument for such an  $N$ -th tensor power would lead to **sharper inequalities** than the inequalities obtained here, it is difficult to see how to obtain such sharper inequalities via a routine generalization of the above argument. In fact, as we shall see in [IUTchIV], these sharper inequalities are known to be **false** [cf. [IUTchIV], Remark 2.3.2, (ii)].

(xii) In the context of the argument of (xi), it is useful to observe the important role played by the **global** realified Frobenioids that appear in the  $\Theta_{\text{LGP}}^{\times\mu}$ -link. That is to say, since ultimately one is only concerned with the computation of *log-volumes*, it might appear, at first glance, that it is possible to dispense with the use of such global Frobenioids and instead work only with the various *local Frobenioids*, for  $\underline{v} \in \underline{\mathbb{V}}$ , that are directly related to the computation of log-volumes. On the other hand, observe that since the isomorphism of [local or global!] Frobenioids arising from the  $\Theta_{\text{LGP}}^{\times\mu}$ -link only preserves **isomorphism classes of objects** of these Frobenioids [cf. the discussion of Remark 3.6.2, (i)], to work only with local Frobenioids means that one must contend with the **indeterminacy** of not knowing whether, for instance, such a local Frobenioid object at some  $\underline{v} \in \underline{\mathbb{V}}^{\text{non}}$  corresponds to a *given open submodule* of the log-shell at  $\underline{v}$  or to, say, the  $p_{\underline{v}}^N$ -*multiple of this submodule*, for  $N \in \mathbb{Z}$ . Put another way, one must contend with the indeterminacy arising from the fact that, unlike the case with the global Frobenioids “ $\mathcal{F}_{\text{MOD}}^{\circledast}$ ”, “ $\mathcal{F}_{\text{MOD}}^{\circledR}$ ”, objects of the various local Frobenioids that arise admit **endomorphisms which are not automorphisms**. This indeterminacy has the effect of rendering *meaningless* any attempt to perform a precise log-volume computation as in (xi).  
○

**Remark 3.12.1.**

(i) In [IUTchIV], we shall be concerned with obtaining *more explicit upper bounds* on  $- |\log(\underline{\underline{\Theta}})|$ , i.e., *estimates* “ $C_{\Theta}$ ” as in the statement of Corollary 3.12.

(ii) It is not difficult to verify that, for  $\lambda \in \mathbb{Q}_{>0}$ , one may obtain a similar theory to the theory developed in the present series of papers for “*generalized  $\Theta_{\text{LGP}}^{\times\mu}$ -links*” of the form

$$\underline{\underline{q}}^{\lambda} \quad \mapsto \quad \underline{\underline{q}}^{\begin{pmatrix} 1^2 \\ \vdots \\ (l^*)^2 \end{pmatrix}}$$



— i.e., so the theory developed in the present series of papers corresponds to the case of  $\lambda = 1$ . This sort of “generalized  $\Theta_{\text{LGP}}^{\times\mu}$ -link” is roughly reminiscent of — but by no means equivalent to! — the sort of issues considered in the discussion of Remark 2.2.2, (i). Here, we observe that raising to the  $\lambda$ -th power on the “ $\underline{q}$  side” *differs quite fundamentally* from raising to the  $\lambda$ -th power on the “ $\underline{q}^{(1^2 \dots (l^*)^2)}$  side”, an issue that is discussed briefly [in the case of  $\lambda = N$ ] in the final portion of Step (xi) of the proof of Corollary 3.12. That is to say, “generalized  $\Theta_{\text{LGP}}^{\times\mu}$ -links” as in the above display *differ fundamentally* both from the situation of Remark 2.2.2, (i), and the situation discussed in the final portion of Step (xi) of the proof of Corollary 3.12 in that the theory of the **first power** of the **étale theta function** is left unchanged [i.e., relative to the theory developed in the present series of papers] — cf. the discussion of Remark 2.2.2, (i); Step (xi) of the proof of Corollary 3.12. At any rate, in the case of “generalized  $\Theta_{\text{LGP}}^{\times\mu}$ -links” as in the above display, one may apply the same arguments as the arguments used to prove Corollary 3.12 to conclude the *inequality*

$$C_{\Theta} \geq -\lambda$$

— i.e., which is *sharper*, for  $\lambda < 1$ , than the inequality obtained in Corollary 3.12 in the case of  $\lambda = 1$ . In fact, however, such sharper inequalities will not be of interest to us, since, in [IUTchIV], our estimates for the upper bound  $C_{\Theta}$  will be *sufficiently rough as to be unaffected* by adding a constant of absolute value  $\leq 1$ .

### Remark 3.12.2.

(i) One of the main themes of the present series of papers is the issue of *dismantling the two underlying combinatorial dimensions* of a number field — cf. Remarks 1.2.2, (vi), of the present paper, as well as [IUTchI], Remarks 3.9.3, 6.12.3, 6.12.6; [IUTchII], Remarks 4.7.5, 4.7.6, 4.11.2, 4.11.3, 4.11.4. The principle examples of this topic may be summarized as follows.

- (a) splittings of various monoids into **unit** and **value group** portions;
- (b) separating the “ $\mathbb{F}_l$ ” arising from the  $l$ -torsion points of the elliptic curve — which may be thought of as a sort of “*finite approximation*” of  $\mathbb{Z}$ ! — into a [*multiplicative*]  $\mathbb{F}_l^*$ -**symmetry** — which may also be thought of as corresponding to the *global arithmetic* portion of the arithmetic fundamental groups involved — and  $a(n)$  [*additive*]  $\mathbb{F}_l^{\times\pm}$ -**symmetry** — which may also be thought of as corresponding to the *geometric* portion of the arithmetic fundamental groups involved;
- (c) separating the ring structures of the various **global number fields** that appear into their respective underlying **additive** structures — which may be related directly to the various *log-shells* that appear — and their respective underlying **multiplicative** structures — which may be related directly to the various *Frobenioids* that appear.

From the point of view of Theorem 3.11, example (a) may be seen in the “**non-interference**” properties that underlie the **log-Kummer correspondences** of

Theorem 3.11, (ii), (b), (c), as well as in the  $\Theta_{\text{LGP}}^{\times\mu}$ -**link compatibility** properties discussed in Theorem 3.11, (ii), (c), (d).

(ii) On the other hand, another important theme of the present §3 consists of the issue of “**reassembling**” *these two dismantled combinatorial dimensions* by means of the **multiradial mono-analytic containers** furnished by the **mono-analytic log-shells** — cf. Fig. 3.4 — i.e., of exhibiting the extent to which these two dismantled combinatorial dimensions **cannot be separated** from one another by describing the “**structure of the intertwining**” between these two dimensions that existed prior to their separation. From this point of view, one may think of the **multiradial representations** discussed in Theorem 3.11, (i) [cf. also Theorem 3.11, (ii), (iii)], as the *final output* of this “reassembling procedure”. Put another way, from the point of view of example (a) of the discussion of (i), this “reassembling procedure” allows one to **compute/estimate the value group portions** of various monoids of arithmetic interest in terms of the **unit group portions** of these monoids. It is precisely these estimates that give rise to the **inequality** obtained in Corollary 3.12.

(iii) One fundamental aspect of the theory that renders possible the “reassembling procedure” discussed in (ii) [cf. the discussion of Step (iv) of the proof of Corollary 3.12] is the “**juggling of  $\boxplus, \boxtimes$** ” [cf. the discussion of Remark 1.2.2, (vi)] effected by the **log-links**, i.e., the **vertical arrows** of the log-theta-lattice. This “juggling of  $\boxplus, \boxtimes$ ” may be thought of as a sort of combinatorial way of representing the **arithmetic holomorphic structure** associated to a **vertical line** of the log-theta-lattice. Indeed, at *archimedean primes*, this juggling amounts essentially to *multiplication by  $\pm i$* , which is a well-known method [cf. the notion of an “*almost complex structure*”!] for representing holomorphic structures in the classical theory of differential manifolds. On the other hand, it is important to recall in this context that this “juggling of  $\boxplus, \boxtimes$ ” is precisely what gives rise to the **upper semi-compatibility** indeterminacy (Ind3) [cf. Proposition 3.5, (ii); Remark 3.10.2, (i)].

**Remark 3.12.3.**

(i) Let  $S$  be a *hyperbolic Riemann surface of finite type* of genus  $g_S$  with  $r_S$  punctures. Write  $\chi_S \stackrel{\text{def}}{=} -(2g_S - 2 + r_S)$  for the *Euler characteristic* of  $S$  and  $d\mu_S$  for the Kähler metric on  $S$  [i.e., the (1,1)-form] determined by the *Poincaré metric* on the upper half-plane. Recall the *analogy* discussed in [IUTchI], Remark 4.3.3, between the theory of **log-shells**, which plays a key role in the theory developed in the present series of papers, and the **classical metric geometry of hyperbolic Riemann surfaces**. Then, relative to this analogy, the **inequality** obtained in Corollary 3.12 may be regarded as corresponding to the inequality

$$\chi_S = - \int_S d\mu_S < 0$$

— i.e., in essence, a statement of the **hyperbolicity** of  $S$  — arising from the classical **Gauss-Bonnet formula**, together with the **positivity** of  $d\mu_S$ . Relative to the analogy between *real analytic Kähler metrics* and *ordinary Frobenius liftings*

discussed in [ $p$ Ord], Introduction, §2 [cf. also the discussion of [ $p$ Teich], Introduction, §0], the *local* property constituted by this positivity of  $d\mu_S$  may be thought of as corresponding to the [*local* property constituted by the] *Kodaira-Spencer isomorphism of an indigenous bundle* — i.e., which gives rise to the *ordinarity* of the corresponding Frobenius lifting on the ordinary locus — in the  $p$ -adic theory. As discussed in [AbsTopIII], §I5, these properties of indigenous bundles in the  $p$ -adic theory may be thought of as corresponding, in the theory of *log-shells*, to the “*maximal incompatibility*” between the various *Kummer isomorphisms* and the *corically constructed data* of the Frobenius-picture of Proposition 1.2, (x). On the other hand, it is just this “maximal incompatibility” that gives rise to the “*upper semi-commutativity*” discussed in Remark 1.2.2, (iii), i.e., [from the point of view of the theory of the present §3] the **upper semi-compatibility** indeterminacy (Ind3) of Theorem 3.11, (ii), that underlies the **inequality** of Corollary 3.12 [cf. Step (x) of the proof of Corollary 3.12].

(ii) The “*metric aspect*” of Corollary 3.12 discussed in (i) is reminiscent of the analogy between the theory of the present series of papers and *classical complex Teichmüller theory* [cf. the discussion of [IUTchI], Remark 3.9.3] in the following sense:

Just as *classical complex Teichmüller theory* is concerned with relating distinct holomorphic structures in a sufficiently **canonical** way as to **minimize** the resulting **volume distortion**, the **canonical** nature of the algorithms discussed in Theorem 3.11 for relating **alien arithmetic holomorphic structures** [cf. Remark 3.11.1] gives rise to a relatively **strong estimate** of the [**log-]volume distortion** [cf. Corollary 3.12] resulting from such a deformation of the arithmetic holomorphic structure.

**Remark 3.12.4.** In light of the discussion of Remark 3.12.3, it is of interest to reconsider the analogy between the theory of the present series of papers and the  $p$ -adic *Teichmüller theory* of [ $p$ Ord], [ $p$ Teich], in the context of Theorem 3.11, Corollary 3.12.

(i) First, we observe that the **splitting monoids** at  $\underline{v} \in \mathbb{V}^{\text{bad}}$  [cf. Theorem 3.11, (i), (b); Theorem 3.11, (ii), (b)] may be regarded as analogous to the **canonical coordinates** of  $p$ -adic Teichmüller theory [cf., e.g., [ $p$ Teich], Introduction, §0.9] that are constructed over the *ordinary locus* of a canonical curve. In particular, it is natural to regard the **bad primes**  $\in \mathbb{V}^{\text{bad}}$  as corresponding to the **ordinary locus** of a canonical curve and the **good primes**  $\in \mathbb{V}^{\text{good}}$  as corresponding to the **supersingular locus** of a canonical curve. This point of view is reminiscent of the discussion of [IUTchII], Remark 4.11.4, (iii).

(ii) On the other hand, the **bi-coric mono-analytic log-shells** — i.e., the various local “ $\mathcal{O}^{\times\mu}$ ” — that appear in the tensor packets of Theorem 3.11, (i), (a); Theorem 3.11, (ii), (a), may be thought of as corresponding to the [**multiplicative!**] **Teichmüller representatives** associated to the various Witt rings that appear in  $p$ -adic Teichmüller theory. Within a *fixed arithmetic holomorphic structure*, these *mono-analytic log-shells* arise from “**local holomorphic units**” — i.e., “ $\mathcal{O}^{\times}$ ” — which are subject to the  $\mathbb{F}_l^{\times\pm}$ -**symmetry**. These “local holomorphic units” may be thought of as corresponding to the **positive characteristic**

**ring structures** on [the positive characteristic reductions of] Teichmüller representatives. Here, the **uniradial**, i.e., “**non-multiradial**”, nature of these “local holomorphic units” [cf. the discussion of [IUTchII], Remark 4.7.4, (ii); [IUTchII], Figs. 4.1, 4.2] may be regarded as corresponding to the *mixed characteristic nature of Witt rings*, i.e., the **incompatibility** of Teichmüller representatives with the **additive structure** of Witt rings.

(iii) The set  $\mathbb{F}_l^*$  of  $l^*$  “**theta value labels**”, which plays an important role in the theory of the present series of papers, may be thought of as corresponding to the “**factor of  $p$** ” that appears in the “*mod  $p/p^2$  portion*”, i.e., the gap separating the “*mod  $p$* ” and “*mod  $p^2$* ” portions, of the rings of Witt vectors that occur in the  $p$ -adic theory. From this point of view, one may think of the *procession-normalized volumes* obtained by taking **averages over  $j \in \mathbb{F}_l^*$**  [cf. Corollary 3.12] as corresponding to the operation of **dividing by  $p$**  to relate the “*mod  $p/p^2$  portion*” of the Witt vectors to the “*mod  $p$  portion*” of the Witt vectors [i.e., the characteristic  $p$  theory]. In this context, the **multiradial representation** of Theorem 3.11, (i), by means of *mono-analytic log-shells labeled by elements of  $\mathbb{F}_l^*$*  may be thought of as corresponding to the **derivative** of the **canonical Frobenius lifting** on a canonical curve in the  $p$ -adic theory [cf. the discussion of [AbsTopIII], §I5] in the sense that this multiradial representation may be regarded as a sort of **comparison** of the **canonical splitting monoids** discussed in (i) to the “**absolute constants**” [cf. the discussion of (ii)] constituted by the **bi-coric mono-analytic log-shells**. This “*absolute comparison*” is precisely what results in the **indeterminacies** (Ind1), (Ind2) of Theorem 3.11, (i).

(iv) In the context of the discussion of (iii), we note that the set of labels  $\mathbb{F}_l^*$  may, alternatively, be thought of as corresponding to the **infinitesimal moduli** of the positive characteristic curve under consideration in the  $p$ -adic theory [cf. the discussion of [IUTchII], Remark 4.11.4, (iii), (d)]. That is to say, the “*deformation dimension*” constituted by the *horizontal dimension* of the log-theta-lattice in the theory of the present series of papers or by the deformations modulo various powers of  $p$  in the  $p$ -adic theory [cf. Remark 1.4.1, (iii); Fig. 1.3] is **highly canonical** in nature, hence may be thought of as being equipped with a natural isomorphism to the “**absolute moduli**” — i.e., so to speak, the “*moduli over  $\mathbb{F}_1$* ” — of the *given number field equipped with an elliptic curve*, in the theory of the present series of papers, or of the *given positive characteristic hyperbolic curve equipped with a nilpotent ordinary indigenous bundle*, in  $p$ -adic Teichmüller theory.

(v) Let  $A$  be a ring of Witt vectors of a perfect field  $k$  of characteristic  $p$ ;  $X$  a *smooth, proper hyperbolic curve* over  $A$  of genus  $g_X$  which is **canonical** in the sense of  $p$ -adic Teichmüller theory;  $\widehat{X}$  the  $p$ -adic formal scheme associated to  $X$ ;  $\widehat{U} \subseteq \widehat{X}$  the *ordinary locus* of  $\widehat{X}$ . Write  $\omega_{X_k}$  for the canonical bundle of  $X_k$ . Then when [cf. the discussion of (iii)] one computes the **derivative** of the **canonical Frobenius lifting**  $\Phi : \widehat{U} \rightarrow \widehat{U}$  on  $\widehat{U}$ , one must contend with “*interference phenomena*” between the various *copies* of some positive characteristic algebraic geometry set-up — i.e., at a more concrete level, the various *Frobenius conjugates* “ $t^p$ ” [where  $t$  is a local coordinate on  $X_k \stackrel{\text{def}}{=} X \times_A k$ ] associated to various  $n \in \mathbb{N}_{\geq 1}$ . In particular, this derivative only yields [upon dividing by  $p$ ] an *inclusion* [i.e., not an isomorphism!] of line bundles

$$\omega_{X_k} \hookrightarrow \Phi^* \omega_{X_k}$$

— also known as the “[**square**] **Hasse invariant**” [cf. [pOrd], Chapter II, Proposition 2.6; the discussion of “generalities on ordinary Frobenius liftings” given in [pOrd], Chapter III, §1]. Thus, at the level of *global degrees of line bundles*, we obtain an *inequality* [i.e., not an equality!]

$$(1 - p)(2g_X - 2) \leq 0$$

<i>inter-universal Teichmüller theory</i>	<i>p-adic Teichmüller theory</i>
<b>splitting monoids</b> at $\underline{v} \in \underline{\mathbb{V}}^{\text{bad}}$	<b>canonical coordinates</b> on the <i>ordinary locus</i>
<b>bad primes</b> $\in \underline{\mathbb{V}}^{\text{bad}}$	<b>ordinary locus</b> of a can. curve
<b>good primes</b> $\in \underline{\mathbb{V}}^{\text{good}}$	<b>supersing.</b> locus of a can. curve
<b>mono-analytic log-shells</b> “ $\mathcal{O}^{\times\mu}$ ”	[ <b>multiplicative!</b> ] <b>Teich. reps.</b>
<b>uniradial</b> “ <b>local hol. units</b> $\mathcal{O}^{\times}$ ” subject to $\mathbb{F}_l^{\times\pm}$ - <b>symmetry</b>	<b>pos. char. ring structures</b> on [pos. char. reductions of] <b>Teich. reps.</b>
set of “ <b>theta value labels</b> ” $\mathbb{F}_l^*$	<b>factor <math>p</math></b> in mod $p/p^2$ portion of Witt vectors
<b>multiradial rep.</b> via $\mathbb{F}_l^*$ -labeled <b>mono-analytic log-shells</b> (cf. (Ind1), (Ind2))	<b>derivative</b> of the <b>canonical Frobenius lifting</b>
set of “ <b>theta value labels</b> ” $\mathbb{F}_l^*$	implicit “ <b>absolute moduli</b> / $\mathbb{F}_1$ ”
<b>inequality</b> arising from <b>upper semi-compatibility</b> (cf. (Ind3))	<b>inequality</b> arising from interference between <b>Frobenius conjugates</b>

Fig. 3.7: The analogy between inter-universal Teichmüller theory and *p*-adic Teichmüller theory

— which may be thought of as being, in essence, a statement of the **hyperbolicity** of  $X$  [cf. the inequality of the display of Remark 3.12.3, (i)]. Since the “*Frobenius conjugate dimension*” [i.e., the “ $n$ ” that appears in “ $t^n$ ”] in the  $p$ -adic theory corresponds to the *vertical dimension* of the log-theta-lattice in the theory of the present series of papers [cf. Remark 1.4.1, (iii); Fig. 1.3], we thus see that the inequality of the above display in the  $p$ -adic case arises from circumstances that are *entirely analogous* to the circumstances — i.e., the **upper semi-compatibility** indeterminacy (Ind3) of Theorem 3.11, (ii) — that underlie the **inequality** of Corollary 3.12 [cf. Step (x) of the proof of Corollary 3.12; the discussion of Remark 3.12.3, (i)].

(vi) The analogies of the above discussion are summarized in Fig. 3.7 above.



**Bibliography**

- [Lang] S. Lang, *Algebraic number theory*, Addison-Wesley Publishing Co. (1970).
- [ $p$ Ord] S. Mochizuki, A Theory of Ordinary  $p$ -adic Curves, *Publ. Res. Inst. Math. Sci.* **32** (1996), pp. 957-1151.
- [ $p$ Teich] S. Mochizuki, *Foundations of  $p$ -adic Teichmüller Theory*, AMS/IP Studies in Advanced Mathematics **11**, American Mathematical Society/International Press (1999).
- [QuCnf] S. Mochizuki, Conformal and quasiconformal categorical representation of hyperbolic Riemann surfaces, *Hiroshima Math. J.* **36** (2006), pp. 405-441.
- [SemiAnbd] S. Mochizuki, Semi-graphs of Anabelioids, *Publ. Res. Inst. Math. Sci.* **42** (2006), pp. 221-322.
- [FrdI] S. Mochizuki, The Geometry of Frobenioids I: The General Theory, *Kyushu J. Math.* **62** (2008), pp. 293-400.
- [FrdII] S. Mochizuki, The Geometry of Frobenioids II: Poly-Frobenioids, *Kyushu J. Math.* **62** (2008), pp. 401-460.
- [EtTh] S. Mochizuki, The Étale Theta Function and its Frobenioid-theoretic Manifestations, *Publ. Res. Inst. Math. Sci.* **45** (2009), pp. 227-349.
- [AbsTopIII] S. Mochizuki, *Topics in Absolute Anabelian Geometry III: Global Reconstruction Algorithms*, RIMS Preprint **1626** (March 2008).
- [IUTchI] S. Mochizuki, *Inter-universal Teichmüller Theory I: Construction of Hodge Theaters*, preprint.
- [IUTchII] S. Mochizuki, *Inter-universal Teichmüller Theory II: Hodge-Arakelov-theoretic Evaluation*, preprint.
- [IUTchIV] S. Mochizuki, *Inter-universal Teichmüller Theory IV: Log-volume Computations and Set-theoretic Foundations*, preprint.