

RIMS-1829

**Harnack inequalities and local central limit
theorem for the polynomial lower tail
random conductance model**

By

O. BOUKHADRA, T. KUMAGAI and P. MATHIEU

June 2015



京都大学 数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

HARNACK INEQUALITIES AND LOCAL CENTRAL LIMIT THEOREM FOR THE POLYNOMIAL LOWER TAIL RANDOM CONDUCTANCE MODEL

O. BOUKHADRA¹, T. KUMAGAI² AND P. MATHIEU³

¹Département de Mathématiques, Université de Constantine 1

boukhadra@umc.edu.dz

²Research Institute for Mathematical Sciences, Kyoto University,
Kyoto 606-8502, Japan

kumagai@kurims.kyoto-u.ac.jp

³Centre de Mathématiques et Informatique

Aix Marseille Université, CNRS, Centrale Marseille, I2M, UMR 7373,
13453 Marseille France

pierre.mathieu@univ-amu.fr

ABSTRACT. *We prove upper bounds on the transition probabilities of random walks with i.i.d. random conductances with a polynomial lower tail near 0. We consider both constant and variable speed models. Our estimates are sharp. As a consequence, we derive local central limit theorems, parabolic Harnack inequalities and Gaussian bounds for the heat kernel. Some of the arguments are robust and applicable for random walks on general graphs. Such results are stated under a general setting.*

keywords : *Markov chains, Random walk, Random environments, Random conductances, Percolation.*

AMS 2000 Subject Classification : *60G50; 60J10; 60K37.*

1. Introduction and Results

The work presented below mainly concerns the Random Conductance Model (RCM) with polynomial lower tail. We shall obtain various heat kernel bounds, Harnack inequalities and a local central-limit theorem for such models under sharp conditions on the fatness of the tail of the conductances near 0. Some of our arguments exploit specific features of the model - mainly some geometric information on the field of conductances and its spectral implications - while other arguments are general properties of random walks on graphs. In the rest of this introduction, we will separate results that are more robust from those that are specific to the RCM. The robust results will be discussed in the first subsection below, and results specific to the RCM and references to the existing literature will be given in the second subsection. Readers who are interested in RCM may start reading this paper from the second subsection.

Notation: We use c or C as generic positive constants.

1.1 Part I: Framework and the results.

In this subsection, we give some sufficient conditions for various heat kernel bounds, Harnack inequalities and a local central-limit theorem on a general graph. The results will be used in the next subsection for a concrete RCM.

Let (G, π) be a weighted graph. That is, G is a countable set and $\omega_{xy} = \omega_{yx} \geq 0$ for each $x, y \in G$. We write $x \sim y$ if and only if $\omega_{xy} > 0$. For $x \neq y$, $\ell(x, y) = \{x_0, x_1, \dots, x_m\}$ is called a path from x to y if $x = x_0, x_1, \dots, x_m = y$ and $x_i \sim x_{i+1}$ for $i = 0, \dots, m-1$. Write $|\ell(x, y)| = m$. Define the graph distance by $d(x, y) = \min\{|\ell(x, y)| : \ell(x, y) \in \mathcal{P}(x, y)\}$ where $\mathcal{P}(x, y)$ is the set of paths from x to y . We define $d(x, x) = 0$ for $x \in G$. We assume (G, π) is connected and it has bounded degree, that is, $d(x, y) < \infty$ for each $x, y \in G$ and there exists $M > 0$ such that $|\{y \in G : \omega_{xy} > 0\}| \leq M$ for each $x \in G$. Write $B(x, R) := \{x \in G : d(x, y) < R\}$ and $\bar{B}(x, R) := \{x \in G : d(x, y) \leq R\}$. For $A \subset G$, define $\pi(A) = \sum_{x \in A} \pi(x)$ where $\pi(x) = \sum_{y \sim x} \omega_{xy}$, and $\nu(A) = \sum_{x \in A} \nu_x$ where $\nu_x \equiv 1$.

We will consider VSRW (variable speed random walk) and CSRW (constant speed random walk) that correspond to (G, π) . Both are continuous time Markov chains whose transition probability from x to y is given by $\omega_{xy}/\pi(x)$. The holding time at x is exponentially distributed with mean $\pi(x)^{-1}$ for VSRW and with mean 1 for CSRW. The corresponding discrete Laplace operator and heat kernel can be written as

$$\mathcal{L}_\theta f(x) = \frac{1}{\theta_x} \sum_y (f(y) - f(x)) \omega_{xy}, \quad p_t^{(\theta)}(x, y) = P^x(X_t^{(\theta)} = y) / \theta_y,$$

where $\theta_x = \theta(x) = \pi(x)$ for CSRW and $\theta_x = 1$ for VSRW. Thus the notation \mathcal{L}_π and $X^{(\pi)}$ correspond to CSRW and \mathcal{L}_ν , $X^{(\nu)}$ correspond to VSRW. We may and will often remove the script when results are valid for both types of random walks.

Let $\tilde{d}(\cdot, \cdot)$ be a metric defined by

$$\tilde{d}(x, y) = \min \left\{ \sum_{i=0}^{m-1} (1 \wedge \omega_{x_i x_{i+1}}^{-1/2}) : \ell(x, y) = \{x_0, x_1, \dots, x_m\} \in \mathcal{P}(x, y) \right\}.$$

Note that by definition, it is clear that $\tilde{d}(x, y) \leq d(x, y)$ for all $x, y \in G$. Write $\tilde{B}(x, R) := \{x \in \mathbb{Z}^d : \tilde{d}(x, y) < R\}$. For $A \subset G$, let $\tau_A = \inf\{t \geq 0 : X_t \notin A\}$.

In the following, we fix θ (which is either π or ν) and consider either CSRW or VSRW.

Assumption 1.1 *Let $x_0 \in G$ be a distinguished point.*

(i) *There exist $\delta > 0, c_1 > 0$ and $T_0(x_0) \in [1, \infty)$ such that*

$$p_t(x, y) \leq c_1 t^{-d/2} \quad \forall x, y \in B(x_0, t^{(1+\delta)/2}), \quad t \geq T_0(x_0). \quad (1.1)$$

(ii) *There exist $\delta > 0, c_2 > 0$ and $R_0(x_0) \in [1, \infty)$ such that the following hold:*

(CSRW case: $\theta = \pi$) $c_2 r^2 \leq E^x[\tau_{B(x, r)}]$ for all $x \in B(x_0, r^{1+\delta})$ and all $r \geq R_0(x_0)$.

(VSRW case: $\theta = \nu$) $c_2 r^2 \leq E^x[\tau_{\tilde{B}(x,r)}]$ for all $x \in \tilde{B}(x_0, r^{1+\delta})$ and all $r \geq \tilde{R}_0(x_0)$.
 (iii) There exist $C_E > 0$ and $R_1(x_0) \in [1, \infty)$ such that if $R \geq R_1(x_0)$ and a positive function $h : \bar{B}(x_0, R) \rightarrow \mathbb{R}_+$ is harmonic on $B = B(x_0, R)$, then writing $B' = B(x_0, R/2)$,

$$\sup_{B'} h \leq C_E \inf_{B'} h. \quad (\text{H})$$

(iv) Let θ be as above. There exist $\delta > 0$, $c_3, c_4 > 0$ and $R_2(x_0) \in [1, \infty)$ such that

$$c_3 R^d \leq \theta(B(x_0, R)) \leq \sup_{x \in B(x_0, R^{1+\delta})} \theta(B(x, R)) \leq c_4 R^d, \quad \text{for all } R \geq R_2(x_0).$$

(v) (CSRW case: $\theta = \pi$) There exist $\kappa > 0$ and $R_3(x_0) \in [1, \infty)$ such that

$$\min_{x \in B(x_0, R)} \pi(x) \geq R^{-\kappa} \quad \text{for all } R \geq R_3(x_0).$$

(VSRW case: $\theta = \nu$) There exist $c_5 > 0$ and $R_4(x_0) \in [1, \infty)$ such that for any $x \in B(x_0, R)$, $R \geq R_4(x_0)$, if $d(x, y) \geq R$ then it holds that

$$\tilde{d}(x, y) \geq c_5 d(x, y).$$

Under the assumption, we have the following.

Heat kernel estimates

Proposition 1.2 Assume Assumption 1.1 and let $\varepsilon \in (0, \delta/(1 + \delta))$. There exist $c_1, \dots, c_5 > 0$ and $R_*(x_0) \in [1, \infty)$ such that for $x, y \in G$ and $t > 0$, if

$$c_1(d(x, y) \vee t^{\frac{1}{2-\varepsilon}}) \geq R_*(x_0), \quad (1.2)$$

and

$$d(x_0, x) \leq c_1(d(x, y) \vee t^{\frac{1}{2-\varepsilon}}), \quad (1.3)$$

hold, then

$$p_t(x, y) \leq c_2 t^{-d/2} \exp\left(-c_3 d(x, y)^2/t\right) \quad \text{for } t > d(x, y), \quad (1.4)$$

$$p_t(x, y) \leq c_4 \exp\left(-c_5 d(x, y)(1 \vee \log(d(x, y)/t))\right) \quad \text{for } t \leq d(x, y). \quad (1.5)$$

Corollary 1.3 Assume Assumption 1.1. There exist $c_1 > 0$ and $R_*(x_0) \in [1, \infty)$ such that if $R \geq R_*(x_0)$, then

$$\sup_{0 < s \leq T} p_s(x, y) \leq c_1 T^{-d/2} \quad \text{for all } x, y \in B(x_0, 2R) \text{ with } d(x, y) \geq R,$$

where $T = R^2$.

For a subset $A \subset G$, let $\{X_t^A\}_{t \geq 0}$ be the process killed on exiting A and define the Dirichlet heat kernel $p_t^A(\cdot, \cdot)$ as

$$p_t^A(x, y) = P^x(X_t^A = y)/\theta_y.$$

Then the following heat kernel lower bound holds.

Proposition 1.4 *Assume Assumption 1.1. Then there exist $c_1, \delta_0 \in (0, 1)$ and $T_1(x_0) \in [1, \infty)$ such that*

$$p_t^{B(x_0, t^{1/2})}(x, y) \geq c_1 t^{-d/2}, \quad \forall x, y \in B(x_0, \delta_0 t^{1/2})$$

for all $t \geq T_1(x_0)$.

Parabolic Harnack inequalities and Hölder continuity of caloric functions

For $x \in G$ and $R, T > 0$, let $C_* \geq 2$, $Q(x, R, T) := (0, 4T] \times B(x, C_* R)$ and define

$$Q_-(x, R, T) := [T, 2T] \times B(x, R), \quad Q_+(x, R, T) := [3T, 4T] \times B(x, R).$$

Let $u(t, x)$ be a function defined on $[0, 4T] \times \bar{B}(x, C_* R)$. We say $u(t, x)$ is caloric on Q if it satisfies the following: for $t \in (0, 4T)$ and $y \in B(x, C_* R)$:

$$\partial_t u(t, y) = \mathcal{L}_\theta u(t, y).$$

We then have the following.

Theorem 1.5 (Parabolic Harnack inequalities)

Assume Assumption 1.1. Then there exist $c_1 > 0, C_ \geq 2$ and $R_5(x_0) \in [1, \infty)$ such that for any $R \geq R_5(x_0)$, and any non-negative function $u = u(t, x)$ which is caloric on $Q(x_0, R, R^2)$, it holds that*

$$\sup_{(t, x) \in Q_-(x_0, R, R^2)} u(t, x) \leq c_1 \inf_{(t, x) \in Q_+(x_0, R, R^2)} u(t, x). \quad (1.6)$$

Corollary 1.6 *Assume Assumption 1.1. Then there exist $c_1, \beta > 0$, $C_* \geq 2$ and $R_6(x_0) \in [1, \infty)$ such that the following holds: For any $R \geq R_6(x_0)$ and $T' \geq R^2 + 1$, let $R' = \sqrt{T'}$ and suppose that u is a positive caloric function on $Q(x_0, R', T')$. Then for any $x_1, x_2 \in B(x_0, R)$ and any $t_1, t_2 \in [4(T' - R^2), 4T']$, we have*

$$|u(t_1, x_1) - u(t_2, x_2)| \leq c_1 (R/T'^{1/2})^\beta \sup_{Q_+(x_0, R', T')} u.$$

Local central limit theorem

In the following, we write the Gaussian heat kernel with covariance matrix Σ (which is a positive definite $d \times d$ matrix) as

$$k_t(x) := \frac{1}{\sqrt{(2\pi t)^d \det \Sigma}} \exp\left(-\frac{x \Sigma^{-1} x}{2t}\right).$$

When $G = \mathbb{Z}^d$, $x_0 = 0$ and $d \geq 2$, if we further assume the invariance principle, we can obtain the following local limit theorem.

Proposition 1.7 *Assume Assumption 1.1 and the following;*

$$\text{There exists } c_1 > 0 \text{ such that } \lim_{R \rightarrow \infty} R^{-d} \pi(B(0, R)) = c_1,$$

$$\lim_{n \rightarrow \infty} P^0(n^{-1/2} X_{[nt]} \in H(y, R)) = \int_{H(y, R)} k_t(z) dz, \quad \forall y \in \mathbb{R}^d, R, t > 0,$$

where $H(y, R) = y + [-R, R]^d$. Then there exist $a > 0$ such that for each $T_1, T_2 > 0$ and each $M > 0$, we have

$$\lim_{n \rightarrow \infty} \sup_{|x| \leq M} \sup_{t \in [T_1, T_2]} |n^d p_{n^2 t}^\omega(0, [nx]) - ak_t(x)| = 0,$$

where we write $[x] = ([x_1], \dots, [x_d])$ for $x = (x_1, \dots, x_d) \in \mathbb{R}^d$.

1.2 Part II: Models and results.

In this subsection, we will consider the specific RCM with polynomial lower tail. In Part I, we consider a general weighted graph, but in Part II we consider $G = \mathbb{Z}^d$ and the conductance is nearest neighbor and random.

Let us first define the model precisely (for more information on the RCM, see Biskup [12] or Kumagai [27]). Consider the d -dimensional hypercubic lattice \mathbb{Z}^d and let \mathbb{E}_d denote the set of (unordered) nearest-neighbor pairs, called edges or bonds, i.e. $\mathbb{E}_d = \{\{x, y\} : x, y \in \mathbb{Z}^d, |x - y| = 1\}$. We use the notation $x \sim y$ if $(x, y) \in \mathbb{E}_d$, and $\omega_e = \omega_{xy} = \omega_{yx}$ to denote the random conductance of an edge e . Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the probability space that governs the randomness of the media. We assume $\{\omega_e : e \in \mathbb{E}_d\}$ to be positive and i.i.d.. We define π , CSRW, VSRW, their Laplace operators and heat kernels etc. as in Part I. Note that we have two sources of randomness for the Markov chain: the randomness of the media and the randomness of the Markov chain. In order to clarify the randomness of the media, we often put $\omega \in \Omega$. For example, we denote by $(P_\omega^x, x \in \mathbb{Z}^d)$ the Markov laws induced by the semigroup $P_\omega^t := e^{t\mathcal{L}_\omega}$, and by $p_t^\omega(x, y) = P_\omega^x(X_t = y)/\theta(y)$ the heat kernel. Let E_ω^x be the expectation with respect to P_ω^x . As in the last subsection, we use the same notation for CSRW and VSRW when it is clear which Markov chain we are talking about.

Our purpose is to investigate the effects of fluctuations in the environment on the behavior of the random walk. We shall in particular get bounds on the long time behavior of the return probability $P_\omega^0(X_t = 0)$.

It is well known that when the conductances are bounded and bounded away from 0 (the uniformly elliptic case), then the decay of the return probability obeys a standard power law with exponent $d/2$: there exist constants c and C such that for all t, x and y , then

$$ct^{-d/2} \leq P_\omega^x(X_t = y) \leq Ct^{-d/2},$$

both for CSRW and VSRW. We refer to Delmotte [20].

The first sharp results for non-uniformly elliptic conductances were obtained in Mathieu and Remy [30] in the case of random walks on super-critical percolation clusters. Then conductances are allowed to take two values only, 0 and 1. We assume that $\mathbb{P}(\omega_b > 0) > p_c(d)$, where $p_c(d)$ is the critical threshold for bond percolation on \mathbb{Z}^d and we condition on the event that the origin belongs to the infinite cluster of positive conductances. Mathieu and Remy showed that there exists a constant C such that, for almost all realizations of

the conductances, for large enough t , we have

$$\sup_y P_\omega^0(X_t = y) \leq Ct^{-d/2}. \quad (1.7)$$

The results in [30] were later improved in [4].

Quite often in statistical mechanics, results in percolation help understanding more general situations through comparison arguments; the present paper is no exception.

The bounds on the return probability in the percolation case eventually lead to the proof of functional central limit theorems and local C.L.T. . We refer to Sidoravicius and Sznitman [31], Berger and Biskup [10] and Mathieu and Piatnitski [29] for the percolation model, and Barlow and Hambly [9] for the local C.L.T., and to Mathieu [28], Biskup and Prescott [14], Barlow and Deuschel [7], Andres, Barlow, Deuschel and Hambly [1] for more general models of random conductances.

In the other direction, examples show that a slow decay of the return probability is possible for random positive conductances. In Fontes and Mathieu [23], the authors computed the annealed return probability for a model of random walk with positive conductances whose law has a power tail near 0. They showed a transition from a classical decay like $t^{-d/2}$ to a slower decay. In [11], Berger, Biskup, Hoffman and Kozma proved that for $d \geq 5$, given any sequence $\lambda_n \uparrow \infty$, there exists a product law \mathbb{P} on $(0, \infty)^{\mathbb{E}_d}$ such that

$$P_\omega^0(X_{n_k} = 0) \geq c(\omega)(\lambda_{n_k} n_k)^{-2}$$

along a deterministic sequence (n_k) , with $c(\omega) > 0$ almost surely. In this construction, although the conductances are almost surely positive, their law has a very heavy tail near 0 of the form $\mathbb{P}(\omega_{xy} < s) \sim |\log(s)|^{-\theta}$, $\theta > 0$. (Here we write $f \sim g$ to mean that $f(t)/g(t) = 1 + o(1)$ for functions f and g .)

One may then ask for what choice of \mathbb{P} does the transition from a classical decay with rate $t^{-d/2}$ to a slower decay happens. A partial answer to this question is in the papers of Boukhadra [16]–[17].

Let us consider positive and bounded conductances, with a power-law tail near zero: let $\gamma > 0$ and assume the following conditions : for any $e \in \mathbb{E}_d$,

$$\omega_e \in [0, 1], \quad \mathbb{P}(\omega_e \leq u) = u^\gamma(1 + o(1)), \quad u \rightarrow 0. \quad (\text{P})$$

It is proved in Boukhadra [17] that, when (P) is satisfied with $\gamma > d/2$, then

$$P_\omega^0(X_t = 0) = t^{-\frac{d}{2} + o(1)}, \quad (1.8)$$

for almost all environments and as t tends to $+\infty$.

On the other hand, it is proved in [16] that, still for an environment satisfying (P), then

$$P_\omega^0(X_n = 0) \geq C(\omega) n^{-(2+\delta)}, \quad (1.9)$$

where $\delta = \delta(\gamma)$ is a constant such that $\delta(\gamma) \rightarrow 0$ as $\gamma \rightarrow 0$.

The next theorem improves upon (1.8) in two respects: first we have extended the domain of admissible values of γ ; secondly and more importantly, we obtain a much sharper upper bound on the return probability, to be compared with (1.7).

In this subsection, we use an equivalent and more appropriate definition of the box:

$$B(x, n) = x + [-n, n]^d \cap \mathbb{Z}^d$$

for all $x \in \mathbb{Z}^d$ and write $B_n = B(0, n)$.

Theorem 1.8 *Let $d \geq 2$ and suppose that the conductances $(\omega_e, e \in \mathbb{E}_d)$ are i.i.d. satisfying (P). Then we have :*

(1) *For the CSRW, for any $\gamma > \frac{1}{8} \frac{d}{d-1/2}$, there exist positive constants $\delta, c_1 > 0$ such that \mathbb{P} -a.s. for all $x, y \in B(0, t^{(1+\delta)/2})$ and for t large enough,*

$$p_t^\omega(x, y) \leq c_1 t^{-d/2}. \quad (1.10)$$

(2) *For the VSRW for any $\gamma > 1/4$, there exist positive constants $\delta', c_2 > 0$ such that \mathbb{P} -a.s. for all $x, y \in B(0, t^{(1+\delta')/2})$ and for t large enough,*

$$p_t^\omega(x, y) \leq c_2 t^{-d/2}. \quad (1.11)$$

Using the results in Part I, we obtain the following.

Theorem 1.9 *Let $\gamma > \frac{1}{8} \frac{d}{d-1/2}$ for CSRW and $\gamma > 1/4$ for VSRW. Then the conclusions of Proposition 1.4 (Heat kernel lower bound), Theorem 1.5 (Parabolic Harnack inequality), Corollary 1.6 (Hölder continuity of caloric functions) and Proposition 1.7 (Local central-limit theorem) hold.*

Remarks 1.10 (1) Let us discuss in what sense the statements in Theorem 1.8 are optimal.

The restrictions on the value of γ are related to trapping effects on the random walk induced by fluctuations of the conductances. These trapping effects depend on the model, CSRW or VSRW.

The CSRW cannot be trapped on a site but it might be trapped on an edge. Indeed, assume there exists in B_n and edge $e = \{x, y\}$ of conductance of order 1 that is surrounded by edges of conductances of order $n^{-\mu}$ for some $\mu > 0$.

Starting at x , the random walk will oscillate between x and y for a time of order n^μ . If we insist that $p_t^\omega(x, x) \leq c_1 t^{-d/2}$ when t is of order n^2 , as in part (1) of Theorem 1.8, this imposes $\mu < 2$.

It is not difficult to see that, under assumption (P), there will \mathbb{P} -a.s exist edges of conductance of order 1 that are surrounded by edges of conductances smaller than $n^{-\mu}$ for all μ such that $\mu\gamma(4d-2) < d$. Thus we deduce that it is not correct that $p_t^\omega(x, x)$ decays faster than $t^{-d/2}$ uniformly on the box $B_{\sqrt{t}}$ when $\gamma < \frac{1}{8} \frac{d}{d-1/2}$.

The VSRW may be trapped on a point: let x be such that all edges containing x have conductances of order $n^{-\mu}$. Then the VSRW will wait for a time of order n^μ before its first jump. Thus the estimate $p_t^\omega(x, x) \leq c_1 t^{-d/2}$ when t is of order n^2 cannot hold unless $\mu < 2$. It is easy to deduce from that fact that statement (1.11) in part (2) of Theorem 1.8 is false when $\gamma < 1/4$.

(2) For $d = 2$ or $d = 3$, the return probability $P_\omega^0(X_t = 0)$ a.s. decays like $t^{-d/2}$ even when our restrictions on γ are not satisfied (in fact for any choice of i.i.d. positive conductances) as was proved in [11].

(3) One may also compare our estimates with the results in [2]. In [2], the authors consider stationary environments of random conductances under some integrability conditions. When applied to i.i.d. conductances satisfying (P), they obtain heat kernel upper bounds as in Theorem 1.8 provided that $\gamma > 1/4$ for both models CSRW and VSRW. (See [2, Proposition 6.3] for CSRW. The same argument also works for VSRW, see the discussion in [2, Remark 1.5].)

Thus statement (2) in Theorem 1.8 is not new but statement (1) improves upon [2]. Observe also that our strategy strongly differs from the one in [2]. The authors of [2] first establish elliptic and parabolic Harnack inequalities from Sobolev inequalities, and then deduce heat kernel bounds. We approach the problem the other way around: we shall first establish Theorem 1.8 using probabilistic arguments (in particular percolation estimates) and deduce the Harnack inequality from Theorem 1.8.

The organization of the paper is as follows. The proofs of the results in Part I and II are given in Sections 2 and 3 respectively. The key tool in the proof of Theorem 1.8 (the main theorem in Part II) is Proposition 3.2, and its proof is given in Section 6. The proof of Proposition 3.2 requires some preliminary percolation results and spectral gap estimates, which are given in Section 4 and 5 respectively. Some relatively standard proof is given in Appendix (Section 7) for completeness.

2. Proof of the results in Part I

In the following three sections, we prove results in Part I. We first give a preliminary lemma.

Lemma 2.1 *(i) Assume Assumption 1.1 (i), (iv). Then there exists $c_1 > 0$ and $R_7(x_0) \in [1, \infty)$ such that*

$$E^y[\tau_{B(x,r)}] \leq c_1 r^2, \quad (2.1)$$

for all $x \in B(x_0, r^{1+\delta}/2)$, all $y \in G$ and all $r \geq R_7(x_0)$.

(ii) Assume Assumption 1.1 (i), (ii), (iv). Then there exist $c_2 > 0, p \in (0, 1)$ and $R_8(x_0) \in [1, \infty)$ such that

$$P^x(\tau_{B(x,r)} \leq t) \leq p + c_2 t/r^2, \quad (2.2)$$

for all $x \in B(x_0, r^{1+\delta})$, $t \geq 0$ and all $r \geq R_8(x_0)$.

Proof. (i) Let $R_7(x_0) := T_0^{1/2}(x_0) \vee R_2(x_0)$. For $R > R_7(x_0)$ and $x \in B(x_0, R^{1+\delta})$, if $y, z \in B(x, R)$ and $t = c_* R^2$ where $c_* \geq 4$ is chosen later, we have $x, y, z \in B(x_0, 2R^{1+\delta}) \subset B(x_0, t^{(1+\delta)/2})$ and $t \geq T_0$. Thus, by Assumption 1.1 (i), (iv), we have

$$P^y(X_t \in B(x, R)) = \sum_{z \in B(x, R)} p(t, y, z) \theta(z) \leq c_1 t^{-d/2} \theta(B(x, R)) \leq c_1 c_4 t^{-d/2} R^d \leq \frac{1}{2},$$

where we chose $c_*^{d/2} \geq 2c_1c_4$. This implies

$$P^y(\tau_{B(x,R)} > t) \leq \frac{1}{2}.$$

By the Markov property, for m a positive integer

$$P^y(\tau_{B(x,R)} > (m+1)t) \leq E^y[P^{Y_{mt}}(\tau_{B(x,R)} > t) : \tau_{B(x,R)} > mt] \leq \frac{1}{2}P^y(\tau_{B(x,R)} > mt).$$

By induction,

$$P^y(\tau_{B(x,R)} > mt) \leq 2^{-m},$$

and we obtain $E^y[\tau_{B(x,R)}] \leq cR^2$. When $y \notin B(x,R)$, clearly $E^y[\tau_{B(x,R)}] = 0$, so the result follows.

(ii) Write $\tau = \tau_{B(x,r)}$. Using (i) and Assumption 1.1 (ii), we have

$$c_2r^2 \leq E^x[\tau] \leq t + E^x[1_{\{\tau > t\}}E^{X_t}[\tau]] \leq t + cr^2P^x(\tau > t) \leq t + cr^2(1 - P^x(\tau \leq t)),$$

for $x \in B(x_0, r^{1+\delta})$, $r \geq R_0(x_0) \vee R_7(x_0) =: R_8(x_0)$. Rewriting, we have

$$P^x(\tau \leq t) \leq 1 - c_2/c + t/(cr^2),$$

and (2.2) is proved. \square

The following lemma is from [6, Lemma 1.1].

Lemma 2.2 *Let $\{\xi_i\}_{i=1}^m, H$ be non-negative random variables such that $H \geq \sum_{i=1}^m \xi_i$. If the following holds for some $p \in (0, 1)$, $a > 0$,*

$$P(\xi_i \leq t | \sigma(\xi_1, \dots, \xi_{i-1})) \leq p + at, \quad t > 0,$$

then

$$\log P(H \leq t) \leq 2(amt/p)^{1/2} - m \log(1/p).$$

Given Lemma 2.1, we have the following.

Proposition 2.3 *Assume Assumption 1.1 (i), (ii), (iv), and let $\varepsilon \in (0, \delta/(1+\delta))$. Then, there exist $c_1, c_2, c_3 > 0$ such that the following holds for $\rho, t > 0$ that satisfy $\rho^{2-\varepsilon} \leq t$ and $t/\rho \geq c_1R_8(x_0)$;*

$$P^x(\tau_{B(x,\rho)} \leq t) \leq c_2 \exp(-c_3\rho^2/t), \quad \text{for all } x \in B(x_0, \rho). \quad (2.3)$$

Proof. The following argument has been often made for heat kernel upper bounds on fractals. We closely follow [4, Proposition 3.7].

Let $r = \lfloor \rho/m \rfloor \geq 1$ where $m \in \mathbb{N}$ is chosen later. Define inductively

$$\sigma_0 = 0, \quad \sigma_i = \inf\{t > \sigma_{i-1} : d(X_{\sigma_{i-1}}, X_t) = r\}, \quad r \geq 1.$$

Let $\xi_i = \sigma_i - \sigma_{i-1}$ and let $\mathcal{F}_t = \sigma(X_s : s \leq t)$ be the filtration of X . By Lemma 2.1, we have

$$P^x(\xi_i < u | \mathcal{F}_{\sigma_{i-1}}) \leq p + c_1u/r^2 \quad (2.4)$$

if $X_{\sigma_{i-1}} \in B(x_0, r^{1+\delta})$, $r \geq R_8(x_0)$ and $u \geq 0$. Note that $d(x, X_{\sigma_m}) = d(X_0, X_{\sigma_m}) \leq mr \leq \rho$ so that $\sigma_m \leq \tau_{B(x, \rho)}$ and $X_{\sigma_i} \in B(x, \rho)$ for $i = 0, 1, \dots, m$. Using Lemma 2.2 with $a = c_1/r^2$, we obtain

$$\begin{aligned} \log P^x(\tau_{B(x, \rho)} \leq t) &\leq \log P^x(\sigma_m \leq t) \leq 2(c_1 m t / (p r^2))^{1/2} - m \log(1/p) \\ &\leq -c_2 m (1 - (c_3 t m / \rho^2)^{1/2}) \end{aligned} \quad (2.5)$$

if

$$x \in B(x_0, r^{1+\delta}/2), \rho \leq r^{1+\delta}/2 \text{ and } r \geq R_8(x_0). \quad (2.6)$$

Let $\lambda = \rho^2/(2c_3 t)$. If $\lambda \leq 1$, then (2.3) is immediate by adjusting c_2 in (2.3) appropriately, so we may assume $\lambda > 1$. If we can choose $m \in \mathbb{N}$ with $\lambda/2 \leq m < \lambda$ and (2.5) hold, then we have the desired estimate. So let us now verify the conditions (2.6). Set $m = \lfloor \lambda/2 \rfloor + 1 \in [\lambda/2, \lambda)$; then since $m \geq 1$, we have $r \leq \rho$. By definition, $r = \lfloor \rho/m \rfloor \geq c_4 t / \rho$ for some $c_4 > 0$, so the assumption implies $r \geq c_5 R_8(x_0)$. The assumption $\rho^{2-\varepsilon} \leq t$ and the fact $\varepsilon \in (0, \delta/(1+\delta))$ implies (noting that one can choose $\rho \geq r$ large) $r^{1+\delta} > 2\rho$. Since $x \in B(x_0, \rho)$, we have verified that (2.3) holds. \square

Let $d_\theta(\cdot, \cdot)$ be a metric that satisfies

$$\theta_x^{-1} \sum_y d_\theta(x, y)^2 \omega_{xy} \leq 1 \text{ for all } x \in G, \quad (2.7)$$

and $d_\theta(x, y) \leq 1$ for all $x \sim y \in G$. The following estimates, which are generalizations of [19, Corollary 11, 12], are given in [22, Theorem 2.1, 2.2].

Proposition 2.4 *There exist $c_1, \dots, c_4 > 0$ such that the following hold for $x, y \in G$;*

$$p_t(x, y) \leq \frac{c_1}{\sqrt{\theta_x \theta_y}} \exp\left(-c_2 d_\theta(x, y)^2 / t\right) \text{ for } t > d_\theta(x, y), \quad (2.8)$$

$$p_t(x, y) \leq \frac{c_3}{\sqrt{\theta_x \theta_y}} \exp\left(-c_4 d_\theta(x, y)(1 \vee \log(d_\theta(x, y)/t))\right) \text{ for } t \leq d_\theta(x, y). \quad (2.9)$$

We are now ready to prove Proposition 1.2.

Proof of Proposition 1.2. We first consider CSRW, namely $\theta_x = \pi(x)$. In this case the graph distance $d(\cdot, \cdot)$ satisfies the condition of d_θ in (2.7). Write $D = d(x, y)$ and $R = d(x_0, x)$.

Case 1: Consider first the case $D^{2-\varepsilon} \geq t$. By (1.2), we have $c_1 D \geq R_*(x_0)$, and by (1.3), $R \leq c_1 D$. So

$$d(x_0, y) \leq d(x_0, x) + d(x, y) = R + D \leq (c_1 + 1)D.$$

Substituting $(c_1 + 1)D$ to R in Assumption 1.1 (v), we have $\min_{x \in B(x_0, (c_1+1)D)} \pi(x) \geq c_2 D^{-\kappa}$ if $(c_1 + 1)D \geq R_3(x_0)$, so taking $R_*(x_0) \geq c_1 R_3(x_0)/(c_1 + 1)$ and plugging this into (2.8) and (2.9) gives the desired estimates by noting

$$D^\kappa t^{d/2} \leq D^{\kappa+d(2-\varepsilon)/2} \leq c_3 \exp(c_4 D^\varepsilon) \leq c_3 \exp(c_4 D^2/t), \text{ for } D^{2-\varepsilon} \geq t,$$

with $c_4 > 0$ smaller than $c_2/2$ in (2.8).

Case 2: Consider the case $D^{2-\varepsilon} < t$ and let $\rho = \lfloor D/2 \rfloor + 1$ if $D \geq 1$, $\rho = 0$ if $D = 0$. Note that $d(x_0, y) \leq (2D) \vee (2R)$. By (1.2), $R_*(x_0) \leq c_1 t^{1/(2-\varepsilon)}$. Also by (1.3), $R \leq c_1 t^{1/(2-\varepsilon)}$, so that $d(x_0, y) \leq c_5 t^{1/(2-\varepsilon)} < t^{(1+\delta)/2}$ by the choice of ε . Since $D^{2-\varepsilon} < t$, $(t/2)/\rho > c_6 t^{(1-\varepsilon)/(2-\varepsilon)}$, which is larger than $c_6 (R_*(x_0)/c_1)^{1-\varepsilon}$. So the assumption for Proposition 2.3 is satisfied by choosing $R_*(x_0) \geq c_* R_8(x_0)^{1/(1-\varepsilon)}$ for large $c_* > 0$. Let $A_x = \{z \in G : d(x, z) \leq d(y, z)\}$ and $A_y = G \setminus A_x$. Then

$$\begin{aligned} p_t(x, y) &= P^x(X_t = y, X_{t/2} \in A_y)/\theta_y + P^x(X_t = y, X_{t/2} \in A_x)/\theta_y \\ &= P^x(X_t = y, X_{t/2} \in A_y)/\theta_y + P^y(X_t = x, X_{t/2} \in A_x)/\theta_x =: I + II. \end{aligned} \quad (2.10)$$

Write $\tau = \tau_{B(x, \rho)}$. Then

$$\begin{aligned} I &= P^x(X_t = y, X_{t/2} \in A_y)/\theta_y = P^x(\tau < t/2, X_t = y, X_{t/2} \in A_y)/\theta_y \\ &\leq P^x(1_{\{\tau < t/2\}} P^{X_\tau}(X_{t-\tau} = y))/\theta_y \\ &\leq P^x(\tau < t/2) \sup_{z \in \partial B(x, \rho), s < t/2} p_{t-s}(z, y) \\ &\leq c_7 \sup_{z \in \partial B(x, \rho), s < t/2} p_{t-s}(z, y) \exp(-c_8 D^2/t), \end{aligned}$$

where Proposition 2.3 is used in the last inequality. Noting that $d(x_0, z) \leq R + \rho \leq c_8 t^{1/(2-\varepsilon)} < t^{(1+\delta)/2}$, we obtain $I \leq c_9 t^{-d/2} \exp(-c_8 D^2/t)$. II can be bounded similarly, so that we obtain (1.4).

We next discuss the VSRW case (i.e. $\theta_x = 1$) briefly. In this case the metric $\tilde{d}(\cdot, \cdot)/\sqrt{M}$, where M is the maximum degree of the vertices, is relevant; indeed it satisfies the condition of d_θ in (2.7). So the conclusion (w.r.t. \tilde{d}) holds if (1.2) and (1.3) hold w.r.t. \tilde{d} . Using Assumption 1.1 (v), it is easy to verify that (1.2) and (1.3) w.r.t. d imply (1.2) and (1.3) w.r.t. \tilde{d} . Finally let us deduce (1.4) and (1.5) for d from those for \tilde{d} . When $t \geq \tilde{d}(x, y)^2$, (1.4) is an on-diagonal estimate, so no distance appears there. When $t < \tilde{d}(x, y)^2$, (1.2) for \tilde{d} implies $R_*(x_0) \leq c_1 \tilde{d}(x, y)^{1/(1-\varepsilon)}$, so by taking $(R_*(x_0)/c_3)^{1-\varepsilon} \geq R_4(x_0)$, we can apply Assumption 1.1 (v) (since $\tilde{d}(x, y) \leq d(x, y)$) and deduce (1.4) and (1.5) for d from those for \tilde{d} . Thus the desired estimates are established. \square

Remark 2.5 In order to obtain similar estimates, [1, Theorem 4.5] uses the approach introduced by Grigor'yan for manifolds (see [22, Theorem 1.3] and [18, Theorem 1.2] for the graph setting close to ours). However, it does not seem that our Assumption 1.1 (i) is strong enough to apply the method – compare [1, Corollary 4.3] with our assumption.

Proof of Corollary 1.3. It is easy to check (1.2) and (1.3), so we can apply Proposition 1.2. If $s \geq R$, then the result follows directly from (1.4). If $s < R$, then (1.5) implies

$$p_s(x, y) \leq c_1 \exp\left(-c_2 R(1 \vee \log(R/s))\right) \leq c_1 \exp(-c_2 R) \leq c_3 R^{-d},$$

so the result holds. \square

We next prepare some propositions in order to prove Proposition 1.4. The idea of the proof is based on that of [24, Theorem 3.1].

A function u is said to be harmonic in a set $A \subset \mathbb{Z}^d$ if u is defined in \bar{A} (that consists of all points in A and all their neighbors) and if $\mathcal{L}u(x) = 0$ for any $x \in A$.

As a first step, we should check the elliptic oscillation inequalities. For any nonempty finite set U and a function u on U , denote

$$\text{osc}_U u := \max_U u - \min_U u.$$

Proposition 2.6 *Assume Assumption 1.1 (iii). Then, for any $\varepsilon > 0$, there exists $\sigma = \sigma(\varepsilon, C_E) < 1$ such that, for any $\sigma R > R_1(x_0)$ and for any function u defined in $\bar{B}(x_0, R)$ and harmonic in $B(x_0, R)$, we have*

$$\text{osc}_{B(x_0, \sigma r)} u \leq \varepsilon \text{osc}_{B(x_0, r)} u, \quad \forall r \in (R_1(x_0)/\sigma, R/2]. \quad (2.11)$$

The proof is standard. For completeness, we give the proof in Section 7.

We write

$$\bar{E}(x, R) := \max_{y \in B(x, R)} E^y[\tau_{B(x, R)}].$$

Then, under Assumption 1.1 (i) and (iv), we have the following due to Lemma 2.1:

$$\bar{E}(x, R) \leq CR^2, \quad \forall x \in B(x_0, R), R \geq R_7(x_0). \quad (2.12)$$

The next proposition can be proved similarly as [25, Proposition 11.2]. For completeness, we give the proof in Section 7.

Proposition 2.7 *Assume Assumption 1.1 (iii) and let $R \geq R_1(x_0)$, u be a function on $B(x_0, R)$ satisfying the equation $\mathcal{L}u = f$ with zero boundary condition. Then, for any positive $r < R/2$ with $\sigma r \geq R_1(x_0)$,*

$$\text{osc}_{B(x_0, \sigma r)} u \leq 2(\bar{E}(x_0, r) + \varepsilon \bar{E}(x_0, R)) \max_{B(x_0, R)} |f|, \quad (2.13)$$

where σ and ε are the same as in Proposition 2.6.

We now give some time derivative properties of the heat kernel.

Proposition 2.8 *Let A be a nonempty finite subset of \mathbb{Z}^d .*

(i) *Let f be a function on A .*

$$u_t(x) = P_t^A f(x).$$

Then, for all $0 < s \leq t$,

$$\|\partial_t u_t\|_2 \leq \frac{1}{s} \|u_{t-s}\|_2. \quad (2.14)$$

(ii) *For all $x, y \in A$,*

$$|\partial_t p_t^A(x, y)| \leq \frac{1}{s} \sqrt{p_{2v}^A(x, x) p_{2(t-s-v)}^A(y, y)} \quad (2.15)$$

for all positive t, s, v such that $s + v \leq t$.

(iii) *Under Assumption 1.1 (i), for all x, y we have*

$$|\partial_t p_t^A(x, y)| \vee |\partial_t p_t(x, y)| \leq C t^{-(\frac{d}{2}+1)}, \quad \forall x, y \in B(x_0, t^{(1+\delta)/2}), t \geq T_0(x_0). \quad (2.16)$$

The proof is an easy modification of the corresponding results in [25] for discrete time. For completeness, we give the proof in Section 7.

We are now ready to prove Proposition 1.4.

Proof of Proposition 1.4. Let $\varepsilon < 1/2$ (we will impose some further bounds of ε later). Let $R = (t/\varepsilon)^{1/2}$, $A = B(x_0, R)$ and for any $x \in B(x_0, \varepsilon R) = B(x_0, (\varepsilon t)^{1/2})$, introduce the function

$$u(y) := p_t^A(x, y).$$

First, we claim that $u(x) \geq ct^{-d/2}$ for large $t > 0$. Let $B = B(x, \varepsilon^{1/4}R)$; we choose ε small enough so that $B \subset A$. Using the Schwarz inequality, we have

$$\begin{aligned} p_t^A(x, x) &\geq p_t^B(x, x) \geq \left(\sum_z p_{t/2}^B(x, z) \theta_z \right)^2 / \theta(B) = (1 - P^x(X_t \notin B))^2 / \theta(B) \\ &\geq (1 - P^x(\tau_{B(x, \varepsilon^{1/4}R)} \leq t))^2 / \theta(B) \geq (1 - p - c_6 \varepsilon^{1/2})^2 / \theta(B) \geq c / \theta(B), \end{aligned}$$

where (2.2) is used in the third inequality and we take $\varepsilon > 0$ small enough. (We take R large so that $\varepsilon^{1/4}R \geq R_8(x_0)$.) So, using Assumption 1.1 (iv), the claim follows.

Now let us show that

$$|u(x) - u(y)| \leq \frac{c}{2} t^{-d/2} \quad (2.17)$$

for all $y \in B(x_0, \varepsilon R)$ so that $d(x, y) \leq 2(\varepsilon t)^{1/2}$, which would imply $u(y) \geq (c/2)t^{-d/2}$ and hence prove the desired result.

u is the function on A that solves $\mathcal{L}^A u(y) = \partial_t u(y)$. Noting that $x \in B(x_0, \varepsilon R) \subset B(x_0, R)$, by Proposition 2.8 (iii),

$$\max_{y \in B(x_0, R)} |\partial_t p_t^A(x, y)| \leq C t^{-(\frac{d}{2}+1)}, \quad \text{for large } t. \quad (2.18)$$

By Proposition 2.7, we have, for any $0 < r < R/3$ and for some $\sigma \in (0, 1)$,

$$\operatorname{osc}_{B(x_0, \sigma r)} u \leq 2(\bar{E}(x_0, r) + \varepsilon^2 \bar{E}(x_0, R)) \max_{y \in B(x_0, R)} |\partial_t p_t^A(x, y)|, \quad (2.19)$$

for all $\sigma r \geq R_1(x_0)$ where ε in Proposition 2.7 is now written as ε^2 .

Estimating $\max |\partial_t p_t^A(x, y)|$ by (2.18) and using (2.12), we obtain, from (2.19),

$$\operatorname{osc}_{B(x_0, \sigma r)} u \leq C \frac{r^2 + \varepsilon^2 R^2}{t^{d/2+1}}, \quad \forall x \in B(x_0, r), \quad t, r \text{ large.}$$

Choosing $r = \varepsilon R$ and noting $t = \varepsilon R^2$, we obtain

$$\operatorname{osc}_{B(x_0, \sigma r)} u \leq 2C \varepsilon t^{-d/2} \leq \frac{c}{2} t^{-d/2} \quad (2.20)$$

provided $\varepsilon \leq c/(4C)$, $x \in B(x_0, (\varepsilon t)^{1/2}) = B(x_0, \varepsilon R)$ and t large.

Note that

$$\sigma r = \sigma \varepsilon R = \sigma \varepsilon \left(\frac{t}{\varepsilon} \right)^{1/2} = \sigma \sqrt{\varepsilon} t^{1/2} = \delta_0 t^{1/2},$$

where $\delta_0 = \sigma \varepsilon^{1/2}$. Hence (2.20) implies (2.17), which was to be proved. \square

Let us briefly mention other proof of the results in Part I.

Proof of Theorem 1.5 is given in Section 7.

Proof of Corollary 1.6. Given Theorem 1.5, the proof is standard and similar to the proof of [8, Corollary 4.2]. (Given Theorem 1.5, one can also modify the proof of [2, Proposition 4.6] and [9, Proposition 3.2].) So we omit the proof. \square

Proof of Proposition 1.7. Given Corollary 1.6, the proof is similar to [2, Theorem 1.11] and [9, Theorem 4.2], so we omit it. \square

3. Proof of the results in Part II

3.1 Strategy and proof of Theorem 1.8.

We now discuss the strategy of the proof of Theorems 1.8 and how one compares random walks with random conductances with random walks on percolation clusters.

Choose a threshold parameter $\xi > 0$ such that $\mathbb{P}(\omega_b \geq \xi) > p_c(d)$ where $p_c(d)$ is the threshold percolation cluster. The i.i.d. nature of the probability measure \mathbb{P} ensures that for \mathbb{P} almost any environment ω , there exists a unique infinite cluster in the graph $(\mathbb{Z}^d, \mathbb{E}_d)$, that we denote by $\mathcal{C}^\xi = \mathcal{C}^\xi(\omega)$.

Provided ξ is small enough, the complement of \mathcal{C}^ξ in \mathbb{Z}^d , here denoted by \mathcal{H}^ξ , is a union of finite connected components that we will refer to as *holes*, see Lemma 4.1. Thus, by definition, holes are connected sub-graphs of the grid. Note that holes may contain edges such that $\omega_b \geq \xi$.

Consider the following additive functional :

$$A(t) = \int_0^t \mathbb{1}_{\{X_s \in \mathcal{C}^\xi\}} ds. \quad (3.1)$$

We shall need to make a time change for the process X to bring us back to the situation that we already know, namely random walks on an infinite percolation cluster.

Recall $A(t)$ from (3.1) and let $A^{-1}(t) = \inf\{s; A(s) > t\}$ be its inverse. Define the corresponding time changed process

$$X_t^\xi := X_{A^{-1}(t)},$$

which is obtained by suppressing in the trajectory of X all the visits to the holes.

For the proof of Theorem 1.8, we need the fact that X^ξ behaves in a standard way in almost any realization of the environment ω (see for eg. [28, Lemma 4.1] or [1, Theorem 4.5]). Recall that we use here the box $B_n = [-n, n]^d \cap \mathbb{Z}^d$.

Lemma 3.1 *There exists a constant c_1 such that \mathbb{P} -a.s. and for t large enough,*

$$\sup_y P_\omega^x(X_t^\xi = y) \leq c_1 t^{-d/2}, \quad (3.2)$$

for all $x \in B_t \cap \mathcal{C}^\xi$.

The key tool in the proof of Theorem 1.8 is the following control on the time spent by the process outside \mathcal{C}^ξ .

Call τ_h the exit time of the random walk X from \mathcal{H}^ξ ; if $X_0 \notin \mathcal{H}^\xi$, then $\tau_h = 0$.

Proposition 3.2 (1) *Let $d \geq 2$ and choose $\varepsilon \in (0, 1)$. Then,*

(1) *For the CSRW, for any $\gamma > \frac{1}{8} \frac{d}{d-1/2}$, there exist positive constants δ, σ and c_1, \dots, c_4 such that for $\xi > 0$ small enough, \mathbb{P} -a.s. for all $x \in B(0, t^{(1+\delta)/2})$ and all t large enough, we have*

$$P_\omega^x(A(t) \leq \varepsilon t) \leq c_1 e^{-c_2 t^\sigma}, \quad (3.3)$$

and

$$P_\omega^x(\tau_h \geq t/2) \leq c_3 e^{-c_4 t^\sigma}. \quad (3.4)$$

(2) *For the VSRW for any $\gamma > 1/4$, there exist positive constants δ', σ' and c_5, \dots, c_8 such that for $\xi > 0$ small enough, \mathbb{P} -a.s. for all $x \in B(0, t^{(1+\delta')/2})$ and all t large enough, we have*

$$P_\omega^x(A(t) \leq \varepsilon t) \leq c_5 e^{-c_6 t^{\sigma'}} \quad \text{and} \quad P_\omega^x(\tau_h \geq t/2) \leq c_7 e^{-c_8 t^{\sigma'}}. \quad (3.5)$$

Proof of Theorem 1.8. Let X be the CSRW with conductances satisfying (P) and assume $\gamma > \frac{1}{8} \frac{d}{d-1/2}$. One can follow the same argument for the VSRW with $\gamma > 1/4$ and with the counting measure instead of π .

We start by reproducing here the same reasoning as in [17]. Let $n = t^{(1+\delta)/2}$ with δ as in Proposition 3.2 and such that $\delta < 1$. Assume first that x belongs to $\mathcal{C}^\xi \cap B_n$. Since the probability of return is decreasing, see for eg. [17, Lemma 3.1], we have

$$P_\omega^x(X_t = x) \leq \frac{2}{t} \int_{t/2}^t P_\omega^x(X_v = x) dv = \frac{2}{t} E_\omega^x \left[\int_{t/2}^t \mathbb{1}_{\{X_v = x\}} dv \right]. \quad (3.6)$$

The additive functional $A(\cdot)$ being a continuous increasing function of the time and null outside the support of the measure $dA(v)$, so by operating a change of variable by setting $u = A(v)$, we get

$$\begin{aligned} E_\omega^x \left[\int_{t/2}^t \mathbb{1}_{\{X_v = x\}} dv \right] &= E_\omega^x \left[\int_{t/2}^t \mathbb{1}_{\{X_v = x\}} \mathbb{1}_{\{X_v \in \mathcal{C}^\xi\}} dv \right] \\ &= E_\omega^x \left[\int_{A(t/2)}^{A(t)} \mathbb{1}_{\{X_u^\xi = x\}} du \right], \end{aligned}$$

which is bounded by

$$E_\omega^x \left[\int_{A(t/2)}^t \mathbb{1}_{\{X_u^\xi = x\}} du \right],$$

since $A(t) \leq t$.

Therefore, for $\varepsilon \in (0, 1)$

$$\begin{aligned} P_\omega^x(X_t = x) &\leq \frac{2}{t} E_\omega^x \left[\int_{A(t/2)}^t \mathbb{1}_{\{A(t/2) \geq \varepsilon t/2\}} \mathbb{1}_{\{X_u^\xi = x\}} du \right] \\ &\quad + \frac{2}{t} E_\omega^x \left[\int_{A(t/2)}^t \mathbb{1}_{\{A(t/2) \leq \varepsilon t/2\}} \mathbb{1}_{\{X_u^\xi = x\}} du \right] \\ &\leq \frac{2}{t} \int_{\varepsilon t/2}^t P_\omega^x(X_u^\xi = x) du + \frac{2}{t} \int_0^t P_\omega^x(A(t/2) \leq \varepsilon t/2) du, \end{aligned}$$

and using Lemma 3.1,

$$\begin{aligned} P_\omega^x(X_t = x) &\leq \frac{2c_1}{t} \int_{\varepsilon t/2}^t u^{-d/2} du + 2P_\omega^x(A(t/2) \leq \varepsilon t/2) \\ &\leq 2c_1(1 - (\varepsilon/2)^{1-d/2}) t^{-d/2} + 2P_\omega^x(A(t/2) \leq \varepsilon t/2), \end{aligned} \quad (3.7)$$

which by virtue of Proposition 3.2 for t large enough, is less than

$$c_2 t^{-d/2} + 2c_3 e^{-c_4 t^\sigma}.$$

Since $\pi(x) > \xi$, we obtain that

$$p_t^\omega(x, x) \leq c_5 t^{-d/2}.$$

Then Cauchy-Schwarz gives

$$p_t^\omega(x, y) \leq \sqrt{p_t^\omega(x, x) p_t^\omega(y, y)} \leq c_6 t^{-d/2}, \quad (3.8)$$

for any $x, y \in B(0, t^{(1+\delta)/2}) \cap \mathcal{C}^\xi$ and all t large enough.

Recall $n = t^{(1+\delta)/2}$. Suppose $x \in \mathcal{H}^\xi \cap B_n$ and $y \in \mathcal{C}^\xi \cap B_n$. Note that x belongs to a hole with a size less than $(\log n)^c$ included in B_{2n} (see Lemma 4.1 below). It implies that $X_{\tau_h} \in \mathcal{C}^\xi \cap B_{2n}$ if $X_0 = x$. Then the strong Markov property gives

$$P_\omega^x(X_t = y) \leq P_\omega^x(\tau_h > t/2) + E_\omega^x \left(\mathbb{1}_{\{\tau_h \leq t/2\}} P_\omega^{X_{\tau_h}}(X_{t-\tau_h} = y) \right) \quad (3.9)$$

which, by (3.8) and (3.4), and for t large enough, is less than

$$c_3 e^{-c_4 t^{\sigma(1+\delta)/2}} + \max_{z \in \mathcal{C}^\xi \cap B_{2n}} \sup_{s \in [t/2, t]} P_\omega^z(X_s = y) \leq c_7 t^{-d/2} \pi(y). \quad (3.10)$$

Since $\pi(y) \geq \xi$, we deduce that

$$p_t^\omega(x, y) \leq c_8 t^{-d/2}. \quad (3.11)$$

Using the reversibility, we also deduce that

$$p_t^\omega(x, y) \leq c_9 t^{-d/2} \quad (3.12)$$

whenever $y \in \mathcal{H}^\xi \cap B_n$ and $x \in \mathcal{C}^\xi \cap B_n$.

Last, suppose $x, y \in \mathcal{H}^\xi \cap B_n$. The strong Markov property yields

$$\frac{P_\omega^x(X_t = y)}{\pi(y)} \leq \frac{P_\omega^x(\tau_h > t/2)}{\pi(y)} + \frac{1}{\pi(y)} P_\omega^x \left(\mathbb{1}_{\{\tau_h \leq t/2\}} P_\omega^{X_{\tau_h}}(X_{t-\tau_h} = y) \right), \quad (3.13)$$

which by (3.4) and (3.12) is less than

$$\frac{c_3}{\pi(y)} e^{-c_4 t^{\sigma(1+\delta)/2}} + \max_{z \in \mathcal{H}^\xi \cap B_{2n}} \sup_{s \in [t/2, t]} p_s(z, y) \leq \frac{c_3}{\pi(y)} e^{-c_4 t^{\sigma(1+\delta)/2}} + c t^{-d/2}. \quad (3.14)$$

Since $1/\pi(y) \leq n^c$ with a constant c depending only on d and γ (cf. Lemma 4.4 below.), the claim follows. \square

The proof of Proposition 3.2 is deferred to Section 6. Section 4 contains some preliminary percolation results, followed by Section 5, which provides some spectral gap estimates necessary to the proof of the proposition.

Although the main strategy is close to the argument in Boukhadra [17], note that the spectral gap estimates we prove here are sharper and their proof involves a much more detailed analysis of the geometry of the percolation cluster.

3.2 Proof of Theorem 1.9.

Proof of Theorem 1.9. It is enough to check Assumption 1.1 with $x_0 = 0$ and the hypothesis in Proposition 1.7. (1.1) is a consequence of Theorem 1.8. Assumption 1.1 (ii) holds since it is true for the time changed process X^ξ as in [1, Proposition 4.7]. (H) is proved in [1, Theorem 7.3]. Note that VSRW and CSRW share the same harmonic functions, so this fact can be used both of them. Assumption 1.1 (iv) will be proved in Lemma 4.5 for the CSRW case (it is trivial for the VSRW case because the reference measure is a uniform measure). Assumption 1.1 (v) for CSRW case is true because of Lemma 4.4 below. We have

$$\min_{x \in B_n} \pi(x) \geq n^{-\kappa} \quad \text{with} \quad \kappa > \frac{d}{\gamma},$$

where γ is the parameter that we see in the law of the environment (P). Assumption 1.1 (ii), (v) for VSRW case is obvious in this case because $\tilde{d}(\cdot, \cdot) = d(\cdot, \cdot)$ in our setting since $\omega_e \leq 1$ for each edge.

The first hypothesis in Proposition 1.7 holds by the law of large numbers, and the second hypothesis is proved in [14, Theorem 2.1] and [28, Theorem 1.3]. \square

4. Percolation

This section contains percolation results necessary to the spectral gap estimates in the following section.

We consider the standard Bernoulli percolation model on the grid \mathbb{Z}^d : we independently assign to edges the value 1 (open) and 0 (closed) with probability p and $q = 1 - p$. Let \mathbb{P} denote the product probability measure thus defined on $\{0, 1\}^{\mathbb{E}^d}$. We assume p is supercritical so that, for \mathbb{P} almost any environment ω , there exists a unique infinite open

cluster that we denote by \mathcal{C} . For p small enough, the complement of \mathcal{C} in \mathbb{Z}^d , denoted by \mathcal{H} , is a union a finite open clusters that are called *holes*.

Let $x \in \mathbb{Z}^d$ and let \mathcal{H}_x be the (possibly empty) set of sites in the finite component of $\mathbb{Z}^d \setminus \mathcal{C}$ containing x .

Lemma 4.1 *Let $d \geq 2$. For p sufficiently close to 1, then there exist constants $C < \infty$ and $c > 0$ such that for all $n \geq 1$*

$$\mathbb{P}(\text{diam } \mathcal{H}_0 > n) \leq C e^{-cn}.$$

Here “diam” is the diameter in the $|\cdot|_\infty$ -distance on \mathbb{Z}^d .

Proof. See Lemma 3.1 in [14]. □

Recall $B_n = [-n, n]^d \cap \mathbb{Z}^d$ the ball in \mathbb{Z}^d centered at 0 and of radius n . We have the following lemma on the proportion of sites belonging to \mathcal{C} in a box B_n .

Lemma 4.2 *Let $\eta \in (0, 1)$. For p sufficiently close to 1, there exists constants $C < \infty$ and $c > 0$ such that for all $n \geq 1$*

$$\mathbb{P}(|B_n \cap \mathcal{C}| \leq \eta |B_n|) \leq C e^{-cn}. \quad (4.1)$$

This estimate which is sufficient for us is probably not optimal. The expected behavior would be an exponential decay in the perimeter of B_n as in dimension 2, [21, Theorem 3].

Proof. Let $\theta_d(p)$ be the bond percolation probability in the grid \mathbb{Z}^d . Note that $\theta_d(p)$ tends to 1 when $p \rightarrow 1$ [cf. [26], Section 1.4]. Call $\mathcal{C}(\mathbb{G})$ the infinite percolation cluster of a (sub) graph $\mathbb{G} \subseteq \mathbb{Z}^d$.

First note that \mathbb{P} -a.s.

$$S_n := \sum_{x \in B_n} \mathbb{1}_{\{x \in \mathcal{C}\}} \geq \sum_{-n \leq \ell \leq n} \sum_{x \in \{\ell\} \times [-n, n]^{d-1}} \mathbb{1}_{x \in \mathcal{C}(\{\ell\} \times \mathbb{Z}^{d-1})} =: \sum_{-n \leq \ell \leq n} S_n(\ell). \quad (4.2)$$

Then repeating the operation we get

$$S_n \geq \sum_{-n \leq \ell_1, \dots, \ell_{d-2} \leq n} S_n(\ell_1, \dots, \ell_{d-2}) \quad (4.3)$$

with

$$S_n(\ell_1, \dots, \ell_{d-2}) := \sum_{x \in \prod_{i=1}^{d-2} \{\ell_i\} \times [-n, n]^2} \mathbb{1}_{x \in \mathcal{C}(\{\ell_1\} \times \dots \times \{\ell_{d-2}\} \times \mathbb{Z}^2)}.$$

The sub-graphs $\{\ell_1\} \times \dots \times \{\ell_{d-2}\} \times \mathbb{Z}^2$ are disjoint copies of \mathbb{Z}^2 in \mathbb{Z}^d .

Now set

$$Y(\ell_1, \dots, \ell_{d-2}) := S_n(\ell_1, \dots, \ell_{d-2}) / (2n + 1)^2.$$

Let $\eta \in (0, 1)$ and choose p sufficiently close to 1 such that $\eta \in (0, \theta_2(p))$. By [21, Theorem 3], for any $\ell_1, \dots, \ell_{d-2} \in [-n, n]$ and for some $c, C > 0$, we have

$$\mathbb{P}(Y(\ell_1, \dots, \ell_{d-2}) \leq \eta) \leq Ce^{-cn}. \quad (4.4)$$

Combined with (4.3), it implies that

$$\begin{aligned} \mathbb{P}(|B_n \cap \mathcal{C}| \leq \eta |B_n|) &\leq \mathbb{P}\left(\sum_{-n \leq \ell_1, \dots, \ell_{d-2} \leq n} Y(\ell_1, \dots, \ell_{d-2}) / (2n+1)^{d-2} \leq \eta\right) \\ &\leq \mathbb{P}\left(\bigcup_{-n \leq \ell_1, \dots, \ell_{d-2} \leq n} \{Y(\ell_1, \dots, \ell_{d-2}) \leq \eta\}\right) \\ &\leq C n^{d-2} e^{-cn}, \end{aligned}$$

which gives (4.1). \square

Write $\mathcal{C}(x)$ for the open cluster containing the point x . Then we have:

Lemma 4.3 *For q small enough, there exists a constant $c_1 > 1$ such that*

$$\mathbb{P}(|\mathcal{C}(0)| < \infty) \leq c_1 q^{2d}, \quad (4.5)$$

and, for all $x \sim 0$,

$$\mathbb{P}(|\mathcal{C}(0)| < \infty \text{ and } |\mathcal{C}(x)| < \infty) \leq c_1 q^{4d-2}. \quad (4.6)$$

Proof. Let us recall some necessary definitions that we can find in [26], Section 1.4. Call a *plaquette* any unit $(d-1)$ -dimensional hypercube in \mathbb{R}^d that is a face of a cube of the form $x + [-\frac{1}{2}, \frac{1}{2}]^d$. Let \mathbb{L}_d be the set of plaquettes. There is a one to one correspondence between edges in \mathbb{E}_d and plaquettes in \mathbb{L}_d . Indeed, for any edge $\{x, y\} \in \mathbb{E}_d$, the segment $[x, y]$ intersects one and only one plaquette.

We couple the percolation process on \mathbb{E}_d with a percolation on \mathbb{L}_d by declaring a plaquette *open* when the corresponding edge is open and declaring it is *closed* otherwise.

Let us suppose that $\mathcal{C}(0)$ is finite. Then there exists a finite *cutset* of closed plaquettes, say ϖ , around the origin. (A cutset around the origin is a connected set of plaquettes \mathbf{c} such that the origin lies in a finite connected component of the complement of \mathbf{c} .)

The number of such cutsets around the origin which contain m plaquettes is at most μ^m , for some constant $\mu = \mu(d)$ depending only on the dimension. The smallest cutset is unique and contains $2d$ plaquettes. Then the usual ‘Peierls argument’ gives that the probability on the left hand side in (4.5) is bounded by

$$\sum_{\varpi, \text{ cutset around } 0} \mathbb{P}(\text{all plaquettes in } \varpi \text{ are closed}) \leq \sum_{m \geq 2d} (\mu q)^m,$$

which converges and is bounded by cq^{2d} for some c provided p is sufficiently close to 1 such that $q\mu < 1$.

As for the second estimate (4.6), we follow the same argument but we find the exponent $4d-2$ since this is the size of the smallest number of plaquettes necessary to form a cutset around both the origin and x . \square

We now describe application of the preceding Lemmas to conductances satisfying assumption **(P)**.

We recall the following result. Call \mathbb{B}_n the set of edges in the box B_n .

Lemma 4.4 *Suppose that the conductances $(\omega_e, e \in \mathbb{E}_d)$ satisfy **(P)**. Then \mathbb{P} -a.s., we have*

$$\lim_{n \rightarrow \infty} \frac{\log \inf_{e \in \mathbb{B}_n} \omega_e}{\log n} = -\frac{d}{\gamma}.$$

Proof. The proof is similar to [23, Lemma 3.6]. \square

The density estimate Lemma 4.2 yields the following volume property for the measure θ .

Lemma 4.5 *Let $\eta \in (0, 1)$ and $\beta \geq 1$. Let \mathbb{P} be a product probability measure satisfying **(P)**. Then for $\xi > 0$ small enough, for \mathbb{P} -a.e. environment, for all $x \in B_{n^\beta}$ and n large enough, we have*

$$\xi \eta |B_n| \leq \pi(B(x, n)) \leq 2d |B_n|. \quad (4.7)$$

Proof. Let $\eta \in (0, 1)$. Recall the infinite cluster \mathcal{C}^ξ introduced in subsection 3.1. The right-hand side inequality in (4.7) comes from the fact that $\pi(x) \leq 2d$. As for the left-hand side inequality, observe that by (4.1) and the i.i.d. character of the conductances,

$$\mathbb{P}\left(\bigcup_{x \in B_{n^\beta}} \{|\mathcal{C}^\xi \cap B(x, n)| < C\eta |B_n|\}\right) \leq |B_{n^\beta}| e^{-cn}. \quad (4.8)$$

By the Borel-Cantelli lemma, we get that for n large enough, for all $x \in B_{n^\beta}$, we have $|\mathcal{C}^\xi \cap B(x, n)| \geq \eta |B_n|$. Since $\pi(x) \geq \xi$ for $x \in \mathcal{C}^\xi$, the claim follows. \square

In next two lemmas we construct sets of ‘good’ paths in the percolation clusters.

Lemma 4.6 *Let \mathbb{P} be a product probability measure satisfying **(P)**.*

(1) *Let $\gamma > \frac{1}{8} \frac{d}{d-1/2}$ and choose $\alpha \in (0, 2)$ such that $\gamma\alpha(4d-2) > d$. For ξ small enough, \mathbb{P} -a.s., for n large enough, for any edge e in B_n , there exists a path of conductances larger than $n^{-\alpha}$ connecting one of the endpoints of e to the frontier ∂B_n .*

(2) *Let $\gamma > 1/4$ and choose $\alpha \in (0, 2)$ such that $\gamma > 1/(2\alpha)$. For ξ small enough, \mathbb{P} -a.s. for n large enough, for any $x \in B_n$, there exists a path of conductances larger than $n^{-\alpha}$ joining x to the frontier ∂B_n .*

Let $\mathcal{H}_n = B_n \cap \mathcal{H}^\xi$ and $\mathcal{C}_n = B_n \cap \mathcal{C}^\xi$.

Lemma 4.7 (1) *Let $\gamma > \frac{1}{8} \frac{d}{d-1/2}$ and choose $\alpha \in (0, 2)$ such that $\gamma\alpha(4d-2) > d$. For ξ small enough, \mathbb{P} -a.s. for n large enough, there exists an injective map φ on \mathcal{H}_n into \mathcal{C}_n such that for all edge $e = \{x, y\}$ with $x \in \mathcal{H}_n$, there exists a path $\ell(e, \varphi(x))$ from one of the endpoints of e to $\varphi(x)$ satisfying*

$$|\ell(e, \varphi(x))| \leq (\log n)^{2d^2} \quad \text{and} \quad \frac{1}{\omega_b} < 4n^\alpha, \quad \forall b \in \ell(e, \varphi(x)). \quad (4.9)$$

(2) Let $\gamma > 1/4$ and choose $\alpha \in (0, 2)$ such that $2\alpha\gamma > 1$. For ξ small enough, \mathbb{P} -a.s. for n large enough, there exists an injective map φ' on \mathcal{H}_n into \mathcal{C}_n such that for all $x \in \mathcal{H}_n$, there exists a path $\ell(x, \varphi'(x))$ from x to $\varphi'(x)$ satisfying

$$|\ell(x, \varphi'(x))| \leq (\log n)^{2d^2} \quad \text{and} \quad \frac{1}{\omega_b} < 4n^\alpha, \quad \forall b \in \ell(x, \varphi'(x)). \quad (4.10)$$

Proof of Lemma 4.6. (1) Let $\alpha\gamma(4d-2) > d$ for some $\alpha \in (0, 2)$. Recall \mathbb{B}_n the set of edges in the box B_n and set $\partial\mathbb{B}_n = \mathbb{B}_n \setminus \mathbb{B}_{n-1}$. Note that $|\mathbb{B}_n| = O(n^{d-1})$.

Let \mathcal{E}_n be the event: there exists an edge $e \in \partial\mathbb{B}_n$ such that none of its endpoints can be joined by a path to $\partial\mathbb{B}_{n+1}$ along edges with conductances larger than $n^{-\alpha}$; this last event is denoted by $\{e \nleftrightarrow \partial\mathbb{B}_{n+1}\}$, i.e.

$$\mathcal{E}_n := \bigcup_{e \in \partial\mathbb{B}_n} \{e \nleftrightarrow \partial\mathbb{B}_{n+1}\}. \quad (4.11)$$

Then Lemma 4.3 with $q = \mathbb{P}(\omega_e < n^{-\alpha\gamma})$ and (P) imply that

$$\mathbb{P}(\mathcal{E}_n) \leq cn^{-1-(\alpha\gamma(4d-2)-d)}, \quad (4.12)$$

By the Borel-Cantelli lemma we then get that there is a finite positive random variable $N = N(\omega)$ such that for any $n \geq N$, for every edge $e \in \partial\mathbb{B}_n$, there exists a path of conductances larger than $n^{-\alpha}$ joining e to $\partial\mathbb{B}_{n+1}$. It implies that there exists a path of conductances larger than $n^{-\alpha}$ joining one of the endpoints of every edge in $\mathbb{B}_n \setminus \mathbb{B}_{N-1}$ to $\partial\mathbb{B}_{n+1}$. Indeed, consider an edge $f \in \partial\mathbb{B}_m$ for some $m \geq N$. From one of its endpoints starts a path of conductances larger than $n^{-\alpha}$ reaching $\partial\mathbb{B}_{m+1}$. Let $e \in \partial\mathbb{B}_{m+1}$ be the last edge of this path. Observe that the conductance of e is larger than $n^{-\alpha}$. There is a path of conductances larger than $n^{-\alpha}$ starting from one of the endpoints of e and reaching $\partial\mathbb{B}_{m+2}$. But since the conductance of e is larger than $n^{-\alpha}$, there is actually a path of conductances larger than $n^{-\alpha}$ starting from any of the endpoints of e and reaching $\partial\mathbb{B}_{m+2}$. Thus we constructed a path from f to $\partial\mathbb{B}_{m+2}$. Iterating this construction, we obtain a path from one endpoint of f to $\partial\mathbb{B}_{n+1}$.

By Lemma 4.4, all conductances in B_N are greater than N^{-c} for some positive constant c depending on d and γ . We can choose n large enough such that $N^{-c} \geq n^{-\alpha}$, which ensures the existence of a path of conductances larger than $n^{-\alpha}$ from one of the endpoints of $e \in \mathbb{B}_n$ to $\partial\mathbb{B}_{n+1}$.

(2) For the second assertion of the lemma, we can follow the same reasoning with a slight adaptation. Let $\gamma > 1/(2\alpha)$ for some $\alpha \in (0, 2)$. Set $\partial B_n = \{x \in B_n : \exists y \notin B_n \text{ s.t. } x \sim y\}$, the frontier of B_n . As before, define \mathcal{E}_n to be the event: there exists a vertex $x \in \partial B_n$ such that any path from x to the boundary ∂B_{n+1} has at least one edge with conductance less than $n^{-\alpha}$. Then we have by Lemma 4.3 and (P) that

$$\mathbb{P}(\mathcal{E}_n) \leq cn^{-1-d(2\alpha\gamma-1)}. \quad (4.13)$$

The rest of the proof is similar. \square

Proof of Lemma 4.7. First let $\gamma > \frac{1}{8} \frac{d}{d-1/2}$ and choose $\alpha \in (0, 2)$ such that $\gamma\alpha(4d-2) > d$.

Let $m \in \mathbb{N}^*$ and set $B_m(z) = (2m+1)z + B_m$ for $z \in \mathbb{Z}^d$. The family $\{B_m(z)\}_{z \in \mathbb{Z}^d}$ constitutes a partition of \mathbb{Z}^d . Note that $|B_m(z)| \geq m^d$. Then choosing $m = \lfloor (\log n)^{d+1} \rfloor$, Lemma 4.2 and the Borel-Cantelli lemma yield that \mathbb{P} -a.s. for n large enough, the vertices in any $B_m(z)$ with $B_m(z) \cap B_n \neq \emptyset$ belong to \mathcal{C}^ξ with a proportion that approaches 1 when ξ is small enough. We choose ξ small enough such that this proportion is larger than $1/2$. Therefore, for any box $B_m(z)$ that intersects B_n , there are sufficiently many sites in $B_m(z) \cap \mathcal{C}_n$ to associate with sites in $B_m(z) \cap \mathcal{H}_n$ (if any) in an injective way. Let φ be an injective map from \mathcal{H}_n into \mathcal{C}_n such that it associates a site in $B_m(z) \cap \mathcal{H}_n$ to a site in $B_m(z) \cap \mathcal{C}_n$.

Let us now construct the path $\ell(e, \varphi(e))$ for some edge $e = \{x, y\}$ and $x \in B_m(z) \cap \mathcal{H}_n$ with $B_m(z) \cap B_n \neq \emptyset$. By Lemma 4.6 (1), \mathbb{P} -a.s. for n large enough, for any e of B_n , there exists a self-avoiding path, say (x^1, x^2, x^3, \dots) with $x^1 = x$ or y , which reaches from e the boundary of B_{2n} with conductances larger than $(2n)^{-\alpha}$. By Lemma 4.1 together with the Borel-Cantelli lemma, \mathbb{P} -a.s. for n large enough, m is larger than the size of a hole. It follows that there is some $k < m$ such that $x^k \in \mathcal{C}^\xi$. This gives us the first part of the path $\ell(e, \varphi(x))$.

Next we claim that it is possible to join x^k with $\varphi(x) \in B_m(z) \cap \mathcal{C}_n$ through a path on \mathcal{C}^ξ inside $A_{4m}(x^k) := x^k + B_{4m}$ (note that x^k and $\varphi(x)$ belong to $A_{4m}(x^k)$ and that $d(\varphi(x), \partial A_{4m}(x^k)) > m$). Indeed, if we suppose that it is not possible to find such a path, there would exist a closed cutset (as seen in Lemma 4.3) of conductances less than ξ and of diameter at least m separating x^k from $\varphi(x)$ in $A_{4m}(x^k)$. But Lemma 4.1 rules out this possibility since m is larger than the possible diameter of a hole. Therefore, there exists a self-avoiding path from e to $\varphi(x)$ through edges with conductances larger than $(2n)^{-\alpha}$ and of length less than $m + (8m)^d \leq (\log n)^{2d^2}$. Note here that this path may leave the box B_n .

(2) The case for which $\gamma > 1/4$ can be treated identically using the assertion (2) of Lemma 4.6. \square

5. Spectral gaps estimates

We work in $L^2(\theta)$, the Hilbert space of functions on \mathbb{Z}^d with scalar product

$$\langle f, g \rangle = \sum_{x \in \mathbb{Z}^d} f(x)g(x)\theta(x),$$

where $\theta(x) = \pi(x)$ in the CSRW and $\theta(x) = 1$ for the VSRW.

We also define the Dirichlet form

$$\mathcal{E}^\omega(f, f) = \frac{1}{2} \sum_{\{x, y\} \in \mathbb{E}_d} (f(x) - f(y))^2 \omega_{xy}. \quad (5.1)$$

For both models, CSRW or VSRW, then \mathcal{E}^ω is always the Dirichlet of the corresponding random walk when acting on $L^2(\theta)$.

Consider the self-adjoint operator

$$\mathcal{G}_\omega(\lambda) := \mathcal{L}_\theta - \lambda \mathcal{M}_\varphi, \quad (5.2)$$

where $\varphi(x) := \mathbb{1}_{\{x \in \mathcal{C}^\xi\}}$ and \mathcal{M}_φ is the multiplicative operator by the function φ , i.e. $\mathcal{M}_\varphi f(x) = \varphi(x)f(x)$. Let $R_\omega^t(\lambda)$ be the semigroup generated by $\mathcal{G}_\omega(\lambda)$. The Feynman-Kac formula (see [17, Proposition 3.3]) reads

$$R_\omega^t(\lambda)f(x) = E_\omega^x \left(f(X_t) e^{-\lambda A(t)} \right), \quad t \geq 0, x \in \mathbb{Z}^d. \quad (5.3)$$

The semigroup of the operator $\mathcal{G}_\omega(\lambda)$ with Dirichlet boundary conditions outside the box B_n is given by

$$(R_n^t f)(x) := E_\omega^x \left[f(X_t) e^{-\lambda A(t)}; \tau_{B_n} > t \right].$$

Note that the operator $-\mathcal{G}_\omega(\lambda)$ with Dirichlet boundary conditions outside B_n is a non-negative symmetric operator with respect to the restriction of the measure θ to B_n . Let $\{\lambda_i, i \in \{1, \dots, |B_n|\}\}$ be the set of its eigenvalues labelled in increasing order, and $\{\psi_i, i \in \{1, \dots, |B_n|\}\}$ the corresponding eigenfunctions with due normalization.

Then, by the min-max Theorem and (5.2), the eigenvalue λ_1 is given by

$$\lambda_1 = \inf_{f \neq 0} \frac{\mathcal{E}^\omega(f, f) + \lambda \sum_{x \in \mathcal{C}_n} f^2(x) \theta(x)}{\sum_{x \in B_n} f^2(x) \theta(x)}, \quad (5.4)$$

where the infimum is taken over functions f vanishing outside the box B_n . Recall the notation $\mathcal{C}_n = B_n \cap \mathcal{C}^\xi$.

First, we want to prove the following key estimates on λ_1 .

Lemma 5.1 (1) *Let X be the CSRW and take $\gamma > \frac{1}{8} \frac{d}{d-1/2}$. Then there exists $\alpha \in (0, 2)$ such that for sufficiently small ξ , for a.e. environment, we have for n large enough*

$$\lambda_1 \geq n^{-\alpha} \quad (5.5)$$

when we choose $\lambda = (1 + 8d/\xi) n^{-\alpha}$.

(2) *For the VSRW, for any $\gamma > 1/4$, there exists $\alpha \in (0, 2)$ such that for ξ small enough, for a.e. environment, for n large enough,*

$$\lambda_1 \geq n^{-\alpha} \quad (5.6)$$

when we choose $\lambda = 3n^{-\alpha}$.

To obtain bounds for the exit time as in Proposition 3.2, we need to estimate another eigenvalue.

Denote by $\mathcal{L}_{\mathcal{H}_n}$ the generator of the random walk with the vanishing Dirichlet boundary condition on $\mathcal{H}_n = B_n \cap \mathcal{H}^\xi$. The associated semigroup is given by $P_{\mathcal{H}_n}^t = e^{t\mathcal{L}_{\mathcal{H}_n}}$.

The operator $-\mathcal{L}_{\mathcal{H}_n}$ is symmetric with respect to the measure θ and has $|\mathcal{H}_n|$ nonnegative eigenvalues that we enumerate in increasing order and denote as follows:

$$\zeta_1 \leq \zeta_2 \leq \dots \leq \zeta_{|\mathcal{H}_n|}. \quad (5.7)$$

$\{\phi_i, i = 1, \dots, |\mathcal{H}_n|\}$ is the set of the associated normalized eigenfunctions.

The spectral gap ζ_1 admits the variational definition

$$\zeta_1 = \inf_{f \neq 0} \frac{\langle -\mathcal{L}_{\mathcal{H}_n} f, f \rangle}{\langle f, f \rangle} = \inf_{f \neq 0} \frac{\mathcal{E}^\omega(f, f)}{\sum_{x \in \mathcal{H}_n} f(x)^2 \theta(x)}, \quad (5.8)$$

where the infimum is taken over functions f vanishing outside \mathcal{H}_n .

Lemma 5.2 (1) *For the CSRW, for any $\gamma > \frac{1}{8} \frac{d}{d-1/2}$, there exists $\alpha \in (0, 2)$ such that for sufficiently small ξ , for a.e. environment, for n large enough,*

$$\zeta_1 \geq n^{-\alpha}. \quad (5.9)$$

(2) *For the VSRW, for any $\gamma > 1/4$, there exists $\alpha \in (0, 2)$ such that for ξ small enough, for a.e. environment, for n large enough,*

$$\zeta_1 \geq n^{-\alpha}. \quad (5.10)$$

Proof of Lemma 5.1. (1) Let $\gamma > \frac{1}{8} \frac{d}{d-1/2}$; then choose $\alpha' \in (0, 2)$ such that $\gamma\alpha'(4d-2) > d$ and α such that $\alpha' < \alpha < 2$. Let f be a function vanishing outside B_n . We use the notation $df(b) := f(a) - f(c)$ for any edge $b = \{a, c\}$.

Let $x \in \mathcal{H}_n$ and call $e = \{x, y\}$ the edge such that $\omega_e = \max_{b \ni x} \omega_b$.

We use the paths ℓ constructed in Lemma 4.7 to get that

$$f(x) = f(x) - f(y) + \sum_{b \in \ell(e, \varphi(x))} df(b) + f(\varphi(x)), \quad (5.11)$$

if the path $\ell(e, \varphi(x))$ starts at y . Otherwise,

$$f(x) = \sum_{b \in \ell(e, \varphi(x))} df(b) + f(\varphi(x)). \quad (5.12)$$

Let us consider the case (5.11) - the case (5.12) can be treated in the same way. Observe that Cauchy-Schwarz inequality gives

$$f(x)^2 \leq 2(f(x) - f(y))^2 + 4|\ell(e, \varphi(x))| \sum_{b \in \ell(e, \varphi(x))} df(b)^2 + 4f(\varphi(x))^2. \quad (5.13)$$

Noting that $\pi(\varphi(x)) \geq \xi$ and

$$\pi(x) \leq 2d\omega_e \leq 2d, \quad (5.14)$$

we obtain that

$$f(x)^2 \pi(x) \leq 4d(f(x) - f(y))^2 \omega_e + 8d|\ell(e, \varphi(x))| \sum_{b \in \ell(e, \varphi(x))} df(b)^2 + \frac{8d}{\xi} f(\varphi(x))^2 \pi(\varphi(x)).$$

Using the bounds from Lemma 4.7 (that we apply with α' rather than α), we get

$$\begin{aligned} & f(x)^2 \pi(x) \\ & \leq 4d(f(x) - f(y))^2 \omega_e + 32dn^{\alpha'} (\log n)^{2d^2} \sum_{b \in \ell(e, \varphi(x))} df(b)^2 \omega_b + \frac{8d}{\xi} f(\varphi(x))^2 \pi(\varphi(x)). \end{aligned}$$

Let us now sum this inequality for $x \in \mathcal{H}_n$. Observe that: - a given edge appears at most $(\log n)^{2d^3}$ (because of the bound on the length of the path), - a given $\varphi(x)$ only appears at most once. So

$$\sum_{x \in \mathcal{H}_n} f(x)^2 \pi(x) \leq 32dn^{\alpha'} (\log n)^{2d^2} (\log n)^{2d^3} \mathcal{E}^\omega(f, f) + \frac{8d}{\xi} \sum_{x \in \mathcal{C}_n} f(x)^2 \pi(x).$$

Choose n big enough so that $32dn^{\alpha'} (\log n)^{2d^2} (\log n)^{2d^3} \leq n^\alpha$. We have obtained the inequality

$$\sum_{x \in \mathcal{H}_n} f(x)^2 \pi(x) \leq n^\alpha \mathcal{E}^\omega(f, f) + \left(\frac{8d}{\xi}\right) \sum_{x \in \mathcal{C}_n} f^2(x) \pi(x).$$

To conclude, use the variational formula (5.4).

(2) The argument is the same and here we just give an outline of the proof. Let $\gamma > 1/4$, choose $\alpha' \in (0, 2)$ such that $\gamma > 1/(2\alpha')$ and α such that $\alpha' < \alpha < 2$.

Let f be a function vanishing outside B_n .

Let $x \in \mathcal{H}_n$. Then Lemma 4.7 implies

$$f(x) = \sum_{b \in \ell(x, \varphi'(x))} df(b) + f(\varphi'(x)), \quad (5.15)$$

which by Cauchy-Schwarz inequality gives

$$f(x)^2 \leq 2|\ell(x, \varphi'(x))| \sum_{b \in \ell(x, \varphi'(x))} df(b)^2 + 2f(\varphi'(x))^2.$$

Summing over \mathcal{H}_n , note that a given edge appears at most $(\log n)^{2d^3}$ (because of the bound on the length of the path), and a given $\varphi(x)$ only appears once. Thus we obtain

$$\sum_{x \in \mathcal{H}_n} f(x)^2 \leq 2(\log n)^{2d^2} (\log n)^{2d^3} n^{\alpha'} \mathcal{E}^\omega(f, f) + 2 \sum_{x \in \mathcal{C}_n} f(x)^2,$$

and hence

$$\sum_{x \in B_n} f(x)^2 \leq n^\alpha \mathcal{E}^\omega(f, f) + 3 \sum_{x \in \mathcal{C}_n} f^2(x),$$

when n is large enough. The variational formula (5.4) then yields the desired estimate. \square

We pass now to the proof of the second spectral gap ζ_1 .

Proof of Lemma 5.2. The argument is the same as in estimating λ_1 .

(1) Let X be the CSRW and assume that $\gamma > \frac{1}{8} \frac{d}{d-1/2}$.

Suppose $x \in \mathcal{H}_n$ and call $e = \{x, y\}$ the edge such that $\omega_e = \max_{b \ni x} \omega_b$. Let f be a function vanishing outside \mathcal{H}_n . Then thanks to Lemma 4.7, there exists a path $\ell(e, \varphi(x))$ connecting e to a site $\varphi(x) \in \mathcal{C}^\xi \cap B_n$. If this path starts at y , write then

$$f(x) = f(x) - f(y) + \sum_{b \in \ell(e, \varphi(x))} df(b). \quad (5.16)$$

Otherwise, write

$$f(x) = \sum_{b \in \ell(e, \varphi(x))} df(b). \quad (5.17)$$

Consider the case (5.16) and do the same thing for the second one.

By Cauchy-Schwarz inequality, (5.16) gives

$$f(x)^2 \leq 2(f(x) - f(y))^2 + 2|\ell(x, \varphi(x))| \sum_{b \in \ell(e, \varphi(x))} df(b)^2. \quad (5.18)$$

Multiply (5.18) by $\pi(x)$ and use (5.14) and (4.9) to obtain

$$\sum_{x \in \mathcal{H}_n} f(x)^2 \pi(x) \leq 4d \sum_{x \in \mathcal{H}_n} (f(x) - f(y))^2 \omega_e + 8d (\log n)^c n^{\alpha'} \mathcal{E}^\omega(f, f), \quad (5.19)$$

where $\alpha' \in (0, 2)$ is chosen such that $\gamma\alpha'(4d - 2) > d$ and we used again the fact that a given edge appears at most $(\log n)^{2d^3}$ (because of the bound on the length of the path). Thus for $\alpha \in (\alpha', 2)$ and n large enough,

$$\text{R.H.S. of 5.19} \leq n^\alpha \mathcal{E}^\omega(f, f).$$

which, using (5.8), gives the lower bound (5.2).

(2) As for the VSRW, instead of (5.16), we have by Lemma 4.7,

$$f(x) = \sum_{b \in \ell(x, \varphi(x))} df(b).$$

The remainder of the proof is the same. \square

6. Proof of Proposition 3.2

With all the necessary tools in hand, we can finally provide the proof of Proposition 3.2.

Proof of Proposition 3.2. (1) Let X be the CSRW. Take $\gamma > \frac{1}{8} \frac{d}{d-1/2}$ and let $\alpha \in (0, 2)$ be as in Lemma 5.1. Choose $\delta > 0$ such that

$$1 - \alpha \frac{(1 + \delta)}{2} > 0. \quad (6.1)$$

Let $n = t^{(1+\delta)/2}$ and suppose $x \in B_{n/2}$. Observe that for any constant $\lambda > 0$ (may be \mathbb{P} -random) and any $\varepsilon \in (0, 1)$, Chebyshev's inequality gives

$$\begin{aligned} P_\omega^x(A(t) \leq \varepsilon t) &= P_\omega^x(A(t) \leq \varepsilon t; \tau_{B_n} > t) + P_\omega^x(A(t) \leq \varepsilon t; \tau_{B_n} \leq t) \\ &\leq P_\omega^x\left(e^{-\lambda A(t)} \geq e^{-\varepsilon \lambda t}; \tau_{B_n} > t\right) + P_\omega^x(\tau_{B_n} \leq t) \\ &\leq e^{\varepsilon \lambda t} E_\omega^x\left(e^{-\lambda A(t)}; \tau_{B_n} > t\right) + P_\omega^x(\tau_{B_n} \leq t). \end{aligned} \quad (6.2)$$

By [1, Proposition 4.7]), we have for t large enough,

$$P_\omega^x(\tau_{B_n} \leq t) \leq C e^{-ct^\delta}, \quad (6.3)$$

where C, c are numerical constants.

Let us look now at the first term of the right hand side of (6.2). Recall the eigenvalues $\{\lambda_i, i \in \{1, \dots, |B_n|\}\}$ of the restricted operator $-\mathcal{G}_\omega(\lambda)$ and their associated normalized eigenfunctions $\{\psi_i, i \in \{1, \dots, |B_n|\}\}$. For $f = \mathbb{1}_{B_n}$, observe first that

$$(R_n^t f)(x) = E_\omega^x(e^{-\lambda A(t)}; \tau_{B_n} > t) = \sum_{i=1}^{|B_n|} e^{-\lambda_i t} \langle f, \psi_i \rangle \psi_i(x). \quad (6.4)$$

Then

$$(R_n^t f)^2(x) \pi(x) \leq \sum_{y \in B_n} (R_n^t f)^2(y) \pi(y) = \sum_i e^{-2\lambda_i t} \langle f, \psi_i \rangle^2 \leq e^{-2\lambda_1 t} \|f\|_2^2,$$

which is less than

$$2d |B_n| e^{-2\lambda_1 t}.$$

Thus by Lemma 5.1 (choosing $\lambda = cn^{-\alpha}$) and using the fact that $1/\pi(x) \leq n^c$, $c > 0$ being a constant that depends only on d and γ (cf. Lemma 4.4), we obtain

$$e^{\varepsilon \lambda t} E_\omega^x(e^{-\lambda A(t)}; \tau_{B_n} > t) \leq C n^{d+c} e^{-(1-\varepsilon)t^{1-\alpha(1+\delta)/2}}. \quad (6.5)$$

According to (6.1) and since $\varepsilon \in (0, 1)$, for large enough t , we have

$$\text{R.H.S. of (6.5)} \leq c e^{-(1-\varepsilon)t^\sigma} \quad (6.6)$$

for any $\sigma < 1 - \alpha(1 + \delta)/2$. Thus (6.2)–(6.3)–(6.6) give the desired upper bound for any σ small enough.

As for the exit time estimate, suppose $x \in \mathcal{H}_n = B_n \cap \mathcal{H}^\xi$ with $n = t^{(1+\delta)/2}$. Recall the eigenvalues $\{\zeta_i, i \in \{1, \dots, |\mathcal{H}_n|\}\}$ of the restricted operator $-\mathcal{L}_{\mathcal{H}_n}$ and their associated normalized eigenfunctions $\{\phi_i, i = 1, \dots, |\mathcal{H}_n|\}$. Let $f = \mathbb{1}_{\mathcal{H}_n}$ and observe that

$$P_\omega^x(\tau_h > t/2) = P_{\mathcal{H}_n}^{t/2} f(x) = \sum_{i=1}^{|\mathcal{H}_n|} e^{-\zeta_i t/2} \langle f, \phi_i \rangle \phi_i(x) \quad (6.7)$$

which, by Lemmas 5.2 – 4.1, yields that

$$P_\omega^x(\tau_h > t/2) \leq \frac{e^{-\zeta_1 t/2}}{\sqrt{\pi(x)}} \|f\|_2 \leq \frac{|B_n|}{\sqrt{\pi(x)}} e^{-\zeta_1 t/2} \leq n^c e^{-\frac{1}{2}t^{1-\alpha(1+\delta)/2}} \quad (6.8)$$

where we used again that $1/\pi(x) \leq n^c$. The claim follows for any $\sigma < 1 - \alpha(1 + \delta)/2$.

(2) Clearly, the above argument for the CSRW holds for the VSRW with $\gamma > 1/4$ and the counting measure instead of π . \square

7. APPENDIX

Here, we give some relatively standard proof for completeness.

Proof of Proposition 2.6. Fix a large ball $B(x_0, R)$, and denote for simplicity $B_r = B(x_0, r)$. Let us prove that, for any $R_1(x_0) \leq r < R/6$,

$$\operatorname{osc}_{B_r} u \leq (1 - \delta) \operatorname{osc}_{B_{3r}} u, \quad (7.1)$$

where $\delta = \delta(C_E) \in (0, 1)$. Then (2.11) follows from (7.1) by iterating.

The function $u - \min_{B_{3r}} u$ is nonnegative in B_{2r} and harmonic in B_{2r} . Applying Assumption 1.1 (iii) to this function, we obtain

$$\max_{B_r} u - \min_{B_{3r}} u \leq C_E (\min_{B_r} u - \min_{B_{3r}} u),$$

for all $R_1(x_0) \leq r \leq R/6$, so

$$\operatorname{osc}_{B_r} u \leq (C_E - 1) (\min_{B_r} u - \min_{B_{3r}} u).$$

Similarly, we have $\operatorname{osc}_{B_r} u \leq (C_E - 1) (\max_{B_{3r}} u - \max_{B_r} u)$. Summing up these two inequalities, we get

$$(1 + C_E) \operatorname{osc}_{B_r} u \leq (C_E - 1) \operatorname{osc}_{B_{3r}} u,$$

whence (7.1) follows. \square

Proof of Proposition 2.7. Denote for simplicity $B_r = B(x_0, r)$. Let

$$g_{B_R}(x, y) = \int_0^\infty p_t^{B_R}(x, y) dt.$$

We then have

$$u(y) = - \sum_{z \in B_R} g_{B_R}(y, z) f(z) \theta_z,$$

and since $E(x_0, R) = \sum_{y \in B_R} g_{B_R}(x_0, y) \theta_y$, we obtain

$$\max_{B_R} |u| \leq \overline{E}(x_0, R) \max_{B_R} |f|.$$

Let v be a function on B_r that solves the Poisson equation $\mathcal{L}v = f$ in B_r . In the same way

$$\max_{B_r} |v| \leq \overline{E}(x, r) \max_{B_r} |f|.$$

The function $w = u - v$ is harmonic in $B_r \subset B_R$ whence, by Proposition 2.6,

$$\operatorname{osc}_{B_{\sigma r}} w \leq \varepsilon \operatorname{osc}_{B_r} w \quad \forall (\sigma r) \geq R_1(x_0).$$

Since $w = u$ on $B_R \setminus B_r$, the maximum principle implies that

$$\operatorname{osc}_{B_r} w \leq \operatorname{osc}_{B_R} w = \operatorname{osc}_{B_R \setminus B_r} w = \operatorname{osc}_{B_R \setminus B_r} u \leq 2 \max_{B_R} |u|.$$

Hence,

$$\operatorname{osc}_{B_{\sigma r}} u \leq \operatorname{osc}_{B_{\sigma r}} v + \operatorname{osc}_{B_{\sigma r}} w \leq 2 \max_{B_{\sigma r}} |v| + 2\varepsilon \max_{B_R} |u| \leq 2(\overline{E}(x_0, r) + \varepsilon \overline{E}(x_0, R)) \max_{B_R} |f|,$$

\square

Proof of Proposition 2.8. (i) Let \mathcal{L}_V^A be the restriction of the operator \mathcal{L}_V on A with Dirichlet boundary conditions outside A and denote by $\{\lambda_i, i \in \{1, \dots, |A|\}\}$ be the set of eigenvalues of the positive symmetric operator $-\mathcal{L}_V^A$ labelled in increasing order, and $\{\psi_i, i \in \{1, \dots, |A|\}\}$ the corresponding eigenfunctions with due normalization. We have

$$u_t = P_t^A f = \sum_i e^{-\lambda_i t} \langle f, \psi_i \rangle \psi_i,$$

which gives

$$-\partial_t u_t = \sum_i \lambda_i e^{-\lambda_i t} \langle f, \psi_i \rangle \psi_i,$$

and thus

$$\|\partial_t u_t\|_2^2 = \sum_i \lambda_i^2 e^{-2\lambda_i t} \langle f, \psi_i \rangle^2.$$

Using the inequality $\lambda_i s \leq e^{\lambda_i s}$, we get

$$\|\partial_t u_t\|_2^2 \leq \frac{1}{s^2} \sum_i e^{-2\lambda_i(t-s)} \langle f, \psi_i \rangle^2 = \frac{1}{s^2} \|u_{t-s}\|_2^2.$$

(ii) We have the semigroup identity

$$p_t^A(x, y) = \sum_z p_v^A(x, z) p_{t-v}^A(z, y) \theta_z,$$

from which we get

$$\partial_t p_t^A(x, y) = \sum_z p_v^A(x, z) \partial_t p_{t-v}^A(z, y) \theta_z,$$

whence

$$|\partial_t p_t^A(x, y)| \leq \|p_v^A(x, \cdot)\|_2 \|\partial_t p_{t-v}^A(y, \cdot)\|_2,$$

By Proposition 2.8 (i),

$$\|\partial_t p_{t-v}^A(y, \cdot)\|_2 \leq \frac{1}{s} \|\partial_t p_{t-v-s}^A(y, \cdot)\|_2$$

for any $s \leq t - v$. Since

$$\|p_v^A(x, \cdot)\|_2^2 = \sum_z p_v^A(x, z)^2 \theta_z = p_{2v}^A(x, x),$$

we obtain (2.15).

(iii) Choose $v \simeq s \simeq t/3$, it follows then from Assumption 1.1 (i) that for any nonempty finite set $A \subset \mathbb{Z}^d$ and t large enough,

$$p_{2v}^A(x, x) \leq C t^{-d/2} \text{ and } p_{2(t-v-s)}^A(y, y) \leq C t^{-d/2}, \quad \forall x, y \in B(x_0, t^{(1+\delta)/2}),$$

when $2t/3 \geq T_0(x_0)$, whence by Proposition 2.8 (ii),

$$|\partial_t p_t^A(x, y)| \leq C t^{-(\frac{d}{2}+1)}.$$

By letting $A \rightarrow \mathbb{Z}^d$, we obtain (2.16). \square

Proof of Theorem 1.5. Given Proposition 1.4, we can use the balayage argument as in the proof of [9, Theorem 3.1]. Note that the statement of [9, Theorem 3.1] includes estimates of very good balls, but as in the proof, we only need the heat kernel estimates.

Let $C_* = \delta_0^{-1}$ and define

$$B = B(x_0, C_*R), \quad B_1 = B(x_0, R), \\ Q = Q(x_0, R, R^2) = (0, 4R^2] \times B, \quad E = (0, 4R^2] \times B_1.$$

Let $u(t, x) \geq 0$ be caloric on Q . Let Z be the space-time process on $\mathbb{R} \times G$ given by $Z_t = (V_0 - t, X_t)$, where X is the Markov chain on G , and V_0 is the initial time. Define the réduite u_E by

$$u_E(t, x) = E^x(u(t - T_E, X_{T_E}); T_E < \tau_Q),$$

where $T_E = \inf\{t \geq 0 : Z_t \in E\}$ and $\tau_Q = \inf\{t \geq 0 : Z_t \notin Q\}$. Clearly, $u_E = u$ on E , $u_E = 0$ on Q^c , and $u_E \leq u$ on $Q - E$. Since a dual process of Z exists and can be written as $(V_0 + t, X_t)$, the balayage formula holds and we can write

$$u_E(t, x) = \int_E p_{t-r}^B(x, y) \nu_E(dr, dy), \quad (t, x) \in Q,$$

for a suitable measure ν_E . Here $p_t^B(x, y)$ is the heat kernel of X , killed on exiting from B . In this case we can write things more explicitly. Set

$$Jf(x) = \begin{cases} \sum_{y \in B} \frac{\omega_{xy}}{\theta(y)} f(y) & \text{if } x \in B_1, \\ 0 & \text{if } x \in B - B_1. \end{cases} \quad (7.2)$$

The balayage formula takes the form

$$u_E(t, x) = \sum_{y \in B_1} p_t^B(x, y) u(0, y) \theta(y) + \sum_{y \in B_1} \int_{(0, T]} p_{t-s}^B(x, y) k(s, y) \theta(y) ds, \quad (7.3)$$

where $k(s, y)$ is zero if $y \in B - B_1$ and

$$k(s, y) = J(u(s, \cdot) - u_E(s, \cdot))(y), \quad y \in B_1. \quad (7.4)$$

(See [3, Proposition 3.3]; See also [9, Appendix] for a self-contained proof of (7.3) and (7.4) for the discrete time case.) Since $u = u_E$ on E , if $s > 0$ then (7.4) implies that $k(r, y) = 0$ unless $y \in \partial(B - B_1)$.

Now let $(t_1, y_1) \in Q_-$ and $(t_2, y_2) \in Q_+$. Note that since $(t_i, y_i) \in E$ for $i = 1, 2$, we have $u_E(t_i, y_i) = u(t_i, y_i)$. Choose $R_5(x_0)$ large enough such that $R_5(x_0) \geq C(R_*(x_0) + \sqrt{T_0(x_0)} + \sqrt{T_1(x_0)})$ for some $C \geq 1$. By Assumption 1.1 (i), Proposition 1.4 and Corollary 1.3, we have, writing $A = \partial(B - B_1)$ and $T = R^2$,

$$\begin{aligned} p_{t_2-s}^B(x, y) &\geq c_1 T^{-d/2} && \text{for } x, y \in B_1, 0 \leq s \leq T, \\ p_s(x, y) &\leq c_2 T^{-d/2} && \text{for } x, y \in B_1, T \leq s \leq 2T, \\ p_{t_1-s}(x, y) &\leq c_2 T^{-d/2} && \text{for } x \in B_1, y \in A, 0 < s \leq t_1. \end{aligned}$$

Substituting these bounds in (7.3), we have

$$\begin{aligned}
u(t_2, y_2) &= \sum_{y \in B_1} p_{t_2}^B(y_2, y) u(0, y) \theta(y) + \sum_{y \in A} \int_0^{t_2} p_{t_2-s}^B(y_2, y) k(s, y) \theta(y) ds \\
&\geq \sum_{y \in B_1} c_1 T^{-d/2} u(0, y) \theta(y) + \sum_{y \in A} \int_0^{t_1} c_1 T^{-d/2} k(s, y) \theta(y) ds \\
&\geq \sum_{y \in B_1} c_1 c_2^{-1} p_{t_1}^B(y_1, y) u(0, y) \theta(y) + \sum_{y \in A} \int_0^{t_1} c_1 c_2^{-1} p_{t_1-s}^B(y_1, y) k(s, y) \theta(y) ds \\
&= c_1 c_2^{-1} u(t_1, y_1),
\end{aligned}$$

which proves (1.6). □

DEDICATION

The authors would dedicate this paper to Professor Kiyosi Itô. Omar Boukhadra would also dedicate it to the memory of his father Youcef Bey.

ACKNOWLEDGMENTS

This research was supported by the french ANR project MEMEMO2 2010 BLAN 0125, and by the Grant-in-Aid for Scientific Research (A) 25247007, Japan.

REFERENCES

- [1] S. Andres, M.T. Barlow, J.-D. Deuschel and B.M. Hambly, Invariance principle for the random conductance model, *Probab. Theory Related Fields* **156** (2013), 535–580.
- [2] S. Andres, J.-D. Deuschel and M. Slowik, Harnack inequalities on weighted graphs and some applications for the random conductance model. *arXiv:1312.5473*
- [3] M.T. Barlow, R.F. Bass and T. Kumagai, Parabolic Harnack inequality and heat kernel estimates for random walks with long range jumps. *Math. Z.* **261** (2009), no. 2, 297–320.
- [4] M. T. Barlow, Random walks on supercritical percolation clusters, *Ann. Probab.* **32** (2004), no. 4, 3024–3084.
- [5] M. T. Barlow, *Diffusions on fractals*, Lecture Notes in Mathematics **1690**, Ecole d’été de probabilités de Saint-Flour XXV–1995, Springer, New York, (1998).
- [6] M.T. Barlow and R.F. Bass, *Construction of Brownian motion on the Sierpinski carpet*. *Ann. Inst. H. Poincaré*, **25** (1989), 225–257.
- [7] M. T. Barlow and J.-D. Deuschel, Invariance principle for the random conductance model with unbounded conductances, *Ann. Probab.* **38** (2010), no. 1, 234–276.
- [8] M. T. Barlow, A. Grigor’yan and T. Kumagai, On the equivalence of parabolic Harnack inequalities and heat kernel estimates, *J. Math. Soc. Japan* **64** (2012), no. 4, 1091–1146.
- [9] M. T. Barlow and B. M. Hambly, Parabolic Harnack inequality and local limit theorem for percolation clusters, *Electron. J. Probab.* **14** (2009), no. 1, 1–27.
- [10] N. Berger and M. Biskup, Quenched invariance principle for simple random walk on percolation clusters, *Probab. Theory Related Fields* **137** (2007), no. 1-2, 83–120.

- [11] N. Berger, M. Biskup, C.E. Hoffman, G. Kozma, Anomalous heat-kernel decay for random walk among bounded random conductances, *Ann. Inst. Henri Poincaré Probab. Stat.* **44** (2008), no. 2, 374–392.
- [12] M. Biskup, Recent progress on the Random Conductance Model, *Prob. Surveys* **8** (2011), 294–373.
- [13] M. Biskup and O. Boukhadra, Subdiffusive heat-kernel decay in four-dimensional i.i.d. random conductance models, *J. London Math. Soc.* **86** (2012), no. 2, 455–481.
- [14] M. Biskup and T. M. Prescott, Functional CLT for random walk among bounded random conductances, *Electron. J. Probab.* **12** (2007), no. 49, 1323–1348.
- [15] M. Biskup, O. Luidor, A. Rozinov and A. Vandenberg-Rodes, Trapping in the random conductance model, *J. Statist. Phys.* **150** (2011), no. 1, 66–87.
- [16] O. Boukhadra, Heat-kernel estimates for random walk among random conductances with heavy tail, *Stochastic Process. Appl.* **120** (2010), no. 2, 182–194.
- [17] O. Boukhadra, Standard spectral dimension for the polynomial lower tail random conductances model, *Electron. J. Probab.* **15** (2010), no. 68, 2069–2086.
- [18] X. Chen, Pointwise upper estimates for transition probability of continuous time random walks on graphs, arXiv:1310.2680
- [19] E.B. Davies, Large deviations for heat kernels on graphs, *J. London Math. Soc.* (2) **47** (1993), 65–72.
- [20] T. Delmotte, Parabolic Harnack inequality and estimates of Markov chains on graphs, *Rev. Mat. Iberoamericana* **15** (1999), no. 1, 181–232.
- [21] R. Durrett and R. H. Schonmann, Large deviations for the contact process and two-dimensional percolation, *Probab. Theory Related Fields* **77** (1988), no. 4, 583–603.
- [22] M. Folz, Gaussian upper bounds for heat kernels of continuous time simple random walks, *Elect. J. Probab.* **16** (2011), 1693–1722.
- [23] L. R. G. Fontes and P. Mathieu, On symmetric random walks with random conductances on \mathbb{Z}^d , *Probab. Theory Related Fields* **134** (2006), no. 4, 565–602.
- [24] A. Grigor'yan and A. Telcs, Harnack inequalities and sub-Gaussian estimates for random walks, *Math. Annalen* **324** (2002), 521–556.
- [25] A. Grigor'yan and A. Telcs, Sub-Gaussian estimates of heat kernels on infinite graphs, *Duke Math. J.* **109** (2001), 451–510.
- [26] G. Grimmett, *Percolation*, second edition, Grundlehren der Mathematischen Wissenschaften, 321, Springer, Berlin, 1999.
- [27] T. Kumagai, *Random Walks on Disordered Media and their Scaling Limits*, Lect. Notes in Math., **2101**, École d'Été de Probabilités de Saint-Flour XL–2010. Springer, New York, (2014).
- [28] P. Mathieu, Quenched invariance principles for random walks with random conductances, *J. Stat. Phys.* **130** (2008), no. 5, 1025–1046.
- [29] P. Mathieu and A. Piatnitski, Quenched invariance principles for random walks on percolation clusters, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **463** (2007), no. 2085, 2287–2307.
- [30] P. Mathieu and E. Remy, Isoperimetry and heat kernel decay on percolation clusters, *Ann. Probab.* **32** (2004), no. 1A, 100–128.
- [31] V. Sidoravicius and A.-S. Sznitman, Quenched invariance principles for walks on clusters of percolation or among random conductances, *Probab. Theory Related Fields* **129** (2004), no. 2, 219–244.