RIMS-1858

Cactus group and Gelfand-Tsetlin group

By

Arkady BERENSTAIN and Anatol N. KIRILLOV

October 2016

京都大学　数理解析研究所

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES

KYOTO UNIVERSITY, Kyoto, Japan

Cactus group and Gelfand-Tsetlin group

ARKADY BERENSTAIN

Department of Mathematics, University of Oregon, Eugene, OR 97403, USA E-mail address: arkadiy@math.uoregon.edu

ANATOL N. KIRILLOV

Research Institute of Mathematical Sciences (RIMS)
Kyoto 606-8502, Japan
URL: http://www.kurims.kyoto-u.ac.jp/~kirillov

and

The Kavli Institute for the Physics and Mathematics of the Universe (IPMU),
5-1-5 Kashiwanoha, Kashiwa, 277-8583, Japan

and

Department of Mathematics, National Research University Higher School of Economics,

7 Vavilova Str., 117312, Moscow, Russia

Abstract

We prove the Etingof conjecture stated that the quotient $Cact_n^{(0)}$ of the Cactus group $Cact_n$ [2],[5],[9] by the relation $(s_{12}s_{13})^6 = 1$ is isomorphic to the Gelfand–Tsetlin group, known also as the BK_n group, which has been introduced and studied in [1]. In particular, a subtraction free birational (as well as piece-wise linear one) action of the (reduced) Cactus group $Cact_n^{(0)}$ on the projective space $\mathbb{P}^{n^2}$ is described. This action is geometric/tropical lift of the combinatorial action of the local Schützenberger transformation on the set of semistandard Young tableaux, cf [1], [6].

2000 Mathematics Subject Classifications: 05E05, 05E10, 05A19.

1 Introduction

The cactus group $Cact_n := \pi_1(\overline{M_{0,n+1}(\mathbb{R})})$ is the fundamental group of the real locus of the Deligne–Mumford compactification $\overline{M_{0,n+1}}$ of the moduli space of genus zero stable curves with $n+1$ marked points, see e.g., [2]. The action of (reduced) cactus group on the set of standard Young tableaux has been described in [5],[4]. On the other hand, a combinatorial action of the Gelfand–Tsetlin/Berenstein–Kirillov group GT_n/BK_n on the set of semistandard Young tableaux had been introduced by A. Lascoux and M.-P. Schützenberger, see e.g., [7]; see also [3], [6].

Based on results obtained in [1],[6], we construct birational/geometric representation of the (reduced) cactus group $Cact_n^{(0)}$, and identify it with the Gelfand–Tsetlin group GT_n. In fact, in [1] the affine extension of the Gelfand–Tsetlin group has been introduced and studied, as well as presented an action of the (affine) group GT_n on the set of semistandard Young tableaux and that of transportation matrices, cf [8]. It will be interesting task to find a geometric interpretations of affine Cactus and affine Gelfand–Tsetlin groups. Another interesting problem we are looking for, is to study of combinatorial and algebraic properties of the higher genus cactus group $Cact_{g,n} := \pi_1(\overline{M_{g,n+1}}(\mathbb{R}))$, work in progress.

2 Cactus group $Cact_n$, [2],[5], [4], [9]

Definition 2.1 *The Cactus group $Cact_n$ is a group (with a unit) generated by elements σ_{ij}, $1 \le i < j \le n$, subject to the set of relations*

- $\sigma_{ij}^2 = 1, \quad if \quad 1 \le i < j \le n,$
- $\sigma_{ij}\ \sigma_{kl} = \sigma_{kl}\ \sigma_{ij}, \quad if \quad j < k,$
- $\sigma_{ij}\ \sigma_{kl}\ \sigma_{ij} = \sigma_{i+j-l,i+j-k}, \quad if \ \ i \le k < l \le j.$

Let us set $\sigma_i := \sigma_{1,i+1}, \ 1 \le i \le n-1$. It is clear that $\sigma_i^2 = 1$, and the elements $\sigma_1, \ldots, \sigma_{n-1}$ generate the Cactus group $Cact_n$. We denote by $Cact_n^{(0)}$ the quotient of the cactus group $Cact_n$ by the normal subgroup generated by the element $((\sigma_1\sigma_2)^6 - 1)$.

3 Gelfand–Tsetlin group, [1], [6]

Definition 3.1 *([1],[6]) The Gelfand–Tsetlin group GT_n, known also as the Berenstein–Kirillov group and denoted by BK_n, is a group (with a unit) generated by the elements $t_1, \ldots, t_{n-1}$ subject to the set of relations*

- $(t_1\ t_2)^6 = 1,$
- $t_i\ t_j = t_j\ t_i, \ \ if\ |i-j| \ge 2,$
- $(t_i\ q_j)^4 = 1 \ \ if\ j - i \ge 2, \quad where$

$$q_i = t_1\ \underbrace{t_2\ t_1}\cdots\underbrace{t_i t_{i-1}\ldots t_1}, \quad 1 \le i < n.$$

Proposition 3.2 *([1],[6]) The following relations in the group BK_n are satisfied*

- $[t_i, q_k\ t_j\ q_k] = 0, \quad if \quad k \geq i+j+1,$
- $[q_i, q_k\ q_j\ q_k] = 0, \quad if \quad k \geq i+j+1,$

where $[a,b] := a\ b - b\ a$ denotes the commutator of elements a and b.

Theorem 3.3 *([1]) The elements of the Gelfand–Tsetlin group GT_n listed below*

$$s_i := q_i\ t_1\ q_i, \quad 1 \leq i < n$$

satisfy the following relations

- $s_i^2 = 1$,
- *(Coxeter relations)* $(s_i\ s_{i+1})^3 = 1$, *if* $1 \leq i < n$,

(*commutativity*) $(s_i\ s_j)^2 = 1$, *if* $|i-j| \geq 2$. *We* **expect** *that the group generated by the elements $s_1, \ldots, s_{n-1}$ is isomorphic to the symmetric group $\mathbb{S}_n$.*

Theorem 3.4 *The maps $\sigma_i \longleftrightarrow q_i$ can be extended to the* **isomorphism** *of groups*

$$Cact_n^{(0)} \cong GT_n.$$

Indeed, it is clear that under the above correspondence one has

$$q_k\ q_j\ q_k = \sigma_{k+1-j,k+1}.$$

Therefore, if $k \geq i+j+1$, then $k-j+1 \geq i+2$, and therefore

$$[\sigma_{1,i+1}, \sigma_{k+1-j,k+1}] = [q_i, q_k\ q_j\ q_k] = 0.$$

Now let us check that the image of relations

$$\sigma_{in}\ \sigma_{jk} = \sigma_{i+n-k,i+n-j}\ \sigma_{in}, \quad 1 \leq j < k \leq n$$

are satisfied in the Gelfand–Tsetlin group. Indeed, we have to show that the relations

$$q_{n-1}\ q_{n-i}\ q_{n-1}\ q_{k-1}\ q_{k-j}\ q_{k-1} = q_{n+i-j-1}\ q_{k-j}\ q_{n+i-j-1}\ q_{n-1}\ q_{n-i}\ q_{n-1}$$

are valid in the group GT_n. By definition, $q_r = q_{r-1}\ p_r$, where $p_r := \underbrace{t_r\ t_{r-1} \cdots t_2\ t_1}$. Therefore,

$$q_{n-1} = q_{i+n-j-1}\ u_{n+i-j}^{(n-1)},$$

where we set $u_b^{(a)} := p_a \ldots p_b$, if $a > b$. Therefore one can rewrite relations we have to prove, in the following form

$$u_{i+n-j}^{(n-1)}\ q_{n-i}\ q_{n-1}q_{k-1}\ q_{k-ju}q_{k-1} = q_{k-j}\ u_{n+i-j}^{(n-1}\ q_{n-i}\ q_{n-1}.$$

Now we use the following deletion $u_{n-1} = q_{n-i}\ u_{n_i+1}^{(n-1)}$,so that we can rewrite the above relations in the form

$$u_{n+i-j}^{(n-1)}\ u_{n-i+1}^{(n-1)}\ q_{k-1} = q_{k-j}\ u_{n+i-j}^{(n-1)}\ u_{n-i+1}^{(n-1)}\ q_{k-1}.$$

Note that the number of terms in the product $u_{n+i-j}^{(n-1)}\ u_{n-i+1}^{(n-1)}$ is equal to $j-1$. Therefore,

$$u_{n+i-j}^{(n-1)}\ u_{n-i+1}^{(n-1)}\ q_{k-1} = q_{k-j} \times A_{j+2},$$

where A_{j+2} is a certain product of generators $t_n, \ldots, t_{j+2}$ only. This statement is clearly seen from the relations

$$p_a\ q_b = t_a\ t_{a-1} \cdots t_{b+1}\ q_{b-1}, \quad if \quad a > b.$$

The relations $s_{ij}\ s_{kl} = s_{kl}\ s_{ij},\ if\ j < k$, can be proved in the same fashion.

Clearly, one has the maps $GT_n \longleftrightarrow Cact_n^{(0)}$ given by

$$q_i \longleftrightarrow s_{1,i+1}$$

. Therefore we came to conclusion that the groups GT_n and $Cact_n^{(0)}$ are isomorphic.

Acknowledgments This note has been written during the second author visit of the Department of Statistic of the University of Warwick and the Clay Mathematical Institute, Oxford during May, 2015. A.K. would like to express deepest thanks to Prof. N. Zygouras for invitation to visit the University of Warwick and the workshop "Random Polymers and Algebraic Combinatorics", May 25–29, at the Clay Mathematical Institute, Oxford. A.K. wants to thanks Professors N.Zygouras and B.Westbury for many interesting discussions concerning tropical/geometric RSK and cactus group.

Remark After this note was written, we was informed about a preprint arXive:1609.2046 " The Berenstein-Kirillov group and cactus groups" by M. Chmutov, M. Glick, P. Pylyavskyy, which also contains another approach to identify the cactus group $Cactus)n^{(0)}$ and that BK_n.

References

[1] A. Berenstein and A. N. Kirillov. *Group generated by involutions, Gelfand-Tsetlin patterns and combinatorics of Young tableaux*, Algebra i Analiz 7 (1995), no.1, 92-152; *translation in* St.Petersburg Math. J. 7 (1996), no.1, 77-127.

[2] P.Etingof, A.Henriques, J. Kamnitzer, E. Rains, Eric, *The cohomology ring of the real locus of the moduli space of stable curves of genus 0 with marked points*, Ann. of Math. (2) **171** (2010), no. 2, 731–777.

[3] E.R. Gansner, *On the equality of two plane paryition correspondences*, Discrete Math, **30** (1980), 121-132.

[4] A. Henriques, *An action of the cactus group*, arXiv:0705.3000 [math.AG].

[5] A. Henriques and J. Kamnitzer, *Crystals and coboundary categories*, Duke Math. J. **132** no. 2 (2006), 191–216.

[6] A.N. Kirillov, *Introduction to Tropical Combinatorics* , Physics and Combinatorics 2000 (Proceedings of the Nagoya 2000 International Workshop on Physics and Combinatorics, Nagoya University, August 21-26, 2000, ed. A.Kirillov and N.Liskova), 82–150, World Scientific, Singapore, 2001

[7] A. Lascoux and M.-P. Schützenberger, *Keys and standard bases*, Invariant Theory and Tableaux (Minneapolis, MN,1988); IMA Vol. Math.Appl. vol **19** (1990), 125-144, Springer, New-York-Berlin.

[8] I. Losev, *Cacti and cells*, arXiv:1506.04400 [math.RT].

[9] L. Rybnikov, *Cactus group and monodromy of Bethe vectors*, arXiv:1409.0131 [math.QA].