

Geometrical formality of solvmanifolds and solvable Lie type geometries

By

HISASHI KASUYA*

Abstract

We show that for a Lie group $G = \mathbb{R}^n \rtimes_{\phi} \mathbb{R}^m$ with a semisimple action ϕ which has a cocompact discrete subgroup Γ , the solvmanifold G/Γ admits a canonical invariant formal (i.e. all products of harmonic forms are again harmonic) metric. We show that a compact oriented aspherical manifold of dimension less than or equal to 4 with the virtually solvable fundamental group admits a formal metric if and only if it is diffeomorphic to a torus or an infra-solvmanifold which is not a nilmanifold.

§ 1. Introduction

Let (M, g) be a compact oriented Riemannian n -manifold. We call g formal if all products of harmonic forms are again harmonic. If a compact oriented manifold admits a formal Riemannian metric, we call it geometrically formal. If g is formal, then the space of the harmonic forms is a subalgebra of the de Rham complex of M and isomorphic to the real cohomology of M . By this, a geometrically formal manifold is a formal space (in the sense of Sullivan [22]). But the converse is not true (see [15] [16]). For very simple examples, closed surfaces with genus ≥ 2 are formal but not geometrically formal. Thus we have one problem of geometrical formality of formal spaces. Kotschick's nice work in [15] stimulates us to consider this problem.

In this paper we prove the following theorem by using computations of the de Rham cohomology of general solvmanifolds given in [14].

Theorem 1.1. *Let $G = \mathbb{R}^n \rtimes_{\phi} \mathbb{R}^m$ with a semisimple action ϕ . Suppose G has a lattice Γ . Then G/Γ admits an invariant formal metric.*

Received September 27, 2011. Revised December 28, 2011.

2000 Mathematics Subject Classification(s): 22E25, 53C20, 58A14

Key Words: solvmanifold, Harmonic form

*Graduate school of mathematical science University of Tokyo Japan

e-mail: khsc@ms.u-tokyo.ac.jp

We also study geometrical formality of low-dimensional aspherical manifolds with the virtually solvable fundamental groups. We consider infra-solvmanifolds which are quotient spaces of simply connected solvable Lie groups by subgroups of the groups of the affine transformations of G satisfying some conditions (see Section 7 for the definition). We classify geometrically formal compact aspherical manifolds of dimension less than or equal to 4 with the virtually solvable fundamental groups.

Theorem 1.2. *Let M be a compact oriented aspherical manifold of dimension less than or equal to 4 with the virtually solvable fundamental group. Then M is geometrically formal if and only if M is diffeomorphic to a torus or an infra-solvmanifold which is not a nilmanifold.*

§ 2. Notation and conventions

Let k be a subfield of \mathbb{C} . A group \mathbf{G} is called a k -algebraic group if \mathbf{G} is a Zariski-closed subgroup of $GL_n(\mathbb{C})$ which is defined by polynomials with coefficients in k . Let $\mathbf{G}(k)$ denote the set of k -points of \mathbf{G} and $\mathbf{U}(\mathbf{G})$ the maximal Zariski-closed unipotent normal k -subgroup of \mathbf{G} called the unipotent radical of \mathbf{G} . Denote $U_n(k)$ the group of k -valued upper triangular unipotent matrices of size n .

§ 3. Unipotent hull of solvable Lie group

Theorem 3.1. ([19]) *Let G be a simply connected solvable Lie group. Then there exists a unique \mathbb{R} -algebraic group \mathbf{H}_G with an injective group homomorphism $\psi : G \rightarrow \mathbf{H}_G(\mathbb{R})$ so that:*

- (1) $\psi(G)$ is Zariski-dense in \mathbf{H}_G .
- (2) The centralizer $Z_{\mathbf{H}_G}(\mathbf{U}(\mathbf{H}_G))$ of $\mathbf{U}(\mathbf{H}_G)$ is contained in $\mathbf{U}(\mathbf{H}_G)$.
- (3) $\dim \mathbf{U}(\mathbf{H}_G) = \dim G$ (resp. $\text{rank } G$).

We denote $\mathbf{U}_G = \mathbf{U}(\mathbf{H}_G)$.

Theorem 3.2. ([13]) *Let G be a simply connected solvable Lie group. Then \mathbf{U}_G is abelian if and only if $G = \mathbb{R}^n \rtimes_{\phi} \mathbb{R}^m$ such that the action $\phi : \mathbb{R}^n \rightarrow \text{Aut}(\mathbb{R}^m)$ is semisimple.*

§ 4. Hodge theory

Let (V, g) be a \mathbb{R} or \mathbb{C} -vector space of dimension n with an inner product g . Let $\bigwedge V = \bigoplus_{p=0} \bigwedge^p V$ be the exterior algebra of V . We extend g to the inner product on

$\wedge V$. Take $vol \in \wedge^n V$ such that $g(vol, vol) = 1$. We define the linear map $*_g : \wedge^p V \rightarrow \wedge^{n-p} V$ as:

$$v \wedge *_g \bar{u} = g(v, u) vol$$

Let $\{\theta_1, \dots, \theta_n\}$ be an orthonormal basis of (V, g) . Then we have

$$*_g(\theta_{i_1} \wedge \dots \wedge \theta_{i_p}) = (\text{sgn} \sigma_{IJ}) \theta_{j_1} \wedge \dots \wedge \theta_{j_{n-p}}$$

where $J = \{j_1, \dots, j_{n-p}\}$ is the complement of $I = \{i_1, \dots, i_p\}$ in $\{1, \dots, n\}$ and σ_{IJ} is the permutation $\begin{pmatrix} 1 \cdots p & p+1 \cdots n \\ i_1 \cdots i_p & j_1 \cdots j_{n-p} \end{pmatrix}$.

Let (M, g) be a compact oriented Riemannian n -manifold. Let $(A^*(M), d)$ be the de Rham complex of M with the exterior derivation d . For $x \in M$ by the inner product g_x on $T_x M$ we define the linear map $*_g : A^p(M) \rightarrow A^{n-p}(M)$ by

$$(*_g(\omega))_x = *_{g_x} \omega_x$$

for $\omega \in A^p(M)$. Define $\delta : A^p(M) \rightarrow A^{p-1}(M)$ by $\delta = (-1)^{np+n+1} *_g d *_g$. We call $\omega \in A^p(M)$ harmonic if $d\omega = 0$ and $\delta\omega = 0$. Let $\mathcal{H}^p(M)$ be the subspace of $A^p(M)$ which consists of harmonic p -forms. Let $\mathcal{H}(M) = \bigoplus \mathcal{H}^p(M)$. It is known that the inclusion $\mathcal{H}(M) \subset A^*(M)$ induces an isomorphism

$$\mathcal{H}^p(M) \cong H^p(M, \mathbb{R}).$$

In general a wedge product of harmonic forms is not harmonic and so $\mathcal{H}^p(M)$ is not a subalgebra of $A^*(M)$.

Definition 4.1. We call a Riemannian metric g formal if all products of harmonic forms are again harmonic. We call an oriented compact manifold M geometrical formal if M admits a formal metric.

§ 5. Invariant forms on solvmanifolds (proof of Theorem 1.1)

Let G be a simply connected solvable Lie group and \mathfrak{g} the Lie algebra which is the space of the left invariant vector fields on G . Consider the exterior algebra $\wedge \mathfrak{g}^*$ of the dual space of \mathfrak{g} . Denote $d : \wedge^1 \mathfrak{g}^* \rightarrow \wedge^2 \mathfrak{g}^*$ the dual map of the Lie bracket of \mathfrak{g} and $d : \wedge^p \mathfrak{g}^* \rightarrow \wedge^{p+1} \mathfrak{g}^*$ the extension of this map. We can identify $(\wedge \mathfrak{g}^*, d)$ with the left invariant forms on G with the exterior derivation. Let $\text{Ad} : G \rightarrow \text{Aut}(\mathfrak{g})$ be the adjoint representation. Let Ad_{sg} be the semi-simple part of $\text{Ad}_g \in \text{Aut}(\mathfrak{g})$ for $g \in G$. Since representations of G are trigonalizable in \mathbb{C} by Lie's theorem, $\text{Ad}_s : G \rightarrow \text{Aut}(\mathfrak{g}_{\mathbb{C}})$ is a diagonalizable representation. Let X_1, \dots, X_n be a basis of $\mathfrak{g}_{\mathbb{C}}$ such that Ad_s is

represented by diagonal matrices. Then we have $\text{Ad}_{sg}X_i = \alpha_i(g)X_i$ for characters α_i of G . Let x_1, \dots, x_n be the dual basis of X_1, \dots, X_n . We assume that G has a lattice Γ . Define the sub-DGA A^* of the de Rham complex $A_{\mathbb{C}}^*(G/\Gamma)$ as

$$A^p = \left\langle \alpha_{i_1 \dots i_p} x_{i_1} \wedge \dots \wedge x_{i_p} \mid \begin{array}{l} 1 \leq i_1 < i_2 < \dots < i_p \leq n, \\ \text{the restriction of } \alpha_{i_1 \dots i_p} \text{ on } \Gamma \text{ is trivial} \end{array} \right\rangle$$

where $\alpha_{i_1 \dots i_p} = \alpha_{i_1} \dots \alpha_{i_p}$.

Theorem 5.1. ([14, v4. Corollary 7.6]) *The inclusion*

$$A^* \subset A_{\mathbb{C}}^*(G/\Gamma)$$

induces a cohomology isomorphism and A^ can be considered as a sub-DGA of $\bigwedge \mathfrak{u}^*$ where \mathfrak{u} is the Lie algebra of \mathbf{U}_G as in Section 3.*

Define g the Hermitian inner product as

$$g(X_i, X_j) = \delta_{ij}.$$

Since Ad_s is an \mathbb{R} -valued representation, the restriction of g on \mathfrak{g} is an inner product on \mathfrak{g} . We consider g as an invariant Riemannian metric on G/Γ .

Theorem 5.2. *If \mathbf{U}_G is abelian, then g is a formal metric on G/Γ .*

Proof. By the assumption, the differential on $\bigwedge \mathfrak{u}^*$ is 0. By Theorem 5.1, the derivation on A^* is 0 and we have an isomorphism

$$A^* \cong H^*(G/\Gamma).$$

Thus it is sufficient to show that all elements of A^* are harmonic. Let $*_g$ be the star operator. Then for $\alpha_{i_1 \dots i_p} x_{i_1} \wedge \dots \wedge x_{i_p} \in A^p$ we have

$$*_g(\alpha_{i_1 \dots i_p} x_{i_1} \wedge \dots \wedge x_{i_p}) = (\text{sgn} \sigma_{IJ}) \bar{\alpha}_{i_1 \dots i_p} x_{j_1} \wedge \dots \wedge x_{j_{n-p}}.$$

Since the restriction of $\alpha_{i_1 \dots i_p}$ on Γ is trivial, the image $\alpha_{i_1 \dots i_p}(G) = \alpha_{i_1 \dots i_p}(G/\Gamma)$ is compact and hence $\alpha_{i_1 \dots i_p}$ is unitary. Since G has a lattice Γ , G is unimodular (see [19, Remark 1.9]) and hence we have

$$\bar{\alpha}_{i_1 \dots i_p} = \alpha_{i_1 \dots i_p}^{-1} = \alpha_{j_1 \dots j_{n-p}}.$$

Hence we have

$$\bar{\alpha}_{i_1 \dots i_p} x_{j_1} \wedge \dots \wedge x_{j_{n-p}} = \alpha_{j_1 \dots j_{n-p}} x_{j_1} \wedge \dots \wedge x_{j_{n-p}} \in A^{n-p}$$

and thus we have $*_g(A^*) \subset A^*$. Since the derivation on A^* is 0, we have $\delta(A^*) = 0$. Hence the theorem follows. \square

By this theorem and Theorem 3.2, we have Theorem 1.1.

Remark. Not every invariant metric on G/Γ in Theorem 1.1 is formal. See the following example.

Example 5.3. Let $H = \mathbb{R} \rtimes_{\phi} \mathbb{R}^2$ such that $\phi(z)(x, y) = (e^z x, e^{-z} y)$. Consider $G = H \times \mathbb{R}$. Then for some non-zero number $a \in \mathbb{R}$, $\phi(a)$ is conjugate to an element of $SL_2(\mathbb{Z})$, and hence G has a lattice $\Gamma = a\mathbb{Z} \rtimes_{\phi} \Gamma' \times \mathbb{Z}$ for a lattice Γ' of \mathbb{R}^2 . Let \mathfrak{g} be the Lie algebra of G and \mathfrak{g}^* the dual of \mathfrak{g} . The cochain complex $(\wedge \mathfrak{g}^*, d)$ is generated by a basis $\{x, y, z, w\}$ such that

$$dx = -z \wedge x, \quad dy = z \wedge y, \quad dz = 0, \quad dw = -z \wedge x.$$

Consider the invariant metric $g = x^2 + y^2 + z^2 + w^2$. Then z and $w - x$ are harmonic for g . But $z \wedge (w - x)$ is not harmonic. Thus g is not formal.

Example 5.4. Let $G = \mathbb{C} \rtimes_{\phi} \mathbb{C}^2$ with $\phi(z)(x, y) = (e^z x, e^{-z} y)$. For some $p, q \in \mathbb{R}$, $\phi(p\mathbb{Z} + \sqrt{-1}q\mathbb{Z})$ is conjugate to a subgroup of $SL_4(\mathbb{Z})$ and hence we have a lattice $\Gamma = (p\mathbb{Z} + \sqrt{-1}q\mathbb{Z}) \rtimes \Gamma''$ for a lattice Γ'' of \mathbb{C}^2 (see [17] and [9]). For any lattice Γ , G/Γ is geometrically formal by Theorem 1.1.

Remark. In [2] for some lattice of G in Example 5.4, it is proved that G/Γ is geometrically formal. But the de Rham cohomology of G/Γ varies according to a choice of a lattice Γ .

Example 5.5. Let K be a finite extension field of \mathbb{Q} with the degree r for positive integers. We assume K admits embeddings $\sigma_1, \dots, \sigma_s, \sigma_{s+1}, \dots, \sigma_{s+2t}$ into \mathbb{C} such that $s + 2t = r$, $\sigma_1, \dots, \sigma_s$ are real embeddings and $\sigma_{s+1}, \dots, \sigma_{s+2t}$ are complex ones satisfying $\sigma_{s+i} = \bar{\sigma}_{s+i+t}$ for $1 \leq i \leq t$. We suppose $s > 0$. Denote \mathcal{O}_K the ring of algebraic integers of K , \mathcal{O}_K^* the group of units in \mathcal{O}_K and

$$\mathcal{O}_K^{*+} = \{a \in \mathcal{O}_K^* : \sigma_i(a) > 0 \text{ for all } 1 \leq i \leq s\}.$$

Define $\sigma : \mathcal{O}_K \rightarrow \mathbb{R}^s \times \mathbb{C}^t$ by

$$\sigma(a) = (\sigma_1(a), \dots, \sigma_s(a), \sigma_{s+1}(a), \dots, \sigma_{s+t}(a))$$

for $a \in \mathcal{O}_K$. We denote

$$\sigma(a) \cdot \sigma(b) = (\sigma_1(a)\sigma_1(b), \dots, \sigma_s(a)\sigma_s(b), \sigma_{s+1}(a)\sigma_{s+1}(b), \dots, \sigma_{s+t}(a)\sigma_{s+t}(b))$$

for $a, b \in \mathcal{O}_K$. Then the image $\sigma(\mathcal{O}_K)$ is a lattice in $\mathbb{R}^s \times \mathbb{C}^t$. Define $l : \mathcal{O}_K^{*+} \rightarrow \mathbb{R}^{s+t}$ by

$$l(a) = (\log |\sigma_1(a)|, \dots, \log |\sigma_s(a)|, 2 \log |\sigma_{s+1}(a)|, \dots, 2 \log |\sigma_{s+t}(a)|)$$

for $a \in \mathcal{O}_K^{*+}$. Then by Dirichlet's units theorem, the image $l(\mathcal{O}_K^{*+})$ is a lattice in the vector space $L = \{x \in \mathbb{R}^{s+t} \mid \sum_{i=1}^{s+t} x_i = 0\}$. By this we have a geometrical representation of the semi-direct product $\mathcal{O}_K^{*+} \rtimes \mathcal{O}_K$ as $l(\mathcal{O}_K^{*+}) \rtimes_{\phi} \sigma(\mathcal{O}_K)$ with

$$\phi(t_1, \dots, t_{s+t})(\sigma(a)) = \sigma(l^{-1}(t_1, \dots, t_{s+t})) \cdot \sigma(a)$$

for $(t_1, \dots, t_{s+t}) \in l(\mathcal{O}_K^{*+})$. Since $l(\mathcal{O}_K^{*+})$ and $\sigma(\mathcal{O}_K)$ are lattices of L and $\mathbb{R}^s \times \mathbb{C}^t$ respectively, we have an extension $\bar{\phi} : L \rightarrow \text{Aut}(\mathbb{R}^s \times \mathbb{C}^t)$ of ϕ and $l(\mathcal{O}_K^{*+}) \rtimes_{\phi} \sigma(\mathcal{O}_K)$ can be seen as a lattice of $L \rtimes_{\bar{\phi}} (\mathbb{R}^s \times \mathbb{C}^t)$. By Theorem 1.1, the solvmanifold $L \rtimes_{\bar{\phi}} (\mathbb{R}^s \times \mathbb{C}^t) / l(\mathcal{O}_K^{*+}) \rtimes_{\phi} \sigma(\mathcal{O}_K)$ is geometrically formal by Theorem 5.4. For a subgroup $U \subset \mathcal{O}_K^{*+}$, we have a Lie group $L' \rtimes_{\bar{\phi}} (\mathbb{R}^s \times \mathbb{C}^t)$ which contains $l(U) \rtimes_{\phi} \sigma(\mathcal{O}_K)$ as a lattice. The solvmanifold $L' \rtimes_{\bar{\phi}} (\mathbb{R}^s \times \mathbb{C}^t) / l(U) \rtimes_{\phi} \sigma(\mathcal{O}_K)$ is also geometrically formal by Theorem 1.1.

Example 5.6. Let $G = \mathbb{R} \rtimes_{\phi} U_3(\mathbb{R})$ such that

$$\phi(t) \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & e^t x & z \\ 0 & 1 & e^{-t} y \\ 0 & 0 & 1 \end{pmatrix}.$$

The left-invariant forms $\wedge \mathfrak{g}^*$ on G is generated by $\{e^{-t} dx, e^t dy, dz - xdy, dt\}$. It is known that G has a lattice Γ (see [20]). By simple computations, we have $H^1(\mathfrak{g}^*) = \langle dt \rangle$, $\dim H^2(\mathfrak{g}^*) = 0$ and $\dim H^3(\mathfrak{g}^*) = 1$. Since G is completely solvable, we have $H^*(G/\Gamma, \mathbb{R}) \cong H^*(\mathfrak{g}^*)$ (see [10]) and hence $\mathcal{H}(\mathfrak{g}) = \mathcal{H}(G/\Gamma)$ where $\mathcal{H}^*(\mathfrak{g})$ is the set of left-invariant harmonic forms. By $d(\wedge^3 \mathfrak{g}^*) = 0$, for any invariant metric g on G , we have:

$$\begin{aligned} \mathcal{H}^1(\mathfrak{g}) &= \langle dt \rangle, \\ \mathcal{H}^2(\mathfrak{g}) &= 0, \\ \mathcal{H}^3(\mathfrak{g}) &= \langle (*_g dt) \rangle. \end{aligned}$$

Thus any invariant metric on G/Γ is formal. Otherwise we have $\mathbf{U}_G = U_3(\mathbb{C}) \times \mathbb{C}$ and hence this solvmanifold is different from examples of geometrically formal solvmanifold given in Theorem 1.1.

§ 6. The extension of Theorem 1.1

Let G be a simply connected solvable Lie group and g an invariant metric which we construct in Section 5. Denote C_g the group of the isometrical automorphisms of (G, g) . Consider $C_g \times G$ and the projection $p : C_g \times G \rightarrow C_g$.

Corollary 6.1. *Suppose $G = \mathbb{R}^n \times_{\phi} \mathbb{R}^m$ with a semi-simple action ϕ . Let $\Gamma \subset C_g \times G$ be a torsion-free discrete subgroup such that G/Γ is compact. Suppose $p(\Gamma)$ is finite.*

Then the metric g given in the last section is a formal metric on G/Γ .

Proof. Let $\Delta = \Gamma \cap G$. Since $\Gamma/\Delta \cong p(\Gamma)$, Δ is a finite index normal subgroup of Γ and G/Δ is compact and hence $\Delta \subset G$ is a lattice. Denote $\mathcal{H}(G/\Gamma)$ and $\mathcal{H}(G/\Delta)$ the sets of the harmonic forms on G/Γ and G/Δ for the metric g . Since we have $A^*(G/\Gamma) = A^*(G/\Delta)^{\Gamma/\Delta}$, we have

$$\mathcal{H}(G/\Gamma) = \mathcal{H}(G/\Delta)^{\Gamma/\Delta}.$$

By Theorem 1.1, $\mathcal{H}(G/\Delta)$ is closed under the wedge product, so is $\mathcal{H}(G/\Delta)^{\Gamma/\Delta}$. Hence the corollary follows. \square

Remark. Not all cocompact discrete subgroup Γ satisfies the assumption of the finiteness of $p(\Gamma)$. See the following example.

Example 6.2. Let $G = \mathbb{R} \times_{\phi} \mathbb{R}^3$ such that $\phi(t) = \begin{pmatrix} e^t & 0 & 0 \\ 0 & e^t & 0 \\ 0 & 0 & e^{-2t} \end{pmatrix}$. Then G has no

lattice (see [11, Chapter 7]). Consider the metric $g = e^{-2t}dx^2 + e^{-2t}dy^2 + e^{4t}dz^2 + dt^2$. Then we have $C_g = O(2) \times O(1)$ acting as rotations and reflections on the (x, y) -coordinates and reflection on the z -coordinate. $C_g \times G$ admits a torsion-free cocompact discrete subgroup Γ . Since $G \cap \Gamma$ is not a lattice of G , $p(\Gamma)$ is not finite. In [11, Chapter 8] it is proved that $\Gamma \cong \mathbb{Z} \times_{\phi} \mathbb{Z}^3$ and for $t \neq 0$ $\phi(t) \in SL_3(\mathbb{Z})$ has a pair of complex conjugate eigenvalues (see [11, Chapter 7]). Hence Γ can be a lattice of a Lie group

$H = \mathbb{R} \times_{\phi} \mathbb{R}^3$ with $\phi(t) = \begin{pmatrix} e^t \cos ct & -e^t \sin ct & 0 \\ e^t \sin ct & e^t \cos ct & 0 \\ 0 & 0 & e^{-2t} \end{pmatrix}$, and $G/\Gamma = H/\Gamma$ is geometrically

formal by Theorem 1.1.

§ 7. Thurston's Geometries and infrasolvmanifold

We say that a compact oriented manifold M admits a geometry (X, g) if $M = X/\Gamma$ where X is a simply connected manifold with a complete Riemannian metric g and Γ is a cocompact discrete subgroup of the group $Isom_g(X)$ of isometries. If (X, g) is a solvable Lie group with an invariant metric g , we call it a solvable Lie type geometry. We consider the following 3-dimensional solvable Lie type geometries.

(3-A) $X = E^3 = \mathbb{R}^3$, $g_{E^3} = dx^2 + dy^2 + dz^2$.

$$(3-B) \ X = Nil^3 = U_3(\mathbb{R}) = \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} : x, y, z \in \mathbb{R} \right\}, \ g_{Nil^3} = dx^2 + dy^2 + (dz - xdy)^2.$$

$$(3-C) \ X = Sol^3 = \mathbb{R} \times_{\phi} \mathbb{R}^2 \text{ with } \phi(z) = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix}, \ g_{Sol^3} = e^{2z}dx^2 + e^{-2z}dy^2 + dz^2.$$

By the theory of geometry and topology of 3-dimensional manifolds we have the following theorem (see [21]).

Theorem 7.1. *A compact aspherical 3-dimensional manifold with the virtually solvable fundamental group admits one of the geometries (3-A~C).*

We also consider the following 4-dimensional solvable Lie type geometries (listed in [24]).

$$(4-A) \ X = E^4 = \mathbb{R}^4, \ g_{E^4} = dx^2 + dy^2 + dz^2 + dt^2.$$

$$(4-B) \ X = Nil^3 \times E = U_3(\mathbb{R}) \times \mathbb{R}, \ g_{Nil^3 \times E} = dx^2 + dy^2 + (dz - xdy)^2 + dt^2.$$

$$(4-C) \ X = Nil^4 = \left\{ \begin{pmatrix} 1 & t & \frac{1}{2}t^2 & z \\ 0 & 1 & t & y \\ 0 & 0 & 1 & x \\ 0 & 0 & 0 & 1 \end{pmatrix} : x, y, z, t \in \mathbb{R} \right\},$$

$$g_{Nil^4} = dx^2 + (dy - tdz)^2 + (dz - tdy + \frac{1}{2}t^2dx)^2 + dt^2$$

$$(4-D) \ X = Sol^3 \times E, \ g_{Sol^3 \times E} = e^{2z}dx^2 + e^{-2z}dy^2 + dz^2 + dt^2.$$

$$(4-E) \ X = Sol_{m,n}^4 = \mathbb{R} \times_{\phi} \mathbb{R}^3 \text{ such that } \phi(t) = \begin{pmatrix} e^{at} & 0 & 0 \\ 0 & e^{bt} & 0 \\ 0 & 0 & e^{ct} \end{pmatrix}, \text{ where } e^a, e^b, e^c \text{ are}$$

distinct roots of $X^3 - mX^2 + nX - 1$ for real numbers $a < b < c$ and integers $m < n$, $g_{Sol_{m,n}^4} = e^{-2at}dx^2 + e^{-2bt}dy^2 + e^{-2ct}dz^2 + dt^2$.

$$(4-F) \ X = Sol_0^4 = \mathbb{R} \times_{\phi} \mathbb{R}^3 \text{ such that } \phi(t) = \begin{pmatrix} e^t & 0 & 0 \\ 0 & e^t & 0 \\ 0 & 0 & e^{-2t} \end{pmatrix},$$

$$g_{Sol_0^4} = e^{-2t}dx^2 + e^{-2t}dy^2 + e^{4t}dz^2 + dt^2.$$

$$(4-G) \ X = Sol_1^4 = \mathbb{R} \times_{\phi} U_3(\mathbb{R}) \text{ such that } \phi(t) \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & e^t x & z \\ 0 & 1 & e^{-t} y \\ 0 & 0 & 1 \end{pmatrix},$$

$$g_{Sol_1^4} = e^{-2t}dx^2 + e^{2t}dy^2 + (dz - xdy)^2 + dt^2.$$

Let G be a simply connected solvable Lie group and g an invariant metric on G . We consider the affine transformation group $\text{Aut}(G) \times G$ and the projection $p : \text{Aut}(G) \times G \rightarrow \text{Aut}(G)$. Let $\Gamma \subset \text{Aut}(G) \times G$ be a torsion-free discrete subgroup such that $p(\Gamma)$ is contained in a compact subgroup of $\text{Aut}(G)$ and the quotient G/Γ is compact. We call G/Γ an infra-solvmanifold. If G is nilpotent, G/Γ is called an infra-nilmanifold. Since $\Gamma \subset \text{Isom}_g(G)$ does not satisfy $\Gamma \subset \text{Aut}(G) \times G$, a compact manifold with a

solvable Lie type geometry is not an infra-solvmanifold in general. Suppose $Isom_g(G) \subset Aut(G) \times G$. Then for an isometry transformation $(\phi, x) \in Aut(G) \times G$, ϕ is an also isometry transformation. By this, for the group C_g of the isometrical automorphisms of G , we have $Isom_g(G) = C_g \times G$. Thus in the assumption $Isom_g(G) \subset Aut(G) \times G$, a compact manifold with a solvable Lie type geometry is an infra-solvmanifold. It is known that for the Euclidian geometry $(E^n, g_{E^n} = dx_1^2 + \dots + dx_n^2)$ we have $Isom_{g_{E^n}} = O(n) \times \mathbb{R}^n$ and the geometries (3-A~C) satisfies $Isom_g(G) \subset Aut(G) \times G$ (see [21]). In [11], Hillman studied the structures of $Isom_g(G)$ of the geometries (4-A~H) and proved $Isom_g(G) \subset Aut(G) \times G$. In [12], Hillman proved the following theorem.

Theorem 7.2. ([12, Theorem 8]) *A 4-dimensional infra-solvmanifold is diffeomorphic to a manifold which admits one of the geometries (4-A~G).*

Remark. By Baues's result in [3], any compact aspherical manifold with the virtually solvable fundamental group is homotopy equivalent to an infra-solvmanifold G/Γ . But for dimension ≥ 4 , there may exist a compact aspherical manifold with virtually solvable fundamental group which is not diffeomorphic to an infra-solvmanifold.

§ 8. Geometrical formality of 3-manifolds

Theorem 8.1. *Let M be a compact oriented aspherical 3-manifold with the virtually solvable fundamental group. If M is a torus or not a nilmanifold, then M is geometrically formal.*

Proof. By Theorem 7.1, it is sufficient to consider the geometries (3-A~C). In the case (3-A), by Corollary 6.1 and the first Bieberbach theorem g_{E^3} is a formal metric on G/Γ .

In the case (3-C), it is known that C_g is isomorphic to the finite dihedral group $D(8)$ (see [21]) and hence by Corollary 6.1 g_{Sol^3} is a formal metric on G/Γ .

Suppose (G, g) is in the case (3-B). Then C_g has two components and the identity component of C_g is isomorphic to a circle S^1 . Let $\Delta = \Gamma \cap G$. By Generalized Bieberbach's theorem (see [1]), Δ is a finite index normal subgroup of Γ . Consider the projection $p : C_g \times G \rightarrow C_g$. If $p(\Gamma)$ is trivial, then $\Gamma \subset G$ is a lattice and G/Γ is a non-toral nilmanifold and hence not formal (see [8]). Suppose $p(\Gamma)$ is non-trivial. By Nomizu's theorem ([18]) we have

$$H^*(G/\Delta, \mathbb{R}) \cong H^*(\mathfrak{g})$$

where \mathfrak{g} is the Lie algebra of G . By this we have

$$H^*(G/\Gamma, \mathbb{R}) \cong H^*(G/\Delta, \mathbb{R})^{\Gamma/\Delta} \cong H^*(\mathfrak{g})^{\Gamma/\Delta}.$$

In [4, Lemma 13.1], it is shown that a non-trivial semisimple automorphism of a nilpotent Lie algebra \mathfrak{g} acts non-trivially on $H^1(\mathfrak{g})$. Since $\Gamma/\Delta \cong p(\Gamma)$ is a nontrivial finite group,

$$H^1(\mathfrak{g})^{\Gamma/\Delta} \neq H^1(\mathfrak{g}).$$

Since $\dim H^1(\mathfrak{g}) = 2$, $\dim H^1(\mathfrak{g})^{\Gamma/\Delta} = 0$ or 1 . If $\dim H^1(\mathfrak{g})^{\Gamma/\Delta} = 0$, then G/Γ is a rational homology sphere and any metric on G/Γ is formal. Suppose $\dim H^1(\mathfrak{g})^{\Gamma/\Delta} = 1$. Then $b_i = 1$ for any $1 \leq i \leq 3$. For any $1 \leq i \leq 3$, we have

$$H^*(G/\Gamma, \mathbb{R}) \cong H^*(\mathfrak{g})^{\Gamma/\Delta} = \bigoplus_{i=1}^3 \langle [\alpha_i] \rangle$$

for non-zero cohomology classes $[\alpha_i] \in H^i(\mathfrak{g})$. We can choose invariant harmonic forms α_i , $i = 1, 2, 3$ for the invariant metric g . Then we have $\mathcal{H}(G/\Gamma) = \bigoplus_{i=1}^3 \langle \alpha_i \rangle$. Since all elements of $\bigwedge^3 \mathfrak{g}^*$ are harmonic, $\alpha_1 \wedge \alpha_2$ is harmonic. For $i < j$ with $(i, j) \neq (1, 2)$, we have $\alpha_i \wedge \alpha_j = 0$. Thus g is a formal metric on G/Γ . This completes the proof of the theorem. \square

Remark. There exists a closed 3-dimensional infra-nilmanifold which is not a nilmanifold. By this theorem such a manifold is geometrically formal.

Example 8.2. Consider $\Gamma = \mathbb{Z} \times_{\phi} \mathbb{Z}^2$ such that $\phi(t) = \begin{pmatrix} (-1)^t & (-1)^t t \\ 0 & (-1)^t \end{pmatrix}$. Then we can embed Γ in $Isom_{g_{Nil^3}}(Nil^3)$ (see [13]). By the direct computation of the lower central series, Γ is non-nilpotent and hence Nil^3/Γ is not a nilmanifold.

§ 9. Aspherical manifolds with the virtually solvable fundamental groups

Theorem 9.1. *Let M be an oriented 4-dimensional infra-solvmanifold. If M is a torus or not a nilmanifold, then M is geometrically formal.*

Proof. In the case (4-A), by Corollary 6.1 and the first Bieberbach theorem g_{E^4} is a formal metric on G/Γ .

In the case (4-D) (resp (4-E)), C_g is isomorphic to the finite group $D(8) \times (\mathbb{Z}/2\mathbb{Z})$ (resp. $(\mathbb{Z}/2\mathbb{Z})^3$) (see [11, Chapter 7]) and hence $g_{Sol^3 \times E}$ (resp. $g_{Sol_{m,n}^4}$) is a formal metric on G/Γ by Corollary 6.1.

As we showed in Example 6.2, in the case (4-F) G/Γ is geometrically formal.

In the case (4-B), the group of all the orientation preserving isomorphisms is $Isom_{g_{Nil^3}} Nil^3 \times \mathbb{R}$ (see [23], [24], or [25]). Thus as the proof of Theorem 8.1 for the case (3-B), if G/Γ is a nilmanifold then G/Γ is not formal, and if G/Γ is an infra-nilmanifold but not a nilmanifold then $g_{Nil^3 \times \mathbb{R}}$ is formal.

In the case (4-C), the group of all the orientation preserving isomorphisms is Nil^4 itself (see [23], [24], or [25]). Thus oriented Nil^4 manifolds are only nilmanifolds and so all oriented Nil^4 manifolds are not formal.

In the case (4-G) we have $Isom_{g_{Sol_1^4}}(Sol_1^4) \cong D(4) \ltimes Sol_1^4$. For any cocompact discrete subgroup $\Gamma \subset Isom_{g_{Sol_1^4}}(Sol_1^4)$, since for the projection $p : \text{Aut}G \ltimes G \rightarrow \text{Aut}(G)$, $p(\Gamma) \subset D(4)$ is finite, we have a subgroup $\Delta \subset \Gamma$ which is a lattice of Sol_1^4 and we have $\mathcal{H}(Sol_1^4/\Gamma) = \mathcal{H}(Sol_1^4/\Delta)^{\Gamma/\Delta}$. In Example 5.6, we showed that the metric $g_{Sol_1^4}$ on the solvmanifold Sol_1^4/Δ is formal. Thus the metric $g_{Sol_1^4}$ on every Sol_1^4 manifold is formal. Hence the theorem follows. □

Finally we prove:

Theorem 9.2. *Let M be a compact oriented aspherical manifold of dimension less than or equal to 4 with the virtually solvable fundamental group. Then M is geometrically formal if and only if M is diffeomorphic to a torus or an infra-solvmanifold which is not a nilmanifold.*

Proof. It is sufficient to show that M is an infra-solvmanifold if M is geometrically formal. If $\dim M \leq 2$, it is obvious. If $\dim M = 3$, it follows from Theorem 7.1. We consider $\dim M = 4$. As Remark 7, M is homotopy equivalent to an infra-solvmanifold G/Γ . It is known that the Euler characteristic $\chi(G/\Gamma)$ of an infra-solvmanifold is 0 (see [11, Chapter 8]). Since G/Γ is an oriented 4-manifold, $\chi(G/\Gamma) = 0$ implies $b_1(G/\Gamma) \neq 0$. Thus we have $b_1(M) \neq 0$. If M is geometrically formal, we have a submersion $M \rightarrow T^{b_1(M)}$ (see [15, Theorem 7]) and hence M is a fiber bundle over a torus $T^{b_1(M)}$. Now we suppose that M is a compact oriented aspherical manifold of dimension 4 with the virtually solvable fundamental group. By the exact sequence of homotopy groups associated by the fiber bundle, the fiber of $M \rightarrow T^{b_1(M)}$ is a compact aspherical manifold of dimension less than or equal 3 with the virtually solvable fundamental group, and hence it is an infra-solvmanifold. Thus M is a fiber bundle whose fiber is an infra-solvmanifold and base space is a torus. By [12, Theorem 7], M is diffeomorphic to an infra-solvmanifold with the fundamental group $\pi_1(M)$. □

Acknowledgements. The author would like to express many thanks to the referee for his careful reading of the earlier version of manuscript with several important remarks, which lead to many improvements in the revised version.

References

- [1] Auslander, L., Bieberbach's theorems on space groups and discrete uniform subgroups of Lie groups, *Ann. of Math. (2)* **71** 1960 579–590.
- [2] de Bartolomeis, P. and Tomassini, A., On solvable generalized Calabi-Yau manifolds, *Ann. Inst. Fourier (Grenoble)* **56** (2006), no. 5, 1281–1296.
- [3] Baues, O., Infra-solvmanifolds and rigidity of subgroups in solvable linear algebraic groups. *Topology* **43** (2004), no. 4, 903–924.
- [4] Baues, O. and Grunewald, O., Automorphism groups of polycyclic-by-finite groups and arithmetic groups, *Publ. Math. Inst. Hautes Etudes Sci.* No. **104** (2006), 213–268.
- [5] Borel, A., *Linear algebraic groups* 2nd enl. ed Springer-verlag (1991).
- [6] Dekimpe, K., Lee, K.-B. and Raymond, F., Bieberbach theorems for solvable Lie groups, *Asian J. Math.* **5** (2001), no. 3, 499–508.
- [7] Deligne, P., Griffiths, P., Morgan, J. and Sullivan, D., Real homotopy theory of Kahler manifolds, *Invent. Math.* **29** (1975), no. 3, 245–274.
- [8] Hasegawa, K., Minimal models of nilmanifolds, *Proc. Amer. Math. Soc.* **106** (1989), no. 1, 65–71.
- [9] Hasegawa, K., Small deformations and non-left-invariant complex structures on six-dimensional compact solvmanifolds, *Differential Geom. Appl.* **28** (2010), no. 2, 220–227.
- [10] Hattori, A., Spectral sequence in the de Rham cohomology of fibre bundles, *J. Fac. Sci. Univ. Tokyo Sect. I* **8** 1960 289–331 (1960).
- [11] Hillman, J.-A., Four-manifolds, geometries and knots, *Topology and Geometry Monographs*, 5.
- [12] Hillman, J.-A., Geometries and infrasolvmanifolds in dimension 4, *Geom. Dedicata* **129** (2007), 57–72.
- [13] Kasuya, H., Formality and hard Lefschetz property of aspherical manifolds, *Osaka J. Math.* **50** (2013) no. 2
- [14] Kasuya, H., Minimal models, formality and hard Lefschetz properties of solvmanifolds with local systems, *J. Differential Geom.* **93**, (2013), 269–298.
- [15] Kotschick, D., On products of harmonic forms. *Duke Math. J.* **107** (2001), no. 3, 521–531.
- [16] Kotschick, D., Terzić, S., On formality of generalized symmetric spaces, *Math. Proc. Cambridge Philos. Soc.* **134** (2003), no. 3, 491–505.
- [17] Nakamura, I., Complex parallelisable manifolds and their small deformations, *J. Differential Geom.* **10** (1975), 85–112.
- [18] Nomizu, K. On the cohomology of compact homogeneous spaces of nilpotent Lie groups, *Ann. of Math. (2)* **59**, (1954). 531–538.
- [19] Raghathan, M.-S., *Discrete subgroups of Lie Groups*, Springer-verlag, New York, 1972.
- [20] Sawai, H., A construction of lattices on certain solvable Lie groups, *Topology Appl.* **154** (2007), no. 18, 3125–3134.
- [21] Scott, P., The geometries of 3-manifolds, *Bull. London Math. Soc.* **15** (1983), no. 5, 401–487
- [22] Sullivan, D., Infinitesimal computations in topology, *Inst. Hautes Études Sci. Publ. Math.* No. **47** (1977), 269–331 (1978).
- [23] Ue, M., Geometric 4-manifolds in the sense of Thurston and Seifert 4-manifolds. I, *J. Math. Soc. Japan* **42** (1990), no. 3, 511–540.
- [24] Wall, C.-T.-C., Geometric structures on compact complex analytic surfaces, *Topology*, **25** (1986), 119–133.
- [25] Wall, C.-T.-C., Geometries and geometric structures in real dimension 4 and complex dimension 2, *In: Proc. of University of Maryland, Geometry and Topology, Lecture Notes*

in Math., **1167**, Springer, 1985, pp. 268–292.