

Asymptotically abelian II_1 -factors

Shōichirō SAKAI

1. Introduction. Recently, physicists (cf. [2], [4], [5], [7], [8], [12], [13], [14]) have introduced the notion of asymptotic abelianess into the theory of operator algebras and obtained various interesting results. In the present paper, we shall extend this notion to finite factors, and by using it, we shall show the existence of a new II_1 -factor. For type III-factors, we shall discuss in another paper.

2. Theorems. First of all, we shall define

Definition 1. Let M be a finite factor and let τ be the unique normalized trace on M . M is called asymptotically abelian, if there exists a sequence of $*$ -automorphisms $\{\rho_n\}$ on M such that $\lim_{n \rightarrow \infty} \|[\rho_n(a), b]\|_2 = 0$ for $a, b \in M$, where $[x, y] = xy - yx$ and $\|x\|_2 = \tau(x^*x)^{1/2}$ for $x, y \in M$.

Let \mathcal{A} be a finite factor, and let φ be the normalized trace on \mathcal{A} . Let $\mathcal{L} = \bigotimes_{n=1}^{\infty} \mathcal{A}_n$ with $\mathcal{A}_n = \mathcal{A}$ be the infinite C^* -tensor product (cf. [3]), and let $\psi = \bigotimes_{n=1}^{\infty} \varphi_n$ with $\varphi_n = \varphi$ be the infinite product trace on \mathcal{L} . Let G be the group of finite permutations of positive integers N , i.e. an element $g \in G$ is an one-to-one mapping of N onto itself which leaves all but a finite number of positive integers fixed.

Then, g will define an $*$ -automorphism, also denoted by g of \mathcal{L} by $g(\Sigma \otimes a_n) = \Sigma \otimes a_{g(n)}$, where $a_n = 1$ for all but a finite number of indices.

For each integer n , we denote by g_n the permutation

$$g_n(k) = \begin{cases} 2^{n-1}+k & \text{if } 1 \leq k \leq 2^{n-1} \\ k-2^{n-1} & \text{if } 2^{n-1} < k \leq 2^n \\ k & \text{if } 2^n < k \end{cases}$$

Then, we can easily show that $\lim_n \|[g_n(a), b]\| = 0$ for $a, b \in \mathcal{L}$. Clearly, the trace ψ on \mathcal{L} is G -invariant --that is, $\psi(g(a)) = \psi(a)$ for $g \in G$ and $a \in \mathcal{L}$.

Let $\{\Pi_\psi, \mathcal{H}_\psi\}$ be the $*$ -representation of \mathcal{L} on a Hilbert space \mathcal{H}_ψ constructed via ψ , then there exists a unitary representation $g \rightarrow U_g$ of G on \mathcal{H}_ψ such that $U_g 1_\psi = 1_\psi$ for $g \in G$ and $\Pi_\psi(g(a)) = U_g \Pi_\psi(a) U_g^*$ for $a \in \mathcal{L}$, where 1_ψ is the image of 1 in \mathcal{H}_ψ . Let \mathcal{M} be the weak closure of $\Pi_\psi(\mathcal{L})$ on \mathcal{H}_ψ , then \mathcal{M} is a finite factor (cf. [3]). The mapping $x \rightarrow U_g x U_g^*$ ($x \in \mathcal{M}$) will define a $*$ -automorphism ρ_g on \mathcal{M} .

Definition 2. The finite factor \mathcal{M} is called the canonical infinite W^* -tensor product of finite factors $\{\mathcal{A}_n\}$ and denoted by $\bigotimes_{n=1}^{\infty} \mathcal{A}_n$.

Proposition 1. $\bigotimes_{n=1}^{\infty} \mathcal{A}_n$ is asymptotically abelian.

Proof. Let τ be the normalized trace on $\bigotimes_{n=1}^{\infty} \mathcal{A}_n$.

We shall identify \mathcal{L} with the image $\Pi_\psi(\mathcal{L})$. Then, $\tau = \psi$ on \mathcal{L} . Let $x, y \in \bigotimes_{n=1}^{\infty} \mathcal{A}_n$, then by Kaplansky's density theorem, there exist two sequences (x_m) and (y_m) in \mathcal{L} such that $\|x_m\| \leq \|x\|$, $\|y_m\| \leq \|y\|$ and $\|x_m - x\|_2 \rightarrow 0$, $\|y_m - y\|_2 \rightarrow 0$ ($m \rightarrow \infty$),

where $\|v\|_2 = \tau(v^*v)^{1/2}$ for $v \in \bigotimes_{n=1}^{\infty} \mathcal{A}_n$.

Then

$$\begin{aligned}
 & \| [\rho_{g_n}(x), y] - [\rho_{g_n}(x_m), y_m] \|_2 \\
 & \leq \| [\rho_{g_n}(x), y] - [\rho_{g_n}(x), y_m] + [\rho_{g_n}(x), y_m] - [\rho_{g_n}(x_m), y_m] \|_2 \\
 & \leq \| [\rho_{g_n}(x), y - y_m] \|_2 + \| [\rho_{g_n}(x) - \rho_{g_n}(x_m), y_m] \|_2 \\
 & \leq \| \rho_{g_n}(x) \cdot (y - y_m) \|_2 + \| (y - y_m) \rho_{g_n}(x) \|_2 + \| \rho_{g_n}(x - x_m) y_m \|_2 + \| y_m \rho_{g_n}(x - x_m) \|_2 \\
 & \leq \| \rho_{g_n}(x) \| \| y - y_m \|_2 + \| y - y_m \|_2 \| \rho_{g_n}(x) \| + \| \rho_{g_n}(x - x_m) \|_2 \| y_m \| \\
 & \quad + \| y_m \| \| \rho_{g_n}(x - x_m) \|_2 \\
 & = 2 \| x \| \| y - y_m \|_2 + 2 \| y_m \| \| x - x_m \|_2 \rightarrow 0 \quad (m \rightarrow \infty).
 \end{aligned}$$

Hence, for arbitrary $\varepsilon > 0$, there exists an m_0 such that

$$| \| [\rho_{g_n}(x), y] \|_2 - \| [\rho_{g_n}(x_{m_0}), y_{m_0}] \|_2 | < \varepsilon \quad \text{for all } n.$$

On the other hand, $\| [\rho_{g_n}(x_{m_0}), y_{m_0}] \|_2 < \varepsilon$ for $n \geq n_0$, where n_0 is some integer; hence $\| [\rho_{g_n}(x), y] \|_2 < 2\varepsilon$ for $n \geq n_0$. This completes the proof.

Now let Φ be a countably discrete group, and let $\mathcal{L}(\Phi)$ be the W^* -algebra generated by the left regular representation of Φ . We shall show examples of asymptotically abelian finite factors.

Example 1. Let \mathcal{L}_1 be the type I_1 -factor, then clearly it is asymptotically abelian.

Example 2. Let Π be the countably discrete group of all finite permutations on the set of all positive integers, then the W^* -algebra $\mathcal{L}(\Pi)$ is a hyperfinite II_1 -factor (cf. [6]). Since all hyperfinite II_1 -factors on separable Hilbert

spaces are *-isomorphic (cf. [6]), $\mathcal{L}(\Pi)$ is *-isomorphic to $\bigotimes_{n=1}^{\infty} \mathcal{L}_{2,n}$, with $\mathcal{L}_{2,n} = \mathcal{L}_2$, where \mathcal{L}_2 is the type I_2 -factor. Since the asymptotic abelianess is preserved under an *-isomorphism, by Proposition 1, $\mathcal{L}(\Pi)$ is asymptotically abelian.

Example 3. Let ϕ_2 be the countably discrete, free group with two generators, then the W^* -algebra $\mathcal{L}(\phi_2)$ is a II_1 -factor (cf. [6]).

Let $\bigotimes_{n=1}^{\infty} \mathcal{D}_n$ with $\mathcal{D}_n = \mathcal{L}(\phi_2)$, then by Proposition 1, $\bigotimes_{n=1}^{\infty} \mathcal{D}_n$ is asymptotically abelian.

Now, we shall show examples of finite factors which are not asymptotically abelian.

Example 4. Let \mathcal{L}_p be the type I_p -factor with $2 \leq p < +\infty$ (p integer), then \mathcal{L}_p is not asymptotically abelian.

Proof. Let (ρ_n) be a sequence of *-automorphisms on \mathcal{L}_p . Let $B(\mathcal{L}_p)$ be the Banach algebra of all bounded operators on \mathcal{L}_p , then $B(\mathcal{L}_p)$ is finite-dimensional; therefore there exists a subsequence (ρ_{n_j}) of (ρ_n) such that $\|\rho_{n_j} - T\| \rightarrow 0$ ($j \rightarrow \infty$), where T is a bounded operator on \mathcal{L}_p . It is easy to show that T is also a *-automorphisms on \mathcal{L}_p ; hence clearly \mathcal{L}_p is not asymptotically abelian.

Example 5. Let ϕ_2 be the countably discrete, free group with two generators, then $\mathcal{L}(\phi_2)$ is not asymptotically abelian.

Proof. Suppose that $\mathcal{L}(\phi_2)$ is asymptotically abelian, and let (ρ_n) be a family of *-automorphisms such that

$$\|[\rho_n(a), b]\|_2 \rightarrow 0 \quad (n \rightarrow \infty) \quad \text{for } a, b \in \mathcal{L}(\phi_2).$$

Clearly, there exists an unitary element u in $\mathcal{L}(\phi_2)$ such that $\tau(u) = 0$, where τ is the normalized trace on $\mathcal{L}(\phi_2)$. Then $\|[\rho_n(u), b]\|_2 = \|\rho_n(u)b - b\rho_n(u)\|_2 = \|\rho_n(u)b\rho_n(u)^* - b\|_2 \rightarrow 0$

$(n \rightarrow \infty)$ for $b \in \mathcal{L}(\Phi_2)$.

Since $\rho_n(u)$ is unitary and $\tau(\rho_n(u)) = \tau(u) = 0$, $\mathcal{L}(\Phi_2)$ has the property Γ ; this is a contradiction (cf. [6]).

Example 6. Let Π be the group of all finite permutations on the set of all positive integers, and let Φ_2 be the free group of two generators, and let $\Phi_2 \times \Pi$ be the direct product group of Φ_2 and Π . Then, $\mathcal{L}(\Phi_2 \times \Pi)$ is $*$ -isomorphic to the W^* -tensor product $\mathcal{L}(\Phi_2) \bar{\otimes} \mathcal{L}(\Pi)$ of $\mathcal{L}(\Phi_2)$ and $\mathcal{L}(\Pi)$ (cf. [6], [9]).

In the following considerations, we shall show that $\mathcal{L}(\Phi_2 \times \Pi)$ is not asymptotically abelian.

Lemma 1. Let ϕ be a group and let E a subset of ϕ . Suppose there exists a subset $F \subset E$ and two elements $g_1, g_2 \in \phi$ such that (i) $F \cup g_1 F g_1^{-1} = E$; (ii) $F, g_2^{-1} F g_2$ and $g_2 F g_2^{-1}$ are contained in E and mutually disjoint. Let $f(g)$ be a complex valued function on ϕ such that $\sum_{g \in \phi} |f(g)|^2 < +\infty$ and $(\sum_{g \in \phi} |f(g_i g g_i^{-1}) - f(g)|^2)^{1/2} < \epsilon$ ($i = 1, 2$). Then, $(\sum_{g \in E} |f(g)|^2)^{1/2} < K\epsilon$, where K does not depend on ϵ and f .

Proof. $v(x) = \sum_{g \in X} |f(g)|^2$ for a subset $X \subset \phi$.

Then,

$$\epsilon > (\sum_{g \in \phi} |f(g_1 g g_1^{-1}) - f(g)|^2)^{1/2} \geq |v(g_1 F g_1^{-1})^{1/2} - v(F)^{1/2}|.$$

Putting $v(E)^{1/2} = s$, then

$$|v(g_1 F g_1^{-1}) - v(F)| = |v(g_1 F g_1^{-1})^{1/2} + v(F)^{1/2}| \cdot |v(g_1 F g_1^{-1})^{1/2} - v(F)^{1/2}|$$

$$\leq 2s\epsilon \text{ and so } v(g_1 F g_1^{-1}) \leq v(F) + 2s\epsilon;$$

hence

$$s^2 \leq v(g_1 F g_1^{-1}) + v(F) < 2(v(F) + s\epsilon),$$

so that

$$v(F) > \frac{s^2}{2} - s\epsilon.$$

Since

$$(\sum_{g \in \phi} |f(g_2 g g_2^{-1}) - f(g)|^2)^{1/2} = (\sum_{g \in \phi} |f(g_2 g_2^{-1} g g_2 g_2^{-1}) - f(g_2^{-1} g g_2)|^2)^{1/2},$$

analogously we have

$$|\nu(g_2 F g_2^{-1}) - \nu(F)| < 2s\epsilon$$

and

$$|\nu(g_2^{-1} F g_2) - \nu(F)| < 2s\epsilon ;$$

hence

$$\nu(g_2 F g_2^{-1}) > \nu(F) - 2s\epsilon > \frac{s^2}{2} - 3s\epsilon$$

and

$$\nu(g_2^{-1} F g_2) > \frac{s^2}{2} - 3s\epsilon .$$

Therefore,

$$s^2 = \nu(E) \geq \nu(F) + \nu(g_2^{-1} F g_2) + \nu(g_2 F g_2^{-1}) > \frac{3}{2}s^2 - 7s\epsilon ;$$

hence

$$s < 14\epsilon .$$

This completes the proof.

Now, let us consider the group $\Phi_2 \times \Pi$. Let k_1, k_2 be the generators of the group Φ_2 , and let F_1 be the set of elements $\in \Phi_2$ which when written as a power of k_1, k_2 of

minimum length end with k_1^n , $n = \pm 1, \pm 2, \dots$. Let $F = F_1 \times \Pi$ and let $a_1 = (k_1, e)$ and $a_2 = (k_2, e)$, where e is the unit of Π .

Then,

$$a_1 F a_1^{-1} = (k_1 F_1 k_1^{-1}, \Pi)$$

and

$$a_2 F a_2^{-1} = (k_2 F_1 k_2^{-1}, \Pi);$$

moreover

$$F \cup a_1 F a_1^{-1} = (F_1, \Pi) \cup (k_1 F_1 k_1^{-1}, \Pi) = (F_1 \cup k_1 F_1 k_1^{-1}, \Pi) = (e, \Pi)^c,$$

where $(\cdot)^c$ is the complement of (\cdot) ; F , $a_1^{-1} F a_1$ and $a_2 F a_2^{-1}$ are contained in $(e, \Pi)^c$ and mutually disjoint. Hence by Lemma 1, we have

Lemma 2. Suppose that (f_n) be a sequence of complex valued functions on $\Phi_2 \times \Pi$ such that $(\sum_{a \in \Phi_2 \times \Pi} |f_n(a)|^2)^{1/2} < +\infty$

and

$$\lim_{n \rightarrow \infty} (\sum_{a \in \Phi_2 \times \Pi} |f_n(a_i a a_i^{-1}) - f_n(a)|^2)^{1/2} = 0 \quad (i = 1, 2).$$

Then,

$$\lim_{n \rightarrow \infty} (\sum_{a \in (e, H)^c} |f_n(a)|^2)^{1/2} = 0.$$

Now we shall show

Theorem 1. $\mathcal{L}(\Phi_2 \times \Pi)$ is not asymptotically abelian.

Proof. Suppose that $\mathcal{L}(\Phi_2 \times \Pi)$ is asymptotically abelian, and let (ρ_n) be a sequence of *-automorphisms on $\mathcal{L}(\Phi_2 \times \Pi)$ such that

$$\|[\rho_n(x), y]\|_2 \rightarrow 0 \quad (n \rightarrow \infty)$$

for $x, y \in \mathcal{L}(\Phi_2 \times \Pi)$.

Let ε_t ($t \in \Phi_2 \times \Pi$) be the unitary element of $\mathcal{L}(\Phi_2 \times \Pi)$ such that $(\varepsilon_t f)(a) = f(t^{-1}a)$ for $f \in \ell^2(\Phi_2 \times \Pi)$ and $a \in \Phi_2 \times \Pi$, where $\ell^2(\Phi_2 \times \Pi)$ is the Hilbert space of all complex valued square summable functions on $\Phi_2 \times \Pi$.

Since all elements of $\mathcal{L}(\Phi_2 \times \Pi)$ are realized as left convolution operators by elements of a subset of $\ell^2(\Phi_2 \times \Pi)$ (cf. [6]), we shall embed $\mathcal{L}(\Phi_2 \times \Pi)$ into $\ell^2(\Phi_2 \times \Pi)$. Then, $x \in \mathcal{L}(\Phi_2 \times \Pi)$ is a complex valued square summable function on $\Phi_2 \times \Pi$.

Now let x_1, x_2, \dots, x_p be a finite subset of elements of $\mathcal{L}(\Phi_2 \times \Pi)$. Then,

$$\|[\rho_n(x_j), \varepsilon_{a_i}]\|_2 \rightarrow 0 \quad (n \rightarrow \infty)$$

for $i = 1, 2$ and $j = 1, 2, \dots, p$.

$$\begin{aligned} \|\rho_n(x_j), \varepsilon_{a_i}\|_2 &= \|\rho_n(x_j)\varepsilon_{a_i} - \varepsilon_{a_i}\rho_n(x_j)\|_2 \\ &= \|\varepsilon_{a_i}^{-1}\rho_n(x_j)\varepsilon_{a_i} - \rho_n(x_j)\|_2 = \left(\sum_{a \in \Phi_2 \times \Pi} |\rho_n(x_j)(a_i a a_i^{-1}) - \rho_n(x_j)(a)|^2 \right)^{1/2}. \end{aligned}$$

Hence, by Lemma 2,

$$\left(\sum_{a \in (e, \Pi)^c} |\rho_n(x_j)(a)|^2 \right)^{1/2} \rightarrow 0 \quad (n \rightarrow \infty).$$

Put

$$f_n(x_j)(a) = \begin{cases} \rho_n(x_j)(a) & \text{if } a \in (e, \Pi) \\ 0 & \text{if } a \notin (e, \Pi) \end{cases}$$

, then we can easily show that $f_n(x_j) \in \mathcal{L}(\Phi_2 \times \Pi)$. Let

$\mathcal{H} = \{\lambda \mid \lambda(a) = 0 \text{ for } a \notin (e, \Pi) \text{ and } \lambda \in \mathcal{L}(\Phi_2 \times \Pi)\}$, then \mathcal{H} is a W^* -subalgebra of $\mathcal{L}(\Phi_2 \times \Pi)$; moreover put $\tilde{\lambda}(h) = \lambda(e, h)$ for $h \in \Pi$ and $\lambda \in \mathcal{H}$, then the mapping $\lambda \rightarrow \tilde{\lambda}$ is an $*$ -isomorphism of \mathcal{H} onto the II_1 -factor $\mathcal{L}(\Pi)$; hence \mathcal{H} is a hyperfinite II_1 -factor.

For arbitrary $\varepsilon > 0$, there exists a positive integer n_0 such that

$$\left(\sum_{a \in (e, \Pi)^c} |\rho_{n_0}(x_j)(a)|^2 \right)^{1/2} < \varepsilon \quad \text{for } j = 1, 2, \dots, p.$$

Then,

$$\|\rho_{n_0}(x_j) - f_{n_0}(x_j)\|_2 < \varepsilon \quad \text{for } j = 1, 2, \dots, p$$

Since \mathcal{H} is a hyperfinite II_1 -factor, there exist a type I_{n_p} subfactor \mathcal{L}_{n_p} of \mathcal{H} and elements r_1, r_2, \dots, r_n such that

$$\|f_{n_0}(x_j) - r_j\|_2 < \varepsilon \quad \text{for } j = 1, 2, \dots, p.$$

Therefore,

$$\|\rho_{n_0}(x_j) - r_j\|_2 < 2\varepsilon \text{ for } j = 1, 2, \dots, p.$$

Since ρ_{n_0} is a $*$ -automorphism, $\|x_j - \rho_{n_0}^{-1}(r_j)\|_2 < 2\varepsilon$ and $\rho_{n_0}^{-1}(r_j) \in \rho_{n_0}^{-1}(\mathcal{L}_{n_p})$ for $j = 1, 2, \dots, p$. $\rho_{n_0}^{-1}(\mathcal{L}_{n_p})$ is a type I_{n_p} factor and $\mathcal{L}(\Phi_2 \times \Pi)$ is a II_1 -factor on a separable Hilbert space; hence by the result of Murray and von Neumann [6], $\mathcal{L}(\Phi_2 \times \Pi)$ is a hyperfinite II_1 -factor.

On the other hand, by Schwartz's theorem [10], $\mathcal{L}(\Phi_2 \times \Pi)$ is not hyperfinite. This is a contradiction and completes the proof.

Now we shall show the existence of the fifth example of II_1 -factors on separable Hilbert spaces.

Corollary 1. $\mathcal{L}(\phi_2 \times \mathbb{I})$ is not *-isomorphic to $\bigotimes_{n=1}^{\infty} \mathcal{D}_n$ with $\mathcal{D}_n = \mathcal{L}(\phi_2)$.

Proof. Clearly the asymptotic abelianess is preserved under *-isomorphisms; hence $\mathcal{L}(\phi_2 \times \mathbb{I})$ is not *-isomorphic to $\bigotimes_{n=1}^{\infty} \mathcal{D}_n$. This completes the proof.

Proposition 2. $\bigotimes_{n=1}^{\infty} \mathcal{D}_n \bar{\otimes} \mathcal{L}(\mathbb{I}) = \bigotimes_{n=1}^{\infty} \mathcal{D}_n$, where $\mathcal{D}_n = \mathcal{L}(\phi_2)$ and $(\cdot) \bar{\otimes} (\cdot)$ is the W*-tensor product of (\cdot) and (\cdot) .

Proof. Since $\mathcal{L}(\phi_2)$ is a II_1 -factor, there exists a type I_2 -factor \mathcal{L}_2 such that $\mathcal{L}(\phi_2) = \mathcal{L}_2 \bar{\otimes} \mathcal{L}'_2$, where \mathcal{L}'_2 is the commutant of \mathcal{L}_2 in $\mathcal{L}(\phi_2)$; hence $\bigotimes_{n=1}^{\infty} \mathcal{D}_n = \bigotimes_{n=1}^{\infty} \mathcal{L}_{2,n} \bar{\otimes} \mathcal{L}'_{2,n}$, where $\mathcal{L}_{2,n} = \mathcal{L}_2$ and $\mathcal{L}'_{2,n} = \mathcal{L}'_2$.

Hence

$$\begin{aligned} \bigotimes_{n=1}^{\infty} \mathcal{D}_n \bar{\otimes} \mathcal{L}(\mathbb{I}) &= \left(\bigotimes_{n=1}^{\infty} \mathcal{L}_{2,n} \right) \bar{\otimes} \left(\bigotimes_{n=1}^{\infty} \mathcal{L}'_{2,n} \bar{\otimes} \mathcal{L}(\mathbb{I}) \right) \\ &= \bigotimes_{n=1}^{\infty} \mathcal{L}_{2,n} \bar{\otimes} \mathcal{L}_{2,n} = \bigotimes_{n=1}^{\infty} \mathcal{D}_n \end{aligned}$$

because $\bigotimes_{n=1}^{\infty} \mathcal{L}_{2,n}$ and $\mathcal{L}(\mathbb{I})$ are hyperfinite and so $\bigotimes_{n=1}^{\infty} \mathcal{L}_{2,n} \bar{\otimes} \mathcal{L}(\mathbb{I})$ is also hyperfinite.

This completes the proof.

The following definition is due to Ching [1]

Definition 2. A finite factor M is said to have property C, if for each sequence $u_n (n=1, 2, \dots)$ of unitary elements in M with the property that

$$\|u_n^* x u_n - x\|_2 \rightarrow 0 \quad (n \rightarrow \infty)$$

for each $x \in M$, there exists a uniformly bounded sequence v_n ($n=1,2,\dots$) of mutually commuting elements in M such that $\|u_n - v_n\|_2 \rightarrow 0$ ($n \rightarrow \infty$).

Then, Ching [1] proved that $\mathcal{L}(\Pi)$ and $\mathcal{L}(\phi_2 \times \Pi)$ have not property C and also there exists a type II_1 -factor M_4 which has both of properties C and Γ .

It is not so difficult to see that $\mathcal{L}(\phi_2)$ has property C, although we do not need it here.

Corollary 2. $\bigotimes_{n=1}^{\infty} \mathcal{A}_n$ with $\mathcal{A}_n = \mathcal{L}(\phi_2)$ has not property C.

Let g_i be the element in Π which permutes i and $i+1$ and leaves all other positive integers fixed, for each $i=1,2,\dots$.

Clearly $\|\varepsilon_{g_i}^* x \varepsilon_{g_i} - x\|_2 \rightarrow 0$ for $x \in \mathcal{L}(\Pi)$.

Hence, let 1 be the unit of $\bigotimes_{n=1}^{\infty} \mathcal{A}_n$, then

$\|1 \otimes \varepsilon_{g_i}^* y 1 \otimes \varepsilon_{g_i} - y\|_2 \rightarrow 0$ for $y \in \bigotimes_{n=1}^{\infty} \mathcal{A}_n \bar{\otimes} \mathcal{L}(\Pi)$. Suppose

$\bigotimes_{n=1}^{\infty} \mathcal{A}_n$ has the property C, then $\bigotimes_{n=1}^{\infty} \mathcal{A}_n \bar{\otimes} \mathcal{L}(\Pi)$ has property C.

Let $\{v_i\}$ be a uniformly bounded sequence of mutually commuting elements in $\bigotimes_{n=1}^{\infty} \mathcal{A}_n \bar{\otimes} \mathcal{L}(\Pi)$ such that $\|1 \otimes \varepsilon_{g_i} - v_i\|_2 \rightarrow 0$ ($i \rightarrow \infty$).

Then, since $g_i g_{i+1} \neq g_{i+1} g_i$,

$$\begin{aligned} \sqrt{2} &= \|1 \otimes \varepsilon_{(g_i g_{i+1})} \otimes \varepsilon_{(g_{i+1} g_i)}\|_2 \\ &= \|1 \otimes \varepsilon_{g_i} 1 \otimes \varepsilon_{g_{i+1}} - 1 \otimes \varepsilon_{g_{i+1}} 1 \otimes \varepsilon_{g_i}\|_2 \\ &\leq \|(1 \otimes \varepsilon_{g_i} - v_i) 1 \otimes \varepsilon_{g_{i+1}}\|_2 + \|v_i (1 \otimes \varepsilon_{g_{i+1}} - v_{i+1})\|_2 \\ &\quad + \|(v_{i+1} 1 \otimes \varepsilon_{g_{i+1}}) v_i\|_2 + \|1 \otimes \varepsilon_{g_{i+1}} (v_i \otimes \varepsilon_{g_i})\|_2 \rightarrow 0 \quad (n \rightarrow \infty). \end{aligned}$$

This is a contradiction and completes the proof.

Proposition 4. There are five examples of II_1 -factors with different algebraical types on separable Hilbert spaces.

Proof. $\bigotimes_{n=1}^{\infty} \mathcal{D}_n$ with $\mathcal{D}_n = \mathcal{L}(\Phi_2) \neq \mathcal{L}(\Pi)$, because $\mathcal{L}(\Pi)$ can not contain the II_1 -factor which is $*$ -isomorphic to $\mathcal{L}(\Phi_2)$ as a W^* -subalgebra (cf. [10]).

Clearly $\bigotimes_{n=1}^{\infty} \mathcal{D}_n \neq \mathcal{L}(\Phi_2)$, because $\bigotimes_{n=1}^{\infty} \mathcal{D}_n = \bigotimes_{n=1}^{\infty} \mathcal{D}_n \otimes \mathcal{L}(\Pi)$ has property Γ ; by Theorem 1 $\bigotimes_{n=1}^{\infty} \mathcal{D}_n \neq \mathcal{L}(\Phi_2 \times \Pi)$; by Proposition 3, $\bigotimes_{n=1}^{\infty} \mathcal{D}_n \neq M_4$.

This completes the proof.

References

1. W. Ching, Non-isomorphic non-hyperfinite factors, to appear in the Canadian Journal of Math.
2. S. Doplicher, R. Kadison, D. Kastler and D. Robinson, Asymptotically abelian systems, Commun. Math. Phys. 6 (1967), 101-120.
3. A. Guichardet, Ann. Scient. Ec. Norm. Sup. 83 (1966), 1-52.
4. R. Haag, D. Kastler and L. Michel, Central decompositions of Ergodic states, to appear.
5. O. Lanford and D. Ruelle, Integral representation of invariant states on R^* -algebras, J. Math. Phys. 8 (1967) 1460-1463.
6. F. Murray and J. von Neumann, On rings of operators IV, Ann of Math. 44 (1943), 716-808.
7. R. Powers, Representations of uniformly hyperfinite algebras and their associated von Neumann ring, Ann. of Math. 86 (1967), 138-171.
8. D. Ruelle, States of Physical systems, Commun. Math. Phys. 3 (1966), 133-150.
9. S. Sakai, The theory of W^* -algebras, Lecture notes Yale University, 1962.
10. S. Sakai, On the hyperfinite II_1 -factor, to appear in the proceeding of Amer. Math. Soc.
11. T. Schwartz, Two finite, non-hyperfinite non-isomorphic factors, Comm. Pure Appl. Math., 16 (1963), 19-26.
12. E. Størmer, Commun. Math. Phys. 5 (1967), 1-22.
13. E. Størmer, Symmetric states of infinite tensor products of C^* -algebras, to appear.

14. H. Araki, On the algebra of all local observables,
Progress of Theoretical Physics, 32 (1964) 844-854.

University of Pennsylvania