

Pontrjagin classes of rational homology manifold

(Report on work by Don Zagier [3])

Lecture by F. Hirzebruch. (Notes by S. Morita)

1. L-classes in the equivariant case.

Let  $X$  be a compact oriented rational homology manifold and assume that a compact Lie group  $G$  acts on  $X$  by orientation preserving simplicial homeomorphisms. Then there is defined the equivariant signature  $\text{sign}(g, X) \in \mathbb{C}$  for any  $g \in G$  as follows (see [1]).

(i) If  $\dim X \equiv 1 \pmod{2}$ , we put  $\text{sign}(g, X) = 0$ .

(ii) If  $\dim X = 4k$ , then the cup product defines a non-degenerate quadratic form  $B$  on  $H^{2k}(X; \mathbb{Q})$ . Let

$$H^{2k}(X; \mathbb{Q}) = V^+ \oplus V^-$$

be an equivariant decomposition of the  $G$ -vector space  $H^{2k}(X; \mathbb{Q})$  such that  $B$  is positive (negative) definite on  $V^+$  ( $V^-$ ). Then we define

$$\text{sign}(g, X) = \text{Trace}(g|V^+) - \text{Trace}(g|V^-)$$

Observe that if  $g$  acts on  $X$  trivially, then

$\text{sign}(g, X) = \text{sign } X$ , where  $\text{sign } X$  is the ordinary signature of  $X$ .

(iii) If  $\dim X = 4k + 2$ , then we can give a complex vector space structure to  $H^{2k+1}(X; \mathbb{R})$  such that the action of  $G$  preserves this structure. We define

$$\text{sign}(g, X) = 2i \text{Im}(\text{Trace } g|H^{2k+1}(X; \mathbb{R})).$$

Now Thom defined the Pontrjagin classes (or equivalently the L-classes) for any rational homology manifold. Then Milnor simplified the Thom's definition by using a  $t$ -regularity argument and the (ordinary) signature.

Recently Zagier has generalized this procedure to the equivariant case.

Precisely, assume that a finite group  $G$  acts on a compact oriented rational homology manifold  $X$ . Then Zagier has defined the "equivariant L-class"

$$L(g, X) \in H^*(X; \mathbb{C})$$

for any  $g \in X$ . This class can be used to calculate the ordinary L-class of the rational homology manifold  $X/G$ , by virtue of the following theorem. This theorem is one of the main results of Zagier.

Theorem 1. Let  $G$  be a finite group and  $X$  a compact oriented rational homology  $G$ -manifold. Let

$$\pi : X \rightarrow X/G$$

be the natural projection. Then

$$\frac{1}{\deg \pi} \pi^* L(X/G) = \frac{1}{|G|} \sum_{g \in G} L(g, X).$$

Here  $\deg \pi$  is the degree of the map  $\pi$  (we do not assume that the action of  $G$  is effective) and  $L(X/G)$  is the Thom-Milnor L-class of the rational homology manifold  $X/G$ .

We will sketch the definition of the class  $L(g, X)$  for the case when  $X$  is a differentiable  $G$ -manifold.

The proof of Theorem 1 in the differentiable case then follows from a calculation depending on Milnor's definition of the L-class  $L(X/G)$  and the Atiyah-Singer  $G$ -signature theorem.

The general case (i.e. the case when  $X$  is only a rational homology  $G$ -manifold) follows from a parallel extension in the equivariant context of Milnor's argument.

Thus let  $X$  be a compact oriented differentiable  $G$ -manifold, where  $G$  is a finite group. Let  $X^g = \{x \in X \mid gx = x\}$ , the fixed point set of  $g$ . Then by Atiyah-Singer [1],

Theorem (G-signature theorem)

$$\text{sign.}(g, X) = L'(g, X) [X^g]$$

for a certain class  $L'(g, X) \in H^*(X^g; \mathbb{C})$ , defined below.

Now the right hand side of the above equation depends only on the top dimensional components of the class  $L'(g, X^g)$ . However to define the equivariant L-class, lower terms of  $L'(g, X)$  are also necessary. Since the "correct" class  $L'(g, X)$  for our purpose differs from the original one defined by Atiyah-Singer [1] by powers of two, we define it explicitly.

Let  $N^g$  be the normal bundle of  $X^g$  in  $X$ . Then  $N^g$  can be decomposed equivariantly as follows,

$$N^g = N^g(-1) \oplus \sum_{0 < \theta < \pi} N^g_\theta$$

where  $N^g(-1)$  is a real bundle over  $X^g$  on which  $g$  acts as  $-1$ .

$N^g_\theta$  is a complex bundle on which  $g$  acts as  $e^{i\theta}$ . We define

$$L_{-1}(N^g(-1)) = e(N^g(-1))L(N^g(-1))^{-1}$$

where  $L(N^g(-1))$  is the L-class of the real bundle  $N^g(-1)$  and

$e(N^g(-1))$  is the Euler class. For the complex part  $N^g_\theta$ , we define

$$L_\theta(N^g_\theta) = (\coth \frac{i\theta}{2})^q \prod_j \frac{\coth(X_j + \frac{i\theta}{2})}{\coth \frac{i\theta}{2}}$$

where  $q = \dim_{\mathbb{C}} N^g_\theta$  and  $X_j$  is the usual formal class such that the Chern classes are the elementary symmetric polynomials in  $x_j$ 's. Now

we define

$$L'(g, X) = L(X^g)L_{-1}(N^g(-1)) \prod_{0 < \theta < \pi} L_\theta(N^g_\theta).$$

We are now prepared to define the equivariant L-class,  $L(g, X)$ . Let

$j: X^g \rightarrow X$  be the inclusion map. Then we simply define

$$L(g, X) = j^*L'(g, X)$$

where  $j!$  is the Gysin homomorphism.

We will give two applications of Theorem 1 in §§ 2, 3.

One is the case of linear actions of complex projective space  $P_n \mathbb{C}$  (§ 2) and the other is the action of the symmetric group of degree  $n$ ,  $S_n$ , on

$$S^n = \underbrace{X \times \cdots \times X}_{n \text{ times}} \quad (\S 3)$$

## 2. Complex projective space

Let  $P_n \mathbb{C} = \{[z_0, z_1, \dots, z_n] \mid z_i \in \mathbb{C}\}$  be  $n$ -dimensional complex projective space. We define a finite group  $G_a$  by

$$G_a = G_{a_0} \times G_{a_1} \times \cdots \times G_{a_n}$$

$$G_{a_j} = \{\lambda \mid \lambda^{a_j} = 1\}.$$

Then  $G_a$  acts on  $P_n \mathbb{C}$  by

$$(\lambda_0, \lambda_1, \dots, \lambda_n)[z_0, z_1, \dots, z_n] = [\lambda_0 z_0, \lambda_1 z_1, \dots, \lambda_n z_n]$$

$$(\lambda_0, \lambda_1, \dots, \lambda_n) \in G_a, \quad [z_0, z_1, \dots, z_n] \in P_n \mathbb{C}.$$

Let  $\pi : P_n \mathbb{C} \rightarrow P_n \mathbb{C}/G_a$  be the natural projection. Then Bott has calculated

Theorem 2 (Bott)

$$\pi^* L(P_n \mathbb{C}/G_a) = \frac{1}{d} \sum_{0 \leq \xi < \pi} \prod_{j=0}^n \frac{a_j x}{\tanh(a_j(x + i\xi))}$$

where  $d$  is the greatest common divisor of the natural numbers  $a_0, a_1, \dots, a_n$  and  $x \in H^2(P_n \mathbb{C})$  is the standard generator.

The sum on the right hand side is taken over all real numbers  $\xi \in [0, \pi)$ . However the product  $\prod_{j=0}^n \frac{a_j x}{\tanh(a_j(x + i\xi))}$  is equal to zero, unless there is at least one  $a_j$  such that  $a_j \xi$  is a multiple of  $\pi$  (because  $x^{n+1} = 0$ ). Therefore the sum is well-defined.

Now this theorem can be obtained from Theorem 1 as follows.

Proof of Theorem 2 using Theorem 1. By Theorem 1, we have

$$(1) \quad \pi^* L(P_n \mathbb{C}/G_a) = \frac{\deg \pi}{|G_a|} \sum_{g \in G_a} L(g, P_n \mathbb{C}).$$

But it is easy to see that

$\frac{\deg \pi}{|G_a|} = \frac{1}{d}$ . Hence we have only to show that

$$(2) \quad \sum_{g \in G_a} L(g, P_n \mathbb{C}) = \sum_{0 \leq \xi < \pi} \prod_{j=0}^n \frac{a_j x}{\tanh(a_j(x+i\xi))}.$$

Let  $g = (\zeta_0, \zeta_1, \dots, \zeta_n)$ ,  $\zeta_j \in G_{a_j}$ . Then

$$(3) \quad P_n \mathbb{C}^{\mathcal{G}} = \{ [z_0, z_1, \dots, z_n] \mid \zeta_j z_j = \zeta z_j \text{ for } j = 0, 1, \dots, n \\ \text{some } \zeta \in S^1 \} = \bigcup_{\zeta \in S^1} X(\zeta)$$

where  $X(\zeta) = \{ [z_0, z_1, \dots, z_n] \in P_n \mathbb{C}^{\mathcal{G}} \mid \zeta_j z_j = \zeta z_j \text{ for all } j \}$ .

Clearly if  $\zeta \notin \{ \zeta_0, \zeta_1, \dots, \zeta_n \}$ . Then  $X(\zeta) = \emptyset$ , while  $X(\zeta_j)$ , is isomorphic to  $P_s \mathbb{C}$ , where  $s+1$  is the number of indices  $i$  with  $\zeta_i = \zeta_j$ .

Now by the definition of the equivariant L-class, we have

$$L(g, P_n \mathbb{C}) = j! L'(g, P_n \mathbb{C}) \quad \text{where } j: P_n \mathbb{C}^{\mathcal{G}} \rightarrow P_n \mathbb{C} \text{ is the inclusion.}$$

Thus we must calculate the class

$$L'(g, P_n \mathbb{C}) \in H^*(P_n \mathbb{C}^{\mathcal{G}}; \mathbb{C}).$$

Let  $L'(g, P_n \mathbb{C})_{\zeta}$  be the component of  $L'(g, P_n \mathbb{C})$  corresponding to the connected component  $X(\zeta) \subset P_n \mathbb{C}^{\mathcal{G}}$ .

As mentioned earlier,  $X(\zeta)$  is isomorphic to  $P_s \mathbb{C}$  and it is easy to check that, to calculate  $L(g, P_n \mathbb{C})_{\zeta}$ , we may assume that

$$X(\zeta) = P_s \mathbb{C} \subset P_n \mathbb{C}, \quad \text{where}$$

$$P_s \mathbb{C} = \{ [z_0, z_1, \dots, z_s, 0, \dots, 0] \in P_n \mathbb{C} \}.$$

Now let  $j: P_s \mathbb{C} \rightarrow P_n \mathbb{C}$  be the inclusion and let  $N$  be the normal bundle of  $P_s \mathbb{C}$  in  $P_n \mathbb{C}$ . Then clearly  $y = j^* x$  is a generator of  $H^2(P_s \mathbb{C})$ .

We study the action of  $g$  on  $N$ . Since

$$g[z_0, z_1, \dots, z_s, z_{s+1}, \dots, z_n] = [\zeta_0 z_0, \zeta_1 z_1, \dots, \zeta_s z_s, \zeta_{s+1} z_{s+1}, \dots, \zeta_n z_n] \\ = [\zeta z_0, \zeta z_1, \dots, \zeta z_s, \zeta_{s+1} z_{s+1}, \dots, \zeta_n z_n]$$

$$= [z_0, z_1, \dots, z_s, \zeta^{-1} \zeta_{s+1} z_{s+1}, \dots, \zeta^{-1} \zeta_n z_n],$$

we have

$$N = \sum_{\theta} N_{\theta},$$

where  $N_{\theta} = 0$  unless  $\theta = \zeta^{-1} \zeta_j$  for some  $j = s+1, \dots, n$  and

$N_{\zeta^{-1} \zeta_j}$  is a complex line bundle over  $P_s \mathbb{C}$ . We obtain

$$(4) \quad L'(g, P_n \mathbb{C})_{\zeta} = L(P_s \mathbb{C}) \prod_{\theta} L_{\theta}(N_{\theta}) = \left( \frac{x}{\tanh y} \right)^{s+1} \prod_{j=s+1}^n \frac{\zeta^{-1} \zeta_j e^{2y+1}}{\zeta^{-1} \zeta_j e^{2y-1}}.$$

Therefore

$$(5) \quad L(g, P_n \mathbb{C})_{\zeta} = j! L'(g, P_n \mathbb{C})_{\zeta} = \left( \frac{x}{\tanh x} \right)^{s+1} \cdot \prod_{j=s+1}^n \frac{\zeta^{-1} \zeta_j e^{2x+1}}{\zeta^{-1} \zeta_j e^{2x-1}} \cdot x^{\bar{n}-s} \\ = \prod_{j=0}^n \left( x \frac{\zeta^{-1} \zeta_j e^{2x+1}}{\zeta^{-1} \zeta_j e^{2x-1}} \right).$$

Observe that the right hand side of (5) is equal to zero unless

$$\zeta \in \{ \zeta_0, \zeta_1, \dots, \zeta_n \}.$$

Now we can show (2) by using the trigonometric identity

$$\sum_{\lambda=1}^a \frac{\lambda z + 1}{\lambda z - 1} = a \frac{z^a + 1}{z^a - 1}.$$

Thus

$$(6) \quad \sum_{g \in G_a} L(g, P_n \mathbb{C}) = \sum_{\zeta_0, \dots, \zeta_n} \sum_{\zeta \in S^1} \prod_{j=0}^n \left( x \frac{\zeta^{-1} \zeta_j e^{2x+1}}{\zeta^{-1} \zeta_j e^{2x-1}} \right) \\ = \sum_{\zeta \in S^1} \prod_{j=0}^n \left( x \cdot \sum_{\zeta_j \in S^1} \frac{\zeta^{-1} \zeta_j e^{2x+1}}{\zeta^{-1} \zeta_j e^{2x-1}} \right) \\ = \sum_{\zeta \in S^1} \prod_{j=0}^n \left( a_j x \frac{\zeta^{-a_j} e^{2a_j x + 1}}{\zeta^{-a_j} e^{2a_j x - 1}} \right) \\ = \sum_{0 \leq \xi < \pi} \prod_{j=0}^n \frac{a_j x}{\tanh(a_j(x+i\xi))}. \quad (\text{Q.E.D.})$$

Now suppose  $a_0, a_1, \dots, a_n$  are mutually relatively prime numbers.

Then by Theorem 2, we have

$$(7) \quad \pi^* L(P_n \mathbb{C}/G_a) = \prod_{j=0}^n \frac{a_j x}{\tanh a_j x} \pmod{x^n}.$$

Therefore, in terms of the total Pontrjagin class  $p$ , we have

$$(8) \quad \tau^* p(P_n \mathbb{C}/G_a) = \prod_{j=0}^n (1 + a_j^2 x^2) \text{ mod } x^n$$

Suppose  $n$  is even, say  $n=2k$ , then there arises a natural question;

Question. Are there values of  $a_0, a_1, \dots, a_{2k}$  such that (8) holds also in the highest term?

Now suppose  $\{a_0, a_1, \dots, a_{2k}\}$  satisfies the requirement of the Question. Then

$$(9) \quad \tau^* p(P_n \mathbb{C}/G_a) = \prod_{j=0}^n (1 + a_j^2 x^2).$$

Since the action of  $G_a$  extends to an action of the torus  $T^{n+1}$ , we have

$$(10) \quad \tau^* : H^*(P_n \mathbb{C}/G_a; \mathbb{Q}) \xrightarrow{\cong} H^*(P_n \mathbb{C}; \mathbb{Q}).$$

Hence

$$(11) \quad \text{sign } P_n \mathbb{C}/G_a = 1.$$

On the other hand,  $P_n \mathbb{C}/G_a$  is a rational homology manifold. Therefore its signature is equal to the L-genus. From (9) and (11), we obtain

$$(12) \quad L_k(p_1, \dots, p_k) = a_0 a_1 \dots a_{2k}$$

where  $p_j$  is the  $j$ -th elementary symmetric polynomial in  $a_j^2$ 's.

Conversely assume that (12) holds. Then it is easy to see that

$\{a_0, a_1, \dots, a_{2k}\}$  satisfies the requirement of the Question. Thus we have obtained

Proposition 3. Let  $a_0, a_1, \dots, a_{2k}$  be mutually relatively prime natural numbers  $\geq 1$ . Then

$$\tau^* p(P_n \mathbb{C}/G_a) = \prod_{j=0}^n (1 + a_j^2 x^2)$$

if and only if  $\{a_j\}$  satisfies the Diophantine equation

$$L_k(p_1, \dots, p_k) = a_0 a_1 \dots a_{2k}.$$

For  $k=1$ , the equation (12) is

$$a_0^2 + a_1^2 + a_2^2 = 3a_0 a_1 a_2$$

and all solutions are known (see [2]). For  $k=2$ , the equation is

$$7(a_0^2 a_1^2 + a_0^2 a_2^2 + \dots + a_3^2 a_4^2) - (a_0^2 + \dots + a_4^2)^2 = 45a_0 a_1 \dots a_4 .$$

Are there infinitely many solutions?

It is easy to check that  $(1, 1, 1, 1, 1)$  and  $(2, 1, 1, 1, 1)$  are solutions. Recently Zagier has found a solution  $(2, 7, 19, 47, 59)$  using a computer. Up to permutation, these are the only solutions in mutually relatively prime natural numbers  $\leq 100$ .

### 3. L-classes of symmetric products.

Let  $X$  be a closed oriented differentiable manifold and let  $X^n$  be the  $n$ -th Cartesian product of  $X$ . Then the symmetric group of degree  $n$ ,  $S_n$ , acts on  $X^n$  by permuting the factors.

Now if  $\dim X$  is even, say  $2s$ , then this action is orientation preserving. Thus we can apply the result of §1.

Let  $X(n) = X^n/S_n$  be the  $n$ -th symmetric product of  $X$ . If we choose a fixed point  $x_0 \in X$ , we have natural inclusions

$$X = X(1) \subset X(2) \subset \dots \subset X(\infty)$$

where  $X(\infty) = \varinjlim_n X(n)$ . We will write  $j$  for any inclusion map  $j: X(n) \rightarrow X(m)$ ,  $\infty \geq m \geq n$ .

Now if we use  $\mathbb{Q}$  for the coefficient of the cohomology, we have

$$(13) \quad H^*(X(n)) \cong H^*(X^n)^{S_n}$$

where the right hand side is the  $S_n$ -invariant subgroup of  $H^*(X^n)$ .

Henceforth we will identify these two groups by the above isomorphism.

It is rather easy to calculate  $H^*(X(n))$ . Let  $\{f_0, f_1, \dots, f_b\}$  be a homogeneous basis for  $H^*(X)$  with  $f_0 = z \in H^{2s}(X)$ , the cohomology fundamental class and  $f_b = 1$ . Let  $n_0, \dots, n_b$  be non-negative integers with  $n_0 + n_1 + \dots + n_b = n$ . We define an element



$\langle n_0 f_0 \dots n_b f_b \rangle \in H^*(X(n))$  as follows.

9

Let  $\sigma \in S_n$ . Then  $\sigma$  acts on  $X^n$  by

$$\sigma(x_1, \dots, x_n) = (x_{\sigma(1)}, \dots, x_{\sigma(n)}).$$

We define an element

$$\langle u_1, u_2, \dots, u_n \rangle \in H^*(X(n)) \quad \text{for } u_j \in H^*(X)$$

$$\text{by } \langle u_1, u_2, \dots, u_n \rangle = \sum_{\sigma \in S_n} \sigma^*(u_1 \times \dots \times u_n) \in H^*(X^n)^{S_n} = H^*(X(n))$$

and we put

$$\langle n_0 f_0 \dots n_b f_b \rangle = \langle \underbrace{f_0, \dots, f_0}_{n_0}, \dots, \underbrace{f_b, \dots, f_b}_{n_b} \rangle.$$

Then it can be shown that

**Proposition 4.** The elements  $\langle n_0 f_0 \dots n_b f_b \rangle$  with  $n_0 + \dots + n_b = n$  and  $n_i \leq 1$  if degree  $f_i$  is odd, form a basis for  $H^*(X(n))$ .

Now we define an element  $[n_0 f_0 \dots n_{b-1} f_{b-1}]_n \in H^*(X(n))$  by

$$[n_0 f_0 \dots n_{b-1} f_{b-1}]_n = \begin{cases} 0 & \text{if } n < (n_0 + \dots + n_{b-1}) \\ (n_b!)^{-1} \langle n_0 f_0 \dots n_b f_b \rangle & \text{if } n_b = n - (n_0 + \dots + n_{b-1}) \geq 0 \end{cases}$$

Then it can be seen that

$$(14) \quad j^* [n_0 f_0 \dots n_{b-1} f_{b-1}]_{n+1} = [n_0 f_0 \dots n_{b-1} f_{b-1}]_n$$

Thus the elements  $[n_0 f_0 \dots n_{b-1} f_{b-1}]_n$  ( $n = 1, 2, \dots$ ) defines an element  $[n_0 f_0 \dots n_{b-1} f_{b-1}] \in H^*(X(\infty))$  so that

$$(15) \quad j^* [n_0 f_0 \dots n_{b-1} f_{b-1}]_n = [n_0 f_0 \dots n_{b-1} f_{b-1}] = [n_0 f_0 \dots n_{b-1} f_{b-1}]_n$$

where  $j: X(n) \rightarrow X(\infty)$ . We write  $\eta$  for the element  $[f_0] \in H^{2s}(X(\infty))$ .

Then  $\eta_n = [f_0]_n = \sum_{i=1}^n \pi_i^* z$ , where  $\pi_i: X^n \rightarrow X$  is the projection on the  $i$ -th factor. Then it can be shown that

$$(16) \quad [n_0 f_0 \dots n_{b-1} f_{b-1}]_n = \eta^{n_0} [n_1 f_1 \dots n_{b-1} f_{b-1}]_n$$

and

**Proposition 5.** The elements

$\eta^{n_0} [n_1 f_1 \dots n_{b-1} f_{b-1}]_n$  form a basis for  $H^*(X(\infty))$  and

$$j^* (\eta^{n_0} [n_1 f_1 \dots n_{b-1} f_{b-1}]_n) = \eta^{n_0} [n_1 f_1 \dots n_{b-1} f_{b-1}]_n,$$

where  $j: X(n) \rightarrow X(\infty)$ .

In terms of these elements of  $H^*(X(\infty))$ , we can write the second main result of Zagier.

Theorem 6. Let  $X$  be a connected closed oriented differentiable manifold of dimension  $2s$ . Let  $j: X(n) \rightarrow X(\infty)$  be the inclusion. Then there is a class  $G \in H^{**}(H(\infty))$  such that

$$L(X(n)) = j^*(Q_s(\eta)^{n+1} G) \quad \text{where } Q_s(t) \text{ is a power series defined by}$$

$$Q_s(t) = \frac{t}{f_s(t)},$$

$$f_s(t) = g_s^{-1}(t), \quad g_s(t) = t + \frac{t^3}{3^s} + \frac{t^5}{5^s} + \frac{t^7}{7^s} + \dots$$

Equivalently, let  $j: X(n) \rightarrow X(n+1)$  be the inclusion, Then

$$j^* L(X(n+1)) = Q_s(\eta_n) \cdot L(X(n)).$$

The proof consists of a rather long and complicated calculation applying Theorem 1. Here we concentrate on the cases when  $X = S^{2s}$  and  $s = 1$  and make some remarks.

Thus assume first that  $X = S^{2s}$ . Then the basis for  $H^*(X)$  is just  $\{z, 1\}$  and the class  $G$  that appeared in Theorem 6 can be simply expressed and the result is

Proposition 7. Let  $X = S^{2s}$ . Then

$$L(X(n)) = \frac{f'_s(\eta)}{1 - f_s(\eta)^2} \left( \frac{\eta}{\tanh \eta} \right)^{n+1}$$

where  $f'$  denotes the derivative of  $f$ .

Now if  $s = 1$ , then  $X(n)$  can <sup>be</sup> naturally identified with  $P_n \mathbb{C}$  and  $\eta_n \in H^2(X(n))$  is the standard generator. In this case Prop. 7 simply says the well-known result

$$L(P_n \mathbb{C}) = \left( \frac{\eta}{\tanh \eta} \right)^{n+1}.$$

Next assume that  $s = 1$ . Thus let  $X$  be a Riemann surface of

genus  $g$ . We choose a basis  $\{\alpha_1, \dots, \alpha_g, \alpha_1', \dots, \alpha_g'\}$  for  $H^1(X)$

$$\begin{aligned} \text{such that } \alpha_i \alpha_j' &= \alpha_i' \alpha_j = 0 \quad (\forall i, j) \\ \alpha_i \alpha_j' &= \alpha_i' \alpha_j \quad (i \neq j) \\ \alpha_i \alpha_i' &= -\alpha_i' \alpha_i = z. \end{aligned}$$

We define elements  $\delta_i$  ( $i = 1, \dots, g$ ) by,

$$\delta_i = [1\alpha_i' \quad 1\alpha_i] \in H^2(X(\infty)).$$

Then we can show

Theorem (Macdonald)

Let  $X$  be a Riemann surface of genus  $g$ . Then

$$L(X(n)) = \left( \frac{\eta}{\tanh \eta} \right)^{n-2g+1} \prod_{i=1}^g \frac{\delta_i}{\tanh \delta_i}.$$

This theorem had been proved by Macdonald by a different method.

Finally we mention that Zagier has also calculated the equivariant

$L$ -classes  $L(g, X(n))$  for the actions on  $X(n)$  which are induced from actions on  $X$ .

#### Bibliography

- [1] M.F. Atiyah and I.M. Singer, The index of elliptic operators III, Ann. of Math. 87 (1968), 546-604.
- [2] F. Hirzebruch, The signature theorem; reminiscences and recreation, Prospects in mathematics, Ann. of Math. Studies 70, Princeton, 1971, 3-31.
- [3] D. Zagier, Some problems in differential topology, Equivariant  $L$ -classes, symmetric products and the  $G$ -signature theorem, Dissertation, Oxford-Bonn, 1971.