

ON CLASS NUMBER of a GALOIS EXTENSION

N. NAKAGOSHI

(TOYAMA Univ.)

Let K be a finite Galois extension of an algebraic number field k . The central extension \hat{K} and the genus field K^* over k are defined in [1], [3]. The central class number $z_{K/k} = (\hat{K} : K)$ and the genus number $g_{K/k} = (K^* : K)$ of K with respect to k are given in [1], [3].

Let $a_{K/k}$ be the ambiguous ideal class number of K respect to k . If K/k is a cyclic extension, or a Galois extension of a prime power degree, we have some relations between the class number h_K of K and $a_{K/k}$ ([2], [3], [5]).

In [4], it is proved:

"Let K/k be a cyclic extension of a prime power degree l^v , and suppose K and the absolute class field \bar{k} of k are disjoint over k , i.e. $K \cap \bar{k} = k$. Then h_K is prime to l if and only if $a_{K/k} = h_K$ and h_k is prime to l ."

We shall generalize this criterion to a Galois extension of a prime power degree. In this note, h_k and \bar{k} for a finite algebraic number field k mean as above.

PROPOSITION. Let k be an algebraic number field and K/k be a Galois extension of degree n . Suppose $K \cap \bar{k} = k$.

- (1) If h_k is prime to n , then $z_{K/k} = g_{K/k} = a_{K/k} = h_K$, and h_K is prime to n .
- (2) Let $n = l^v$ be a prime power. Then h_K is prime to l if and only if $z_{K/k} = g_{K/k} = a_{K/k} = h_K$ and h_k is prime to l .

Proof. (1) Since $\bar{z}_{K/R} = (\hat{K} : K^*) \cdot g_{K/R}$, $g_{K/R} = (K^* : K \cdot \bar{K}) h_{K/R}$, $h_{K/R}$, $\bar{z}_{K/R}$ and $g_{K/R}$ are divisible by $h_{K/R}$. In the decomposition

$$\frac{h_{K/R}}{h_{K/R}} = \frac{g_{K/R}}{h_{K/R}} \frac{h_{K/R}}{g_{K/R}} = \frac{\bar{z}_{K/R}}{h_{K/R}} \frac{h_{K/R}}{\bar{z}_{K/R}}$$

if $h_{K/R}$ is prime to π , then $\frac{g_{K/R}}{h_{K/R}}$, $\frac{\bar{z}_{K/R}}{h_{K/R}}$ are prime to π .

From the genus number formula and the central class number formula, we have

$$\frac{g_{K/R}}{h_{K/R}} = \frac{T_{\mathfrak{p}} T_{\mathfrak{q}} e_{\mathfrak{q}}}{(K_0 : k) (E_R : E_R \cap N_{K/R} \cup K)}$$

$$\frac{\bar{z}_{K/R}}{h_{K/R}} = \frac{T_{\mathfrak{p}} T_{\mathfrak{q}} \cdot (K^x \cap N_{K/R} J_K : N_{K/R} K^x)}{(K_0 : k) (E_R : E_R \cap N_{K/R} K^x)}$$

(notations in these formulas are defined in [1], [3]).

$$E_R \cap N_{K/R} \cup K \supset E_R^n = \{ \varepsilon^n = N_{K/R} \varepsilon \mid \varepsilon \in E_R \},$$

$$E_R \cap N_{K/R} K^x \supset E_R^n$$

and it is well-known

$$(K^x \cap N_{K/R} J_K) / N_{K/R} K^x \cong H^{-3}(G, \mathbb{Z}) / F$$

where G is a Galois group of K over k and F is the subgroup of TATE cohomology group $H^{-3}(G, \mathbb{Z})$, generated by $I_{\mathfrak{q}} G_{\mathfrak{q}} / G H^{-3}(G_{\mathfrak{q}}, \mathbb{Z})$ for all the infinite and the finite prime divisors \mathfrak{q} and its decomposition group $G_{\mathfrak{q}}$.

Therefore, the prime factors of $\frac{g_{K/R}}{h_{K/R}}$ and $\frac{\bar{z}_{K/R}}{h_{K/R}}$ are those of π , also

$$\frac{g_{K/R}}{h_{K/R}} = \frac{\bar{z}_{K/R}}{h_{K/R}} = 1; \bar{z}_{K/R} = g_{K/R} = h_{K/R}.$$

Let I_K be the ideal group of K and P_K be the principal ideal group of K . Then

$$A = \{ \alpha \in I_K \mid \alpha^{1-\sigma} \in P_K \text{ for any } \sigma \in G \},$$

$$H = \{ \alpha \in I_K \mid N_{K/k} \alpha \in P_k \}$$

are the subgroups of I_K , and

$$A \cap H \subset \{ \alpha \in I_K \mid \alpha^n \in P_K \}.$$

It follows for the ambiguous ideal class number $a_{K/k}$ of K respect to k

$$a_{K/k} = (A : P_K) = (A : A \cap H) (A \cap H : P_K),$$

where the prime factors of $(A \cap H : P_K)$ are also those of n . If

$f_{K/k}$ is prime to n , then $a_{K/k}$ is prime to n , hence

$$(A \cap H : P_K) = 1; \quad A \cap H = P_K.$$

By $f_{K/k} = f_{K^*/k}$, $K^* = K \cdot \bar{k}$ is the genus field of K over k , i.e. the class field corresponding to H over K . So we have

$$\begin{aligned} f_{K/k} &= (K^* : K) = (I_{K^*} : H) = (I_K : A \cap H) (A \cap H : H) \\ &= (I_K : A \cap H) (A : A \cap H) = (I_K : A \cap H) \cdot a_{K/k}. \end{aligned}$$

On the other hand, the number $a_{K/k}^{(0)}$ of ideal classes represented by an ambiguous ideal in K/k is given by [5]:

$$a_{K/k}^{(0)} = \frac{f_{K/k} \cdot |G| \cdot e_g}{|H^1(G, E_K)|}.$$

If $f_{K/k}$ is prime to n , then $a_{K/k}, a_{K/k}^{(0)}$ and $f_{K/k}$ are prime to n , that is, $a_{K/k}^{(0)} = f_{K/k}$. Since $f_{K/k} = f_{K/k} \equiv 0 \pmod{a_{K/k}}$, and $a_{K/k} \equiv 0 \pmod{a_{K/k}^{(0)}}$, hence $f_{K/k} = f_{K/k} = a_{K/k}$.

(2) In case of $n = l^v$, we know in [3]

$$f_{K/k} \equiv f_{K/k} \pmod{l}.$$

References

- [1] Furuta, Y., Nagoya Math. J., vol.29(1967), 281-285.
- [2] ——— Nagoya Math. J., vol.37(1970), 197-200
- [3] ——— J. Number Theory, vol. 3(1971), 318-322.
- [4] Furuya, H., Tohoku Math. J., vol.23(1971), 207-218.
- [5] Yokoi, H., Nagoya Math. J., vol,29(1967), 31-44.
- [6] ——— J. Math. Soc., Japan, vol.20(1968), 411-418.

12, June, 1973, Kanazawa.