

Stefan problems with the unilateral boundary condition
on the fixed boundary

Shoji Yotsutani

§0. Introduction.

In this note we consider the following one-dimensional two-phase Stefan problem with the unilateral boundary condition on the fixed boundary : Given the data, ϕ and ℓ , find two functions $s = s(t)$ and $u = u(x,t)$ such that the pair (s, u) satisfies

$$\begin{array}{l}
 (0.1) \quad s(0) = \ell, \quad 0 < s(t) < 1 \quad (0 \leq t \leq T), \\
 (0.2) \quad u_{xx} - c_0 u_t = 0 \quad (0 < x < s(t), 0 < t \leq T), \\
 (0.3) \quad u_{xx} - c_1 u_t = 0 \quad (s(t) < x < 1, 0 < t \leq T), \\
 (0.4) \quad \begin{array}{l}
 (a) \quad u_x(0,t) \in \gamma_0(u(0,t)) \quad (0 < t \leq T), \\
 (b) \quad -u_x(1,t) \in \gamma_1(u(1,t)) \quad (0 < t \leq T),
 \end{array} \\
 (0.5) \quad \begin{array}{l}
 (a) \quad u(x,0) = \phi(x) \quad (0 \leq x \leq \ell), \\
 (b) \quad u(x,0) = \phi(x) \quad (\ell \leq x \leq 1),
 \end{array} \\
 (0.6) \quad u(s(t),t) = 0 \quad (0 \leq t \leq T), \\
 \text{and the free boundary condition} \\
 (0.7) \quad b \dot{s}(t) = u_x(s(t)+0,t) - u_x(s(t)-0,t) \quad (0 < t \leq T)
 \end{array}$$

The quantities c_0, c_1 and b are positive physical parameters of the problem. T is a positive constant to be discussed later, and the function ϕ and the value $\ell, 0 < \ell < 1$, are the initial data for (S).

Each γ_i , $i = 0, 1$, is a maximal monotone graph in \mathbb{R}^2 with $\gamma_i(H_i) \ni 0$, where H_i is a constant satisfying $(-1)^i H_i \geq 0$. We put the assumptions of signs for H_0 and H_1 from the physical reasoning. (0.4) are the unilateral boundary conditions and (0.7) is the so-called Stefan's condition.

The system (0.1)-(0.7) is a simple model of a heat-conduction system consisting of two phases (e. g. liquid and solid) of the same substance which are in perfect thermal contact at an interface. $u(x,t)$ represents the temperature distribution in the system, and the curve $s(t)$ represents the position of the interface which varies with time t as solid melts or liquid freezes. The unilateral boundary conditions (0.4) model several physical situations, including the temperature control through the boundary [7, Ch. 1] and the heat flow subject to the nonlinear cooling by the radiation on the boundary [9, Ch. 7]. The boundary conditions at the interface ((0.6), (0.7)) reflect respectively the facts that the temperature at the interface must be equal to the melting temperature (taken to be zero) and that the rate of melting is proportional to the rate of absorption of the heat energy at the interface. In formulating (0.7), we have assumed, without loss of generality, that the thermal conductivity in both phases is 1.

The problems of this type with linear boundary conditions on the fixed boundary have been considered by many authors (Rubinstein [15], Kamenomostkaja [10], Friedman [8, 9], Brézis [1], Cannon-Primicerio [4, 5], Cannon-Henry-Kotlow [6], Nogi [13], etc.). On the other hand Bénilan [22] has treated this type's Stefan problem of n -dimensional case by using the theory of nonlinear contraction semigroups in Banach space L^1 . He got an integral solution. However we do not know the differentiability of the Benilan's integral solution. One-phase problem of this type was recently studied by Yotsutani [20].

Our purpose is to study the global existence and uniqueness of the classical solution.

In §1 we state the main results. In §2 we explain the difference scheme.

§1. Statements of main results.

We suppose that $0 < \ell < 1$, and $\phi(x)$ satisfies the following condition (A) for simplicity.

(A) $\phi(x)$ is a Lipschitz continuous function on $[0,1]$ satisfying $\phi(0) \in D(\gamma_0)$, $\phi(1) \in D(\gamma_1)$ and $(\ell - x) \phi(x) \geq 0$ ($0 \leq x \leq 1$).

DEFINITION. The pair (s, u) is a solution of (S) if $s \in C([0, T]) \cap C^\infty(]0, T])$, $u \in C(\bar{D}_T)$, $u|_{D_{i,T}} \in C^\infty(D_{i,T})$ ($i = 0, 1$), $\int_0^T \int_0^1 |u_{xx}|^2 dx dt < +\infty$, and (1)-(7) hold, where $D_T = D_{0,T} \cup D_{1,T}$, $D_{i,T} = \{(x, t) ; (-1)^i i < (-1)^i x \leq (-1)^i s(t)\}$ ($i = 0, 1$).

THEOREM 1. Let $\phi(x)$ satisfy (A). Then there exist a critical time T^* ($0 < T^* \leq +\infty$) and a pair of functions (s, u) defined on $[0, T^*[$ such that (s, u) is a solution of (S) on $[0, T]$ for any T ($0 < T < T^*$).

Further if $T^* < +\infty$, then the following relations hold,

$$\lim_{t \rightarrow T^*} s(t) = 0 \text{ or } 1, \quad 0 < s(t) < 1 \quad (0 < t < T^*).$$

REMARK. In particular, if γ_i ($i = 0, 1$) is a single valued maximal monotone, then $u_x|_{\tilde{D}_{i,T}} \in C(\tilde{D}_{i,T})$ ($\tilde{D}_{i,T} = D_{i,T} \cup \{x=i\}$) for any T ,

$$(-1)^i u_x(i, t) = \gamma_i(u(i, t)) \quad (0 < t < T^*).$$

THEOREM 2. Under the same assumption of Theorem 1, the critical time T^* and the pair (s, u) on $[0, T^*[$ are uniquely determined.

REMARK. We can loosen the assumption (A) a little.

We construct a solution by using a primitive implicit difference scheme with only a device of capturing a free boundary explicitly through step-by-step process in time (see §2). Uniqueness is based upon the maximum principle, its strong form, a parabolic version of Hopf's lemma and the comparison theorems for the unilateral problem associated with the heat equation. The proofs of our theorems is given in [21].

§2. Difference scheme

Let ℓ be a rational number with $0 < \ell < 1$.

We use a net of rectangular meshes with uniform space width h and variable time step $\{k_n\}$ ($n = 1, 2, 3, \dots$). Time steps $\{k_n\}$ are assumed to be unknown and they are determined by the rule that h/k_n gives gradient of a desired free boundary at each time $t = t_n$, so that the free boundary crosses each mesh lines just at each corresponding mesh points. Let us introduce discrete coordinates

$$x_j = jh \quad (j = 0, 1, 2, \dots, M),$$

$$t_n = \sum_{p=1}^n k_p \quad (n = 1, 2, 3, \dots), \quad t_0 = 0,$$

where h varies in such a way that $\ell/h = J_0$ and $1/h = M$ are integers, and net functions s_n and u_j^n which correspond to $s(t_n)$ and $u(x_j, t_n)$ respectively. By our rule we can put

$$(2.1) \quad s_n = J_n \cdot h \quad (J_n : \text{interger, } n = 0, 1, 2, \dots).$$

Further we introduce usual divided differences :

$$(2.2) \quad u_{jx}^n = \frac{1}{h}(u_{j+1}^n - u_j^n), \quad u_{j\bar{x}}^n = \frac{1}{h}(u_j^n - u_{j-1}^n),$$

$$(2.3) \quad u_{jxx}^n = \frac{1}{h^2}(u_{j+1}^n - 2u_j^n + u_{j-1}^n), \quad u_{j\bar{\tau}}^n = \frac{1}{k_n}(u_j^n - u_j^{n-1}), \quad \text{etc.}$$

In our scheme the heat equation are replaced by pure implicit difference equations,

$$(2.4) \quad u_{jxx}^n - c_0 u_{j\bar{\tau}}^n = 0 \quad (1 \leq j \leq J_n - 1),$$

$$(2.5) \quad u_{jxx}^n - c_1 u_{j\bar{\tau}}^n = 0 \quad (J_n + 1 \leq j \leq M - 1).$$

The boundary and initial conditions are put in the following forms,

$$(2.6) \quad (a) \quad u_{0x}^n \in \gamma_0(u_0^n),$$

$$(b) \quad -u_{M\bar{x}}^n \in \gamma_1(u_M^n),$$

$$(2.7) \quad (a) \quad u_j^0 = \phi_j \equiv \phi(x_j) \quad (0 \leq j \leq J_0 - 1),$$

$$(b) \quad u_j^0 = \phi_j \equiv \phi(x_j) \quad (J_0 + 1 \leq j \leq M),$$

$$(2.8) \quad u_{J_n}^n = 0.$$

The Stefan's condition is replaced by an explicit formula

$$(2.9) \quad \pm b \frac{h}{k_n} = u_{J_{n-1}x}^{n-1} - u_{J_{n-1}\bar{x}}^{n-1}$$

where + or - sign correspond to the cases of positive heat flow

to the interface or negative one at $t = t_n$ respectively.

Our algorithm is the following. β : a positive constant.

$$1^{\circ} \quad u_j^0 = \phi_j \quad (0 \leq j \leq J_0 - 1), \quad u_{J_0}^0 = 0, \quad u_j^0 = \phi_j \quad (J_0 + 1 \leq j \leq M),$$

$$s_0 = J_0 \cdot h = \ell.$$

For $n = 1, 2, 3, \dots$ successively,

$$2.1^{\circ} \quad \text{if } u_{J_{n-1}x}^{n-1} - u_{J_{n-1}\bar{x}}^{n-1} > \beta h^{\frac{1}{2}}, \quad \text{then we take } J_n = J_{n-1} + 1$$

and get k_n from

$$b \frac{h}{k_n} = u_{J_{n-1}x}^{n-1} - u_{J_{n-1}\bar{x}}^{n-1},$$

$$2.2^{\circ} \quad \text{if } u_{J_{n-1}x}^{n-1} - u_{J_{n-1}\bar{x}}^{n-1} < -\beta h^{\frac{1}{2}}, \quad \text{then we take } J_n = J_{n-1} - 1$$

and get k_n from

$$-b \frac{h}{k_n} = u_{J_{n-1}x}^{n-1} - u_{J_{n-1}\bar{x}}^{n-1},$$

$$2.3^{\circ} \quad \text{if } |u_{J_{n-1}x}^{n-1} - u_{J_{n-1}\bar{x}}^{n-1}| \leq \beta h^{\frac{1}{2}}, \quad \text{then we take}$$

$$J_n = J_{n-1} \quad \text{and} \quad k_n = \beta h^{\frac{1}{2}},$$

3° solve the system of difference equations (2.4) and (2.5)

for $\{u_j^n\}_j$ under the boundary conditions (2.6) and (2.8)

with the initial condition $\{u_j^{n-1}\}_j$,

4° if $J_n < J_{n-1} > J_{n-2}$ or $J_n > J_{n-1} < J_{n-2}$, then J_n and

k_n are revised as $J_n = J_{n-1}$ and $k_n = bh^{\frac{1}{2}}/\beta$, and then return again to the step 3^o (as $n = 1$, we consider $J_{-1} = J_0$).

5^o if $J_n = 1$ or $M - 1$, then stop computing.

REMARK 2.1. Step 3^o is well-defined by Lemma 4.1 of Yotsutani [20].

Acknowledgement. The author would like to express his hearty thanks to Prof. H. Brézis for his valuable advices.

References

- [1] H. Brézis, On some degenerate nonlinear parabolic equations, in "Nonlinear Functional Analysis, Proc. Symp. Pure Math. XVIII," (F. Browder, e.d.), Amer. Math. Soc., Providence, R. I., 1970, 28 -38.
- [2] H. Brézis, Problemes unilatéraux, J. Math. Pures Appl., 51 (1972), 1 - 164.
- [3] H. Brézis, Operateurs maximaux monotones et semigroups de contractions dans les espaces de Hilbert, Math. Studies, 5, North Holland, 1973.
- [4] J. R. Cannon and M. Primicerio, A two phase Stefan problem with temperature boundary conditions, Ann. Mat. Pura Appl. (IV), 88 (1971), 177 - 192.
- [5] J. R. Cannon and M. Primicerio, A two phase Stefan problem with flux boundary conditions, Ann. Mat. Pura Appl. (IV), 88 (1971), 193 - 205.
- [6] J. R. Cannon, D. B. Henry, D. B. Kotlow, Classical solutions of the one-dimensional, two-phase Stefan problem, Ann. Mat. pura Appl. (IV), 107 (1975), 311 - 341.
- [7] G. Duvaut and J. L. Lions, Inequalities in mechanics and physics, Springer, 1976.

- [8] A. Friedman, One dimensional Stefan problems with nonmonotone free boundary, *Trans. Amer. Math. Soc.*, 133 (1968), 89 -114.
- [9] A. Friedman, *Partial differential equations of parabolic type*, Prentice - Hall, 1964.
- [10] S. L. Kamenomostkaja, On Stefan's problem, *Math. Sbornik*, 53 (95), (1965), 485 - 514.
- [11] N. Kenmochi, Free boundary problems for a class of nonlinear parabolic equations : an approach by the theory of subdifferential operators, to appear.
- [12] L. Nirenberg, A strong maximum principle for parabolic equations, *Comm. Pure Appl. Math.*, 6 (1953), 167 - 177.
- [13] T. Nogi, A difference scheme for solving the 2-phase Stefan problem of heat equation, in *Functional Analysis and Numerical Analysis, Japan-France Seminar, Tokyo and Kyoto, 1976*: H. Fujita (ed.), Japan Society for Promotion of Science, (1978), 361 - 382.
- [14] I. G. Petrowsky, *Partial differential equations*, originally published in Moscow in 1961 by Fizmatgiz under the title *Lektsii ob uravneniyah s chastnumi proizvodnumi*.
- [15] L. I. Rubinstein, *The Stefan problem*, translation of mathematical monographs 27, American Mathematical Society, 1972.
- [16] D. G. Schaeffer, A new proof of the infinite differentiability of the free boundary in the Stefan problem, *J. Diff. Eqs.*, 20 (1976), 266 - 267.

- [17] D. G. Wilson, A. D. Solomon and P. T. Boggs (Ed.), Moving boundary problems, Academic Press, 1978.
- [18] M. Yamaguchi and T. Nogi, The Stefan problem, Sangyo-Tosho, 1977, (in Japanese).
- [19] S. Yotsutani, Evolution equations associated with the subdifferentials, J. Math. Soc. Japan, 31 (1979), 624 - 646.
- [20] S. Yotsutani, Stefan problems with the unilateral boundary condition on the fixed boundary I, pre-print.
- [21] S. Yotsutani, Stefan problems with the unilateral boundary condition on the fixed boundary II, pre-print
- [22] P. Bénéilan, Opérateurs accréatifs et semi-groupes dans les espaces L^p ($1 \leq p \leq \infty$), in Functional Analysis and Numerical Analysis, Japan-France Seminar, Tokyo and Kyoto, 1976: H. Fujita (ed.), Japan Society for Promotion of Science, (1978), 15 - 53.
- [23] A. Damlamian, Some results on the multi-phase Stefan problem, Comm. in Partial Differential Equations, 2(10), (1977), 1017 - 1044.

Shoji YOTSUTANI

Department of Applied Science

Faculty of Engineering

Miyazaki University

Miyazaki 880

Japan