

Shape of compactifications

Yukihiro Kodama

(Department of Mathematics, University of Tsukuba)

All of spaces are assumed to be metrizable and hence compactifications are metrizable ones.

General questions are as follows.

- 1) Has a good space always a good compactification ?
- 2) Does there exist a good space every compactification of which is bad ?
- 3) Does there exist a bad space having a good compactification ?

Here by "good" we mean "with good shape property".

Example 1. Let X be a countable discrete space. Then for every compactum Y there is a compactification αX such that the remainder $\alpha X - X \approx Y$ and Y is a retract of αX .

Example 2. There are compactifications $\alpha_i R$, $i = 1, 2, 3$, of the real line R :

- 1) $\alpha_1 R$ is an FAR and $\alpha_1 R - R$ consists of two segments.
- 2) $\alpha_2 R$ is an FANR and $\alpha_2 R - R$ consists of two circles.
- 3) $\alpha_3 R$ is not pointed 1-movable. To construct $\alpha_3 R$, let M be a solenoid and let f be a 1-1 map from R onto one component of M . Put $X = \{(f(x), 1/|x|+1) : x \in R\} \subset M \times I$ and $\alpha_3 R = Cl_{M \times I} X$. Then $X \approx R$ and $\alpha_3 R$ is a compactification of R . Since M is a retract of $\alpha_3 R$, $\alpha_3 R$ is not pointed 1-movable.

Let us use the following notations.

$R_+ = [0, \infty)$; $E^* = \sqrt{\beta R_+}$, where βR_+ is the Stone-Čech compactification of R_+ ; $I^* = \beta(I \times R_+) - I \times R_+$. E^* is said to be a Čech 0-cell. I^* contains two Čech 0-cells $E_0^* = (\text{Cl}_{\beta(I \times R_+)} \{0\} \times R_+) \cap I^*$ and $E_1^* = (\text{Cl}_{\beta(I \times R_+)} \{1\} \times R_+) \cap I^*$.

A space X is said to be Čech path connected if for $x_0, x_1 \in X$ there is a map $f : I^* \rightarrow X$ (called a Čech path) such that $f(E_i^*) = x_i$, $i = 0, 1$. X is said to be locally Čech path connected if for $x \in X$ and a neighborhood U of x there is a neighborhood $V \subset U$ of x such that any two points of V are connected by a Čech path in U . The following characterization of pointed 1-movability is given in [1].

Theorem 1. The followings are equivalent for a continuum X .

- (1) X is pointed 1-movable.
- (2) X is Čech path connected.
- (3) Every map $f : E_0^* \cup E_1^* \rightarrow X$ is extendable over I^* .

As a consequence we have the following Krasinkiewicz's theorem.

Corollary (Krasinkiewicz [3]) A continuous image of a pointed 1-movable continuum is pointed 1-movable.

As an application of Theorem 1 we have the following theorem for compactifications of pointed 1-movable spaces.

Theorem 2 [2] Let X be a connected, locally Čech path connected, locally compact space. If αX is a compactification of X such that each component of the remainder $\alpha X - X$ is pointed 1-movable, then αX is pointed 1-movable.

Corollary 1. Let X be a space in Theorem 2. Then the Freudenthal compactification and the one point compactification of X are pointed 1-movable.

Corollary 2. Let X be a locally connected locally compact space. If αX is a compactification of X such that every component of the remainder $\alpha X - X$ is pointed 1-movable, then αX is pointed 1-movable.

In Theorem 2, Corollaries 1 and 2, as shown by Example 2, we can not omit pointed 1-movability of the components of the remainder. Also, we can not replace pointed 1-movability by pointed r -movability ($r \geq 1$).

Example 3. Let $r > 0$. Let $\{S_i, f_{i,i+1} : i=1,2,\dots\}$ be the inverse sequence of r -sphere S_i with bonding map $f_{i,i+1}$ of fixed degree > 1 . Let X' be the telescope associated with $\{S_i\}$. Put $X = C(S_1) \cup X'$, where $C(S_1)$ is the cone over S_1 . Then the one point compactification (= the Freudenthal compactification) of X is not r -movable.

Problem 1. Does there exist a locally compact ANR X such that every compactification of X is not movable ?

Problem 2. Has every (pointed r -) movable locally compact space a (pointed r -) movable compactification ?

Problem 3. Has every locally compact metrizable ANSE a movable compactification ?

Problem 4. Has every locally compact metrizable ASE a compactification which is ASE ?

References

- [1] Y.Kodama, On fine shape theory III.
- [2] ———, Compactification of pointed 1-movable spaces.
- [3] J.Krasinkiewicz, Continuous images of continua and 1-movability, *Fund.Math.*, 98(1978), 141-164.