

A supersymmetric extension of
infinite dimensional Lie algebras

by

Kimio UENO *) (上野 喜三雄)

and

Hirofumi YAMADA **) (山田 裕史)

*) Department of Mathematics, Yokohama City University,
Yokohama 236, Japan

**) Department of Mathematics, Hiroshima University,
Hiróshima 730, Japan

0. Introduction

The transformation theory of the KP (Kadomtsev-Petviashvili) hierarchy was completed in the beautiful work of Date, Jimbo, Kashiwara and Miwa (see e.g., [2]). They have constructed the Fock representation of the Lie algebra $\mathfrak{gl}(\infty)$ and, using the Bose-Fermi correspondence, realized the irreducible components on the polynomial space of infinitely many variables. The Lie algebra $\mathfrak{gl}(\infty)$ has many subalgebras, corresponding ^{to} various type of solutions of the KP hierarchy, for example, the Kac-Moody (affine) algebra of type $A_\ell^{(1)}$ and the Virasoro algebra.

Our main object is the "supersymmetric extension" of above theory. As the first step we introduce in this note the Lie superalgebra $\mathfrak{gl}(\infty|\infty)$ by making use of the free field operators. We show that the "super Kac-Moody algebras" and the "super Virasoro algebra" are contained as the subsuperalgebra of $\mathfrak{gl}(\infty|\infty)$.

Recently the authors got a preprint of Manin and Radul [5]. They have introduced a supersymmetric extension of the KP hierarchy. We shall discuss the relationship between their theory and $\mathfrak{gl}(\infty|\infty)$ in the subsequent paper.

1. Lie superalgebras

We first define the notion of Lie superalgebras.

A \mathbb{Z}_2 -graded complex vector space $\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1$ is called a Lie superalgebra if there is a bilinear bracket product $[,]$ on \mathcal{G} satisfying the following conditions. If $x \in \mathcal{G}_\alpha$ and $y \in \mathcal{G}_\beta$ ($\alpha, \beta = 0, 1$), then 1) $[x, y] \in \mathcal{G}_{\alpha+\beta \pmod{2}}$, 2) $[x, y] = -(-)^{\alpha\beta} [y, x]$ and 3) $[x, [y, z]] = [[x, y], z] + (-)^{\alpha\beta} [y, [x, z]]$. The last relation is referred to as "super Jacobi identity". The space \mathcal{G}_0 (resp. \mathcal{G}_1) is called the even (resp. odd) part. Remark that a Lie superalgebra is not a Lie algebra.

The simplest example of the Lie superalgebras is constructed by the following manner. Let $N = m + n$ be a positive integer, and let

$$\mathcal{G}_0 = \left\{ \begin{bmatrix} A & 0 \\ 0 & D \end{bmatrix} ; A \text{ is } m \times m, D \text{ is } n \times n \right\},$$

$$\mathcal{G}_1 = \left\{ \begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix} ; B \text{ is } m \times n, C \text{ is } n \times m \right\} \text{ so that } \mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1$$

is the space of all $N \times N$ complex matrices. For $X \in \mathcal{G}_\alpha$, $Y \in \mathcal{G}_\beta$ we define $[X, Y] = XY - (-)^{\alpha\beta} YX$. Then the space \mathcal{G} is a Lie superalgebra which is denoted by $\mathfrak{gl}(m|n)$. For $X = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathcal{G}$ we define the "supertrace" by $\text{str } X = \text{tr } A - \text{tr } D$. Supertraceless elements of $\mathfrak{gl}(m|n)$ make a Lie subsuperalgebra $\mathfrak{sl}(m|n)$.

See [1] and [3] for other concepts and examples of Lie superalgebras.

2. The Lie superalgebra $\mathfrak{gl}(\infty|\infty)$

We consider the vector space $V = \mathbb{C}[t, t^{-1}, \xi] / (\xi^2)$ with basis $e_j^{(0)} = t^{-j}$, $e_j^{(1)} = t^{-j} \xi$ ($j \in \mathbb{Z}$). Denote V_0 (resp. V_1) the space spanned by $e_j^{(0)}$'s (resp. $e_j^{(1)}$'s). Let $E_{ij}^{(\alpha\beta)}$ ($\alpha, \beta = 0, 1; i, j \in \mathbb{Z}$) be the endomorphism on V such that $E_{ij}^{(\alpha\beta)} e_k^{(\gamma)} = \delta^{\beta\gamma} \delta_{jk} e_j^{(\alpha)}$. If we define the bracket product for $E_{ij}^{(\alpha\beta)}$'s by

$$[E_{ij}^{(\alpha\beta)}, E_{i'j'}^{(\alpha'\beta')}] = \delta^{\beta\alpha'} \delta_{ji'} E_{ij'}^{(\alpha\beta')} - (-)^{(1-\delta^{\alpha\beta})(1-\delta^{\alpha'\beta'})} \delta^{\beta'\alpha} \delta_{j'i} E_{i'j}^{(\alpha'\beta)},$$

then the space $\mathfrak{gl}(2\infty) = \left\{ \sum a_{ij}^{(\alpha\beta)} E_{ij}^{(\alpha\beta)} ; a_{ij}^{(\alpha\beta)} = 0 \text{ if } |i-j| \gg 1 \right\}$ has the structure of Lie superalgebra. The even (resp. odd) part is the space of linear combinations of $E_{ij}^{(\alpha\beta)}$'s with $\alpha = \beta$ (resp. $\alpha \neq \beta$).

We can construct the Lie superalgebra $\mathfrak{gl}(2\infty)$ by making use of the "free field operators". Let A be the Clifford algebra over \mathbb{C} with generators $\psi_j^{(\alpha)}, \psi_j^{(\alpha)*}$ ($\alpha = 0, 1; i, j \in \mathbb{Z}$), satisfying the defining relations:

$$[\psi_i^{(0)}, \psi_j^{(0)}]_+ = [\psi_i^{(0)*}, \psi_j^{(0)*}]_+ = 0, \quad [\psi_i^{(0)}, \psi_j^{(0)*}]_+ = \delta_{ij},$$

$$[\psi_i^{(1)}, \psi_j^{(1)}] = [\psi_i^{(1)*}, \psi_j^{(1)*}] = 0, \quad [\psi_i^{(1)}, \psi_j^{(1)*}] = -\delta_{ij},$$

$$[\psi_i^{(0)}, \psi_j^{(1)}] = [\psi_i^{(0)}, \psi_j^{(1)*}] = [\psi_i^{(0)*}, \psi_j^{(1)}] = [\psi_i^{(0)*}, \psi_j^{(1)*}] = 0.$$

An element of $W^{(0)} = (\bigoplus_{j \in \mathbb{Z}} \mathbb{C}\psi_j^{(0)}) \oplus (\bigoplus_{j \in \mathbb{Z}} \mathbb{C}\psi_j^{(0)*})$ (resp.

$W^{(1)} = (\bigoplus_{j \in \mathbb{Z}} \mathbb{C}\psi_j^{(1)}) \oplus (\bigoplus_{j \in \mathbb{Z}} \mathbb{C}\psi_j^{(1)*})$) is referred to as a free

fermion (resp. free boson). The following proposition is easy to see.

Proposition 1. The application $E_{ij}^{(\alpha\beta)} \longmapsto \psi_i^{(\alpha)} \psi_j^{(\beta)*}$ defines a representation of the Lie superalgebra $\mathfrak{gl}(2\infty)$.

Next we define the central extension of $\mathfrak{gl}(2\infty)$. First we define the "vacuum expectation value" for quadratic elements in A . Set the linear form by

$$\langle \psi_i^{(\alpha)} \psi_j^{(\beta)} \rangle = \langle \psi_i^{(\alpha)*} \psi_j^{(\beta)*} \rangle = 0,$$

$$\langle \psi_i^{(0)} \psi_j^{(0)*} \rangle = \begin{cases} 1 & i = j < 0 \\ 0 & \text{otherwise,} \end{cases} \quad \langle \psi_j^{(0)*} \psi_i^{(0)} \rangle = \begin{cases} 1 & i = j \geq 0 \\ 0 & \text{otherwise,} \end{cases}$$

$$\langle \psi_i^{(1)} \psi_j^{(1)*} \rangle = \begin{cases} -1 & i = j < 0 \\ 0 & \text{otherwise,} \end{cases} \quad \langle \psi_j^{(1)*} \psi_i^{(1)} \rangle = \begin{cases} 1 & i = j \geq 0 \\ 0 & \text{otherwise,} \end{cases}$$

and the normalization condition $\langle 1 \rangle = 1$. We put

$$:\psi_i^{(\alpha)}\psi_j^{(\beta)*}: = \psi_i^{(\alpha)}\psi_j^{(\beta)*} - \langle \psi_i^{(\alpha)}\psi_j^{(\beta)*} \rangle, \text{ the normal product.}$$

Proposition 2. If we put $Z_{ij}^{(\alpha\beta)} = :\psi_i^{(\alpha)}\psi_j^{(\beta)*}:$, then the following commutation and anti-commutation relations hold:

- 1) $[Z_{ij}^{(00)}, Z_{i'j'}^{(00)}] = \delta_{ji'}Z_{ij'}^{(00)} - \delta_{j'i}Z_{i'j}^{(00)} + \delta_{ji'}\delta_{j'i}(Y_+(j) - Y_+(i)),$
- 2) $[Z_{ij}^{(11)}, Z_{i'j'}^{(11)}] = \delta_{ji'}Z_{ij'}^{(11)} - \delta_{j'i}Z_{i'j}^{(11)} - \delta_{ji'}\delta_{j'i}(Y_+(j) - Y_+(i)),$
- 3) $[Z_{ij}^{(00)}, Z_{i'j'}^{(11)}] = 0,$
- 4) $[Z_{ij}^{(00)}, Z_{i'j'}^{(01)}] = \delta_{ji'}Z_{ij'}^{(01)},$
- 5) $[Z_{ij}^{(00)}, Z_{i'j'}^{(10)}] = -\delta_{j'i}Z_{i'j}^{(10)},$
- 6) $[Z_{ij}^{(11)}, Z_{i'j'}^{(01)}] = -\delta_{j'i}Z_{i'j}^{(01)},$
- 7) $[Z_{ij}^{(11)}, Z_{i'j'}^{(10)}] = \delta_{ji'}Z_{ij'}^{(10)},$
- 8) $[Z_{ij}^{(01)}, Z_{i'j'}^{(01)}]_+ = 0,$
- 9) $[Z_{ij}^{(10)}, Z_{i'j'}^{(10)}]_+ = 0,$
- 10) $[Z_{ij}^{(01)}, Z_{i'j'}^{(10)}]_+ = \delta_{ji'}Z_{ij'}^{(00)} + \delta_{j'i}Z_{i'j}^{(11)} + \delta_{ji'}\delta_{j'i}(Y_+(j) - Y_+(i)),$

$$\text{where } Y_+(j) = \begin{cases} 1 & j \geq 0 \\ 0 & j < 0. \end{cases}$$

By Proposition 2 the space

$$\left\{ \sum a_{ij}^{(\alpha\beta)} z_{ij}^{(\alpha\beta)} ; a_{ij}^{(\alpha\beta)} = 0 \text{ for } |i-j| \gg 1 \right\} \oplus \mathbb{C} \cdot 1$$

has the Lie superalgebra structure which is the one dimensional central extension of $\mathfrak{gl}(2\infty)$. We denote this Lie superalgebra by $\mathfrak{gl}(\infty|\infty)$.

3. Subalgebras

In this section we give some Lie subsuperalgebras of $\mathfrak{gl}(\infty|\infty)$.

The Lie algebra $\mathfrak{gl}(\infty)$ in the sense of [2] is, of course, a subalgebra of the even part of $\mathfrak{gl}(\infty|\infty)$. In fact, the space

$$\left\{ \sum a_{ij}^{(00)} z_{ij}^{(00)} ; a_{ij}^{(00)} = 0 \text{ for } |i-j| \gg 1 \right\} \oplus \mathbb{C} \cdot 1 \quad \text{is}$$

isomorphic to $\mathfrak{gl}(\infty)$. Define the elements $L_m^{(0)} = - \sum_{j \in \mathbb{Z}} j z_{j+m}^{(00)}$

for $m \in \mathbb{Z}$. Then we have the commutation relation

$$[L_m^{(0)}, L_n^{(0)}] = (m - n)L_{m+n}^{(0)} + \frac{1}{6}(m^3 - m) \delta_{m+n} 0.$$

Hence $\bigoplus_{m \in \mathbb{Z}} \mathbb{C} L_m^{(0)} \oplus \mathbb{C} \cdot 1$ is a Lie subalgebra of $\mathfrak{gl}(\infty)$. This

algebra is called the "Virasoro algebra" [4].

There are two manners for the supersymmetric extension of the Virasoro algebra, namely the "Ramond algebra" and the "Neveu-Schwartz algebra" [4]. The Ramond (resp. Neveu-Schwartz) algebra is the complex Lie superalgebra $\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1$, where the even part \mathcal{G}_0 has the basis $\{l_m, c; m \in \mathbb{Z}\}$, and the odd part \mathcal{G}_1 has the basis $\{g_k; k \in \mathbb{Z}\}$ (resp. $\{g_k; k \in \mathbb{Z} + 1/2\}$) satisfying the following bracket relations:

$$[l_m, l_n] = (m - n)l_{m+n} + \frac{1}{8}(m^3 - m) \delta_{m+n} c,$$

$$[l_m, g_k] = (\frac{1}{2}m - k)g_{m+k},$$

$$[g_j, g_k] = 2l_{j+k} + \frac{1}{2}(j^2 - \frac{1}{4}) \delta_{j+k} c \quad \text{and}$$

$$[\mathcal{G}, c] = \{0\}.$$

Proposition 3. Define the elements

$$L_m = -\sum_{j \in \mathbb{Z}} j (Z_{m+j}^{(00)} + Z_{m+j}^{(11)}) - \frac{m}{2} \sum_{j \in \mathbb{Z}} Z_{m+j}^{(11)} + \frac{1}{8} \delta_{m,0} \quad \text{and}$$

$$G_m = -\sqrt{-1} \sum_{j \in \mathbb{Z}} (Z_{m+j}^{(01)} + j Z_{m+j}^{(10)}) \quad \text{for } m \in \mathbb{Z}.$$

Then $l_m \longrightarrow L_m, g_m \longrightarrow G_m, c \longrightarrow 2$ is a representation of the Ramond algebra.

Proposition 4. Define the elements

$$L_m = - \sum_{j \in \mathbb{Z}} j (z_{m+j}^{(00)} + z_{m+j}^{(11)}) - \frac{m-1}{2} \sum_{j \in \mathbb{Z}} z_{m+j}^{(11)} + \frac{1}{2} \delta_{m,0} \quad \text{and}$$

$$G_{m+1/2} = -\sqrt{-1} \sum_{j \in \mathbb{Z}} (z_{m+j}^{(01)} + j z_{m+j+1}^{(10)}) \quad \text{for } m \in \mathbb{Z}.$$

Then $l_m \mapsto L_m$, $g_{m+1/2} \mapsto G_{m+1/2}$, $c \mapsto 2$ is a representation of the Neveu-Schwartz algebra.

In both cases a slight modification gives the more general representation, in which the central element c corresponds to arbitrarily given complex number.

Elements of $\mathfrak{gl}(\infty|\infty)$ are written as $\sum a_{ij}^{(\alpha\beta)} z_{ij}^{(\alpha\beta)} + a$. Consider the following conditions for the coefficients $a_{ij}^{(\alpha\beta)}$:

$$1) \quad a_{i+m}^{(\alpha)} z_{j+m}^{(\beta)} = a_{ij}^{(\alpha\beta)},$$

$$2) \quad \sum_{i=0}^{m^{(0)}-1} a_{i \ i+jm}^{(00)} - \sum_{i=0}^{m^{(1)}-1} a_{i \ i+jm}^{(11)} = 0 \quad \text{for any } j \in \mathbb{Z}.$$

For the sake of simplicity we assume that $m^{(0)} \neq m^{(1)}$.

References

- [1] L. Corwin, Y. Ne'eman and S. Sternberg: Graded Lie algebras in mathematics and physics (Bose-Fermi symmetry), Rev. Mod. Phys. 47 (1975), 573-603.
- [2] E. Date, M. Jimbo, M. Kashiwara and T. Miwa: Transformation groups for soliton equations, Proc. RIMS Symp. "Non-linear Integrable Systems — Classical Theory and Quantum Theory—" M. Jimbo and T. Miwa ed., World Scientific Publishing Co., 1983, 39-119.
- [3] V. G. Kac: Lie superalgebras, Advances in Math. 26 (1977), 8-96.
- [4] S. Mandelstam: Dual resonance models, Phys. Rep. 13 (1974), 259-353.
- [5] Yu. I. Manin and A. O. Radul: A supersymmetric extension of the Kadomtsev-Petviashvili hierarchy, preprint.