

On Gevrey Singularities of solutions of equations with non symplectic characteristics

Tsutomu SAKURAI 桜井 力
(Saitama Univ. Urawa 338) (埼玉大. 理)

In this note we shall construct parametrices for a specific class of differential operators with non symplectic characteristics and clarify the structure of Gevrey singularities of solutions of the corresponding equations using constructed parametrices.

0. Notation and preliminaries

If X is an open set of \mathbb{R}^N and $\nu \geq 1$, the Gevrey class of order ν ; which we denote by $G^\nu(X)$, is the set of all $u \in C^\infty(X)$ such that for every compact set $K \subset X$ there is a constant C_K with

$$|\partial_x^\alpha u(x)| \leq C_K^{|\alpha|+1} (\alpha!)^\nu, \quad x \in K,$$

for all multi-indices $\alpha \in \mathbb{N}^N$.

We use the following definition of the Gevrey wave front set given by Hörmander [14].

Definition 0.1. If $X \subset \mathbb{R}^N$ and $u \in \mathcal{D}'(X)$ we denote by $WF_\nu(u)$ the complement in $T^*(X) \setminus 0$ of the set of $(\dot{x}, \dot{\xi})$ such that there exist a neighborhood $U \subset X$ of \dot{x} , a conic neighborhood $V \subset \mathbb{R}^N \setminus 0$ of $\dot{\xi}$ and a bounded sequence $u_k \in \mathcal{D}'(X)$ which is equal to u in U and satisfies

$$|\hat{u}_k(\xi)| \leq C^{k+1} (k^\nu / |\xi|)^k, \quad k = 1, 2, \dots$$

for some constant C when $\xi \in V$, where \hat{u}_k denotes the Fourier transform of u_k .

$WF_1(u)$ is also denoted by $WF_A(u)$ since this is one of the

definition of the analytic wave front set known to be equivalent to the others; see e.g. Bony [3].

If π denotes the canonical projection of $T^*(X) \setminus 0$ on X then $u \in C^V(X \setminus \pi(WF_V(u)))$ and for a differential operator P with analytic coefficients, we have

$$WF_V(Pu) \subset WF_V(u) \subset \text{Char } P \cup WF_V(Pu),$$

where $\text{Char } P$ denotes the characteristic set of P . We say that P is C^V microhypoelliptic at $(\dot{x}, \dot{\xi})$ if there is a conic neighborhood $V \subset T^*(X) \setminus 0$ of $(\dot{x}, \dot{\xi})$ such that

$$WF_V(Pu) \cap V = WF_V(u) \cap V.$$

1. Statement of the results

Let Σ be the submanifold in $T^*(\mathbb{R}^N) \setminus 0$ of codimension $2d+d'$ given by

$$\Sigma = \{(x, \xi) \in T^*(\mathbb{R}^N) \setminus 0; x_1 = \dots = x_d = 0, \xi_1 = \dots = \xi_{d+d'} = 0\},$$

where $0 < d < d+d' < N$. With this Σ we set

$$\mathbb{R}^N_x = \mathbb{R}_t^d \times \mathbb{R}_y^n = \mathbb{R}_t^d \times \mathbb{R}_{y'}^{d'} \times \mathbb{R}_{y''}^{d''} \quad (d+n=N, d'+d''=n)$$

and denote by $\xi = (\tau, \eta) = (\tau, \eta', \eta'')$ the dual variables of $x = (t, y) = (t, y', y'') \in \mathbb{R}_t^d \times \mathbb{R}_{y'}^{d'} \times \mathbb{R}_{y''}^{d''}$. (In this coordinate $\Sigma = \{(t, y, \tau, \eta', \eta''); t = \tau = \eta' = 0, \eta'' \neq 0\}$.)

For a fixed integer $h \geq 1$ we shall consider a differential operator of order m with polynomial coefficients of the form:

$$(1.1) \quad P = p(t, D_t, D_y) = \sum_{\substack{|\alpha|+|\beta| \leq m \\ |\gamma| = |\alpha|+|\beta'|+(1+h)|\beta''|-m}} a_{\alpha\beta\gamma} t^\gamma D_y^\beta D_t^\alpha,$$

where $(\alpha, \beta, \gamma) = (\alpha, \beta', \beta'', \gamma) \in \mathbb{N}^d \times \mathbb{N}^{d'} \times \mathbb{N}^{d''} \times \mathbb{N}^d$ and $(D_t, D_y) = (-i\partial_t, -i\partial_y)$. Note that the symbol $p(t, \tau, \eta)$ has the following quasi-homogeneity:

$$(1.2) \quad \rho(t/\lambda^\rho, \lambda\tau, \lambda^\rho\eta', \lambda^\rho\eta'') = \lambda^{\rho m} \rho(t, \tau, \eta', \eta''), \quad \lambda > 0$$

with $\rho = 1/(1+h)$.

Let ρ_0 denote the principal symbol given by

$$(1.3) \quad \rho_0(t, \tau, \eta) = \sum_{\substack{|\alpha|+|\beta|=m \\ |\gamma|=h|\beta|}} a_{\alpha\beta\gamma} t^\gamma \eta^\beta \tau^\alpha.$$

For a point $(\dot{x}, \dot{\xi}) = (0, \dot{y}; 0, 0, \dot{\eta}'') \in \Sigma$ ($|\dot{\eta}''| \neq 0$) we suppose:

(H-1) There exists a constant $c > 0$ such that

$$|\rho_0(t, \tau, \eta', \dot{\eta}'')| \geq c(|\tau| + |\eta'| + |t|)^m, \quad (t, \tau, \eta') \in \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^{d'}.$$

We also consider the following condition due to Grushin.

(H-2) For all $\eta' \in \mathbb{R}^{d'}$, $\text{Ker } p(t, D_t, \eta', \dot{\eta}'') \cap \mathcal{G}(\mathbb{R}_t^d) = \{0\}$.

Here $p(t, D_t, \eta', \dot{\eta}'')$ is considered as an operator acting on $\mathcal{G}(\mathbb{R}_t^d)$ with a parameter $\eta' \in \mathbb{R}^{d'}$.

Remark. If $h = 1$, (H-2) is known to be equivalent to C^∞ micro-hypoellipticity with loss of $m/2$ derivatives; see e.g. Boutet de Monvel-Grigis-Helffer [4], see also Grushin [10],[11],[12] and the other authors [8],[15],[28].

Theorem I. Let P be an operator of the form (1.1) satisfying (H-1), (H-2) for $(\dot{x}, \dot{\xi}) \in \Sigma$. If $\nu \geq 1+h$ then P is G^ν micro-hypoelliptic at $(\dot{x}, \dot{\xi})$.

The condition $\nu \geq 1+h$ is the best in the sense that

Theorem II. Let P be an operator of the form:

$$(1.4) \quad P = p'(t, D_t, D_{y''}) + q(D_{y'}) \\ = \sum_{\substack{|\alpha|+|\beta''| \leq m \\ |\gamma| = |\alpha| + (1+h)|\beta''| - m}} a_{\alpha\beta''\gamma} t^\gamma D_{y''}^{\beta''} D_t^\alpha + \sum_{|\beta'|=m} b_{\beta'} D_{y'}^{\beta'}$$

satisfying (H-1) for $(0, \dot{\xi}) \in \Sigma$. Then one can find a neighborhood U of the origin in \mathbb{R}^N and a solution $u \in C^\infty(U)$ of $Pu = 0$ in U such that for every $\nu < 1+h$

$$(1.5) \quad (0, \dot{\xi}) \in WF_\nu(u) \subset WF_A(u) \subset \{(x, \lambda \dot{\xi}); x \in U, \lambda > 0\}.$$

If $1 \leq \nu < 1+h$ we can get a result on propagation of singularities of solutions for these operators.

Let Λ be the involutive submanifold of $T^*(\mathbb{R}^N) \setminus 0$ containing Σ given by

$$\Lambda = \{(t, y; \tau, \eta', \eta'') \in T^*(\mathbb{R}^N) \setminus 0; \eta' = 0\}.$$

Then in the canonical way Λ defines a bicharacteristic foliation in Σ as well as in Λ ; that is, each leaf Γ_0 is an integral submanifold of dimension d' of the vector fields generated by $\{\partial_{y_1}, \dots, \partial_{y_{d'}}\}$. (Note that $T_\rho(\Gamma_0) = T_\rho(\Sigma) \cap T_\rho(\Sigma)^\perp$ for all $\rho \in \Gamma_0$.)

Theorem III. Let Γ_0 be the bicharacteristic leaf passing through $(\dot{x}, \dot{\xi}) \in \Sigma$ defined as above and W be an open set containing $(\dot{x}, \dot{\xi})$ such that $\Gamma_0 \cap W$ is connected. Suppose that P is an operator of the form (1.1) satisfying (H-1) for $(\dot{x}, \dot{\xi})$ and that $1 \leq \nu < 1+h$. If $u \in \mathcal{D}'(\mathbb{R}^N)$ and $WF_\nu(Pu) \cap \Gamma_0 \cap W = \emptyset$ then either $\Gamma_0 \cap W \cap WF_\nu(u) = \emptyset$ or $\Gamma_0 \cap W \subset WF_\nu(u)$.

Remark. If $h = 1$ and $\nu = 1$ this is a special case of Theorem 2 in Grigis-Schapira-Sjöstrand [9]. See also Sjöstrand [29], [30] and Hasegawa [13] in this connexion.

Example. Let

$$(1.6) \quad P = \sum_{j=1}^{N-1} \partial_{x_j}^2 + \sum_{j=1}^d x_j^{2h} \partial_{x_N}^2 + h \sum_{j=1}^d c_j x_j^{h-1} \partial_{x_N},$$

where $c = (c_1, \dots, c_d) \in \mathbb{C}^d$. If

$$(1.7) \quad \sup_{\substack{\langle \sigma, \text{Im} c \rangle = 0 \\ |\sigma|_1 = 1}} |\langle \sigma, \text{Im} c \rangle| < 1 \quad (\sigma \in \mathbb{R}^d, |\sigma|_1 = |\sigma_1| + \dots + |\sigma_d|),$$

then P is G^{1+h} microhypoelliptic at every point in $T^*(\mathbb{R}^N) \setminus 0$.

In fact, noticing that $hx_j^{h-1} \partial_{x_N} = [\partial_{x_j}, x_j^h \partial_{x_N}]$ we get by Theorem 1' of Rothschild-Stein [25]

$$(1.8) \quad \sum_{j=1}^{N-1} \|\partial_{x_j} u\|^2 + \sum_{j=1}^d \|x_j^h \partial_{x_N} u\|^2 \leq C |(Pu, u)|$$

if (1.7) is fulfilled. This implies (H-2) while (H-1) is evident.

2. A study of the Grusin operator

We shall construct a right parametrix K for a self-adjoint operator $Q = (P^*P)^k$ with $2km \geq d+1$. (Note that the quasi-homogeneity (1.1), (1.2) and the conditions (H-1), (H-2) are preserved for Q with the order m replaced by $M = 2km$.) Then clearly $K^*(P^*P)P^*$ is a left parametrix of P and the microhypoellipticity of P follows immediately from that of Q .

In the construction of the parametrix we follow closely Métivier [21] and Ōkaji [22]. In this section we shall derive the estimates for the inverse of $\hat{Q} = \mathcal{F}_y Q \mathcal{F}_y^{-1}$.

2.1. Grusin operator. Let $Q = q(t, D_t, D_y)$ be an operator of the form (1.1) satisfying (H-1), (H-2) for $(\dot{x}, \dot{\xi}) \in \Sigma$. We may assume $(\dot{x}, \dot{\xi}) = (0, e_N) = (0; 0, \dots, 0, 1)$ without loss of generality; henceforth we let $\dot{\xi} = (0, \dot{\eta}) = (0, 0, \dot{\eta}'') = (0, \dots, 0, 1) \in \mathbb{R}^N$.

By the Fourier transform in y , we consider the equation:

$$(2.1) \quad q(t, D_t, \eta) v(t, \eta) = u(t, \eta)$$

in a conic neighborhood $U_\varepsilon \times V_\varepsilon$ of $(0; \dot{\eta}) \in \mathbb{R}^d \times (\mathbb{R}^n \setminus 0)$ given by

$$(2.2) \quad U_\varepsilon = \{t \in \mathbb{R}^d; |t| < 1\},$$

$$V_\varepsilon = \{\eta = (\eta', \eta'') \in \mathbb{R}^n \setminus 0; |\eta'| < \varepsilon \eta_n, |\eta'' - \dot{\eta}'' \eta_n| < \varepsilon \eta_n\}.$$

$q(t, D_t, \eta)$ is essentially the same operator that was studied by Grušin [12]; so we call it *Grušin operator*.

Now we shall start with the following lemma due to Grušin (: Lemma 3.4 in [12]).

Lemma 2.1. Let $Q = q(t, D_t, D_y)$ be an operator of order M of the form (1.1) satisfying (H-1) and (H-2) for $\xi = (0, 0, \dot{\eta}'')$. Then there exist a conic neighborhood V'' of $\dot{\eta}''$ and a constant C such that for all $\eta = (\eta', \eta'') \in \mathbb{R}^{d'} \times V''$

$$(2.3) \quad \sum_{|\beta| \leq M} \int |(|\eta''|^\rho + |\eta'| + |t|^h |\eta''|)|^{M-|\beta|} |D_t^\beta v(t)|^2 dt \\ \leq C \int |q(t, D_t, \eta) v(t)|^2 dt$$

for $v \in \mathcal{G}(\mathbb{R}_t^d)$, where $\rho = 1/(1+h)$.

Let us introduce new variables

$$\bar{t} = t \eta_n^\rho, \quad \bar{\tau} = \tau / \eta_n^\rho, \quad \bar{\eta}' = \eta' / \eta_n^\rho, \quad \bar{\eta}'' = \eta'' / \eta_n, \quad (\eta_n > 0)$$

and set

$$\bar{v}(\bar{t}, \bar{\eta}, \eta_n) = v(\bar{t} / \eta_n, \bar{\eta}' \eta_n, \bar{\eta}'' / \eta_n).$$

Then in view of (1.2) we have

$$(2.4) \quad q(\bar{t}, D_{\bar{t}}, \bar{\eta}) \bar{v}(\bar{t}, \bar{\eta}, \eta_n) = \eta_n^{-\rho M} q(t, D_t, \eta) v(t, \eta),$$

and the conic neighborhood $U_\varepsilon \times V_\varepsilon$ blows up into $\mathbb{R}_{\bar{\eta}}^d \times \bar{V}_\varepsilon$, where $\bar{V}_\varepsilon = \mathbb{R}_{\bar{\eta}}^{d'} \times \{\bar{\eta}'' \in \mathbb{R}^{d''}; |\bar{\eta}'' - \dot{\eta}''| < \varepsilon\}$.

By multiplying $\eta_n^{-\rho(M-d)}$, (2.3) becomes

$$(2.5) \quad \sum_{|\beta| \leq M} \int |(1 + |\bar{\eta}'| + |\bar{t}|^h |\bar{\eta}''|)|^{M-|\beta|} |D_{\bar{t}}^\beta \bar{v}(\bar{t}, \eta_n)|^2 d\bar{t} \\ \leq C \int |q(\bar{t}, D_{\bar{t}}, \bar{\eta}) \bar{v}(\bar{t}, \eta_n)|^2 d\bar{t}.$$

Moreover, we have

Proposition 2.2. *Let*

$$\bar{V}_\varepsilon^{\mathbb{C}} = \{ \bar{\eta} = (\bar{\eta}', \bar{\eta}'') \in \mathbb{C}^{d'} \times \mathbb{C}^{d''}; |\operatorname{Im} \bar{\eta}'| < \varepsilon(1 + |\operatorname{Re} \bar{\eta}'|), |\bar{\eta}'' - \bar{\eta}''| < \varepsilon \}.$$

If ε is chosen sufficiently small then for all $\bar{\eta} \in \bar{V}_\varepsilon^{\mathbb{C}}$ we have (2.5) with another constant C and there exists a left inverse $\bar{K}(\bar{\eta})$ of $q(\bar{t}, D_{\bar{t}}, \bar{\eta})$ depending holomorphically on $\bar{\eta} \in \bar{V}_\varepsilon^{\mathbb{C}}$.

2.2 Commutator estimates. We consider the operators

$$T_j = \partial_{\bar{t}_j} \quad \text{and} \quad T_{-j} = i\bar{t}_j \quad (j = 1, 2, \dots, d).$$

For a sequence $I = (j_1, \dots, j_k) \in \{\pm 1, \dots, \pm d\}^k$ we denote by T_I the operator

$$T_I = T_{j_1} T_{j_2} \dots T_{j_k}$$

and $\langle I \rangle = |I_+| + (1/h)|I_-| = \#\{j_l > 0\} + \#\{j_l < 0\}$.

We define the space

$$B^k(\bar{\eta}) = \{ u \in L^2(\mathbb{R}^d); \forall I, \langle I \rangle \leq k, T_I u \in L^2(\mathbb{R}^d) \}$$

for $k \in \mathbb{N}/h$ equipped with the norm:

$$\|u\|_{k, \bar{\eta}} = \max_{\langle I \rangle + j \leq k} (1 + |\bar{\eta}'|)^j \|T_I u\|_{L^2(\mathbb{R}^d)}$$

depending on $\bar{\eta} \in \bar{V}_\varepsilon^{\mathbb{C}}$. Note that $\|u\|_{0, \bar{\eta}}$ is the usual L^2 norm independent of $\bar{\eta}$; hence denoted by $\|u\|_0$. We also define $B^{-k}(\bar{\eta})$ the dual space of $B^k(\bar{\eta})$.

If L is an operator acting from $\mathcal{G}(\mathbb{R}^d)$ into $\mathcal{G}'(\mathbb{R}^d)$ we set

$$(\operatorname{ad} T_j)(L) = [T_j, L] = T_j L - L T_j \quad (j = \pm 1, \dots, \pm d)$$

and because the $\operatorname{ad} T_j$'s commute, we denote for a multi-index $\alpha =$

$$(\alpha_+, \alpha_-) = (\alpha_1, \dots, \alpha_d; \alpha_{-1}, \dots, \alpha_{-d}) \in \mathbb{N}^d \times \mathbb{N}^d$$

$$(\operatorname{ad} T)^\alpha = \prod_j (\operatorname{ad} T_j)^{\alpha_j}.$$

If the operator L from $\mathcal{S}(\mathbb{R}^d)$ into $\mathcal{S}'(\mathbb{R}^d)$ can be extended as a bounded operator in $L^2(\mathbb{R}^d)$ we denote by $\|L\|_0$ the norm of this extension, otherwise we agree with $\|L\|_0 = +\infty$.

At last, we introduce the norm:

$$\|L\|_{k, \bar{\eta}} = \text{Max}_{\langle I \rangle + \langle J \rangle + j \leq k} (1 + |\bar{\eta}'|)^j \|T_I L T_J\|_0$$

for $k \in \mathbb{N}/h$, then $\|L\|_{k, \bar{\eta}} < +\infty$ only means that L is bounded from $B^{-p}(\bar{\eta})$ to $B^{-p+k}(\bar{\eta})$ for all $p = 0, 1/h, 2/h, \dots, k$.

Now let $\bar{Q} = \bar{Q}(\bar{\eta}) = q(\bar{t}, D_{\bar{t}}, \bar{\eta})$. Then we can write

$$(2.6) \quad \bar{Q}(\bar{\eta}) = \sum_{\langle I \rangle + |\beta| \leq M} b_{I, \beta} \bar{\eta}^\beta T_I$$

and (2.5) by:

$$(2.7) \quad \|u\|_{M, \bar{\eta}} \leq C_0 \|\bar{Q}u\|_0 \quad \text{for } \bar{\eta} \in \bar{V}_\varepsilon^{\mathbb{C}}$$

We obtain as in Ōkaji [22]

Lemma 2.3. *If Q is a self-adjoint operator satisfying (2.7) then there exists a constant C_1 such that*

$$(2.8) \quad \|L\|_{M, \bar{\eta}} \leq C_1 (\|\bar{Q}L\|_0 + \|L\bar{Q}\|_0) \quad \text{for } \bar{\eta} \in \bar{V}_\varepsilon^{\mathbb{C}}$$

Let p be an integer. For real $R > 1$, $\mathcal{L}_R^p(\bar{V}_\varepsilon^{\mathbb{C}})$ denotes the space of operators L for which there is a constant C such that for all $\alpha = (\alpha_+, \alpha_-) \in \mathbb{N}^d \times \mathbb{N}^d$ and $\bar{\eta} \in \bar{V}_\varepsilon^{\mathbb{C}}$

$$\|(\text{ad } T)^\alpha(L)\|_{\langle \alpha \rangle + p, \bar{\eta}} \leq C |\alpha|! R^{|\alpha|},$$

where $\langle \alpha \rangle = (1/h)|\alpha_+| + |\alpha_-|$. Then $\mathcal{L}_R^p(\bar{V}_\varepsilon^{\mathbb{C}})$ becomes a Banach space in an obvious way.

Lemma 2.4. *Let \bar{Q} be as in Lemma 2.3. Then there are constants R_0 and C_2 depending only on C_1 and $\text{Max } |b_{I, \beta}|$ such that if both $\bar{Q}L$ and $L\bar{Q}$ are in $\mathcal{L}_R^0(\bar{V}_\varepsilon^{\mathbb{C}})$ then L is in $\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})$, moreover*

$$(2.9) \quad \|L\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})} \leq C_2 (\|\bar{Q}L\|_{\mathcal{L}_R^0(\bar{V}_\varepsilon^{\mathbb{C}})} + \|L\bar{Q}\|_{\mathcal{L}_R^0(\bar{V}_\varepsilon^{\mathbb{C}})}).$$

Proof is parallel to that of Métivier [21] Proposition 2.3 and Okaji [22] Lemma 7.2 and will be found in [27]. The following proposition is just a consequence of this lemma.

Proposition 2.5. *Let \bar{Q} be a self-adjoint operator satisfying (2.7) and let \bar{K} be the inverse of \bar{Q} such that $\bar{K}\bar{Q} = \bar{Q}\bar{K} = Id$. Then, if R is large enough, \bar{K} is in $\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})$.*

2.3. Kernel of the inverse. For an operator K from $\mathcal{G}(\mathbb{R}^d)$ to $\mathcal{G}'(\mathbb{R}^d)$, we denote by $K(\bar{t}, \bar{s})$ its distribution kernel.

Lemma 2.6. *If K is in $\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})$ with $M \geq d+1$ then $K(\bar{t}, \bar{s})$ is in $L^2(\mathbb{R}^d \times \mathbb{R}^d)$, moreover there exist constants \bar{C} and \bar{R} such that for all $\alpha = (\alpha_+, \alpha_-) \in \mathbb{N}^d \times \mathbb{N}^d$*

$$(2.10) \quad \|(\bar{t}-\bar{s})^{\alpha_-} (\partial_{\bar{t}} + \partial_{\bar{s}})^{\alpha_+} K(\bar{t}, \bar{s})\|_{L^2} \leq \bar{C} \|K\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})} \bar{R}^{|\alpha|} (\alpha_+!)^{1-\rho} (\alpha_-!)^\rho,$$

where $\rho = 1/(1+h)$.

Proof. Note that if K and K^* are bounded from $L^2(\mathbb{R}^d)$ into $B^{d+1}(\bar{\eta})$ then K is a Hilbert-Schmidt operator with the continuous kernel such that

$$\|K(\bar{t}, \bar{s})\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)} \leq C \|K\|_{d+1, \bar{\eta}}.$$

To prove (2.10) we consider

$$(2.11) \quad \begin{aligned} & \left((\bar{t}-\bar{s})^{\alpha_-} (\partial_{\bar{t}} + \partial_{\bar{s}})^{\alpha_+} \right)^{1+h} K(\bar{t}, \bar{s}) \\ &= \sum (\pm) \bar{t}^{\beta'_-} \bar{s}^{\beta''_-} \partial_{\bar{t}}^{\beta'_+} \partial_{\bar{s}}^{\beta''_+} (\bar{t}-\bar{s})^{\alpha_-} (\partial_{\bar{t}} + \partial_{\bar{s}})^{h\alpha_+} K(\bar{t}, \bar{s}), \end{aligned}$$

where the sum consists of $2^{h|\alpha_-|+|\alpha_+|}$ terms of the coefficients 1 or -1 with the multi-indices $\beta'_-, \beta''_-, \beta'_+, \beta''_+$ such that $\beta'_- + \beta''_- = h\alpha_-$, $\beta'_+ + \beta''_+ = \alpha_+$.

Now

$$(2.12) \quad \frac{\beta'_- \beta''_-}{\bar{t} \bar{s}} \frac{\beta'_+ \beta''_+}{\partial_{\bar{t}}^+ \partial_{\bar{s}}^+} (\bar{t} - \bar{s})^{\alpha_-} (\partial_{\bar{t}}^+ + \partial_{\bar{s}}^+)^{h\alpha_+} K(\bar{t}, \bar{s})$$

is the distribution kernel of

$$\frac{\beta'_- \beta''_-}{T_- T_+} (\text{ad } T_-)^{\alpha_-} (\text{ad } T_+)^{\alpha_+} (K) T_- T_+;$$

which is bounded from $L^2(\mathbb{R}^d)$ into $B^M(\bar{\eta})$ together with its adjoint. Since $M \geq d+1$ we know (2.12) is a continuous function with L^2 norm bounded by

$$C \|K\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon)} R^{|\alpha_-| + h|\alpha_+|} (|\alpha_-| + h|\alpha_+|)!.$$

Adding up these estimates we have

$$(2.13) \quad \begin{aligned} & \|((\bar{t} - \bar{s})^{\alpha_-} (\partial_{\bar{t}}^+ + \partial_{\bar{s}}^+)^{\alpha_+})^{1+h} K(\bar{t}, \bar{s})\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)} \\ & \leq C \|K\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon)} \bar{R}^{|\alpha_-|} (|\alpha_-| + h|\alpha_+|)! \end{aligned}$$

provided that $\bar{R} \geq (2R)^h$. Also we have

$$(2.14) \quad \|K(\bar{t}, \bar{s})\|_{L^2(\mathbb{R}^d \times \mathbb{R}^d)} \leq C \|K\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon)}.$$

Then a simple interpolation argument yields (2.10) in view of the Stirling formula. \square

2.4. Symbol of the inverse. We write the operator K of kernel $K(\bar{t}, \bar{s})$ with a symbol $k = \sigma(K)$ in the way that

$$(2.15) \quad K(\bar{t}, \bar{s}) = (2\pi)^{-d} \int e^{i\langle \bar{t} - \bar{s}, \bar{\tau} \rangle} k(\bar{t}, \bar{\tau}) d\bar{\tau}.$$

That is, k is the distribution on \mathbb{R}^{2d} given by

$$(2.16) \quad k(z^+, z^-) = \int e^{i\langle u, z^- \rangle} K(z^+, z^+ + u) du.$$

Here and below we use the notation $z = (z^+, z^-) = (z_1, \dots, z_d; z_{-1}, \dots, z_{-d}) \in \mathbb{R}^{2d}$.

Since (2.15), (2.16) have a sense as the partial Fourier transform

the mapping σ is clearly an isomorphism between $L^2(\mathbb{R}_t^d \times \mathbb{R}_s^d)$ and $L^2(\mathbb{R}_z^{2d})$. Also by the definition of σ we have

$$\sigma((\text{ad } T_j)(K)) = \partial_{z_j} \sigma(K).$$

hence Lemma 2.6 is restated as

Lemma 2.7. Let $k = k(\bar{\eta}) = \sigma(K(\bar{\eta}))$: the symbol of $K(\bar{\eta}) \in \mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})$ with $M \geq d+1$. Then there exist constants \bar{C}, \bar{R} such that for all $\alpha = (\alpha_+, \alpha_-) \in \mathbb{N}^d \times \mathbb{N}^d$ and $\bar{\eta} \in \bar{V}_\varepsilon^{\mathbb{C}}$

$$(2.17) \quad \|\partial_z^\alpha k(\bar{\eta})\|_{L^2(\mathbb{R}_z^{2d})} \leq \bar{C} \|K\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})} \bar{R}^{|\alpha|} (\alpha_+!)^{1-\rho} (\alpha_-!)^\rho$$

where $\rho = 1/(1+h)$.

Now suppose that $K(\bar{\eta}) \in \mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})$ ($M \geq d+1$) depends holomorphically on $\bar{\eta}$. Then we have

Proposition 2.8. Let $K(\bar{\eta})$ be as above and let $k(z, \bar{\eta}) = \sigma(K(\bar{\eta}))(z)$. Then there exists a constant C such that for $(z, \bar{\eta}) \in \mathbb{R}^{2d} \times \bar{V}_\varepsilon^{\mathbb{C}}$ with $0 < \varepsilon' < \varepsilon$ and for all $(\alpha, \beta) = (\alpha_+, \alpha_-, \beta', \beta'') \in \mathbb{N}^d \times \mathbb{N}^d \times \mathbb{N}^{d'} \times \mathbb{N}^{d''}$

$$(2.18) \quad \left| \partial_z^\alpha \partial_{\bar{\eta}}^\beta k(z, \bar{\eta}) \right| \leq C^{|\alpha|+|\beta|+1} \left(\frac{1}{\varepsilon-\varepsilon'} \right)^{|\beta|} (\alpha_+!)^{1-\rho} (\alpha_-!)^\rho \beta! (1+|\bar{\eta}'|)^{-|\beta'|},$$

where $\rho = 1/(1+h)$.

Proof. Recall that

$$\bar{V}_\varepsilon^{\mathbb{C}} = \{ \bar{\eta} = (\bar{\eta}', \bar{\eta}'') \in \mathbb{C}^{d'} \times \mathbb{C}^{d''}; |\text{Im } \bar{\eta}'| < \varepsilon(1+|\text{Re } \bar{\eta}'|), |\bar{\eta}'' - \bar{\eta}''| < \varepsilon \}.$$

Then we use the Cauchy inequality to obtain

$$\left\| \partial_{\bar{\eta}}^\beta K \right\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})} \leq \|K\|_{\mathcal{L}_R^M(\bar{V}_\varepsilon^{\mathbb{C}})} \left(\frac{\eta}{\varepsilon-\varepsilon'} \right)^{|\beta|} \beta! (1+|\bar{\eta}'|)^{-|\beta'|}.$$

Applying Lemma 2.7 to $\partial_{\bar{\eta}}^\beta K(\bar{\eta})$ we get (2.18) by means of the Sobolev

lemma. \square

3. Parametrix; proof of Theorem I

In Section 2 we have showed that there is the inverse $\bar{K}(\bar{\eta})$ of $\bar{Q}(\bar{\eta}) = q(\bar{t}, D_{\bar{t}}, \bar{\eta}) = (P^*P)^k(t, D_t, \eta)$ ($2km \geq d+1$) for $\bar{\eta} \in \bar{V}_\varepsilon^C$ such that

$$q(\bar{t}, D_{\bar{t}}, \bar{\eta}) \bar{K}(\bar{t}, \bar{s}, \bar{\eta}) d\bar{s} = \delta(\bar{t}-\bar{s}) d\bar{s}$$

with the kernel

$$\bar{K}(\bar{t}, \bar{s}, \bar{\eta}) = (2\pi)^{-d} \int e^{i\langle \bar{t}-\bar{s}, \bar{\tau} \rangle} \bar{k}(\bar{t}, \bar{\tau}, \bar{\eta}) d\bar{\tau},$$

where \bar{k} satisfies (2.18) in $\mathbb{R}_{\bar{t}}^d \times \mathbb{R}_{\bar{\tau}}^d \times (\bar{V}_\varepsilon^C \cap \mathbb{R}_{\bar{\eta}}^{d'})$ for $0 < \varepsilon' < \varepsilon$.

Now we return to the original variables:

$$t = \bar{t}/\eta_n^\rho, \quad \tau = \bar{\tau}/\eta_n^\rho, \quad \eta' = \bar{\eta}'/\eta_n^\rho, \quad \eta'' = \bar{\eta}''/\eta_n^\rho, \quad (\eta_n > 0)$$

and set

$$\hat{K}(t, s, \eta) = (2\pi)^{-d} \int e^{i\langle t-s, \tau \rangle} \hat{k}(t, \tau, \eta) d\tau,$$

where

$$\begin{aligned} (3.1) \quad \hat{K}(t, \tau, \eta) &= \eta_n^{-2\rho km} \bar{k}(\bar{t}, \bar{\tau}, \bar{\eta}) \\ &= \eta_n^{-2\rho km} \bar{k}(t\eta_n^\rho, \tau/\eta_n^\rho, \eta'/\eta_n^\rho, \eta''/\eta_n^\rho). \end{aligned}$$

Then in view of (2.4)

$$q(t, D_t, \eta) \hat{K}(t, s, \eta) ds = \delta(t-s) ds$$

for $\eta \in V_\varepsilon = \{(\eta', \eta'') \in \mathbb{R}^n \setminus 0; |\eta'| < \varepsilon\eta_n, |\eta'' - \hat{\eta}''\eta_n| < \varepsilon\eta_n\}$.

Let us introduce a cut off function given by Métivier:

Lemma 3.1. For given two cones $V_1 \subset\subset V_2 \subset \mathbb{R}^N \setminus 0$ and $0 < \rho < 1$ there exist $g \in C^\infty(\mathbb{R}^N)$ and C such that

$$(3.2) \quad g(\xi) = 0 \quad \text{for } \xi \in V_2 \quad \text{or} \quad |\xi| \leq 1$$

$$g(\xi) = 1 \quad \text{for } \xi \in V_1 \text{ and } |\xi| \geq 2$$

and

$$(3.3) \quad |\partial_\xi^\alpha g(\xi)| \leq C^{|\alpha|+1} \left(\frac{|\alpha|}{|\xi|}\right)^{\rho|\alpha|}$$

for all α, ξ such that $|\alpha| \leq |\xi|$. (Lemma 3.1 in [21].)

With $\rho = 1/(1+h)$ and $\xi \in V_1 \subset V_2 = \{\xi = (\tau, \eta) \in \mathbb{R}^d \times \mathbb{R}^n; |\tau| < \varepsilon', \eta \in V_\varepsilon\}$, we take $g(\xi) = g(\tau, \xi)$ as above and set $k_g(t, \tau, \eta) = \hat{\kappa}(t, \tau, \eta)g(\tau, \eta)$. Then

Proposition 3.2. There exists a constant C_0 such that

$$(3.4) \quad |\partial_\eta^\beta \partial_\tau^{\alpha_-} \partial_t^{\alpha_+} k_g(t, \tau, \eta)| \leq C_0^{|\alpha|+|\beta|+1} (1+|t|)^{|\beta|} (|\alpha_+|^{1-\rho} |\xi|^\rho)^{|\alpha_+|} \\ \times \left(\frac{|\alpha_-|}{|\xi|}\right)^{\rho|\alpha_-|} \left(\frac{|\beta'|}{|\xi|^{\rho+|\eta'|}} + \chi_g(\xi) \left(\frac{|\beta'|}{|\xi|}\right)^\rho\right)^{|\beta'|} \left(\frac{|\beta''|}{|\xi|}\right)^{\rho|\beta''|}$$

for $|\alpha_-|+|\beta| \leq |\xi|$, where $\xi = (\tau, \eta) = (\tau, \eta', \eta'') \in \mathbb{R}^N$, $(\alpha, \beta) = (\alpha_+, \alpha_-, \beta', \beta'') \in \mathbb{N}^d \times \mathbb{N}^d \times \mathbb{N}^{d'} \times \mathbb{N}^{d''}$, $\rho = 1/(1+h)$ and χ_g is the characteristic function of the support of $\nabla_\eta g$.

Now let $K_g = k_g(t, D_t, D_y) = \text{Op}(k_g)$; that is, the operator with the kernel:

$$(3.5) \quad K_g(t, y, s, w) = (2\pi)^{-N} \int e^{i\langle t-s, \tau \rangle + i\langle y-w, \eta \rangle} k_g(t, \tau, \eta) d\tau d\eta.$$

Then we have

$$(3.6) \quad QK_g = K_g^*Q = g(D_t, D_y) = \text{Op}(g)$$

and the following

Proposition 3.3.

$$(3.7) \quad \text{WF}_A(K_g) \subset \{(t, y, t, w; \tau, \eta, -\tau, -\eta) \in T^*(\mathbb{R}^{2N}) \setminus 0; y'' = w'', (\tau, \eta) \in V_2\}.$$

$$(3.8) \quad WF_{1+h}(K_g) \subset \{(t, y, t, y; \tau, \eta, -\tau, -\eta) \in T^*(\mathbb{R}^{2N}) \setminus 0; (\tau, \eta) \in V_2\}.$$

Proof. By Lemma 3.3 and Remark 3.4 in Métivier [21] we obtain (3.7). Hence to prove (3.8) it suffices to show that K is in G^{1+h} for $y' \neq 0$. Using the vector field $(1/|y'|^2)\langle y', D_{\eta'} \rangle$ for integrating by parts we can prove this as in the case 2 in the proof of Lemma 3.3 in [21]. \square

For any set V we write $\text{diag}(V) = \{(\rho, \rho) \in V \times V\}$. We have therefore proved the following; from which Theorem I follows immediately.

Theorem 3.4. *Let P be an operator of the form (1.1) satisfying (H-1), (H-2) for $(\dot{x}, \dot{\xi}) \in \Sigma$ and let $Q = (P^*P)^k$ with $2km \geq d+1$. Then there are a conic neighborhood $V \subset \mathbb{R}^N \setminus 0$ of $\dot{\xi}$ and an operator $K: \mathcal{S}'(\mathbb{R}^N)/C_0^\infty(\mathbb{R}^N) \longrightarrow \mathcal{D}'(\mathbb{R}^N)/C^\infty(\mathbb{R}^N)$ such that for every $u \in \mathcal{S}'(\mathbb{R}^N)$*

$$(3.9) \quad WF_A(QKu - u) \cap (\mathbb{R}^N \times V) = \emptyset,$$

$$(3.10) \quad WF_A(K^*Qu - u) \cap (\mathbb{R}^N \times V) = \emptyset$$

and that

$$(3.11) \quad WF'_{1+h}(K) \subset \text{diag}(T^*(\mathbb{R}^N) \setminus 0),$$

where $WF'_{1+h}(K) = \{(x, \xi; \tilde{x}, \tilde{\xi}); (x, \tilde{x}; \xi, -\tilde{\xi}) \in WF_{1+h}(K)\}$.

4. Proof of Theorem II

Let $\dot{\xi} = (0, 0, \dot{\eta}''')$ with $\dot{\eta}''' \neq 0$. We consider the operator $\rho'(t, D_t, \dot{\eta}''')$; which is precisely the same one that was studied by Grušin [10].

From the result of Grušin [10] we can take $c \in \mathbb{C}$ and $0 \equiv v \in \mathcal{G}(\mathbb{R}^N)$ such that

$$(4.1) \quad \rho'(t, D_t, \dot{\eta}''')v(t) = -c^m q(\bar{\eta}'),$$

where $\bar{\eta}' \in \mathbb{R}^{d'}$ is fixed with $|\bar{\eta}'| = 1$. Then

$$u_\lambda(t, y) = \exp(i\lambda^\rho \langle y', \bar{\eta}' \rangle + i\lambda \langle y'', \bar{\eta}'' \rangle) v(t\lambda^\rho), \quad \rho = 1/(1+h)$$

is a solution of $pu = 0$ for every $\lambda \geq 0$. Hence

$$u(t, y) = \int_0^{+\infty} u_\lambda(t, y) e^{-\lambda^\rho} d\lambda$$

is a C^∞ solution in $U = \{(t, y', y'') \in \mathbb{R}^N; |\operatorname{Im} z| |y'| < 1\}$.

By Lemma 3.7 in Ōkaji [22], v satisfies the estimate

$$|\partial_t^\alpha v(t)| \leq C^{|\alpha|+1} (\alpha!)^{1-\rho}.$$

Hence we have

$$(4.2) \quad WF_A(u) \subset \{(t, y; 0, 0, \lambda \bar{\eta}'') \in T^*(\mathbb{R}^N) \setminus 0; \lambda > 0\}$$

in the same way as (3.7).

On the other hand, since v is analytic, $\partial_t^\alpha v(0) \neq 0$ for some $\alpha \in \mathbb{N}^d$. Therefore,

$$(4.3) \quad |\langle \eta'', D_{y''} \rangle^k \partial_t^\alpha u(0, 0)| = \int_0^{+\infty} |\bar{\eta}''|^{2k} \lambda^{\rho|\alpha|+k} |\partial_t^\alpha v(0)| e^{-\lambda^\rho} d\lambda \\ = C |\bar{\eta}''|^{2k} \Gamma((k+1)/\rho + |\alpha|), \quad (C > 0).$$

This combined with (4.2) implies $(0; 0, 0, \bar{\eta}'') \in WF_v(u)$ for every $v < 1+h$, and proof is now complete. \square

5. Second microlocalization in Gevrey class

Following Sjöstrand [29] we introduce the Fourier-Bros-Iagolnitzer transform (F.B.I. tr.):

$$(5.1) \quad T^{(1)} f(z, \lambda) = \int e^{-\lambda(z-x)^2/2} f(x) dx, \quad (f \in \mathcal{G}'(\mathbb{R}^N))$$

associated to $\kappa: T^*(\mathbb{R}^N) \setminus 0 \ni (x, \xi) \longrightarrow x - i\xi \in \mathbb{C}_z^N$.

$T^{(1)} f$ is defined on $\mathbb{C}_z^N \times \mathbb{R}_\lambda^+$, holomorphic with respect to z and bounded by $C e^{\lambda |\operatorname{Im} z|^2/2} (\lambda + |y|)^k$ for some C, k real.

In terms of the F.B.I. tr. we can characterize the Gevrey wave

front set as follows: For $f \in \mathcal{G}'(\mathbb{R}^N)$, $(\dot{x}, \dot{\xi}) \in WF_\nu(f)$ if and only if there are constants $C, c > 0$ such that

$$(5.2) \quad |T^{(1)}f(z, \lambda)| \leq C e^{\frac{\lambda}{2} |\operatorname{Im} z|^2 - c\lambda} \quad \text{for } |z - (\dot{x} - i\dot{\xi})| < c.$$

Let Λ be the involutive submanifold of $T^*(\mathbb{R}^N)$:

$$\Lambda = \{(x, \xi) \in T^*(\mathbb{R}^N); \xi_1 = \dots = \xi_{d'} = 0\} \quad (1 \leq d' < N),$$

and Γ_0 be the bicharacteristic leaf pathing through $(\dot{x}, \dot{\xi}) \in \Lambda$.

Then Λ and Γ_0 can be identified with $\kappa(\Lambda) = \{z \in \mathbb{C}^N; \operatorname{Im} z' = 0\}$ and $\kappa(\Gamma_0) = \{z \in \mathbb{C}^N; \operatorname{Im} z' = 0, z'' = (\dot{x}'' - i\dot{\xi}'')\}$ respectively, where $z = (z', z'') \in \mathbb{C}^{d'} \times \mathbb{C}^{N-d'}$.

We set $\varphi_\Lambda(z) = |\operatorname{Im} z''|/2$; which is the pluri-subharmonic function canonically associated to Λ . If Ω is a neighborhood of $\dot{z} \in \kappa(\Lambda)$, we denote by $H_\Lambda^{\nu, \text{loc}}(\Omega)$ the space of holomorphic functions $u(z, \lambda)$ in Ω with a parameter $\lambda > 0$ such that for all $K \subset\subset \Omega$ and $\varepsilon > 0$ there exists $C_{K, \varepsilon}$ with the estimate:

$$(5.3) \quad |u(z, \lambda)| \leq C_{K, \varepsilon} e^{\lambda \varphi_\Lambda + \varepsilon \lambda^{1/\nu}} \quad \text{for } z \in K, \lambda \geq 1.$$

For $\dot{z} \in \Lambda$ we also use the notation: $u \in H_{\Lambda, \dot{z}}^\nu$ if there is a neighborhood $\omega_{\dot{z}}^\circ$ of \dot{z} such that $u \in H_\Lambda^\nu(\omega_{\dot{z}}^\circ)$.

If $u \in H_\Lambda^{\nu, \text{loc}}(\Omega)$ we denote by $S_\Lambda^\nu(u)$ the subset in Ω defined by:

$$(5.4) \quad \dot{z} \in S_\Lambda^\nu(u) \quad \text{if and only if there exist a neighborhood } \omega_{\dot{z}}^\circ \text{ of } \dot{z} \text{ and constants } C, c > 0 \text{ such that}$$

$$|u(z, \lambda)| \leq C e^{\lambda \varphi_\Lambda - c\lambda^{1/\nu}} \quad \text{for } z \in \omega_{\dot{z}}^\circ, \lambda \geq 1.$$

By applying the maximum principle to $z' \mapsto \lambda^{-1/\nu} (\log |u(z, \lambda)| - \lambda |\operatorname{Im} z''|)$ it can be seen easily the following two lemmas.

Lemma 5.1. *Let Γ_0 be a bicharacteristic leaf in Λ and ω be a connected open set in Γ_0 containing $(\dot{x}, \dot{\xi})$. If $u \in H_{\Lambda, \dot{z}}^\nu$ for all*

$z \in \kappa(\omega)$ and $\kappa(\dot{x}, \dot{\xi}) = \dot{x} - i\dot{\xi} \notin S_{\Lambda}^{\nu}(u)$ then $\kappa(\omega) \cap S_{\Lambda}^{\nu}(u) = \emptyset$.

Lemma 5.2. Let $(\dot{x}, \dot{\xi}) \in \Lambda$, $f \in \mathcal{G}'(\mathbb{R}^N)$. If $(\dot{x}, \dot{\xi}) \notin WF_{\nu}(f)$ and $T_{\Lambda}^{(1)}f \in H_{\Lambda, \dot{x}-i\dot{\xi}}^{\nu}$ then $\dot{x}-i\dot{\xi} \notin S_{\Lambda}^{\nu}(T^{(1)}f)$.

Let us introduce the F.B.I. tr. of second kind along Λ following Lebeau [19]:

$$(5.5) \quad T_{\Lambda}^{(2)}f(w, \mu, \lambda) = \int e^{-\lambda(w''-x'')^2/2 - \lambda\mu(w'-x')^2/2} f(x) dx \quad (f \in \mathcal{G}'(\mathbb{R}^N)).$$

Then $T_{\Lambda}^{(2)}f(w, \mu, \lambda)$ is a holomorphic function with respect to $w \in \mathbb{C}^N$ with the bound:

$$|T_{\Lambda}^{(2)}f(w, \mu, \lambda)| \leq C e^{\frac{\lambda}{2} |\text{Im} w''|^2 + \frac{\lambda\mu}{2} |\text{Im} w'|^2} (\lambda + |w|)^k.$$

It was shown in [20] and [2] that the relation between $T^{(1)}f$ and $T^{(2)}f$ is

$$(5.6) \quad T^{(2)}f(w, \mu, \lambda) = \left(\frac{\lambda}{2\pi(1-\mu)} \right)^{\frac{d'}{2}} \int_{\mathbb{R}^{d'}} e^{-\lambda\rho(w'-x')^2/2} T^{(1)}f(x', w'', \lambda) dx',$$

where $\rho = \mu/(1-\mu)$ with the inversion formula:

$$(5.7) \quad T^{(1)}f(z, \lambda) = \frac{1}{2} \left(\frac{1}{2\pi\lambda} \right)^{\frac{d'}{2}} \int_{\mathbb{R}^{d'}_{\xi}} e^{-\lambda R|\xi'|/2} \left(1 - i \frac{\langle \xi', \nabla' \rangle}{\lambda |\xi'|} \right) T^{(2)}f\left(z' - i \frac{R\xi'}{\lambda |\xi'|}, z'', \mu, \lambda\right) \frac{R d\xi'}{R + |\xi'|},$$

where $\mu = |\xi'|/(R + |\xi'|)$.

Now we define the second wave front set adapted to the Gevrey class. (See also Esser [7].)

Definition 5.3. If $1 \leq \nu < +\infty$ and $f \in \mathcal{G}'(\mathbb{R}^N)$, the second wave front set along Λ of f ; denoted by $WF_{\Lambda, \nu}^{(2)}(f)$, is the subset in $T_{\Lambda}(T^*(\mathbb{R}^N) \setminus 0)$ defined by the following condition:

$$(5.8) \quad (\dot{x}, 0, \dot{\xi}''; \dot{\sigma}') \notin WF_{\Lambda, \nu}^{(2)}(f)$$

if and only if there exist $C, c > 0, 0 < \mu_0 < 1$ and a decreasing function $o(\lambda)$ with $\lim_{\lambda \rightarrow +\infty} o(\lambda) = 0$ such that

$$(5.9) \quad |T_{\Lambda}^{(2)} f(w, \mu, \lambda)| \leq C e^{\frac{\lambda}{2} |\operatorname{Im} w''|^2 + \frac{\lambda}{2} \mu |\operatorname{Im} w'|^2 - c \lambda \mu}$$

for

$$(5.10) \quad 0 < \mu < \mu_0, \lambda \mu > o(\lambda) \lambda^{1/\nu}, |w' - (\dot{x}' - i\dot{\sigma}')| + |w'' - (\dot{x}'' - i\dot{\xi}'')| < c.$$

Using (5.6) and (5.7) we can show the following:

Lemma 5.4. *Let $(\dot{x}, \dot{\xi}) \in \Lambda$ and $f \in \mathcal{G}'(\mathbb{R}^N)$. Then $T^{(1)} f \in H_{\Lambda, \dot{x} - i\dot{\xi}}^{\nu}$ if and only if $\pi_{\Lambda}^{-1}(\dot{x}, \dot{\xi}) \cap WF_{\Lambda, \nu}^{(2)}(f) = \emptyset$, where $\pi_{\Lambda}: T_{\Lambda}(T^*(\mathbb{R}^N) \setminus 0) \rightarrow \Lambda$ is the canonical projection.*

At last, we introduce the space of the partially holomorphic Gevrey functions $G_{\mathcal{A}_x}^{\nu}$ as follows: $f(x) \in G_{\mathcal{A}_x}^{\nu}(\Omega)$ if and only if for every compact set $K \subset \subset \Omega$ there is a constant C such that

$$(5.11) \quad |\partial_x^{\alpha'} \partial_x^{\alpha''} f(x)| \leq C^{|\alpha|+1} \alpha'! (\alpha''!)^{\nu} \quad \text{for } x \in K.$$

We have

Lemma 5.5. *If $f \in \mathcal{G}'(\mathbb{R}^N) \cap G_{\mathcal{A}_x}^{\nu}(\Omega)$ and $1 \leq \nu' < \nu$ then $T^{(1)} f \in H_{\Lambda, z}^{\nu'}$ for every $z \in \kappa(\pi^{-1}(\Omega) \cap \Lambda)$.*

6. Proof of Theorem III

As in Section 2 we suppose that $\dot{x} = 0, \dot{\xi} = (0, 0, \dot{\eta}'') = (0, \dots, 0, 1) \in \mathbb{R}^N \setminus 0$ and set $Q = (P^*P)^k$ with $2km \geq d+1$. Here we also introduce the pseudo-differential operator:

$$(6.1) \quad \operatorname{Op}(r) = \operatorname{Op}(\eta_n^{2km/(1+h)} e^{-|\eta'|^{2l(1+h)}/\eta_n^{2l}},$$

where l is a positive integer to be determined. Then r has the same quasi-homogeneity in its symbol as Q has.

Consider the operator $Q + \operatorname{Op}(r)$. Then it satisfies (H-2) since Q is non negative self-adjoint operator at $\dot{\xi}$. We also note that

though not being polynomial, r is holomorphic with the uniform bound $O(|\xi|^{2km/(1+h)})$ in a small quasi-homogeneous neighborhood of ξ of the form:

$$V_\varepsilon^{\mathbb{C}} = \{(\eta', \eta'') \in \mathbb{C}^{d'} \times \mathbb{C}^{d''}; |\operatorname{Im} \eta'| < \varepsilon(|\eta_n|^{1/(1+h)} + |\operatorname{Re} \eta'|), |\eta''/\eta_n - \dot{\eta}''| < \varepsilon\}.$$

Now all the results in Section 2 are remain valid for $Q + \operatorname{Op}(r)$ and we get the symbol $k_g(t, \tau, \eta)$ satisfying (3.4) such that

$$(6.2) \quad \operatorname{Op}(k_g)^*(Q + \operatorname{Op}(r)) = \operatorname{Op}(g).$$

Here g is an arbitrary cut off function satisfying (3.3) for $\rho = 1/(1+h)$ with its support in

$$(6.3) \quad V_{\varepsilon_0} = \{(\tau, \eta) \in T^*(\mathbb{R}^N) \setminus 0; |\tau| < \varepsilon_0 \eta_n, |\eta'| < \varepsilon_0 \eta_n, |\eta''/\eta_n - \dot{\eta}''| < \varepsilon_0\}.$$

If $(\dot{x}, \dot{\xi}) = (0; 0, 0, \dot{\eta}'') \in \Sigma$ then the bicharacteristic leaf is $\Gamma_0 = \{(0, y', 0; 0, 0, \dot{\eta}''); y' \in \mathbb{R}^{d'}\}$. For any compact set $F \subset \pi(\Gamma_0 \cap W)$ there exist a neighborhood $U \subset\subset O_R = \{x \in \mathbb{R}^N; |x| < R\}$ of F and a conic neighborhood V of $\dot{\xi}$ such that

$$(6.4) \quad WF_V(Pu) \cap \bar{U} \times (\bar{V} \setminus 0) = \emptyset,$$

where \bar{U}, \bar{V} denote the closures of U, V respectively.

After replacing u by φu with a suitable $\varphi \in C_0^\infty(O_R)$ we can suppose $u \in \mathcal{S}'(O_R)$ with no influence on (6.4).

We fix a conic neighborhood V_2 of $\dot{\xi}$ with $V_2 \subset\subset V \cap V_{\varepsilon_0}$. If we choose another conic neighborhood V_1 of $\dot{\xi}$ sufficiently small then the cut off function g in Lemma 3.1 can be taken in the form: $g(\xi) = g'(\eta', \eta_n)g''(\tau, \eta'')$ so that $\operatorname{supp} \nabla_\eta g \subset \{(t, \eta', \eta''); |\eta'| > \delta|\xi|\}$ for some $\delta > 0$.

As in Proposition 3.3 one can see the following:

Proposition 6.1. *If k_g satisfies (3.4) with $\chi_g(\xi) = 0$ for $|\eta'| < \delta|\xi|$ ($\delta > 0$), then*

$$(6.5) \quad K_g(t, y, s, w) \in G^{1+h}_{y', w}, ((\mathbb{R}^N \times \mathbb{R}^N) \setminus \operatorname{diag}(\mathbb{R}^N)),$$

where K_g denotes the distribution kernel of k_g .

Now we let g be taken as above and write for $u \in \mathcal{S}'(O_R)$

$$(6.6) \quad \begin{aligned} \text{Op}(g)u &= \text{Op}(k_g)^*Qu + \text{Op}(k_g)^*\text{Op}(r)u \\ &= \text{Op}(k_g)^*Qu + \text{Op}(r)\text{Op}(k_g)^*u. \end{aligned}$$

We shall apply the theory of second microlocalization along the involutive submanifold:

$$\Lambda = \{(t, y; \tau, \eta', \eta'') \in T^*(\mathbb{R}^N) \setminus 0; \eta' = 0\}.$$

Hereafter, we also denote the coordinate in $T^*(\mathbb{R}^N)$ by

$$x' = y', \quad x'' = (t, y'') \quad \text{and} \quad \xi' = \eta', \quad \xi'' = (\tau, \eta'')$$

and use the notation in Section 4 without mentioning it.

First we study $\text{Op}(r)\text{Op}(k_g)^*u$, where

$$r(\xi) = \frac{2km/(1+h)}{\eta_n} e^{-|\eta'|^{2l(1+h)}/\eta_n^{2l}}$$

was given in (6.1). Now we choose l so that $(1+h)-1/2l > \nu$. Then

$$(6.7) \quad |\eta'|^{2l(1+h)}/\eta_n^{2l} \geq |\eta'| \quad \text{for} \quad |\eta'| \geq \eta_n^{-\varepsilon} \eta_n^{1/\nu}, \quad \eta_n > 0,$$

where $\varepsilon = (1/\nu) - (2l/(2l(1+h)-1)) > 0$. We can see easily the following:

Lemma 6.2. *If $r = O(e^{-c|\eta'|})$, $c > 0$ for $|\eta'| \geq \eta_n^{-\varepsilon} \eta_n^{1/\nu}$, $\eta_n > 0$ then for every $u \in \mathcal{S}'(\mathbb{R}^N)$*

$$(6.8) \quad \text{WF}_{\Lambda, \nu}^{(2)}(\text{Op}(r)u) \cap \pi_{\Lambda}^{-1}(\Gamma_0) = \emptyset.$$

Since $\text{Op}(k_g)(\mathcal{S}) \subset \mathcal{S}$; equivalently $\text{Op}(k_g)^*(\mathcal{S}') \subset \mathcal{S}'$, (6.8) holds for $\text{Op}(r)\text{Op}(k_g)^*u$. Therefore we have

$$(6.9) \quad \Gamma^{(1)}(\text{Op}(r)\text{Op}(k_g)^*u) \in H_{\Lambda, z}^{\nu} \quad \text{for all} \quad z \in \kappa(\Gamma_0)$$

in view of Lemma 5.4.

Next we study $\text{Op}(k_g)^*Qu$. Let \tilde{g} be another cut off function given by Lemma 3.1 with two cones \tilde{V}_1, \tilde{V}_2 such that

$$V_2 \subset\subset \tilde{V}_1 \subset\subset \tilde{V}_2 = V.$$

Noticing that $WF_V(Qu) \subset WF_V(Pu)$, we then get by (6.4)

$$(6.10) \quad WF_V(\text{Op}(\tilde{g})Qu) \subset WF_V(Pu) \cap (\mathbb{R}^N \times V_2) \subset \pi^{-1}(O_R \setminus U),$$

$$(6.11) \quad WF_V(\text{Op}(1-\tilde{g})Qu) \subset WF_V(Pu) \setminus (\mathbb{R}^N \times \tilde{V}_1) \subset \pi^{-1}(O_R) \setminus (\mathbb{R}^N \times \bar{V}_2).$$

Hence we can write

$$(6.12) \quad Qu = \chi_{F_\varepsilon} \text{Op}(\tilde{g})Qu + \chi_{O_R} (1-\chi_{F_\varepsilon}) \text{Op}(\tilde{g})Qu + \chi_{O_R} \text{Op}(1-\tilde{g})Qu \\ (\equiv v_1 + v_2 + v_3),$$

where χ_B denotes the characteristic function of each set B and

$$F_\varepsilon = \{(x', x'') \in \mathbb{R}^N; (x', 0) \in F, |x''| \leq \varepsilon\}$$

with $\varepsilon > 0$ so small that $F_\varepsilon \subset U$.

In the following we assume farther that

$$(6.13) \quad F \text{ is convex with an analytic boundary in } \pi(\Gamma_0),$$

By (6.10) we see that

$$WF_V(v_1) \subset \{(x, \xi); (x', \xi') \in T_{\partial F}^*(\pi(\Gamma_0)), |x''| < \varepsilon\} \cup \pi^{-1}(\{x; |x''| \geq \varepsilon\}).$$

Hence by (3.7)

$$(6.14) \quad \text{Op}(k_g)^*v_1 \in G^{\nu}(\text{Int}(F_\varepsilon)),$$

where $\text{Int}(F_\varepsilon)$ denotes the interior of F_ε .

Since $\text{supp}(v_2) \subset \bar{O}_R \setminus F_\varepsilon$, it follows by Proposition 6.1

$$(6.15) \quad \text{Op}(k_g)^*v_2 \in G^{1+h_{x'}}(\text{Int}(F_\varepsilon)).$$

Thus by Lemma 5.5 we have

$$(6.16) \quad T^{(1)}(\text{Op}(k_g)^*v_2) \in H_{\Lambda, z}^v \quad \text{for all } z \in \kappa(\pi^{-1}(\text{Int}(F_\varepsilon) \cap \Lambda)).$$

In view of (6.11) we have

$$WF_v(v_3) \subset O_R \times (\mathbb{R}^N \setminus \bar{V}_1) \cup T_{\partial O_R}^*(\mathbb{R}^N).$$

Again by (3.7) this yields

$$(6.17) \quad \text{Op}(k_g)^*v_3 \in G^v(\text{Int}(F_\varepsilon)).$$

Consequently, by (6.9) and (6.14)–(6.17), we have

$$(6.18) \quad \text{Op}(g)u = u_1 + u_2,$$

where

$$u_1 = \text{Op}(k_g)^*(v_1 + v_2) \in G^v(\text{Int}(F_\varepsilon))$$

and

$$u_2 = \text{Op}(k_g)^*v_2 + \text{Op}(r)\text{Op}(k_g)^*u$$

with

$$T^{(1)}(u_2) \in H_{\Lambda, z}^v \quad \text{for all } z \in \kappa(\pi^{-1}(\text{Int}(F_\varepsilon)) \cap \Gamma_0).$$

Now we apply Lemma 5.1, 5.2 and obtain

$$(6.18) \quad \text{If } (\dot{x}, \dot{\xi}) \in \pi^{-1}(\text{Int}(F_\varepsilon)) \cap \Gamma_0 \text{ and } (\dot{x}, \dot{\xi}) \notin WF_v(u_2) \\ \text{then } \pi^{-1}(\text{Int}(F_\varepsilon)) \cap \Gamma_0 \cap WF_v(u_2) = \emptyset.$$

Because $g \equiv 1$ in the neighborhood V_1 of $\dot{\xi}$,

$$WF_v(u_2) \cap \pi^{-1}(\text{Int}(F_\varepsilon)) \cap \Gamma_0 = WF_v(u) \cap \pi^{-1}(\text{Int}(F_\varepsilon)) \cap \Gamma_0.$$

Therefore (6.18) implies Theorem III for $\hat{W} = \pi^{-1}(\text{Int}(F_\varepsilon))$.

Since any compact set in $\Gamma_0 \cap W$ can be covered by a finite number of such \hat{W} 's we have actually proved Theorem III. \square

7. Remarks

The problem to determine the Gevrey class in which certain C^∞ hypoelliptic operators still remain hypoelliptic, has its origin in

the celebrated example given by Baouendi-Goulaouic [1]:

$$p_1 = \partial_t^2 + \partial_x^2 + t^2 \partial_y^2;$$

which has a solution u of $p_1 u = 0$ in a neighborhood of the origin only belonging to G^2 .

Deridj-Zuily [5] and Durand [6] have studied Gevrey hypoellipticity for second order operators and proved, for example, G^{1+h+0} and G^{1+h} hypoellipticity of the operators in Example 1 respectively.

However, as was showed by Parenti-Rodino [24], hypoellipticity does not always imply microlocal one. In this respect, Iwasaki [17] proved among others G^2 microhypoellipticity for double characteristic operators. Our Theorem I is an extension of this in some sense, though the operators are much restricted.

Recently, Kajitani-Wakabayashi also studied Gevrey microhypoellipticity in [18] but for more general classes of operators and obtained the results including our Theorem I as a special case.

However our proof by constructing parametrices reveals how the quasi-homogeneity of the operators relate to the lowest order of Gevrey class in which operators being hypoelliptic and gives a more precise information on the singularities of solutions (: Proposition 6.1 and Theorem III).

Moreover present method can be applied to the operator:

$$p_2 = \partial_t^2 + t^2 \partial_x^2 + t^4 \partial_y^2$$

given by Oleinic-Radkevič [23] and one can show that p_2 is hypoelliptic in $G^{3/2}$; while Durand's results gives only G^3 hypoellipticity of p_2 . This will be given in the future publication together with the complete description of the results in Section 5.

References

- [1] Baouendi M.S. and Goulaouic C.: Non analytic hypoellipticity for

- some degenerate operators, Bull. Amer. Math. Soc. **78**(1972) 483-486.
- [2] Bardos C., Lebeau G. and Rauch J.: Scattering frequencies and Gevrey 3 singularities, preprint.
- [3] Bony J.M.: Equivalence des diverses notions de spectre singulier analytique, Sém. Goulaouic-Schwartz **III**(1976-1977).
- [4] Boutet de Monvel L., Grigis A. and Helffer B.: Paramétrixes d'opérateurs pseudodifférentiels à caractéristiques multiples, Astérisque **34-35**(1976) 93-121.
- [5] Derridj M. and Zuily C.: Sur la régularité Gevrey des opérateurs de Hörmander, J. Math. pures et appl. **52**(1973) 309-336.
- [6] Durand M.: Régularité Gevrey d'une classe d'opérateurs hypoelliptiques, J. Math. pure et appl. **57**(1978) 323-360.
- [7] Esser P.: preprint.
- [8] Grigis A.: Hypoellipticité et paramétrix pour des opérateurs pseudodifférentiels à caractéristiques doubles, Astérisque **34-35**(1976)183-204.
- [9] Grigis A., Schapira P. and Sjöstrand J.: Propagation des singularités analytiques pour des opérateurs à caractéristiques multiples, C.R.A.S. **293,I**(1981) 397-400.
- [10] Grušin V.V.: on a class of hypoelliptic operators, Mat. Sb. **83** = Math. U.S.S.R. Sb. **12**(1970) 458-476.
- [11] ———— : On a class of elliptic pseudodifferential operators degenerate on submanifold, Mat. Sb. **84** = Math. U.S.S.R. Sb. **13**(1972) 155-185.
- [12] ———— : Hypoelliptic differential equations and pseudodifferential operators with operator valued symbols, Mat. Sb. **88** = Math. U.S.S.R. Sb. **27**(1972) 497-513.
- [13] Hasegawa K.: Propagation of micro-analyticities of solutions to some class of linear partial differential equations with non-involutive double characteristics, to appear in J. Fac. Sci. Univ. Tokyo.
- [14] Hörmander L.: Uniqueness theorems and wave front set for

- solutions of linear differential equations with analytic coefficients, *Comm. Pure Appl. Math.* 24(1971) 671-704.
- [15] ————— : A class of hypoelliptic pseudo-differential operators with double characteristics, *Math. Ann.* 217(1975) 165-188.
- [16] ————— : *The analysis of linear partial differential operators*, Springer, 1983.
- [17] Iwasaki C.: Gevrey hypoellipticity and pseudo-differential operators on Gevrey class, *Lecture Notes in Math.* 1256(1987) 281-293.
- [18] Kajitani K. and Wakabayashi S.: Hypoelliptic operators in Gevrey class, preprint.
- [19] Lebeau G.: Deuxièm microlocalisation à croissance, *Sém. Goulaouic-Meyer-Schwartz XV*(1982-1983).
- [20] ————— : Deuxièm microlocalisation sur les sous-variétés isotropes, *Ann. Inst. Fourier* 35(1985) 145-216.
- [21] Métivier G.: Analytic hypoellipticity for operators with multiple characteristics, *Comm. PDE* 6(1980) 1-90.
- [22] Ōkaji T.: Analytic hypoellipticity for operators with symplectic characteristics, *J. Math. Kyoto Univ.* 25(1985) 489-514.
- [23] Oleinik O.A. and Radkevič E.V.: On the analyticity of solutions for linear partial differential equations, *Mat. Sb.* 90 = *Math. U.S.S.R. Sb.* 19(1973) 581-596.
- [24] Parenti C. and Rodino L.: Examples of hypoelliptic operators which are not micro-hypoelliptic, *Bolletino U.M.I.* 17-B(1980) 390-409.
- [25] Rothschild L.P. and Stein E.M.: Hypoelliptic differential operators and nilpotent groups, *Acta Math.* 137(1977) 247-320.
- [26] Sakurai T.: Tempered second wave front sets and quasi-homogeneous partial differential operators, *Proc. Conf. on spectral and scattering theory for differential operators*, 1986.
- [27] ————— : Gevrey hypoellipticity for operators with non symplectic characteristics, to appear in *Saitama Math. J.*

- 5(1987) 7-25.
- [28] Sjöstrand J.: Parametrices for pseudo-differential operators with multiple characteristics, Ark. Mat. 12(1974) 85-130.
- [29] ————— : Singularites analytiques microlocales, Astérisque 95(1982) 1-166.
- [30] ————— : Analytic wave front sets and operators with multiple characteristics, Hokkaido Math. J. XII(1983) 392-433.