

Topics on Hilbert's 16th problem

by Goo ISHIKAWA (Faculty of Sci., Hokkaido Univ.; Japan)

石川 剛郎 北大・理

Introduction. The first half of Hilbert's 16th problem is concerned with the relative position of a non-singular real algebraic curve (resp. surface) in  $\mathbb{R}P^2$  (resp.  $\mathbb{R}P^3$ ) of a fixed degree. In fact, Hilbert asked about the topology of sixth degree curves and fourth degree surfaces. Though this concrete problem was solved completely by several mathematicians in the last decade, it is still stimulating the investigation of "topology of real algebraic manifolds" or "enumerative real algebraic geometry" from the viewpoint of contemporary mathematics.

Let  $H_{n,d}$  denote the  $\mathbb{R}$ -vector space of homogeneous polynomials  $F(x_0, x_1, \dots, x_n)$  of degree  $d$  with real coefficients. By a real hypersurface, we mean an element of  $P(H_{n,d}) = (H_{n,d} - 0)/\mathbb{R}^\times$ . Let  $K$  denotes  $\mathbb{R}$  or  $\mathbb{C}$ . For  $[F] \in P(H_{n,d})$ , set  $KZ_F = \{[x] \in KP^n \mid F(x) = 0\}$ . We call  $[F]$  non-singular if  $CZ_F$  is non-singular.

Two non-singular real hypersurfaces  $[F], [G] \in P(H_{n,d})$  are called isotopic if there exists a continuous family  $\sigma_t$  of homeomorphisms of  $\mathbb{R}P^n$  such that  $\sigma_0 = \text{identity}$  and  $\sigma_1(RZ_F) = RZ_G$ . They are called rigidly isotopic if there exists a continuous family  $[F]_t \in P(H_{n,d})$  of non-singular real hypersurfaces such that  $[F]_0 = [F]$  and  $[F]_1 = [G]$ . We see that, if  $[F]$  and  $[G]$  are rigidly isotopic, then they are isotopic.

# 10

The following tables illustrate the known isotopy and rigid isotopy classifications for curves: (Notations will be explained)

d Rigid isotopy classification for curves of degree  $\leq 6$

1  $\langle J \rangle_I$

2  $\langle 1 \rangle_I, \langle 0 \rangle_{II}$

3  $\langle J, 1^- \rangle_I, \langle J \rangle_{II}$

4  $\langle 4 \rangle_I, \langle 3 \rangle_{II}, \langle 1 \langle 1^- \rangle \rangle_I, \langle 2 \rangle_{II}, \langle 1 \rangle_{II}, \langle 0 \rangle_{II}$

5  $\langle J, 3^+, 3^- \rangle_I, \langle J, 5 \rangle_{II}, \langle J, 1^+, 3^- \rangle_I, \langle J, 4 \rangle_{II}, \langle J, 3 \rangle_{II},$

$\langle J, 1^- \langle 1^- \rangle \rangle_I, \langle J, 2 \rangle_{II}, \langle J, 1 \rangle_{II}, \langle J \rangle_{II}$

6 Isotopy classification

(i)  $\langle 1, 1 \langle 9 \rangle \rangle_I, \langle 5, 1 \langle 5 \rangle \rangle_I, \langle 9, 1 \langle 1 \rangle \rangle_I$

(ii)  $\langle 1 \langle 9 \rangle \rangle, \langle 1, 1 \langle 8 \rangle \rangle, \langle 4, 1 \langle 5 \rangle \rangle, \langle 5, 1 \langle 4 \rangle \rangle, \langle 8, 1 \langle 1 \rangle \rangle, \langle 10 \rangle$

(iii)  $\langle a, 1 \langle b \rangle \rangle, (0 \leq a \leq 7, 1 \leq b \leq 8, a + b \leq 8)$

(iv)  $\langle a \rangle, (0 \leq a \leq 9)$  (v)  $\langle 1 \langle 1 \langle 1 \rangle \rangle \rangle_I:$

The eight classes  $\langle 9 \rangle, \langle 4, 1 \langle 4 \rangle \rangle, \langle 1 \langle 8 \rangle \rangle, \langle 5, 1 \langle 1 \rangle \rangle, \langle 3, 1 \langle 3 \rangle \rangle, \langle 1, 1 \langle 5 \rangle \rangle, \langle 2, 1 \langle 2 \rangle \rangle,$  and  $\langle 1 \langle 4 \rangle \rangle$  correspond to exactly two rigid isotopy classes respectively.

Isotopy classification for curves of degree 7

7 (i)  $\langle J, a, 1 \langle b \rangle \rangle, (0 \leq a \leq 13, 1 \leq b \leq 13, a + b \leq 14)$

(ii)  $\langle J, a \rangle, (0 \leq a \leq 15)$  (iii)  $\langle J, \langle 1 \langle 1 \langle 1 \rangle \rangle \rangle$

In general, to obtain the concrete classifications, we need two types of results: (A) Restriction theorems (Estimates), (B) Existence theorems (Explicit constructions and Abstract existence theorems).

In this talk, after mentioning some restriction theorems, we

intend to review the following two important topics related to (B):

(I) Local and global bifurcations of singular real plane curves.

(II) Real structures on polarized K-3 surfaces.

1) Let  $C \subset \mathbb{R}P^2$  be a non-singular plane curve of degree  $d$ .

Denote by  $\ell$  the number of connected components of  $C$ . By

Harnak(1875),  $\ell \leq (1/2)(d-1)(d-2) + 1 (= g + 1)$ . Following

Prtrovskii, we call  $C$  is an M-curve if  $\ell = g + 1$ .

2)The image of an embedding  $i : S^1 \rightarrow \mathbb{R}P^2$  is called an oval

(resp. a pseudo-line) if  $i$  is null-homotopic (resp.

non null-homotopic). An oval divides  $\mathbb{R}P^2$  to a region

diffeomorphic to the open disk, which is called the interior, and to

a region diffeomorphic to the open Möbius band, which is called the

exterior.

3) According to Viro, we symbolize isotopy types of non-singular

real plane curves as follows:

The symbol  $\langle a \rangle$  means (the isotopy type) of union of  $a$ -ovals


outside in each other.  $\langle J \rangle$  means the pseudo-line. For a symbol

$\langle A \rangle$ ,  $\langle 1 \langle A \rangle \rangle$  means the union of  $\langle A \rangle$  and an oval surrounding  $\langle A \rangle$ .

For a sequence of symbols  $\langle A_1 \rangle, \langle A_2 \rangle, \dots, \langle A_r \rangle$ , by  $\langle A_1, A_2, \dots, A_r \rangle$ ,

we mean union of them lying outside in each other.

For example,  $\langle J, 13, 1 \langle 1 \rangle \rangle$  means the isotopy type of the curve



We see that a symbol determines the isotopy type of a non-singular plane curve.

4) An oval  $E$  of a curve  $C$  is called even (resp. odd) if  $E$  is included in the even (resp. odd) number of other ovals of  $C$ .

Set  $p = \#(\text{even ovals})$  and  $n = \#(\text{odd ovals})$ .

**Theorem** (Rokhlin 1972). For an  $M$ -curve of even degree  $d$ ,

$$p - n \equiv (d/2)^2, \pmod{8}.$$

Example ( $M$ -curves of degree 6). Let  $d = 6$  and  $l = 11$ . Then we have  $p + n = 11$  and  $p - n \equiv 1 \pmod{8}$ . By Bézout's theorem, the possible isotopy types are  $\langle \alpha, 1 \langle \beta \rangle \rangle$  with  $\alpha, \beta$  are non-negative integers. Then we have  $p = \alpha + 1$  and  $n = \beta$ . Thus  $\alpha + \beta = 10$  and  $\alpha - \beta \equiv 0 \pmod{8}$ . Therefore  $(\alpha, \beta) = (1, 9), (5, 5), (9, 1)$ . The possible isotopy types of  $M$ -curves of degree 6 are

$$\langle 1, 1 \langle 9 \rangle \rangle, \langle 5, 1 \langle 5 \rangle \rangle \text{ and } \langle 9, 1 \langle 1 \rangle \rangle.$$

In fact these are realized by non-singular curves of degree 6.

5) The isotopy classification of curves of degree  $d \leq 5$  is classically known. The classification for  $d = 6$  and  $d = 7$  are established by Gudkov(1971) and by Viro(1980) respectively.

Consider on  $M$ -curves of degree 8;  $d = 8$  and  $l = 22$ . Then  $p+n = 22$ ,  $p-n \equiv 0 \pmod{8}$ . Thus  $(p, n) = (3, 19), (7, 15), (11, 11), (15, 7), (19, 3)$ . By Bézout's theorem, the possible isotopy types

are  $\langle \alpha, 1 \langle \beta \rangle \rangle$ ,  $\langle \alpha, 1 \langle \beta \rangle, 1 \langle \gamma \rangle \rangle$ ,  $\langle \alpha, 1 \langle \beta \rangle, 1 \langle \gamma \rangle, 1 \langle \delta \rangle \rangle$ , and  $\langle \alpha, 1 \langle \beta, 1 \langle \gamma \rangle \rangle \rangle$ .

**Theorem (Viro 1980).** For an M-curve of degree 8 of type  $\langle \alpha, 1 \langle \beta \rangle, 1 \langle \gamma \rangle, 1 \langle \delta \rangle \rangle$ , if  $\beta\gamma\delta \neq 0$ , then  $\beta$ ,  $\gamma$  and  $\delta$  are all odd.

Nowaday, we have the following table:

d	g+1	$\nu$
2	1	1
4	4	1
6	11	3
8	22	$58 \leq ? \leq 104$

Here,  $\nu = \#(\text{isotopy classes of M-curves of degree } d)$ .

**Conjecture (Viro).** For an M-curves of degree 8, if  $\gamma \neq 0$ , then  $\beta$  and  $\gamma$  are odd.

If Conjecture is true, then we have  $\nu \leq 79$  for  $d = 8$ .

**Problem.** Classify M-cueves of degree 8.

For M-curves of degree 10, there are results of Chislenko and Shustin. For curves in  $\mathbb{RP}^1 \times \mathbb{RP}^1$ , the are results of Hilbert, Gudkov and Matsuoka.

6) Rigid isotopy. Let  $\text{PH}_{n,d}$  be as in introduction. Denote

by  $\Sigma$  the set of singular hypersurfaces in  $\mathbb{P}H_{n,d}$ . Then we have the natural bijection between the set of rigid isotopy classes of non-singular hypersurfaces of degree  $d$  in  $\mathbb{R}P^n$  and  $\pi_0(\mathbb{P}H_{n,d} - \Sigma)$ .


7) Plane curve ( $n = 2$ ). Let  $[F] \in \mathbb{P}H_{2,d}$ . Consider the involution on the Riemann surface  $\mathbb{C}Z_F$  defined by the complex conjugation. The fixed point set is the real locus  $\mathbb{R}Z_F$ .

According to Klein,  $[F]$  is called of type I (resp. II) if  $\mathbb{C}Z_F - \mathbb{R}Z_F$  is not connected (resp. connected).

If  $[F]$  is of type I, the orientation of  $\mathbb{C}Z_F$  induces an orientation, so called the complex orientation, of  $\mathbb{R}Z_F$  (up to simultaneous reversing).

The type and the complex orientation in the case of type I are rigid isotopy invariants.

For example,  $\langle J, 1^+, 3^- \rangle_I$  means a rigid isotopy class of type I with the following complex orientation:



There exist the rigid isotopy classifications for  $d \leq 6$ .


8) Construction. One of methods to construct non-singular curves in prescribed isotopy classes is perturbation of singular

curves: 

Singular curves $\xrightarrow{\text{perturb}}$ Non-singular curves
--

In the classical method of Harnack, Brusotti and so on, only


curves with  $A_1$ -type singularities are considered:



Viro treats the perturbation of  $J_{10}$ -singularity defined by locally  $f(x,y) = (y - \alpha_1 x^2)(y - \alpha_2 x^2)(y - \alpha_3 x^2) = 0$ .

**Theorem (Viro 1982).** There exist exactly 31-classes of perturbations of  $J_{10}$ . (See [1986, Viro]).

By this and the gluing method of Viro, we can construct an M-curve of degree 6 of type  $\langle 5, 1 \langle 5 \rangle \rangle$ :



**Problem.** Let  $F : \mathbb{R}^2 \times \mathbb{R}^{10}, 0 \longrightarrow \mathbb{R}, 0$  be the versal deformation of  $J_{10}$ -singularity and  $\Sigma \subset \mathbb{R}^{10}$  be the bifurcation locus. Then  $\#(\text{local connected components of } \mathbb{R}^{10} - \Sigma \text{ at } 0) = 31?$

9) Space surface ( $n = 3$ ). The rigid isotopy classifications of space surfaces of degree  $\leq 2$  is classically well-known and that of space surfaces of degree 3 is done by Klein. For example, the number of rigid isotopy classes is equals to 1, 3, 5 in the case  $d =$

# 16

1, 2, 3 respectively.

Now consider the rigid isotopy classification problem of space surfaces of degree 4.

Set  $S = \mathbb{C}Z_F \subset \mathbb{C}P^3$ , for  $[F] \in \text{PH}_{3,4} - \Sigma$ . Then the first betti number  $b_1(S) = 0$  and  $\Lambda^2 T^*S$  is trivial, that is,  $S$  is a K-3 surface. As is well-known,  $L = H^2(S, \mathbb{Z})$  is a free  $\mathbb{Z}$ -module of rank 22. With respect to the cup-product, the signature of  $L$  is

$(3, 19)$ , and  $L \cong \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^{\oplus 3} \oplus (-E_8)^{\oplus 2}$ . Denote by  $\tau : S \rightarrow S$  the complex conjugation and set  $\varphi = \tau^* : L \rightarrow L$ . Let  $h \in L$  be the Poincaré dual of the hyperplane section. Then we have  $\varphi(h) = -h$ .

Consider the triple  $(L, \varphi, h)$ . Set  $L^\varphi = \{x \in L \mid \varphi(x) = x\}$  and denote by  $(t_+, t_-)$  the signature of  $L^\varphi$ . Then we have  $t_+ = 1$  and the Euler characteristic  $\chi(RS) = 2t_- - 18$ .

For the natural action of  $\text{PGL}(4)$  on  $\text{PH}_{3,4} - \Sigma$ , we have

Theorem (Nikulin 1979). The construction in above induces a bijection

$$\pi_0((\text{PH}_{3,4} - \Sigma)/\text{PGL}(4)) \longleftrightarrow \left\{ (L, \varphi, h) \left| \begin{array}{l} L \text{ is free of sign } (3, 19), \\ \varphi \text{ is an involution of } L, \\ h^2 = 4, h \text{ is primitive,} \\ \varphi(h) = -h \text{ and } t_+ = 1 \end{array} \right. \right\} / \text{isom..}$$

The right hand side has 134 elements. 72 of them correspond to surfaces with null-homotopic real part  $RS \subset \mathbb{R}P^3$ .

Since  $\pi_0(\text{PGL}(4)) = 2$ , the natural mapping

$$\pi : \pi_0(\text{PH}_{3,4} - \Sigma) \longrightarrow \pi_0((\text{PH}_{3,4} - \Sigma)/\text{PGL}(4))$$

is 2:1 or 1:1 over each point of the target.



Let  $r : \mathbb{R}P^3 \longrightarrow \mathbb{R}P^3$  be the reflection along a plane in  $\mathbb{R}P^3$ , for instance,  $r[x_0, x_1, x_2, x_3] = [-x_0, x_1, x_2, x_3]$ .

We call a rigid isotopy class  $[F]$  is amphicheiral if  $[F \circ r]$  is rigidly isotopic to  $[F]$ .

Then we see that  $\pi$  is one to one over  $\pi[F]$  if and only if  $[F]$  is amphicheiral.

The next result is the criterion of amphicheirality described by only on real loci  $RS$ .

**Theorem** (Kharlamov 1984). Denote by  $p$  the number of spherical components of  $RS$ , and by  $q$  the half of dimension of  $H_1(RS, \mathbb{R})$ . Consider the following four cases:

- (a)  $RS$  is null-homotopic and  $p \geq 3$ ,  $q \geq 3$ ,
- (b)  $RS$  is null-homotopic and  $p < 3$  or  $q < 3$ ,
- (c)  $RS$  is non null-homotopic and  $p \geq 4$ ,
- (d)  $RS$  is non null homotopic and  $p < 4$ .

If (a) or (c), then  $[F]$  is not amphicheiral. If (b) or (d), then  $[F]$  is amphicheiral.

As a result, there exist exactly 170 rigid isotopy classes of non-singular surfaces in  $\mathbb{R}P^3$  of degree 4;

$$\#(\pi_0(\text{PH}_{3,4} - \Sigma)) = 170.$$

### Bibliography on classical theory of real algebraic varieties

This is a collection of papers concerning to the first half of

## 18

the Hilbert's 16th problem, complex manifolds with conjugations, singularities defined over the real and so on.

Of course, there may be many important papers which do not appear in this bibliography.

### 1875

A. Harnack, Über die Vieltheiligkeit der ebenen algebraischen Curven, Math. Ann., 10, 189-199.

### 1891

D. Hilbert, Über die reellen Züge algebraischen Curven, Math. Ann., 38, 115-138.

### 1906

V. Ragsdale, On the arrangement of the real branches of plane algebraic curves, Amer. J. Math., 28, 377-404.

### 1913

K. Rohn, Die Maximalzahl und Anordnung der Ovale bei der ebenen Kurve 6. Ordnung und bei der Fläche 4. Ordnung, Math. Ann., 73, 177-229.

### 1931

A. Comessatti, Reelle Fragen in der algebraischen Geometrie, Jb Deutsche Math. Ver., 41, 107-134.

R. Courant, D. Hilbert, Methoden der mathematischen Physik, I, Springer-Verlag, Berlin.

### 1932

A. Comessatti, Sulla connessione e sui numeri base delle superficie algebrische reali, Rend. Semin. Mat. Univ. Padova, 3, 141-162.

1933

I.G. Petrovsky, Sur la topologie des courbes réelle et algébriques, C. R. Acad. Sci. Paris, , 1270-1271.

1938

I.G. Petrovskii, On the topology of real plane algebraic curves, Ann. of Math., (2) 39, 189-209.

1939

L. Bieberbach, Über die reellen Züge der algebraischen Gegilde, Deutsche Math., 4, 348-360.

1944

M.F. Châtelet, Variations sur un thème de H.Poincaré, Ann. Ec. Norm. Sup., 61, 249-300.

1949

O.A. Oleinik, Some estimates for Betti numbers of real algebraic surfaces, Dokl. Akad. Nauk SSSR, 67, 425-426.

O.A. Oleinik, I.G. Petrovskii, On the topology of real algebraic surfaces, Dokl. Akad. Nauk SSSR, 67, 31-32.

\_\_\_\_\_ , On the topology of real algebraic surfaces, Izv. Akad. Nauk SSSR. Ser. Mat., 13, 389-402. = Amer. Math. Soc. Transl. 7(1952), 399-417.

I.G. Petrovskii, On the topological properties of algebraic lines and surfaces, Vestnik Moskov. Gos. Univ., 11, 23-27.

1950

O.A. Oleinik, On the topology of real algebraic space curves, Dokl. Akad. Nauk SSSR., 70, 13-14.

1951

O.A. Oleinik, On algebraic curves on an algebraic surface, Usp.

Mat. Nauk, 6-4, 209-210.

\_\_\_\_\_, On the topology of real algebraic curves on an algebraic surface, Mat. Sb., 29, 133-156.

\_\_\_\_\_, Estimates of the Betti numbers of real algebraic hypersurfaces, Mat. Sb., 28, 635-640.

O.A. Oleinik, I.G. Petrovskii, On the topology of real algebraic surfaces, Usp. Mat. Nauk, 6-4, 208-209.

#### 1964

J. Milnor, On the betti number of real varieties, Proc. Amer. Math. Soc., 15, 275-280.

#### 1965

R. Thom, Sur l'homologie des varietes algebriques reelles, Differential and combinational topology (A symposium in honor of M.Morse), Princeton Univ. Press, N.J., 255-265.

#### 1966

D.A. Gudkov, G.A. Utkin, M.A. Tai, A complete classification of irreducible curves of the fourth order, Mat. Sb., 69, 222-256.

#### 1968

W. Nuij, A note on hyperbolic polynomials, Math. Scand., 23, 69-72.

#### 1969

D.A. Gudkov, G.A. Utkin, The topology of curves of the sixth order and surfaces of the fourth order, Gor'kov. Gos. Univ. Uchen. Zap., 87, 3-213 = Nine papers on Hilbert's 16th problem, Amer. Math. Soc. Transl., (2) 112(1978).

#### 1971

N.L. Alling, N. Greenleaf, Foundation of the theory of Klein

surfaces, Lect. Notes in Math., 219, Springer.

V.I. Arnol'd, The arrangement of the ovals of real plane algebraic curves, involutions of four-dimensional smooth manifolds, and the arithmetic of integral quadratic forms, Funkt. Anal. Prilozhen., 5-3, 1-9. = Funct. Anal. Appl., 5, 169-176.

### 1972

V.M. Kharlamov, Maximal number of components of the fourth degree in  $\mathbb{R}P^3$ , Funkt. Anal. Prilozhen., 6-4, 101. = Funct. Anal. Appl., 6, 345-346.

V.A. Rokhlin, Proof of Gudkov conjecture, Funkt. Anal. Prilozhen., 6-2, 62-64. = Funct. Anal. Appl., 6, 62-64.

V.A. Rokhlin, Congruences modulo 16 in Hilbert's sixteenth problem I, Funkt. Anal. Prilozhen., 6-4, 58-64. = Funct. Anal. Appl., 6, 301-306.

### 1973

V.I. Arnol'd, The topology of real algebraic curves (the works of Petrovskii and their development), Usp. Mat. Nauk, 28-5, 260-262.

D.A. Gudkov, Construction of a curve of the sixth order of type  $\frac{5}{1}5$ , Izv. Vyssh. Uchebn. Zaved. Mat., 3 (130), 28-36.

D.A. Gudkov, A.D. Krakhnov, Periodicity of the Euler characteristic of real algebraic  $(M-1)$ -varieties, Funkt. Anal. Prilozhen., 7-2, 15-19. = Funct. Anal. Appl., 7, 98-102.

V.M. Kharlamov, New congruences for the Euler characteristic of real algebraic varieties, Funkt. Anal. Prilozhen., 7-2, 74-78. = Funct. Anal. Appl., 7, 147-150.

W. Massey, The quotient space of the complex projective plane under conjugation is a 4 sphere, Geom. Dedicata 2., 371-374.

V.A. Rokhlin, Congruences modulo 16 in Hilbert's sixteenth problem II, *Funkt. Anal. Prilozhen.*, 7-2, 91-92. = *Funct. Anal. Appl.*, 7, 163-164.

## 1974

D.A. Gudkov, The topology of real projective algebraic varieties, *Usp. Mat. Nauk*, 29-4, 3-79. = *Russ. Math. Surveys*, 29-4, 1-79.

V.M. Kharlamov, The generalized Petrovskii inequality, *Funkt. Anal. Prilozhen.*, 8-2, 50-56. = *Funct. Anal. Appl.*, 8, 132-137.

V.A. Rokhlin, Complex orientations of real algebraic curves, *Funkt. Anal. Prilozhen.*, 8-4, 71-75. = *Funct. Anal. Appl.*, 8, 331-334

## 1975

V.M. Kharlamov, Additional congruences for the Euler characteristic of real algebraic manifolds of even dimensions, *Funkt. Anal. Prilozhen.*, 9-2, 51-60. = *Funct. Anal. Appl.*, 9, 134-141.

\_\_\_\_\_ , A generalized Petrovskii inequality II, *Funkt. Anal. Prilozhen.*, 9-3, 93-94. = *Funct. Anal. Appl.*, 9, 266-268.

N.M. Mishachev, Complex orientations of plane M-curves of odd degree, *Funkt. Anal. Prilozhen.*, 9-4, 77-78. = *Funct. Anal. Appl.*, 9, 342-343.

S.M. Natanzon, Moduli of real algebraic curves, *Usp. Mat. Nauk* 30-1, 251-252.

G. Shimura, On the real points of an arithmetic quotient of a bounded symmetric domain, *Math. Ann.*, 215, 135-164.

V.I. Zvonilov, Kharlamov's inequalities and Petrovskii-Oleinik inequalities, *Funkt. Anal. Prilozhen.*, 9-2, 69-70. = *Funct. Anal. Appl.*, 9, 152-153.

## 1976

A.L. Ceponkus, On nests of real plane algebraic curves, Litov.

Mat. Sbornik, 16-4, 239-243.

V.M. Kharlomov, The topological type of nonsingular surfaces in

$\mathbb{R}P^3$  of degree four, Funkt. Anal. Prilozhen., 10-4, 55-68. = Funct.

Anal. Appl., 10, 295-305.

V.A. Krasnov, A generalized Petrovskii inequality for

odd-degree curves, Funkt. Anal. Prilozhen., 10-2, 41-48. = Funct.

Anal. Appl., 10, 116-121.

## 1977

V.I. Arnol'd, Modern development of the work of I.G. Petrovskii

in the topology of real algebraic manifolds, Usp. Mat. Nauk, 32-3,

215-216.

G.M. Polotovskii, A catalogue of M-decomposing curves of sixth

order, Dokl. Akad. Nauk SSSR, 236-3, 548-551. = Soviet Math. Dokl.,

18-5, 1241-1245. (See also Correction, Dokl. Akad. Nauk SSSR, 240,

(1978), 1272.)

## 1978

N. A'campo, Sur la première partie du seizième problème de

Hilbert, Seminaire Bourbaki, 31<sup>e</sup> année, 1978/79, n° 537. =

Lecture Notes in Math., 770, Springer(1980), pp. 208-227.

V.I. Arnol'd, Index of a singular point of a vector field, the

Petrovskii-Oleinik inequality, and mixed Hodge structures, Funkt.

Anal. Prilozhen., 12-1, 1-14. = Funct. Anal. Appl., 12, 1-12.

M.R. Gabel, Algebraic vector fields on  $S^2$ , J. Pure Appl.

Algebra 11(1977), 29-40.

L.V. Golubina, K.K. Tai, in [LA,1978], pp. 130-136.

V.M. Kharlamov, Isotopy types of nonsingular surfaces of fourth degree in  $\mathbb{R}P^3$ , Funkt. Anal. Prilozhen., 12-1, 86-87. = Funct. Anal. Appl., 12, 68-69.

\_\_\_\_\_, Real algebraic surfaces, Proc. Internat. Congr. Math., (Helsinki, 1978), I, 421-428. = Amer. Math. Soc. Transl., (2)-117(1981), 15-21.

A.B. Korchagin, New possibility on the method of Brusotti for the construction of  $M$ -curves of an order  $\geq 8$ , in [LA,1978], pp. 149-159.

R.S. Kulkarni, P.B. Shalen, A note on real cubic surfaces, Bull. Calcutta Math. Soc., 70, 375-377.

[LA]: E.A. Leontovich-Andronova, ed., Methods of qualitative theory of differential equations, Interuniv. Collect. Gor'ki.

E. Looijenga, The discriminant of a real simple singularity, Composit. Math., 37-1, 51-62.

S.M. Natanzon, On the order of a finite group of homeomorphisms of a surface onto itself and the number of real forms of a complex algebraic curve, Dokl. Akad. Nauk SSSR, 242, 765-768. = Sov. Math. Dokl., 19, 1195-1199.

\_\_\_\_\_, Automorphism of the Riemann surface of an  $M$ -curve, Funkt. Anal. Prilozhen., 12-3, 82-83. = Funct. Anal. Appl., 12-3, 228-229.

G.M. Polotovskii,  $(M-1)$  and  $(M-2)$ -ramified curves of sixth order, in [LA,1978], pp. 130-148.

V.A. Rokhlin, Complex topological characteristics of real algebraic curves, Usp. Math. Nauk, 33-5, 77-89. = Russ. Math. Surveys, 33-5, 85-98.



R. Silhol, Géométrie algébrique sur un corps non algébriquement clos, *Comm. Algebra*, 6-11, 1131-1155.

O.Ya. Viro, Generalized Petrovskii and Arnol'd inequalities and curves with singularities, *Usp. Mat. Nauk*, 33-3, 145-146.

G. Wilson, Hilbert's sixteenth problem, *Topology*, 17, 53-73.

### 1979

V.I. Arnol'd, O.A. Oleinik, Topology of real algebraic manifolds, *Vestnik Moskov. Univ. Mat.*, 34-6, 7-17. = *Moskow Univ. Math. Bull.*, 34, 5-17.

E.S. Berezovskaya, Index of a stationary point of a vector field on the plane, *Funkt. Anal. Prilozhen.*, 13-2, 77-77. = *Funct. Anal. Appl.*, 13-2, 137-137.

D.A. Gudkov, On the topology of algebraic curves on a hyperboloid, *Usp. Mat. Nauk*, 34-6, 26-32. = *Russ. Math. Surveys*, 34-6, 27-35.

A.G. Khovanskii, Index of a polynomial field, *Funkt. Anal. Prilozhen.*, 13-1, 49-58. = *Funct. Anal. Appl.*, 13, 38-45.

S.M. Natanzon, Automorphisms and real forms of a class of complex algebraic curves, *Funkt. Anal. Prilozhen.*, 13-2, 89-90. = *Funct. Anal. Appl.*, 13, 148-150.

V.V. Nikulin, Integral symmetric bilinear forms and some of their applications, *Izv. Akad. Nauk SSSR Ser. Mat.*, 43-1, 111-177. = *Math. USSR Izv.*, 14-1, 103-167.

T.M. Rassais, On Hilbert's sixteenth problem, *C. R. Math. Rep. Acad. Sci. Canada*, 1, 203-205.

O. Ya. Viro, Construction of M-surfaces, *Funkt. Anal. Prilozhen.*, 13-3, 71-72. = *Funct. Anal. Appl.*, 13, 212-213.

\_\_\_\_\_ , Construction of multicomponent real algebraic surfaces, Dokl. Akad. Nauk SSSR, 248-2, 279-282. = Soviet Math. Dokl. 20-5, 991-995.

V.I. Zvonilov, Refinement of the Petrovskii-Arnol'd inequality for curves of odd degree, Funkt. Anal. Prilozhen., 13-4, 31-39. = Funct. Anal. Appl., 13, 262-268.

### 1980

B. Chevalier, Sur une manière de construire des courbes réelles dans  $P^2(\mathbb{R})$ , in [R,1980], pp. 27-50.

D.A. Gudkov, Generalization of a theorem of Brusotti for curves on a surface of second order, Funkt. Anal. Prilozhen., 14-1, 20-24. = Funct. Anal. Appl., 14, 15-18.

H. Jaffee, Real algebraic curves, Topology, 19, 81-87.

A. Marin, Quelques remarques sur les courbes algébriques planes réelles, in [R,1980], pp. 51-68.

S.M. Natanzon, On the sum-number of ovals of four complex isomorphic real algebraic curves, Usp. Mat. Nauk, 35-4, 184.

S. Ochanine, Sur l'utilisation de la théorie des groupes formels dans le 16<sup>ème</sup> problème de Hilbert, in [R,1980], pp. 81-100.

[R]: J.J. Risler, ed., Séminaire Sur La Géométrie Algébrique Réelle, Publications mathématiques de l'Université Paris VII, No.9.

J.J. Risler, Sur le 16<sup>ème</sup> problème de Hilbert: Un résumé et quelques questions, in [R,1980], pp. 11-25.

V.A. Rokhlin, New inequalities in the topology of real planer algebraic curves, Funkt. Anal. Prilozhen., 14-1, 37-43. = Funct. Anal. Appl., 14, 29-33.

O.Ya. Viro, Curves of degree 7, curves of degree 8 and the

Ragsdale conjecture, Dokl. Akad. Nauk SSSR, 254-6, 1306-1310. =  
Soviet. Math. Dokl., 22-2, 566-570.

C.T.C. Wall, Affine cubic functions III, The real place, Math.  
Proc. Camb. Phil. Soc., 87-1, 1-14.

### 1981

T. Abiko, Classification of cubic forms with three variables,  
Hokkaido Math. J., 10, 239-248.

N.L. Alling, Real Elliptic Curves, Math. Studies 54,  
North-Holland.

D.S. Aron, Automatic analysis of real algebraic curves, SIGSAM  
Bull., 15-4, 3-9.

B. Berron, Le naeud <<huit>> est algébrique réel, C. R. Acad.  
Sc. Paris, 292, 679-682.

J.L. Colliot-Thélène, F. Ischeback, L'équivalence rationnelle  
sur les cycles de dimension zéro des variétés algébriques réelles, C.  
R. Acad. Sc. Paris, 292-15, 723-725.

B.H. Gross, Topological invariants of real algebraic curves,  
Sémin. Théor. Nombres, Univ. Grenoble I, 1980-1981, Exposé 2.

B.H. Gross, J. Harris, Real algebraic curves, Ann. Sci. Éc.  
Norm. Sup., (4), 14, 157-182.

V.M. Karpushkin, The number of components of the complement to  
the level hypersurface of a harmonic polynomial, Vestnik Moscow Univ.  
Mat., 34-6, 3-4.

V.M. Kharlamov, Rigid isotopic classification of real planer  
curves of degree 5, Funkt. Anal. Prilozhen., 15-1, 88-89. = Funct.  
Anal. Appl., 15, 73-74.

\_\_\_\_\_ , Real algebraic surfaces, Amer. Math. Soc.

Transl., (2), 117, 15-21.

J.S. Milne, K.-y. Shih, The action of complex conjugation on a Shimura variety, *Ann. Math.*, (II)113, 569-599.

G.M. Polotovskii, *Usp. Mat. Nauk.*, 36-4, 235-236.

### 1982

Yu.S. Chislenko, M-curves of degree 10, *Zap. Nauchn. Sem. LOMI* 122, 146-161. = *J. Soviet Math.*, 26, 1689-1699.

T. Fidler, Pencils of lines and the topology of real algebraic curves, *Izv. Akad. Nauk SSSR Ser. Mat.*, 46-4, 853-863. = *Math. USSR Izv.*, 21-1 (1983), 161-170.

S.M. Finashin, The topology of the complement of a real algebraic curve in  $CP^2$ , *Zap. Nauchn. Sem. LOMI*, 122, 137-145. = *J. Soviet Math.*, 26(1984), 1684-1689.

P. Gianni, Sulle disequaglianze di Arnold-Wilson per le curve algebriche reali, *Ann. Univ. Ferrara, Nuova Ser., Sez. VII* 28, 115-126

L.V. Golubina, D.A. Gudkov, Classification of sets of singular points of unicursal plane fifth-order curves, *Differential and Integral Equations* 6, Gor'kov Gos. Univ., Gorki, pp. 126-132.

D.A. Gudkov, A curve of fifth order with five cusps, *Funk. Ana Prilozhen.*, 16-3, 54-55. = *Funct. Anal. Appl.*, 16, 201-202.

V.A. Krasnov, Albanese mapping for real algebraic varieties, *Mat. Zemetki* 32, 365-374. = *Math. Notes Acad. Sci. SSSR.*, 32-3, 661-666.

V.A. Lipnitskii, Norms in the fields of real algebraic functions and the reduced Whitehead group, *Dokl. Akad. Nauk SSSR*, 26 585-588.

M. Seppälä, Real structure of Teichmüller space, *Manuscripta*

Math., 40, 79-86.

R. Silhol, Real abelian varieties and the theory of Comessatti,  
Math. Z., 181, 345-364.

G. Gonzalez-Sprinberg, Nodes of the Laplacian and topology of  
real algebraic varieties, Topology Proc. Spec. Semin. México, 4,  
99-115.

V.I. Zvonilov, Complex topological characteristics of real  
algebraic curves on surfaces, Funkt. Anal. Prilozhen., 16-3, 56-57. =  
Funct. Anal. Appl., 16, 202-204.

O.Ya Viro, Complex orientations of real algebraic surfaces, Usp.  
Mat. Nauk 37-4, 93-93.

### 1983

G. Faltings, Real projective structures on Riemann surfaces,  
Composit. Math., 48-2, 223-269.

T. Fidler, New congruences in the topology of real plane  
algebraic curves, Dokl. Akad. Nauk SSSR, 270-1. = Soviet Math. Dokl.,  
27-3, 566-568.

G. Ishikawa, On degrees of non-roughness of real projective  
varieties, Proc. Japan Acad., 59-6, 277-280.

V.A. Krasnov, Harnack-Thom inequalities for mappings of real  
algebraic varieties, Izv. Acad. Nauk SSSR, Ser. Mat., 47-2. = Math.  
USSR Izv., 22-2, 1984.

V.V. Nikulin, Involutions of integral quadratic forms and their  
applications to real algebraic geometry, Izv. Acad. Nauk SSSR, 47,  
109-158. = Math. USSR Izv. 22-1(1984), 99-172.

A.P. Ogg, Real points on Shimura curves, Arithmetic and  
Geometry, Prog. Math., 35, Birkhäuser, pp. 277-307.

J.J. Risler, On the Bézout theorem in the real case, Monograf. Memor. Mat. 32, Publ. Inst. J. Juan, Madrid.

M. Reggero, On linear systems and the divisor class group of a real variety, Ann. Mat. Pura Appl., IV 135, 349-362.

R. Silhol, Diviseurs sur les varietes algebriques réelles, Boll. U. M. I..

O.Ya. Viro, Real plane curves of degree 7 and 8: New prohibitions, Izv. Akad. Nauk SSSR Ser. Mat., 47-5, 1135-1150. = Math. USSR Izv. 23-2(1984), 409-422.

\_\_\_\_\_ , Progress over the last five years in the topology of real algebraic varieties, Proc. International Conference of Mathematicians, Warsaw, 525-611.

\_\_\_\_\_ , Gluing algebraic hypersurfaces, removing of singularities and constructions of curves, Topology conference, Proc. Collect. Rep., Leningrad 1982, pp.149-197.

C.T.C. Wall, Pencils of real binary cubics, Math. Proc. camb. Phil. soc., 93, 477-484.

#### 1984

W.A. Adkins, A Harnack estimate for real normal surface singularities, Pacific J. Math., 114-2, 257-265.

C. Andradas, J.M. Gamgoa, On projections of real algebraic varieties, C. R. Math. Rep. Acad. Sci. Canada, 6-2, 49-54.

D.S. Arnon, S.A. McCallum, A polynomial-time algorithm for the topological type of real algebraic curves - extended abstract, Rocky Mt. J. Math., 14, 847-850.

L. Beretta, A Bézout theorem in real algebraic geometry, Rend. Sci. Mat. Appl., A 115, 99-111.

J.W. Bruce, C.G. Gibson, Distance genericity for real algebraic hypersurfaces, *Can. J. Math.*, 36, 374-384.

E. Bujalance, J.J. Etoyo, J.M. Gamboa, Real algebraic curves and Klein surfaces, *Rev. R. Acad. Cienc. Exactas Fís. Nat. Madr.*, 78, 221-227.

A.H. Durfee, Four characterizations of real rational double points, *Ensign. Math.* II-30, 1-6.

P. Gianni, Traverso, Shape determination for real curves and surfaces, *Ann. Univ. Ferrara, Nuova Ser., Sez. VII* 29, 87-109.

G. Gozalez-Sprinberg, V. Ruggiero, Petites deformations de droites dans le plan projectif reel, *Ann. Univ. Ferrara, Nuova Ser., Sez. VII* 29, 179-210.

V.M. Kharlamov, Classification of non-singular surfaces of degree 4 in  $\mathbb{R}P^3$  with respect to rigid isotopies, *Funkt. Anal. Prilozhen.*, 18-1, 49-56. = *Funct. Anal. Appl.*, 18, 39-45.

A.G. Khovanskii, Real analytic varieties with the finiteness property and complex abelian integrals, *Funkt. Anal. Prilozhen.*, 18-2, 40-50. = *Funct. Anal. Appl.*, 18, 119-127.

\_\_\_\_\_ , Cycles of dynamical systems on the plane and Rolle's theorem, *Sibir. Mat. Zhurnal*, 25-3, 198-203.

M. Letizia, Quotients by complex conjugation of nonsingular quadrics and cubics in  $P_{\mathbb{C}}^3$  defined over  $\mathbb{R}$ , *Pacific J. Math.*, 110-2, 307-314.

M.G. Marinari, M. Reimondo, On complete intersection real curves, *Rocky Mt. J. Math.*, 14, 919-920.

S.M. Natanzon, Spaces of real meromorphic functions on real algebraic curves, *Sov. Math. Dokl.*, 30, 724-726.

J.-J. Risler, Complexité et géométrie réelle (d'après A.Khovansky), Séminaire Bourbaki 37<sup>e</sup> année 1984-85, n<sup>o</sup> 637.

M. Seppälä, On moduli of real curves, Rocky Mount. J. Math., 14-4, 969-970.

H.W. Schulting, Systems of real varieties, Rocky Mt. J. Math., 14, 977-980.

R. Silhol, Towards a classification of real algebraic surfaces Rocky Mount. J. Math., 14-4, 903-904.

\_\_\_\_\_, Real algebraic surfaces with rational or elliptic fiberings, Math. Z., 186, 465-499.

### 1985

E. Bujalance, D. Singerman, The symmetry type of a Riemann surface, Proc. London Math. Soc., 51, 501-519.

B. Chevallier, Sur les courbes maximales de Harnack, 300, (I)4 109-114.

\_\_\_\_\_, Courbes maximales et courbes non réduites, C. R Acad. Sc. Paris, 301, (I)2, 35-40.

T. Fidler, Additional inequalities in the topology of real plane algebraic curves, Izv. Akad. Nauk SSSR, 49-4, 874-883. = Math. USSR Izv., 27-1(1986), 183-192.

G. Ishikawa, The number of singular points in a pencil of real plane algebraic curves, Japan. J. Math., 11-2, 387-401.

V.N. Karpushkin, Precise order bounds for the number of component of the zeros of harmonic polynomials, Funkt. Anal. Prilozhen., 19-4, 55-60. = Funct. Anal. Appl., 19, 295-299.

V.A. Krasnov, Degenerations of real algebraic varieties, Izv. Akad. Nauk SSSR, 49-4. = Math. USSR Izv., 27-1(1986), 115-140.



V.V. Nikulin, Filtrations of 2-elementary forms and involutions of integral symmetric and skew-symmetric bilinear forms, *Izv. Akad. Nauk SSSR*, 49-4. = *Math. USSR Izv.*, 27-1, 159-182.

J.J. Risler, Type topologique des surfaces algébriques réelles de degré 4 dans  $\mathbb{R}P^3$ , *Astérisque*, 126, 153-168.

E.I. Shustin, Counterexamples to a conjecture of Rokhlin, *Funkt. Anal. Prilozhen.*, 19-2, 94-95. = *Funct. Anal. Appl.*, 19, 162-163.

\_\_\_\_\_ , Estimates for the number of singular points of algebraic curves, *Differential and integral equation*, Gos. Univ. Gorki., pp. 81-83.

\_\_\_\_\_ , The Hilbert-Roon method and removals of singular points of real algebraic curves, *Dokl. Akad. Nauk SSSR*, 281, 33-36. = *Sov. Math. Dokl.*, 31, 282-285.

O.Ya. Viro, Topological problems concerning lines and points of three-dimensional space, *Dokl. Akad. Nauk SSSR*, 284-5, 1049-1052. = *Soviet Math. Dokl.*, 32-2, 528-531.

### 1986

R. Benedetti, Finiteness for the topology of semi-algebraic sets of bounded complexity - Some examples, in [R, 1986], pp.1-27.

E. Bujalance, Automorphism groups of compact Klein surfaces with one boundary component, *Math. Scand.*, 59, 45-58.

E. Bujalance, J.J. Etayo, J.M. Gamboa, Automorphism groups of Klein surfaces without involutions I, *Math. Contributions*, Mon. D. F. Botella Ráduan, pp. 33-75.

B. Chevallier, A propos des courbes de degré 6, *C. R. Acad. Sc, Paris*, 302-(I)1, 33-38.

\_\_\_\_\_ , Courbes maximales de Harnack et discriminant,

in [R,1986], pp. 41-65.

T. Fidler, New congruences in the topology of singular real algebraic plane curves, Dokl. Akad. Nauk SSSR, 286, 1075-1079. = Sov. Math. Dokl., 33, 262-266.

[GM]: L. Guillou, A. Marin, ed., A la Recherche de la Topologie Perdue, Prog. in Math., 62, Birkhäuser.

V.M. Kharlamov (Harlamov), Surfaces non amphichérales de degré 4 dans  $RP^3$ , in [R,1986], pp. 67-72.

A.B. Korchagin, M-curves of degree 9: New prohibitions, Mat. Zametki, 39-2, 277-283.

F. Loeser, Un analogue local de l'inégalité de Petrowsky-Oleinik, in [R,1986], pp. 73-83.

G.M. Polotovskii, Connection between the rigid isotopy class of a fifth-order nonsingular curves in  $RP^2$  and its disposition with respect to a line, Funkt. Anal. Prilozhen., 20-4, 87-88. = Funct. Anal. Appl., 20, 330-332.

[R]: J-J. Risler, ed., Séminaire sur la géométrie algébrique réelle II, Publ. Math. l'Univ. Paris VII.

J-J. Risler, Propriétés métriques des ensembles semi-algébriques réels et complexité additive, in [R,1986], pp. 29-40.

M. Seppälä, On moduli of real curves, in [R,1986], 85-95.

R. Silhol, Bounds for the connected components and the first Betti number mod two of a real algebraic surface, Compositio Math., 60, 53-63.

\_\_\_\_\_, Sur la topologie des surfaces algébriques réelles, in [R,1986], 97-109.

Z. Szafraniec, On the Euler characteristic of analytic and algebraic sets, *Topology*, 25-4, 411-414.

V.A. Vasil'ev, Sharpness and the local Petrovskii condition for strictly hyperbolic operators with constant coefficients, *Izv. Akad. Nauk SSSR*, 50-2. = *Math. USSR Izv.*, 28-2(1987), 233-273.

O.Ya. Viro, Progress in the topology of real algebraic varieties over the last six years, *Usp. Mat. Nauk* 41-3, 45-67. = *Russ. Math. Surveys* 41-3, 55-82.

C.T.C. Wall, Real forms of cusp singularities, *Math. Proc. Camb. Philos. Soc.*, 99, 213-232.

### 1987

[ASV]: J-M. Aroca, T. Sanchez-Giralda, J-L. Vincente, ed., *Géométrie algébrique et applications III, Géométrie réelle, Systèmes différentiels et Théorie de Hodge*, *Travaux en cours*, 24, Hermann.

E. Bujalance, J.J. Etoya, J.M. Gamboa, Double coverings of hyperelliptic Klein surfaces, *C. R. Math. Acad. Sci., Soc. R. Can.*, 9, 107-112.

S.M. Finashin, M. Kreck, O.Ya. Viro, Exotic knottings of surfaces in the 4-sphere, *Bull. Amer. Math. Soc.*, 17-2, 287-290.

E.J. Kani, Unramified double covers of hyperelliptic Klein surfaces, *C. R. Math. Acad. Sci., Soc. R. Can.*, 9, 133-138.

H. Knörrer, T. Miller, Topologische Typen reeller kubischer Flächen, *Math. Z.*, 195, 51-67.

W. Kucharz, Real algebraic curves as complete intersections, *Math. Z.*, 194, 259-266.

S.M. Natanzon, Real meromorphic functions on real algebraic curves, *Dokl. Akad. Nauk SSSR*, 297-1, 40-43.

E.I. Shustin, Versal deformations in the space of planar curves of fixed degree, *Funkt. Anal. Prilozhen.*, 21-1, 90-91. = *Funct. Anal Appl.*, 21, 82-84.

\_\_\_\_\_, A new M-curves of degree eight, *Mat. Zametki*, 42-2, 180-186.

R. Silhol, Cohomologie de Galois et cohomologie des variétés algébriques réelles; applications aux surfaces rationnelles. *Bull. Soc. Math. France* 115-1, 107-125.

\_\_\_\_\_, in [ASV,1987], pp. 149-163.

C.T.C. Wall, Real forms of cusp singularities II, *Math. Proc. Camb. Philos. Soc.* 102, 193-201.

\_\_\_\_\_, Real forms of smooth del Pezzo surfaces, *J. Reine Angew. Math.*, 375/376, 47-66.

### 1988

S. Akbulut, H.C. King, Polynomial equations of immersed surfaces, *Pacific J. Math.*, 131-2, 209-217.

K. Kurdyka, K. Rusek, Polynomial-rational bijections on  $\mathbb{R}^n$ , *Proc. Amer. Math. Soc.*, 102-4, 804-808.

M. Seppälä, Complex algebraic curves with real moduli, *J. Reine Angew. Math.*, 387, 209-220.

1989. 1. 24.