

Residual currents and tensor products of holonomic systems

新潟大教養 田島 慎一

Shinichi TAJIMA

The primary purpose of this note is to provide an elementary introduction to the theory of \mathcal{D}_X -Modules. This note is not meant to be a survey. We try to explain some basic notions in the theory of \mathcal{D}_X -modules. We mainly focus our attention on concrete examples of regular holonomic \mathcal{D}_X -Modules which are related with residual currents. It is hoped that some of the basic features of holonomic systems emerge.

§ 0. Products of residual currents.

To begin with let us recall the notion of residual current.

Let (X, \mathcal{O}_X) be a complex manifold and let $f \in \Gamma(X, \mathcal{O}_X)$ be a holomorphic function. Dolbeault (6) and Herrera-Liebermann (10) showed, by making use of a desingularization of the hypersurface defined by $f = 0$, that for any compactly supported smooth test differential form ϕ , the limits

$$\lim_{\varepsilon \rightarrow +0} \int_{|f| > \varepsilon} \frac{\phi}{f} \quad \text{and} \quad \lim_{\varepsilon \rightarrow +0} \int_{|f| = \varepsilon} \frac{\phi}{f}$$

exist.

The linear functional $[\frac{1}{f}]$ defined by

$$\left[\frac{1}{f}\right] : \phi \longrightarrow \lim_{\varepsilon \rightarrow +0} \int_{|f| > \varepsilon} \frac{\phi}{f}$$

is called the principal value of $\frac{1}{f}$. And the current defined by

$$\phi \longrightarrow \lim_{\varepsilon \rightarrow +0} \int_{|f| = \varepsilon} \frac{\phi}{f}$$

is called a residual current. Note that

$$\bar{\partial} \left[\frac{1}{f} \right] (\phi) = \lim_{\varepsilon \rightarrow 0} \int_{|f| = \varepsilon} \frac{\phi}{f}.$$

Recently Passare [22] defined and intensively studied "products of residual currents". Let r be a product of residual currents of the form

$$r = \bar{\partial} \left[\frac{1}{f_1} \right] \wedge \bar{\partial} \left[\frac{1}{f_2} \right] \wedge \cdots \wedge \bar{\partial} \left[\frac{1}{f_k} \right], \quad f_1, f_2, \dots, f_k \in \Gamma(X, \mathcal{O}_X).$$

Passare showed in particular the following fact.

Theorem(local version). Assume that $f_1 = f_2 = \cdots = f_k = 0$ is a complete intersection at x . Then a germ of holomorphic function $h \in \mathcal{O}_x$ annuls the residual current r if and only if the holomorphic function h belongs to the ideal generated by f_1, f_2, \dots, f_k i.e. $h \in \mathcal{O}_x(f_1, f_2, \dots, f_k)$. Here \mathcal{O}_x denotes the stalk at x of the sheaf of holomorphic functions.

For instance, in $\mathbb{C}^2 = \{(x, y) \mid x, y \in \mathbb{C}\}$, let us consider the

following residual current.

$$r = \bar{\partial}\left[\frac{1}{y^2}\right] \wedge \bar{\partial}\left[\frac{1}{y-x^2}\right]$$

Since $y^2 r = 0$ and $(y-x^2)r = 0$, you get $x^4 r = 0$. Hence in particular the current r is supported at the origin. Therefore r is a linear combination of derivatives of Dirac's delta function. In fact, one can verify the following equality.

$$\left(\bar{\partial}\left[\frac{1}{y^2}\right] \wedge \bar{\partial}\left[\frac{1}{y-x^2}\right] dx \wedge dy\right)\phi = -(2\pi i)^2 \left\{ \frac{1}{6} \frac{\partial^3 \phi}{\partial x^3}(0, 0) + \frac{\partial^2 \phi}{\partial x \partial y}(0, 0) \right\}.$$

But it seems difficult, in general, to get such an explicit formula. We are thus interested in the following problem.

How to calculate or characterize a residual current?

Now let us explain our basic idea.

Let r be a residual current in the sense of Passare (22). Let \mathcal{D}_X be the sheaf on X of rings of holomorphic linear partial differential operators with holomorphic coefficients. We write $\mathcal{I}_r \subset \mathcal{D}_X$ the sheaf of annihilator ideals of the residual current r .

$$\mathcal{I}_{x,x} = \{P \in \mathcal{D}_{x,x} \mid Pm = 0\},$$

where $\mathcal{D}_{x,x}$ denotes the stalk at x of \mathcal{D}_X . Then the current r can be regarded as a distribution solution for the system of linear partial differential equations : $Pr = 0$ for any $P \in \mathcal{I}_r$.

If the annihilator ideal \mathcal{I}_r is determined, one can employ the theory of linear partial differential equations to study the residual current r . And

if in particular the dimension of the vector space of distribution solutions of this system above is equal to one, the ideal \mathcal{I}_r characterizes the residual current r .

In this report we will restrict ourselves mainly to the two dimensional case and we will derive a regular holonomic system and its generator whose distribution solutions may equal to a constant multiple of the residual current r . In other words we will derive a left \mathcal{D}_X -ideal which may equal to the annihilator ideal of the residual current r .

We will use the notions of

- (i) \mathcal{D}_X -Modules and algebraic local cohomologies,
- (ii) regular holonomic distributions,
- (iii) tensor products of holonomic systems,

and

- (iv) blow-up and blow-down of holonomic systems.

§ 1. \mathcal{D}_X -Modules and algebraic local cohomologies.

Let us recall some basic notions and fix our notation.

We start with a linear partial differential equation with an unknown function u : $Pu = 0$, where P is a holomorphic linear partial differential operator with holomorphic coefficients. Notice that $QPu = 0$ holds for any linear partial differential operator Q . Let us consider the left \mathcal{D}_X -linear homomorphism from \mathcal{D}_X to \mathcal{D}_X defined by P :

$$\mathcal{D}_X \ni QP \longleftarrow \xrightarrow{P} Q \in \mathcal{D}_X$$

Here \mathcal{D}_X denotes the sheaf on X of holomorphic linear partial differential operators with holomorphic coefficients.

If we denote by \mathcal{I} the sheaf of left \mathcal{D}_x -ideals generated by P , we have $\text{Ker}(P : \mathcal{D}_x \longrightarrow \mathcal{D}_x) = 0$ and $\text{Im}(P : \mathcal{D}_x \longrightarrow \mathcal{D}_x) = \mathcal{I}$.

If we set

$$\mathcal{M} = \text{Coker}(P : \mathcal{D}_x \longrightarrow \mathcal{D}_x) = \mathcal{D}_x / \mathcal{I},$$

then \mathcal{M} becomes a left \mathcal{D}_x -Module. And the exact sequence

$$0 \longleftarrow \mathcal{M} \longleftarrow \mathcal{D}_x \xleftarrow{P} \mathcal{D}_x \longleftarrow 0.$$

can be regarded as a finite presentation of the left \mathcal{D}_x -Module \mathcal{M} . In this sense \mathcal{M} is an intrinsic object.

Notice that \mathcal{D}_x is a coherent sheaf of rings and the each stalk $\mathcal{D}_{x,x}$ is a Noetherian ring.

Let \mathcal{F} be a left \mathcal{D}_x -Module. The solutions of the equation $Pu = 0$ belonging to \mathcal{F} can be considered as follows. From the exact sequence above we get an exact sequence

$$\begin{array}{ccccc} 0 \longrightarrow \text{Hom}_{\mathcal{D}_x}(\mathcal{M}, \mathcal{F}) & \longrightarrow & \text{Hom}_{\mathcal{D}_x}(\mathcal{D}_x, \mathcal{F}) & \xrightarrow{P} & \text{Hom}_{\mathcal{D}_x}(\mathcal{D}_x, \mathcal{F}) \\ & & \parallel & & \parallel \\ & & \mathcal{F} \ni g & \longrightarrow & Pg \in \mathcal{F}. \end{array}$$

It follows that

$$\text{Hom}_{\mathcal{D}_x}(\mathcal{M}, \mathcal{F})_x = \text{Ker}(P : \mathcal{F} \longrightarrow \mathcal{F})_x = \{g \in \mathcal{F}_x \mid Pg = 0\}.$$

Therefore determining the solutions of $Pu = 0$ is equivalent to determining

the sheaf $\mathcal{H}om_{\mathcal{D}_x}(\mathcal{M}, \mathcal{F}_1)$. Incidentally, we have

$$\text{Ext}_{\mathcal{D}_x}^1(\mathcal{M}, \mathcal{F}_1) = \text{Coker} (P : \mathcal{F}_1 \rightarrow \mathcal{F}_1),$$

in the sense of homological algebra.

More generally let

$$\sum_{j=1}^N P_{ij} u_j = 0 \quad i = 1, 2, \dots, N_1$$

be a system of linear partial differential equations, where P_{ij} denote linear partial differential operators and u_j denote unknown functions. We can associate to this system a coherent left \mathcal{D}_x -Module \mathcal{M} with finite presentation :

$$0 \longleftarrow \mathcal{M} \longleftarrow \mathcal{D}_x^N \longleftarrow \mathcal{D}_x^{N_1}$$

Let \mathcal{F}_1 be a left \mathcal{D}_x -Module. We can consider the sheaf of \mathcal{D}_x -homomorphisms from \mathcal{M} to \mathcal{F}_1 and its extensions :

$$\mathcal{H}om_{\mathcal{D}_x}(\mathcal{M}, \mathcal{F}_1) \quad \text{and} \quad \text{Ext}_{\mathcal{D}_x}^k(\mathcal{M}, \mathcal{F}_1) \quad k \geq 1.$$

Let

$$0 \longleftarrow \mathcal{M} \longleftarrow \mathcal{D}_x^N \xleftarrow{P} \mathcal{D}_x^{N_1} \xleftarrow{P_1} \mathcal{D}_x^{N_2} \xleftarrow{P_2} \dots$$

be a (local) projective resolution of \mathcal{M} , where P_j denote matrices of linear partial differential operators. It is easy to see that $\mathcal{H}om_{\mathcal{D}_x}(\mathcal{M}, \mathcal{F}_1)$ is the solutions sheaf and each cohomology group $\text{Ext}_{\mathcal{D}_x}^k(\mathcal{M}, \mathcal{F}_1)$ gives the

obstructions of the solvability of P_{k-1} with compatibility condition given by P_k .

For this reason we mean by a system of linear partial differential equations a coherent left \mathcal{D}_X -Module.

Example 1.

The structure sheaf \mathcal{O}_X is naturally endowed with a structure of left \mathcal{D}_X -Module by differentiation. For instance, let $X = \{(x, y) \mid x, y \in \mathbb{C}\} = \mathbb{C}^2$. For any germ $f \in \mathcal{O}_{X, x}$ of holomorphic function we have

$$\frac{\partial}{\partial x} f = \frac{\partial f}{\partial x} \in \mathcal{O}_{X, x} \quad \text{and} \quad \frac{\partial}{\partial y} f = \frac{\partial f}{\partial y} \in \mathcal{O}_{X, x}.$$

But if you regard f as an element of $\mathcal{D}_{X, x}$, f becomes a linear partial differential operator of order zero and

$$\frac{\partial}{\partial x} f = \frac{\partial f}{\partial x} + f \frac{\partial}{\partial x}, \quad \text{etc.}$$

Hence we have $\mathcal{O}_X = \mathcal{D}_X / (\mathcal{D}_X \frac{\partial}{\partial x} + \mathcal{D}_X \frac{\partial}{\partial y})$.

We also have

$$\text{Hom}_{\mathcal{D}_X}(\mathcal{O}_X, \mathcal{O}_X) = \mathbb{C}_X \quad \text{and} \quad \text{Ext}_{\mathcal{D}_X}^k(\mathcal{O}_X, \mathcal{O}_X) = 0 \quad \text{for } k \geq 1.$$

Let us recall the notion of algebraic local cohomology.

Let $Y = \{f = 0\}$ be a complex hypersurface, f denotes a defining holomorphic function. We define the algebraic local cohomology sheaf by the inductive limit

$$\mathcal{H}_{[Y]}^1(\mathcal{O}_X) = \varinjlim_k \mathcal{E}_{\text{xt}}^1_{\mathcal{O}_X}(\mathcal{O}_X / (f)^k, \mathcal{O}_X)$$

where $(f)^k$ denotes the \mathcal{O}_X -Ideal generated by f^k . We also consider

$$\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X) = \varinjlim_k \mathcal{H}om_{\mathcal{O}_X}((f)^k, \mathcal{O}_X),$$

hence have a exact sequence

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{H}_{[X|Y]}^0(\mathcal{O}_X) \longrightarrow \mathcal{H}_{[Y]}^1(\mathcal{O}_X) \longrightarrow 0.$$

If we denote by ϕ_k be a section of $\mathcal{H}om_{\mathcal{O}_X}((f)^k, \mathcal{O}_X)_x$ defined by $\phi_k(f^k) = 1$, then ϕ_k can be identified with $\frac{1}{f^k}$. Hence a germ of

$\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X)$ can be represented by

$$h_0 + \frac{h_1}{f} + \frac{h_2}{f^2} + \cdots + \frac{h_k}{f^k}, \quad \text{where } h_0, \dots, h_k \in \mathcal{O}_{X, x}.$$

This means that $\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X)$ is the sheaf of meromorphic functions with pole on Y .

A germ of $\mathcal{H}_{[Y]}^1(\mathcal{O}_X)$ can be represented by

$$\frac{h_1}{f} + \frac{h_2}{f^2} + \cdots + \frac{h_k}{f^k} \pmod{\mathcal{O}_{X, x}}, \quad \text{where } h_0, \dots, h_k \in \mathcal{O}_{X, x}.$$

It is thus easy to verify that these two sheaves have the structure of left \mathcal{D}_X -Module.

Example 2.

Let $X = \mathbb{C}^2 = \{(x, y) \mid x, y \in \mathbb{C}\}$, $Y = \{(x, y) \in X \mid x = 0\}$.

If we set $n = \frac{1}{x}$, then n generates the sheaf $\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X)$ over \mathcal{D}_X .

Since $(x \frac{\partial}{\partial x} + 1)n = 0$ and $\frac{\partial}{\partial y} n = 0$, we have

$$\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X) \cong \mathcal{D}_X n = \mathcal{D}_X / (\mathcal{D}_X(x \frac{\partial}{\partial x} + 1) + \mathcal{D}_X \frac{\partial}{\partial y}).$$

Hence $\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X)$ is a coherent left \mathcal{D}_X -Module.

It is easy to verify the followings.

$$\text{Hom}_{\mathcal{D}_X}(\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X), \mathcal{O}_X) = \mathbb{C}_{X-Y} = j_* \mathbb{C}_{X-Y}$$

and

$$\text{Ext}_{\mathcal{D}_X}^k(\mathcal{H}_{[X|Y]}^0(\mathcal{O}_X), \mathcal{O}_X) = 0 \quad \text{for } k \geq 1,$$

where $j : X-Y \hookrightarrow X$ is the natural open inclusion map.

Example 3.

Let $X = \mathbb{C}^2 = \{(x, y) \mid x, y \in \mathbb{C}\}$, $Y = \{(x, y) \in X \mid x = 0\}$.

If we set $m = \frac{1}{x} \text{ mod } \mathcal{O}_X$, then m generates the \mathcal{D}_X -Module $\mathcal{H}_{[Y]}^1(\mathcal{O}_X)$.

Since $xm = 1 \in \mathcal{O}_X$ and $\frac{\partial}{\partial y} m = 0$, we have

$$\mathcal{H}_{[Y]}^1(\mathcal{O}_X) = \mathcal{D}_X m = \mathcal{D}_X / (\mathcal{D}_X x + \mathcal{D}_X \frac{\partial}{\partial y}).$$

We also have

$$\text{Hom}_{\mathcal{D}_X}(\mathcal{H}_{[Y]}^1(\mathcal{O}_X), \mathcal{O}_X) = 0,$$

$$\text{Ext}_{\mathcal{D}_X}^1(\mathcal{H}_{[Y]}^1(\mathcal{O}_X), \mathcal{O}_X) = \mathbb{C}_Y$$

and

$$\text{Ext}_{\mathcal{D}_X}^k(\mathcal{H}_{[Y]}^1(\mathcal{O}_X), \mathcal{O}_X) = 0 \quad \text{for } k \geq 2.$$

Note. Set $m' = \frac{\partial}{\partial x} m$. Then $m' \in \mathcal{D}_X m$. Hence $\mathcal{D}_X m' \subseteq \mathcal{D}_X m$.

Since $m = -xm'$, we have $\mathcal{D}_X m \subseteq \mathcal{D}_X m'$. Therefore we have

$\mathcal{D}_X m = \mathcal{D}_X m'$ as set. But the annihilator ideal of m' is equal

to

$$\mathcal{D}_X x^2 + \mathcal{D}_X (x \frac{\partial}{\partial x} + 2) + \mathcal{D}_X \frac{\partial}{\partial y}.$$

In what follows we say that a \mathcal{D}_X -Module $\mathcal{D}_X m_1$ generated by m_1 is equal to a \mathcal{D}_X -Module $\mathcal{D}_X m_2$ generated by m_2 if the annihilator ideal of m_1 is equal to the annihilator ideal of m_2 .

Notation.

We write $\mathcal{M}_1 \cong \mathcal{M}_2$ if the \mathcal{D}_X -Module \mathcal{M}_1 is isomorphic to the

\mathcal{D}_X -Module \mathcal{M}_2 . We write $\mathcal{D}_X m_1 = \mathcal{D}_X m_2$ if the \mathcal{D}_X -Module $\mathcal{D}_X m_1$ is

equal to $\mathcal{D}_x^{m_2}$.

Let \mathcal{I} be a coherent \mathcal{O}_x -Ideal and Y the support of $\mathcal{O}_x/\mathcal{I}$. We define the q -th algebraic local cohomology sheaf by the inductive limit

$$\mathcal{H}_{[Y]}^q(\mathcal{O}_x) = \varinjlim_k \text{Ext}_{\mathcal{O}_x}^q(\mathcal{O}_x/\mathcal{I}^k, \mathcal{O}_x).$$

We also set

$$\mathcal{H}_{[X|Y]}^q(\mathcal{O}_x) = \varinjlim_k \text{Ext}_{\mathcal{O}_x}^q(\mathcal{I}^k, \mathcal{O}_x).$$

We thus have a long exact sequence

$$0 \rightarrow \mathcal{H}_{[Y]}^0(\mathcal{O}_x) \rightarrow \mathcal{O}_x \rightarrow \mathcal{H}_{[X|Y]}^0(\mathcal{O}_x) \rightarrow \mathcal{H}_{[Y]}^1(\mathcal{O}_x) \rightarrow 0.$$

and isomorphisms

$$\mathcal{H}_{[X|Y]}^q(\mathcal{O}_x) = \mathcal{H}_{[Y]}^{q+1}(\mathcal{O}_x) \quad q \geq 1.$$

It is easy to see that these are left \mathcal{D}_x -Modules.

Recall that a left coherent \mathcal{D}_x -Module whose characteristic variety is Lagrangian is called a holonomic system.

Following result is known.

Theorem (Kashiwara (13), Mebkhout (20)).

(i) $\mathcal{H}_{[Y]}^q(\mathcal{O}_x)$ is coherent.

(ii) $\mathcal{H}_{[Y]}^q(\mathcal{O}_x)$ is a regular holonomic system

For the notion of regular singularity we refer to (16).

Theorem (Mebkhout (20)).

$$R\mathcal{H}om_{\mathcal{D}_X} (R\Gamma_{[Y]}(\mathcal{O}_X), \mathcal{O}_X) = C_Y,$$

where $R\Gamma_{[Y]}$ denote the right derived functor of $\Gamma_{[Y]}$ and $R\mathcal{H}om_{\mathcal{D}_X}(\cdot, \mathcal{O}_X)$ denotes the right derived functor of $\mathcal{H}om_{\mathcal{D}_X}(\cdot, \mathcal{O}_X)$.

To end this section let us recall that the sheaf \mathcal{D}_X has the following description :

$$\mathcal{D}_X = \mathcal{H}_{[X]}^{\dim X} (\mathcal{O}_{X \times X}^{(0, \dim X)})$$

where $\mathcal{O}_{X \times X}^{(0, \dim X)}$ is the sheaf of $(\dim X)$ -holomorphic forms in the second variables with holomorphic functions as coefficients and X is identified with the diagonal of $X \times X$. Note also that the sheaf \mathcal{D}_X^∞ of rings of linear partial differential operators of infinite order can be described by

$$\mathcal{D}_X^\infty = \mathcal{H}_X^{\dim X} (\mathcal{O}_{X \times X}^{(0, \dim X)}).$$

Let X and Y be complex manifolds and ψ a holomorphic map from Y to X . Kashiwara introduced the sheaf $\mathcal{D}_{Y \rightarrow X}$ and $\mathcal{D}_{X \leftarrow Y}$ by

$$\mathcal{D}_{Y \rightarrow X} = \mathcal{H}_{[Y]}^{\dim X} (\mathcal{O}_{Y \times X} \otimes_{\mathcal{O}_X} \Omega_X^{\dim X}),$$

and

$$\mathcal{D}_{X \leftarrow Y} = \mathcal{H}_{[Y]}^{\dim X} (\Omega_Y^{\dim Y} \otimes_{\mathcal{O}_Y} \mathcal{O}_{Y \times X}),$$

respectively, where $\Omega_X^{\dim X}$ (resp. $\Omega_Y^{\dim Y}$) signifies the sheaf of the holomorphic $(\dim X)$ -forms (resp. $(\dim Y)$ -forms).

§ 2. Regular holonomic distributions and residual currents.

In this section we try to illustrate the Reconstruction theorem for holonomic systems (16). We shall content ourselves with discussing the most simplest case which is related to residual currents.

Let us first examine the one dimensional case. Let $X = \{z \mid z \in \mathbb{C}\}$, $Y = \{0\}$. We set

$$\mathcal{M}^- = \mathcal{D}_X / \mathcal{D}_X \frac{d}{dz} \cong \mathcal{O}_X,$$

$$\mathcal{M} = \mathcal{D}_X / \mathcal{D}_X (z \frac{d}{dz} + 1) \cong \mathcal{H}_{[X|Y]}^0(\mathcal{O}_X),$$

and

$$\mathcal{M}^- = \mathcal{D}_X / \mathcal{D}_X z \cong \mathcal{H}_{[Y]}^1(\mathcal{O}_X).$$

We thus have a short exact sequence of \mathcal{D}_X -Modules.

$$(*) \quad 0 \longrightarrow \mathcal{M}^- \longrightarrow \mathcal{M} \longrightarrow \mathcal{M}^- \longrightarrow 0$$

By applying the derived functor $R\mathcal{H}om_{\mathcal{D}_X}(*, \mathcal{O}_X)$ to the exact

sequence above we get an exact sequence

$$\begin{aligned}
 0 &\longrightarrow \mathcal{H}om_{\mathcal{D}_X}(M^-, \mathcal{O}_X) \longrightarrow \mathcal{H}om_{\mathcal{D}_X}(M, \mathcal{O}_X) \longrightarrow \mathcal{H}om_{\mathcal{D}_X}(M^+, \mathcal{O}_X) \\
 &\longrightarrow \mathcal{E}xt_{\mathcal{D}_X}^1(M^-, \mathcal{O}_X) \longrightarrow \mathcal{E}xt_{\mathcal{D}_X}^1(M, \mathcal{O}_X) \longrightarrow \mathcal{E}xt_{\mathcal{D}_X}^1(M^+, \mathcal{O}_X) \longrightarrow 0.
 \end{aligned}$$

Since $\mathcal{H}om_{\mathcal{D}_X}(M, \mathcal{O}_X)_Y = \mathcal{E}xt_{\mathcal{D}_X}^1(M, \mathcal{O}_X)_Y = 0$, we thus have

$$(**) \quad 0 \longrightarrow C_{X-Y} \longrightarrow C_X \longrightarrow C_Y \longrightarrow 0$$

Let us recall the following quasi-isomorphisms.

$$R\mathcal{H}om_{\mathcal{C}}(C_{X-Y}, \mathcal{O}_X) = [0 \rightarrow \mathcal{B}_{X-Y} \xrightarrow{\bar{\partial}} \mathcal{B}_{X-Y} \rightarrow 0],$$

$$R\mathcal{H}om_{\mathcal{C}}(C_X, \mathcal{O}_X) = [0 \rightarrow \mathcal{B}_X \xrightarrow{\bar{\partial}} \mathcal{B}_X \rightarrow 0],$$

and

$$R\mathcal{H}om_{\mathcal{C}}(C_Y, \mathcal{O}_X) = [0 \rightarrow \Gamma_Y \mathcal{B}_X \xrightarrow{\bar{\partial}} \Gamma_Y \mathcal{B}_X \rightarrow 0].$$

where \mathcal{B}_X denotes the sheaf of hyperfunctions and $\Gamma_Y \mathcal{B}_X$ denotes the sheaf

of hyperfunctions with supports in Y . It follows that

$$R\mathcal{H}om_{\mathcal{C}}(C_{X-Y}, \mathcal{O}_X) = j_* j^{-1} \mathcal{O}_X,$$

$$R\mathcal{H}om_{\mathcal{C}}(C_X, \mathcal{O}_X) = \mathcal{O}_X,$$

and

$$R\mathcal{H}om_c(\mathcal{C}_Y, \mathcal{O}_X) = \mathcal{H}_Y^1(\mathcal{O}_X)[-1]$$

where j denotes the natural open inclusion map $j : X - Y \hookrightarrow X$. We thus verified the following results.

$$R\mathcal{H}om_c(R\mathcal{H}om_{\mathcal{D}_X}(M', \mathcal{O}_X)) = M'$$

$$R\mathcal{H}om_c(R\mathcal{H}om_{\mathcal{D}_X}(M, \mathcal{O}_X)) = \mathcal{D}_X^\infty \otimes_{\mathcal{D}_X} M$$

and

$$R\mathcal{H}om_c(R\mathcal{H}om_{\mathcal{D}_X}(M'', \mathcal{O}_X)) = \mathcal{D}_X^\infty \otimes_{\mathcal{D}_X} M''$$

where \mathcal{D}_X^∞ denotes the sheaf of rings of linear partial differential operators of infinite orders.

Let us reexamine the result above by using the following double complex associated with the exact sequence (**):

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Gamma_Y \mathcal{B}_X & \longrightarrow & \mathcal{B}_X & \longrightarrow & \mathcal{B}_{X-Y} \longrightarrow 0 \\
 & & \downarrow \bar{\partial} & & \downarrow \bar{\partial} & & \downarrow \bar{\partial} \\
 0 & \longrightarrow & \Gamma_Y \mathcal{B}_X & \longrightarrow & \mathcal{B}_X & \longrightarrow & \mathcal{B}_{X-Y} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

Let start with the meromorphic function $\frac{1}{z} \in \Gamma(X-Y, \mathcal{B}_{X-Y})$. Since the sheaf \mathcal{B}_X is flabby, the sheaf morphism $\mathcal{B}_X \rightarrow \mathcal{B}_{X-Y}$ is surjective.

In fact the principal value hyperfunction $(\frac{1}{z}) \in \Gamma(X, \mathcal{B}_X)$ satisfies

$$[\frac{1}{z}] = \frac{1}{z} \text{ on } X - Y \text{ thus in particular that } \bar{\partial}[\frac{1}{z}] \text{ belongs to}$$

$\Gamma_Y \mathcal{B}_X$. Hence the coboundary morphism

$$\text{Ker}(\bar{\partial} : \mathcal{B}_{X-Y} \rightarrow \mathcal{B}_{X-Y}) \rightarrow \text{Coker}(\bar{\partial} : \Gamma_Y \mathcal{B}_X \rightarrow \Gamma_Y \mathcal{B}_X),$$

maps $\frac{1}{z}$ to $\bar{\partial}[\frac{1}{z}]$.

We thus have the following exact sequence of \mathcal{D}_X^∞ -Modules :

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{D}_X^\infty(\frac{1}{z}) \longrightarrow \mathcal{D}_X^\infty(\bar{\partial}[\frac{1}{z}]) \longrightarrow 0,$$

which is isomorphic to

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{D}_X^\infty \otimes \mathcal{H}_{[X|Y]}^0(\mathcal{O}_X) \longrightarrow \mathcal{D}_X^\infty \otimes \mathcal{H}_{[Y]}^1(\mathcal{O}_X) \longrightarrow 0.$$

We also have the following exact sequence of coherent left \mathcal{D}_X -Modules :

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{D}_X(\frac{1}{z}) \longrightarrow \mathcal{D}_X(\bar{\partial}[\frac{1}{z}]) \longrightarrow 0.$$

It is easy to verify that the exact sequence above is isomorphic to the following one :

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{H}_{[X|Y]}^0(\mathcal{O}_X) \longrightarrow \mathcal{H}_{[Y]}^1(\mathcal{O}_X) \longrightarrow 0.$$

It is hoped that the explanation above illustrate some aspects of the subjects. We refer to (16) and (14) for the reconstruction theorem for holonomic systems and for the Riemann-Hilbert correspondence for regular holonomic systems.

Nextly we briefly examine the two dimensional case.

Let f be a holomorphic function defined in a domain X in \mathbb{C}^2 . We set $F = \{f = 0\}$. Let us denote by $\delta(f)$ the section of $\mathcal{H}_{[F]}^1(\mathcal{O}_X)$ defined by

$\frac{1}{f} \pmod{\mathcal{O}_X}$. If the function f is analytically irreducible, the section

$\delta(f)$ generates the \mathcal{D}_X -Module $\mathcal{H}_{[F]}^1(\mathcal{O}_X)$, for $\mathcal{H}_{[F]}^1(\mathcal{O}_X)$ is simple as

\mathcal{D}_X -Module ((8)).

We have the following result (cf. (3), (7) and (19)).

Theorem. 4

Assume that the holomorphic function f is analytically irreducible.

Then $\mathcal{D}_X(\delta(f)) = \mathcal{D}_X(\bar{\partial}[\frac{1}{f}])$ holds. More precisely the annihilator ideal

of $\delta(f)$ and that of the residual current $\bar{\partial}[\frac{1}{f}]$ coincides.

In particular the residual current $\bar{\partial}[\frac{1}{f}]$ is a regular holonomic distribution ((15)).

§ 3. Tensor products of holonomic systems.

In this section we examine tensor products of holonomic systems supported on plane curves.

Let us recall the notion of tensor product.

Let \mathcal{M}_1 and \mathcal{M}_2 be two \mathcal{D}_X -modules. Let X_1 and X_2 be two copies of the complex manifold X . Let p_1 (resp. p_2) be the projection from $X_1 \times X_2$ to X_1 (resp. X_2). We set

$$\mathcal{M}_1 \widehat{\otimes} \mathcal{M}_2 = \mathcal{D}_{X_1 \times X_2} \otimes_{\mathcal{D}_{X_1} \otimes \mathcal{D}_{X_2}} (p_1^{-1} \mathcal{M}_1 \otimes p_2^{-1} \mathcal{M}_2).$$

Let us denote by \otimes^L the left derived functor of \otimes .

We have the following result.

Proposition (Kashiwara (13)).

For two \mathcal{D}_X -Modules \mathcal{M}_1 and \mathcal{M}_2 , we have

$$\mathcal{M}_1 \otimes_x^L \mathcal{M}_2 = \mathcal{D}_{X \rightarrow X_1 \times X_2} \otimes_{\mathcal{D}_{X_1 \times X_2}}^L (\mathcal{M}_1 \widehat{\otimes} \mathcal{M}_2).$$

Now let us examine the tensor products of two algebraic local cohomologies with support in plane curves.

Let X be a domain in $\mathbb{C}^2 = \{(x, y) \mid x, y \in \mathbb{C}\}$ containing the origin $P = (0, 0)$. Let F and G be two analytic plane curves (defined in X) passing through P . Let f (resp. g) be a defining holomorphic function of the curve F (resp. G). We set :

$$\mathcal{L} = \mathcal{H}_{[F]}^1(\mathcal{O}_{X_1}) \widehat{\otimes} \mathcal{H}_{[G]}^1(\mathcal{O}_{X_2}).$$

We identify X with the diagonal of $X_1 \times X_2$. We choose the natural coordinates (x_1, y_1, x_2, y_2) on $X_1 \times X_2$ such that $X = \{(x_1, y_1, x_2, y_2) \mid x_1 = x_2, y_1 = y_2\}$. Since

$$\mathcal{D}_{X \rightarrow X_1 \times X_2} = \mathcal{D}_{X_1 \times X_2} / (x_1 - x_2) \mathcal{D}_{X_1 \times X_2} + (y_1 - y_2) \mathcal{D}_{X_1 \times X_2},$$

$\mathcal{D}_{X \rightarrow X_1 \times X_2} \otimes^L \mathcal{L}$ is quasi-isomorphic to the complex

$$0 \longleftarrow \mathcal{L} \xleftarrow{\chi_1} \begin{array}{c} \mathcal{L} \\ \oplus \\ \mathcal{L} \end{array} \xleftarrow{\chi_2} \mathcal{L} \longleftarrow 0.$$

where $\chi_1 = (x_1 - x_2, y_1 - y_2)$ and $\chi_2 = \begin{pmatrix} y_2 - y_1 \\ x_1 - x_2 \end{pmatrix}$.

If we denote by $1_{X \rightarrow X_1 \times X_2}$ the canonical section of $\mathcal{D}_{X \rightarrow X_1 \times X_2}$

(23), we have the following:

$$x \cdot 1_{X \rightarrow X_1 \times X_2} = 1_{X \rightarrow X_1 \times X_2} \left(\frac{1}{2} (x_1 + x_2) \right),$$

$$\frac{\partial}{\partial x} 1_{X \rightarrow X_1 \times X_2} = 1_{X \rightarrow X_1 \times X_2} \left(\frac{\partial}{\partial x_1} + \frac{\partial}{\partial x_2} \right), \text{ etc.}$$

It is easy to verify the following result.

Proposition 5.

Let F and G be two analytically irreducible plane curves containing the origin P . Assume that F and G meet properly at P i.e. $F \cap G = P$. Then we have the following results.

$$(i) \tau_{\mathcal{O}_x}^k(\mathcal{H}_{[F]}^1(\mathcal{O}_x), \mathcal{H}_{[G]}^1(\mathcal{O}_x)) = \tau_{\mathcal{O}_k}^k(\mathcal{D}_{x \rightarrow x_1 \times x_2}(\mathcal{L})) = 0$$

for any $k \geq 1$.

$$(ii) \text{ The tensor product } \mathcal{H}_{[F]}^1(\mathcal{O}_x) \otimes_{\mathcal{O}_x} \mathcal{H}_{[G]}^1(\mathcal{O}_x) = \mathcal{D}_{x \rightarrow x_1 \times x_2} \otimes_{\mathcal{D}_{x_1 \times x_2}} \mathcal{L}$$

is isomorphic to the simple regular holonomic \mathcal{D}_x -Module $\mathcal{H}_{[P]}^2(\mathcal{O}_x)$

supported by the origin P .

Let us denote by $\delta(f)$ (resp. by $\delta(g)$) the section of the sheaf

$$\mathcal{H}_{[F]}^1(\mathcal{O}_x) \text{ (resp. } \mathcal{H}_{[G]}^1(\mathcal{O}_x)) \text{ defined by } \frac{1}{f} \bmod \mathcal{O}_x \text{ (resp. } \frac{1}{g} \bmod \mathcal{O}_x).$$

We set :

$$\mathfrak{m} = \mathcal{D}_{x \rightarrow x_1 \times x_2} \otimes (\delta(f) \widehat{\otimes} \delta(g)).$$

The following result is an immediate consequence of Proposition 5.

Proposition 6.

Under the assumption of Proposition 5, we have the following equality.

$$\mathcal{D}_x^{\mathfrak{m}} \cong \mathcal{D}_{x \rightarrow x_1 \times x_2} \otimes_{\mathcal{D}_{x_1 \times x_2}} \mathcal{L}.$$

We can use the result above to calculate the annihilator ideal of the generator m .

Example 7 ([24]).

Set $X = \mathbb{C}^2 = \{(x, y) \mid x, y \in \mathbb{C}\}$, $F = \{(x, y) \mid y = 0\}$ and $G = \{(x, y) \mid y - x^2 = 0\}$. Let us denote by $\delta(y)$ (resp. $\delta(y - x^2)$) the canonical generator of $\mathcal{H}_{[F]}^1(\mathcal{O}_X)$ (resp. $\mathcal{H}_{[G]}^1(\mathcal{O}_X)$). We set :

$$m = 1_{x \rightarrow x_1 \times x_2} \otimes (\delta(y_1) \widehat{\otimes} \delta(y_2 - x_2^2)).$$

Then we have

$$\mathcal{D}_X m \cong \mathcal{D}_{x \rightarrow x_1 \times x_2} \otimes_{\mathcal{D}_{x_1 \times x_2}} (\mathcal{H}_{[F]}^1(\mathcal{O}_{x_1}) \widehat{\otimes} \mathcal{H}_{[G]}^1(\mathcal{O}_{x_2})),$$

and

$$\begin{aligned} \mathcal{D}_X m &= \mathcal{D}_X / (\mathcal{D}_X x^2 + \mathcal{D}_X (x \frac{\partial}{\partial x} + 2) + \mathcal{D}_X y) \\ &= \mathcal{D}_X \left(-\frac{\partial}{\partial x} \delta(x, y) \right), \end{aligned}$$

where $\delta(x, y)$ denotes Dirac's delta-function at the origin.

For the detailed account, we refer the reader to [25].

Example 8.

Put $F = \{(x, y) \in \mathbb{C}^2 \mid y = 0\}$ and $G = \{(x, y) \in \mathbb{C}^2 \mid y^2 - x^3 = 0\}$.

If we set

$$m = 1_{x \rightarrow x_1, x_2} \otimes (\delta(y_1) \widehat{\otimes} \delta(y_2^2 - x_2^3)),$$

then we have

$$\mathcal{D}_x m \cong \mathcal{H}_{[F]}^1(\mathcal{O}_x) \otimes_{\mathcal{O}_x} \mathcal{H}_{[G]}^1(\mathcal{O}_x).$$

By direct calculation we have

$$\begin{aligned} \mathcal{D}_x m &= \mathcal{D}_x / (\mathcal{D}_x x^3 + \mathcal{D}_x (x \frac{\partial}{\partial x} + 3) + \mathcal{D}_x y) \\ &= \mathcal{D}_x \left(\frac{\partial^2}{\partial x^2} \delta(x, y) \right). \end{aligned}$$

Let us state a conjecture.

Conjecture.

Let F and G be analytically irreducible plane curves intersecting properly at the origin. Let f and g be holomorphic defining function of F and G respectively. If we set

$$m = 1_{x \rightarrow x_1, x_2} \otimes (\delta(f) \widehat{\otimes} \delta(g))$$

then we have $\mathcal{D}_x m = \mathcal{D}_x r$, where r denotes the residual current

$$r = \bar{\partial} \left[\frac{1}{f} \right] \wedge \bar{\partial} \left[\frac{1}{g} \right].$$

This means that the annihilator ideal of m and that of r coincides.

§ 4. Blow-up and blow-down of holonomic systems.

In order to calculate tensor products of holonomic \mathcal{D}_X -Modules we use the notion of blow-up and blow-down of \mathcal{D}_X -Modules.

Let X and Z be two complex manifolds. Let ψ be a proper holomorphic map from Z to X . For any coherent \mathcal{D}_X -Module \mathcal{M} , we set :

$$L\psi^* \mathcal{M} = \mathcal{D}_{Z \rightarrow X} \otimes_{\psi^{-1} \mathcal{D}_X}^L \psi^{-1} \mathcal{M}.$$

For any coherent \mathcal{D}_Z -Module \mathcal{N} we set :

$$\int_{\psi} \mathcal{N} = R\psi_* (\mathcal{D}_{X \leftarrow Z} \otimes_{\mathcal{D}_Z}^L \mathcal{N}).$$

Here $R\psi_*$ is the right derived functor of ψ_* .

Example 9.

Let $X = \mathbb{C}^2$ and let $\pi : \tilde{X} \rightarrow X$ be the blowing-up of X at the origin $P = (0, 0)$. Then we have

$$\int_{\pi} \mathcal{O}_{\tilde{X}} = \mathcal{O}_X \oplus \mathcal{R}_{[P]}^2(\mathcal{O}_X).$$

We have the following projection formula which we will use later.

Projection Formula (cf. (11), (25)).

Let ψ be a proper holomorphic map from Z to X . For any coherent \mathcal{D}_Z -Module \mathcal{N} , and for any coherent \mathcal{D}_X -Module \mathcal{M} , we have

$$\int_{\psi} (\mathcal{M} \otimes_{\mathcal{D}_Z}^L L\psi^* \mathcal{M}) = \left(\int_{\psi} \mathcal{M} \right) \otimes_{\mathcal{D}_X}^L \mathcal{M}.$$

Note. If we set $\mathcal{M} = \mathcal{O}_Z$ then we have

$$\int_{\psi} L\psi^* \mathcal{M} = \left(\int_{\psi} \mathcal{O}_Z \right) \otimes_{\mathcal{D}_X}^L \mathcal{M}.$$

Now let us return to the two dimensional case. Let X be a domain in \mathbb{C}^2 and let F be a plane curve on X with a holomorphic defining function f . Let $\pi: \tilde{X} \rightarrow X$ be the blow-up of X at the origin. Let us denote by \tilde{F} the total transform of F , i.e. $\tilde{F} = \{f \circ \pi = 0\}$.

It is easy to verify the following result.

Proposition 10.

$$(i) \quad \text{Tor}_k^{\mathcal{D}_X} (\mathcal{D}_{\tilde{X} \rightarrow X}, \mathcal{H}_{[F]}^1(\mathcal{O}_X)) = 0 \quad \text{for } k \geq 1.$$

$$(ii) \quad \mathcal{D}_{\tilde{X} \rightarrow X} \otimes \mathcal{H}_{[F]}^1(\mathcal{O}_X) = \mathcal{H}_{[\tilde{F}]}^1(\mathcal{O}_{\tilde{X}}).$$

It means that the total transform of the sheaf of algebraic local cohomology supported in F is equal to the sheaf of algebraic local cohomology supported in the total transform of F .

Note that if we set $\tilde{f} = f \circ \pi$ then $\delta(\tilde{f}) = 1_{\tilde{X} \rightarrow X} \otimes \delta(f)$ generates the $\mathcal{D}_{\tilde{X}}$ -Module $\mathcal{H}_{[\tilde{F}]}^1(\mathcal{O}_{\tilde{X}})$.

Example 11.

We calculate the annihilator ideal of $1_{x \rightarrow x_1 x_2} \otimes (\delta(y_1) \hat{\otimes} \delta(y_2^2 - x_2^3))$

by making use of the projection formula.

Set $F = \{(x, y) \in \mathbb{C}^2 \mid f(x, y) = y = 0\}$ and $G = \{(x, y) \in \mathbb{C}^2 \mid g(x, y) = y^2 - x^3 = 0\}$. Let \mathcal{M} be the sheaf of algebraic local cohomology with support in the cusp G :

$$\mathcal{M} = \mathcal{D}_x \delta(g) \cong \mathcal{H}_{[G]}^1(\mathcal{O}_x)$$

Let (u, v) be local coordinates on X which satisfy $x = v$ and $y = uv$. Since $\tilde{g} = g \circ \pi = v^2(u^2 - v)$, we have

$$\begin{aligned} \pi^* \mathcal{M} &= \mathcal{D}_x \delta(\tilde{g}) \\ &= \mathcal{D}_x / (\mathcal{D}_x(v^2(u^2 - v)) + \mathcal{D}_x(u \frac{\partial}{\partial u} + 2v \frac{\partial}{\partial v} + 6) + \mathcal{D}_x(u^2 \frac{\partial}{\partial u} - v \frac{\partial}{\partial u} + 2u)). \end{aligned}$$

If we set $\mathcal{N} = \mathcal{D}_x \delta(u) = \mathcal{D}_x / (\mathcal{D}_x u + \mathcal{D}_x \frac{\partial}{\partial v})$, then we have

$$\int_{\pi} \mathcal{M} = \mathcal{D}_x / (\mathcal{D}_x y + \mathcal{D}_x \frac{\partial}{\partial x}).$$

It is easy to verify that

$$\begin{aligned} \mathcal{N} \otimes_{\mathcal{O}_x} \pi^* \mathcal{M} &= \mathcal{D}_x / (\mathcal{D}_x u + \mathcal{D}_x v^3 + \mathcal{D}_x(v \frac{\partial}{\partial v} + 3)), \\ &= \mathcal{D}_x \left(\frac{\partial^2}{\partial v^2} \delta(u, v) \right) \end{aligned}$$

and

$$\int_{\pi} \mathcal{D}_x \left(\frac{\partial^2}{\partial v^2} \delta(u, v) \right) = \mathcal{D}_x / \left(\mathcal{D}_x x^3 + \mathcal{D}_x \left(x \frac{\partial}{\partial x} + 3 \right) + \mathcal{D}_x y \right).$$

Therefore we have

$$\begin{aligned} \mathcal{D}_x m &= \mathcal{D}_x / \left(\mathcal{D}_x x^3 + \mathcal{D}_x \left(x \frac{\partial}{\partial x} + 3 \right) + \mathcal{D}_x y \right) \\ &= \mathcal{D}_x \left(\frac{\partial^2}{\partial x^2} \delta(x, y) \right), \end{aligned}$$

where $m = 1_{x \rightarrow x_1 \times x_2} \otimes (\delta(y_1) \widehat{\otimes} \delta(y_2^2 - x_2^3))$.

References

- (1) Andronikof, E. : The Kashiwara conjugation and wave-front sets of regular holonomic distributions on a complex manifold, preprint.
- (2) Barlet, D. et Kashiwara, M. : Le réseau L^2 d'un système holonome régulier, *Invent. math.*, 86, 35–62 (1986).
- (3) Bjork, J. -E. : Intersection cohomology and regular holonomic D-Modules, Université Louis Pasteur, Strasbourg.
- (4) Coleff, N. R. et Herrera, M. E. : Les courants résiduels associés à une forme méromorphe, *Springer Lecture Notes in Math.*, 633 (1978).
- (5) Dickenstein, A. and Sessa, C. : Canonical representatives in moderate cohomology, *Invent. math.*, 80, 417–434 (1985).
- (6) Dolbeault, P. : Résidues et courants. Questions on algebraic varieties, *C.I.M.E.*, septembre 1960.
- (7) ————— : Valeurs principales et opérateurs différentiels semi-holomorphes, *Fonctions analytiques de plusieurs variables et analyse complexe*, C.N.R.S., 35–50 (1974).

- [8] van Doorn, M. G. M. and van den Essen, A. R. P. : D_n -Modules with support on a curve, Publ. RIMS, Kyoto Univ., 23, 937–953 (1987).
- [9] Hartshorne, R. : Residues and duality, Springer Lecture Notes in Math., 20 (1966).
- [10] Herrera, M. E. and Lieberman, D. I. : Residues and principal values on complex spaces, Math. Ann., 194, 259–294 (1971).
- [11] Hotta, R. : Introduction to D-Modules, Lecture Notes, Institute of Math. Sci., Madras (1987).
- [12] Iversen, B. : Cohomology of sheaves, Lecture Notes Series 55, Aarhus Univ. (1984).
- [13] Kashiwara, M. : On the holonomic systems of linear differential equations II, Invent. math., 49, 121–135 (1978).
- [14] _____ : The Riemann-Hilbert problem for holonomic systems, Publ. RIMS, Kyoto Univ., 20, 319–365 (1984).
- [15] _____ : Regular holonomic D-Modules and distributions on complex manifolds, Advanced Studies in Pure Math. 8, Complex Analytic Singularities, 199–206 (1986).
- [16] Kashiwara, M. and Kawai, T. : On the holonomic systems of micro-differential equations III, Publ. RIMS, Kyoto Univ., 17, 813–979 (1981).
- [17] Kashiwara, M. and Schapira, P. : An introduction to microlocal analysis, Notas de curso, Univ. Federal de Pernambuco, Brasil (1984).
- [18] Lê D. T. and Mebkhout, Z. : Introduction to linear differential systems, Proc. Symp. Pure Math. AMS, 40 (2), 31–63 (1983).
- [19] Mebkhout, Z. : La valeur principale et le résidu simple des formes à singularités essentielles, Springer Lecture Notes in Math., 482, 190–215 (1975).
- [20] _____ : Local cohomology of analytic spaces, Publ. RIMS, Kyoto Univ., 12 suppl., 247–256 (1977).

- [21] _____ : Une autre équivalence de catégories, *Compositio Math.*, 51, 63–88 (1984).
- [22] Passare, M. : Residues, currents, and their relation to ideals of holomorphic functions, *Math. Scand.*, 62, 75–152 (1988).
- [23] Schapira, P. : *Microdifferential Systems in the Complex Domain*, Springer (1985).
- [24] Tajima, S. : A calculus of the tensor product of two holonomic systems with support on non-singular plane curves, *Proc. Japan Acad.*, 63, 390–391 (1987).
- [25] _____ : On the projection formula for tensor products of holonomic systems and its applications, preprint. 数理解析研究所講究録
- [26] Tajima, S. and Uchida, M. : 特異点除去に対応する de Rham系の積分の計算, 数理解析研究所講究録 693, 41–68 (1989).
- [27] Vilonen, K. : Intersection homology D-Module on local complete intersections with isolated singularities, *Invent. math.*, 81, 107–114 (1985).
- [28] Yger, A. : *Formules de division et prolongement méromorphe*, Springer Lecture Notes in Math., 1295, 226–283 (1987).