

Periodic solutions of a singular Hamiltonian system of 2-body type

Kazunaga Tanaka (田中和永)

Department of Mathematics, School of Science, Nagoya University
 Chikusa-ku, Nagoya 464, JAPAN

0. Introduction and results

In this short note, we study the existence of periodic solutions of a Hamiltonian system

$$\ddot{q} + \nabla V(q, t) = 0, \quad (HS)$$

where $q = (q_1, \dots, q_N) \in \mathbf{R}^N$ ($N \geq 3$) and $V(q, t) : \mathbf{R}^N \times \mathbf{R} \rightarrow \mathbf{R}$ is a given potential. We deal with the case where a potential has a singularity and is related to 2-body problem.

More precisely, we assume $V(q, t)$ satisfies

- (V1) $V(q, t) \in C^2((\mathbf{R}^N \setminus \{0\}) \times \mathbf{R}, \mathbf{R})$ is T -periodic in t ;
- (V2) $V(q, t) < 0$ and $V(q, t), \nabla V(q, t) \rightarrow 0$ as $|q| \rightarrow \infty$ uniformly in t ;
- (V3) $V(q, t)$ is of a form:

$$V(q, t) = -\frac{1}{|q|^\alpha} + W(q, t),$$

where $\alpha > 0$ and $W(q, t) \in C^2((\mathbf{R}^N \setminus \{0\}) \times \mathbf{R}, \mathbf{R})$ satisfies

$$\begin{aligned} &|q|^\alpha W(q, t), |q|^{\alpha+1} \nabla W(q, t), |q|^{\alpha+2} \nabla^2 W(q, t), \\ &|q|^\alpha W_t(q, t) \rightarrow 0 \quad \text{as } q \rightarrow 0 \text{ uniformly in } t. \end{aligned}$$

We consider the following two problems:

- (i) *Prescribed Period Problem (PP)*: For a given $T > 0$, we study the existence of T -periodic solutions of (HS), i.e., solutions of (HS) such that

$$q(t+T) = q(t) \quad \text{for all } t. \quad (HS.P)$$

- (ii) *Prescribed Energy Problem (PE)*: Assume V is independent of t . For a given $H \in \mathbf{R}$, we study the existence of periodic solutions of (HS) such that

$$\frac{1}{2} |\dot{q}(t)|^2 + V(q(t)) = H \quad \text{for all } t. \quad (HS.E)$$

(Here we do not fix the period of $q(t)$.)

We study via variational methods these problems. Recently it is observed that the order α of the singularity of $V(q, t)$ at $q = 0$ plays an important role for the existence of

periodic solutions for both of problems. We consider the following cases separately; (i) the *strong force* case $\alpha \geq 2$ for (PP) and $\alpha > 2$ for (PE) (ii) the *weak force* case $\alpha \in (0, 2)$.

For the Prescribed Period problem (PP), we use the following variational formulation. Let $E = \{q \in H_{loc}^1(\mathbf{R}, \mathbf{R}^N); q(t) \text{ is } T\text{-periodic in } t\}$ is a space of T -periodic functions with norm $\|q\|_E^2 = \int_0^T [|\dot{q}(t)|^2 + |q(t)|^2] dt$ and set

$$\Lambda = \{q \in E; q(t) \neq 0 \text{ for all } t\}.$$

We define the functional $I(q) : \Lambda \rightarrow \mathbf{R}$ by

$$I(q) = \int_0^T \left[\frac{1}{2} |\dot{q}|^2 - V(q(t), t) \right] dt.$$

Then there is one-to-one correspondence between critical points $q \in \Lambda$ of $I(q)$ and T -periodic solutions of (HS), (HS.P). Therefore we try to find critical points of $I(q)$.

If (V1)–(V3) holds and $\alpha \geq 2$, more generally, under the conditions of (V2)–(V3) and (V1') $V(q, t) \in C^1((\mathbf{R}^N \setminus \{0\}) \times \mathbf{R}, \mathbf{R})$ is T -periodic in t and the following strong force condition (SF) of Gordon [Go]:

(SF) there is a neighborhood Ω of 0 and $U(q) \in C^1(\Omega \setminus \{0\}, \mathbf{R})$ such that

$$\begin{aligned} U(q) &\rightarrow \infty, & q &\rightarrow 0, \\ -V(q, t) &\geq |\nabla U(q)|^2 & \text{for all } q \in \Omega \setminus \{0\} \text{ and } t, \end{aligned}$$

we can see the functional $I(q)$ satisfies the Palais-Smale condition and we can apply min-max methods to obtain critical points of $I(q)$. We refer to [BR, Gr, AC1]. Our main purpose is to study the weak force case $\alpha \in (0, 2)$. We remark that the Palais-Smale condition does not hold in this case. Our result is as follows:

Theorem 0.1 ([T2]). *Suppose (V1)–(V3) and $\alpha \in (1, 2)$. Then the prescribed period problem (HS), (HS.P) possesses at least one periodic solution.*

For the Prescribed energy problem (PE), we can expect the existence of periodic solutions only under the situations

- (i) $H > 0$ if $\alpha > 2$, or
- (i) $H < 0$ if $\alpha \in (0, 2)$.

Actually, if $V(q) = -\frac{1}{|q|^\alpha}$, we can easily see that periodic solutions of (HS), (HS.E) exist if and only if (i) or (ii) holds. In the strong force case $\alpha > 2$, the Palais-Smale condition holds under additional assumptions and we refer to Ambrosetti and Coti Zelati [AC2] for the existence result. We study the case $\alpha \in (0, 2)$. Here we assume

(V4) there is $\bar{\alpha} \in (0, \alpha]$ such that

$$\nabla V(q)q \geq -\bar{\alpha}V(q) \quad \text{for all } q \in \mathbf{R}^N \setminus \{0\}$$

in addition to (V1)–(V3).

Theorem 0.2 ([T3]). Suppose V is independent of t , $H < 0$ and (V1)–(V4). Moreover assume $\alpha \in (1, 2)$ if $N \geq 4$ and $\alpha \in (4/3, 2)$ if $N = 3$. Then the prescribed energy problem (HS), (HS.E) possesses at least one periodic solution.

We remark that in case of weak force the existence of *generalized solutions*, which may enter the singularity 0, is obtained by [BR] for the prescribed period problem (PP) and by [AC2] for the prescribed energy problem (PE). We also remark the result very closely related to Theorem 0.1 is obtained by Coti Zelati and Serra [CS] independently.

In what follows, we sketch outline of the proof of Theorem 0.1. The proof of Theorem 0.2 is done essentially in a same way (but more complicated) to Theorem 0.1 and we refer to [T3].

1. Perturbed functionals

We take the following approach, which is used by [BR] first time.

1° First we introduce a perturbed potential $V_\epsilon(q, t) = V(q, t) - \frac{\epsilon}{|q|^2}$. The corresponding functional

$$\begin{aligned} I_\epsilon(q) &= \int_0^T \left[\frac{1}{2} |q|^2 - V_\epsilon(q, t) \right] dt \\ &= \int_0^T \left[\frac{1}{2} |q|^2 - V(q, t) + \frac{\epsilon}{|q|^2} \right] dt \end{aligned}$$

satisfies a variant of the Palais-Smale condition and we can apply a minimax method of [BR] to get approximate solution $q_\epsilon(t)$ for each $\epsilon \in (0, 1]$.

2° Second we try to pass to the limit as $\epsilon \rightarrow 0$ and we try to obtain a solution as a limit of $q_\epsilon(t)$

More precisely, we use the following minimax method; we set

$$\Gamma = \{\gamma \in C(S^{N-2}, \Lambda); \deg \tilde{\gamma} \neq 0\} \quad (1.1)$$

where $\tilde{\gamma} : S^1 \times S^{N-2} \simeq ([0, T]/\{0, T\}) \times S^{N-2} \rightarrow S^{N-1}$ is defined by

$$\tilde{\gamma}(t, x) = \frac{\gamma(x)(t)}{|\gamma(x)(t)|}$$

and $\deg \tilde{\gamma}$ denote the Brower degree of $\tilde{\gamma}$. We define

$$b_\epsilon = \inf_{\gamma \in \Gamma} \max_{x \in S^{N-2}} I_\epsilon(\gamma(x)). \quad (1.2)$$

Then we have

Proposition 1.1 ([BR]). For any $\epsilon \in (0, 1]$, b_ϵ is a critical value of $I_\epsilon(q)$. That is, there is a critical point $q_\epsilon(t)$ of $I_\epsilon(q)$ such that $I_\epsilon(q_\epsilon) = b_\epsilon$. Moreover, there are constants $M, m > 0$ independent of $\epsilon \in (0, 1]$ such that

$$m \leq b_\epsilon \leq M \quad \text{for all } \epsilon \in (0, 1]. \quad (1.3)$$

Using the uniform estimate (1.3), we can get

Proposition 1.2 ([BR]). There is a constant $C > 0$ independent of $\epsilon \in (0, 1]$ such that

$$\|q_\epsilon\|_E \leq C \quad \text{for all } \epsilon \in (0, 1].$$

Therefore we can choose a subsequence — still we denote by $\epsilon \rightarrow 0$ — such that $q_\epsilon \rightarrow q_0 \in E$ weakly in E and strongly in L^∞ . If $q_0(t) \neq 0$ for all t , in other words, if $q_0 \in \Lambda$, we can easily see $q_0(t)$ is a periodic solution of (HS), (HS.P). The difficulty is to prove $q_0 \in \Lambda$.

Even if $q_0 \notin \Lambda$, we can see

- (i) Set $D = \{t; q_0(t) = 0\}$. Then $\text{meas } D = 0$;
- (ii) $q_0(t) \in C^2(\mathbf{R} \setminus D, \mathbf{R}^N) \cap C(\mathbf{R}, \mathbf{R}^N)$;
- (iii) $q_0(t)$ satisfies (HS) in $\mathbf{R} \setminus D$.

Bahri and Rabinowitz [BR] called such a limit function $q_0(t)$ *generalized solution* of (HS), (HS.P). They constructed generalized solutions under the conditions (V1'), (V2) and (V3') $V(q, t) \rightarrow -\infty$ as $q \rightarrow 0$ uniformly in t .

To prove $q_0(t)$ does not enter the singularity 0, we use a combination of a re-scaling argument and an estimate of Morse indices.

2. Re-scaling argument

Suppose $q_0(t)$ enters the singularity 0 at $t_0 \in (0, T]$, i.e., $q_0(t_0) = 0$. Then there is a sequence $t_\epsilon \in (0, T]$ such that

- 1° $t_\epsilon \rightarrow t_0$;
- 2° $|q_\epsilon(t)|$ takes its local minimum at t_ϵ .

Case 1: First we study the behavior of $q_\epsilon(t)$ near the singularity 0 more precisely via a re-scaling argument. We set

$$\begin{aligned} \delta_\epsilon &= |q_\epsilon(t_\epsilon)|, \\ x_\epsilon(s) &= \delta_\epsilon^{-1} q_\epsilon(\delta_\epsilon^{(\alpha+2)/2} s + t_\epsilon). \end{aligned}$$

Then $x_\epsilon(s)$ satisfies $|x_\epsilon(0)| = 1$ and

$$\ddot{x}_\epsilon + \frac{\alpha x_\epsilon}{|x_\epsilon|^{\alpha+2}} + \delta_\epsilon^{\alpha+1} \nabla W(\delta_\epsilon x_\epsilon(s), \delta_\epsilon^{(\alpha+2)/2} s + t_\epsilon) + \frac{2\epsilon}{\delta_\epsilon^{2-\alpha}} \frac{x_\epsilon}{|x_\epsilon|^4} = 0.$$

We study the behavior of $x_\epsilon(s)$ instead of $q_\epsilon(t)$.

After taking a suitable subsequence — still we denote by ϵ —, we may assume that

$$d = \lim_{\epsilon \rightarrow 0} \frac{\epsilon}{\delta_\epsilon^{2-\alpha}} \in [0, \infty] \quad (2.1)$$

exists. We consider the following two cases separately.

Case 1: $d < \infty$;

Case 2: $d = \infty$.

Case 1: First we deal with Case 1.

Proposition 2.1. *Suppose $d < \infty$. After taking a subsequence — still denoted by ϵ —, $x_\epsilon(s)$ converges to a function $y_{\alpha,d}(s)$ in $C_{loc}^2(\mathbf{R}, \mathbf{R}^N)$, where $y_{\alpha,d}(s)$ is the solution of*

$$\begin{aligned} \ddot{y} + \frac{\alpha y}{|y|^{\alpha+2}} + \frac{2dy}{|y|^4} &= 0, \quad \text{in } \mathbf{R}, \\ y(0) &= e_1, \\ \dot{y}(0) &= \sqrt{2(1+d)}e_2. \end{aligned} \quad (2.2)$$

Here, $e_1, e_2, \dots, e_N \in \mathbf{R}^N$ are vectors satisfying $e_i \cdot e_j = \delta_{ij}$. ■

Case 2: In this case, we introduce another re-scaled function

$$z_\epsilon(s) = \delta_\epsilon^{-1} q_\epsilon(\epsilon^{-1/2} \delta_\epsilon^2 s + t_\epsilon).$$

Then $z_\epsilon(s)$ satisfies

$$\ddot{z}_\epsilon + \frac{\alpha \delta_\epsilon^{2-\alpha}}{\epsilon} \frac{z_\epsilon}{|z_\epsilon|^{\alpha+2}} + \frac{\alpha \delta_\epsilon^{2-\alpha}}{\epsilon} \delta_\epsilon^{\alpha+1} \nabla W(\delta_\epsilon z_\epsilon, \epsilon^{-1/2} \delta_\epsilon^2 s + t_\epsilon) + \frac{2z_\epsilon}{|z_\epsilon|^4} = 0.$$

We have

Proposition 2.2. *Suppose $d = \infty$. Then, after taking a subsequence — still denoted by ϵ —, we have*

$$z_\epsilon(s) \rightarrow z_0(s) = e_1 \cos \sqrt{2}s + e_2 \sin \sqrt{2}s \quad \text{in } C_{loc}^2(\mathbf{R}, \mathbf{R}^N).$$

Here, $e_1, e_2, \dots, e_N \in \mathbf{R}^N$ are vectors satisfying $e_i \cdot e_j = \delta_{ij}$. ■

We remark $z_0(s)$ is a solution of

$$\begin{aligned} \ddot{z} + \frac{2z}{|z|^4} &= 0, \quad \text{in } \mathbf{R}, \\ z(0) &= e_1, \\ \dot{z}(0) &= \sqrt{2}e_2. \end{aligned}$$

3. Estimates of Morse index

Using the propositions 2.1 and 2.2, we have the following estimate of Morse indices.

Proposition 3.1. *Suppose $q_0(t)$ enters the singularity 0 and set*

$$\nu = \#\{t \in (0, t]; q_0(t) = 0\}.$$

Then

$$\liminf_{\epsilon \rightarrow 0} \text{index } I'_\epsilon(q_\epsilon) \geq (N - 2)i(\alpha)\nu \quad (3.1)$$

where

$$i(\alpha) = \max\{k \in \mathbf{N}; k < \frac{2}{2 - \alpha}\}. \quad \blacksquare$$

Before we sketch the proof of Proposition 3.1, we give a proof of Theorem 0.1.

Proof of Theorem 0.1. First we remark that the following estimate of Morse index follows from the minimax characterization (1.1)–(1.2) of b_ϵ .

Proposition 3.2 (c.f.[BL, LS, T1]). *$q_\epsilon(t) \in \Lambda$ satisfies*

$$\text{index } I'_\epsilon(q_\epsilon) \leq N - 2 \quad \text{for all } \epsilon \in (0, 1]. \quad (3.2)$$

Comparing (3.1) and (3.2), we have

$$i(\alpha)\nu \leq 1. \quad (3.3)$$

Since $i(\alpha) \geq 2$ for $\alpha \in (1, 2)$ and $i(\alpha) = 1$ for $\alpha \in (0, 1]$, we find

$$\begin{aligned} \nu &= 0, & \text{if } \alpha \in (1, 2), \\ \nu &\leq 1, & \text{if } \alpha \in (0, 1]. \end{aligned}$$

Therefore in case $\alpha \in (1, 2)$, we obtain $q_0(t) \neq 0$ for all t and it is a classical solution. ■

Sketch of the proof of Proposition 3.1. Suppose $q_0(t_0) = 0$ and choose $t_\epsilon \in (0, T]$ as above. We deal with only the Case 1: $d < \infty$. The Case 2: $d = 0$ can be treated similarly. For $L > 0$, $\varphi(s) \in H_0^1(-L, L; \mathbf{R})$ and $j = 1, 2, \dots, N$, we set

$$h_{\epsilon,j}(t) = \delta_\epsilon \varphi(\delta_\epsilon^{-(\alpha+2)/2}(t - t_\epsilon))e_j.$$

After the change of variable, we take a limit as $\epsilon \rightarrow 0$ and obtain

$$\begin{aligned} & \delta_\epsilon^{-(2-\alpha)/2} I_\epsilon''(q_\epsilon)(h_{\epsilon,j}, h_{\epsilon,j}) \\ & \rightarrow \int_{-L}^L \left[|\dot{\varphi}|^2 - \frac{\alpha |\varphi|^2}{|y_{\alpha,d}|^{\alpha+2}} + \frac{\alpha(\alpha+2)(y_{\alpha,d}, e_j)^2 |\varphi|^2}{|y_{\alpha,d}|^{\alpha+4}} \right. \\ & \quad \left. - \frac{2d |\varphi|^2}{|y_{\alpha,d}|^4} + \frac{8d(y_{\alpha,d}, e_j)^2 |\varphi|^2}{|y_{\alpha,d}|^6} \right] ds. \end{aligned}$$

Recalling $y_{\alpha,d}(s) \in \text{span}\{e_1, e_2\}$ for all s , we can see

$$\liminf_{\epsilon \rightarrow 0} \text{index } I_\epsilon''(q_\epsilon) \geq (N-2)i(\alpha, d) \quad (3.4)$$

where

$$i(\alpha, d) = \sup_{L>0} \left(\begin{array}{l} \text{the number of negative eigenvalues} \\ \text{of the following eigenvalue problem:} \\ -\ddot{\varphi} - \left(\frac{\alpha}{|y_{\alpha,d}|^{\alpha+2}} + \frac{2d}{|y_{\alpha,d}|^4} \right) \varphi = 0, \\ \varphi(L) = \varphi(-L) = 0. \end{array} \right).$$

We repeat the above argument at all other $t'_0 \in (0, T]$ such that $q_0(t'_0) = 0$ and we find

$$\liminf_{\epsilon \rightarrow 0} \text{index } I_\epsilon''(q_\epsilon) \geq (N-2)i(\alpha)\nu$$

where

$$i(\alpha) = \min_{d \geq 0} i(\alpha, d).$$

Now Proposition 3.1 follows from the following proposition. ■

Proposition 3.3.

$$i(\alpha, d) = \max\{k \in \mathbf{N}; k < \frac{2\sqrt{1+d}}{2-\alpha}\}. \quad (3.5)$$

Thus $i(\alpha) = \max\{k \in \mathbf{N}; k < \frac{2}{2-\alpha}\}$.

Proof. The case $d = 0$ is proved in [T2]. The case $d > 0$ is proved similarly. The key of the proof is the Sturm comparison theorem and the following property of $y_{\alpha,d}(s)$. We use the polar coordinate and write

$$y_{\alpha,d}(s) = r(s)(e_1 \cos \theta(s) + e_2 \sin \theta(s))$$

where $r(s) > 0$ and $\theta(s) \in \mathbf{R}$ with $\theta(0) = 0$. Then we have

- (i) $s \dot{r}(s) > 0$ for all $s \neq 0$ and $r(s) \rightarrow \infty$ and $s \rightarrow \pm\infty$;
- (ii) $\dot{\theta}(s) > 0$ for all s ;
- (iii) $\theta(s) \rightarrow \pm \frac{2\pi\sqrt{1+d}}{2-\alpha}$ as $s \rightarrow \pm\infty$.

■

4. Remarks

In case $\alpha \in (0, 1]$, it seems that the existence of classical periodic solutions is not known. However by (3.3) we can see there is a generalized solution of (HS), (HS.P) that enters at most one time in its period. By (3.4) and (3.5), we also have

$$d \leq (2 - \alpha)^2 - 1 \quad (4.1)$$

where d is defined in (2.1).

We get the following additional information under slightly stronger conditions: (V1), (V2) and

(V3'') $V(q, t)$ is of a form:

$$V(q, t) = -\frac{1}{|q|^\alpha} + W(q, t),$$

where $\alpha > 0$ and $W(q, t) \in C^2((\mathbf{R}^N \setminus \{0\}) \times \mathbf{R}, \mathbf{R})$ satisfies

$$\begin{aligned} & |q|^{\alpha-\rho} W(q, t), |q|^{\alpha-\rho+1} \nabla W(q, t), |q|^{\alpha-\rho+2} \nabla^2 W(q, t), \\ & |q|^{\alpha-\rho} W_t(q, t) \rightarrow 0 \quad \text{as } q \rightarrow 0 \text{ uniformly in } t \end{aligned}$$

for some $\rho \in (0, \alpha)$.

We assume $q_0(t)$ is a generalized solution such that $q_0(t_0) = 0$. Beaulieu [B] proved that the limits

$$a_\pm = \lim_{t \rightarrow t_0 \pm 0} \frac{q_0(t)}{|q_0(t)|} \in S^{N-1}$$

exist. We have

Theorem 4.1 ([T4]). Assume (V1), (V2), (V3'') and let $q_\epsilon(t)$ be a critical point of $I_\epsilon(q)$ which is obtained through a minimax method (1.1)–(1.2). Suppose $q_0(t) = \lim_{\epsilon \rightarrow 0} q_\epsilon(t)$ is a generalized solution such that $q_0(t_0) = 0$ and let $a_\pm = \lim_{t \rightarrow t_0 \pm 0} \frac{q_0(t)}{|q_0(t)|} \in S^{N-1}$. Then we have

$$\text{the angle between } a_+ \text{ and } a_- = \frac{2\pi\sqrt{1+d}}{2-\alpha} \text{ modulo } 2\pi$$

where $d \in [0, (2 - \alpha)^2 - 1]$ is defined in (2.1). ■

In particular, in case $\alpha = 1$ we have $d = 0$ and $a_+ = a_-$.

References

- [AC1] A. Ambrosetti and V. Coti Zelati, Critical points with lack of compactness and applications to singular dynamical system, *Annali Mat. Pura Appl.* **149** (1987), 237–259.
- [AC2] — and —, Closed orbits of fixed energy for singular Hamiltonian systems, *Arch. Rat. Mech. Anal.* **112** (1990), 339–362.
- [B] A. Beaulieu, Etude de solutions generalisees pour un systeme hamiltonien avec potentiel singulier, *preprint*.
- [BL] A. Bahri and P.L. Lions, Morse index of some min-max critical points. I. Application to multiplicity results, *Comm. Pure Appl. Math.* **41** (1988), 1027–1037.
- [BR] A. Bahri and P.H. Rabinowitz, A minimax method for a class of Hamiltonian systems with singular potentials, *J. Funct. Anal.* **82** (1989), 412–428.
- [C] V. Coti Zelati, Periodic solutions for a class of planar, singular dynamical systems, *J. Math. Pure Appl.* **68** (1989), 109–119.
- [CS1] V. Coti Zelati and E. Serra, Collisions and non-collisions solutions for a class of Keplerian-like dynamical systems, *preprint*.
- [CS2] — and —, Multiple brake orbits for some classes of Hamiltonian systems, *preprint*.
- [Go] W.B. Gordon, Conservative dynamical systems involving strong forces, *Trans. Amer. Math. Soc.* **204** (1975), 113–135.
- [Gr] C. Greco, Periodic solutions of a class of singular Hamiltonian systems, *Nonlinear Analysis; T.M.A.* **12** (1988), 259–269.
- [LS] A. C. Lazer and S. Solimini, Nontrivial solutions of operator equations and Morse indices of critical points of min-max type, *Nonlinear Analysis: T. M. A.* **12** (1988), 761–775.
- [R] P. H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, *CBMS Regional Conf. Ser. in Math.* **65** Amer. Math. Soc., Providence 1986.
- [ST] E. Serra and S. Terracini, Noncollision solutions to some singular minimization problems with Keplerian-like potentials, *preprint*.
- [T1] K. Tanaka, Morse indices at critical points related to the symmetric mountain pass theorem and applications, *Comm. Partial Diff. Eq.* **14** (1989), 99–128.
- [T2] —, Non-collision solutions for a second order singular Hamiltonian system with weak force, *Ann. Inst. Henri Poincaré, Analyse nonlinéaire*, (to appear).
- [T3] —, A prescribed energy problem for a singular Hamiltonian system with a weak force, *J. Funct. Anal.*, (to appear).
- [T4] —, *preprint*.