

## ON MAXIMUM MODULUS OF ANALYTIC FUNCTIONS

SHIGEYOSHI OWA (近畿大学・理工)

WEI-QI YANG (北京理工大学)

ABSTRACT. Let  $A$  be the class of functions  $f(z)$  analytic in  $|z| < R$  with  $f(0) = 1$ . The object of the present paper is to investigate analyticity conditions of  $M(r)$  which is the maximum modulus of  $f(z)$  in  $A$ . Our results proved here provide a generalization of some results due to W. K. Hayman (J. Analyse Math. 1(1951), 135 - 154).

1. INTRODUCTION. Let  $A$  be the class of functions  $f(z)$  analytic in the disk  $|z| < R$  with  $f(0) = 1$ . The maximum modulus  $M(r)$  given by

$$(1.1) \quad M(r) = \max\{f(z) \in A: z = re^{i\theta}, 0 < r < R, 0 \leq \theta \leq 2\pi\}$$

was investigated by Hadamard [1], Hayman [3], et al. (see [2], [4]).

The curves from the origin to  $|z| = R$  consisting of all points  $z = re^{i\theta}$  which satisfy

$$(1.2) \quad \frac{\partial |f(re^{i\theta})|}{\partial \theta} = 0$$

are called quasi-extremal curves of  $f(z) \in A$ . Some of quasi-extremal curves are called extremal curves of  $f(z) \in A$  if

$$(1.3) \quad |f(re^{i\theta})| = M(r).$$

In the present paper, we show analyticity conditions of the maximum modulus  $M(r)$  of  $f(z) \in A$ .

2. MAIN RESULTS. In order to discuss our problems, we have to use the following lemmas.

LEMMA 1 (Hayman [3]). Let  $f(z)$  be analytic on the circle  $|z| = r$ . If it satisfies

$$(2.1) \quad \frac{\partial |f(re^{i\theta_0})|}{\partial \theta} = 0, \quad f(re^{i\theta_0}) \neq 0,$$

then  $g(z) = zf'(z)/f(z)$  is real at the point  $z_0 = re^{i\theta_0}$  and

$$(2.2) \quad g(z_0) = r \frac{d}{dr} \log |f(\alpha(r))|,$$

where  $\alpha(t)$  is the parametric representation with  $t = |\alpha(t)|$  of any given curve which is analytic at a point  $z_0$  and non-tangential with the circle  $|z| = r$ .

LEMMA 2. Let  $\Gamma$  be an analytic curve and intersect every circle  $|z| = r$  ( $0 < r \leq R$ ) at only one point. Then  $\Gamma$  is of the form

$$(2.3) \quad \Gamma = \{z: z = re^{i\theta(r)}, 0 \leq r \leq R\},$$

where  $\theta(r)$  is a real analytic function.

PROOF. Since  $\Gamma$  is a simple analytic curve, there exists a parametric representation of  $\Gamma$

$$(2.4) \quad \Gamma = \{z: z = \alpha(t), 0 \leq t \leq 1\}$$

such that  $\alpha(t)$  is an analytic injection with  $\alpha(0) = 0$ . Letting

$$(2.5) \quad \log \frac{\alpha(t)}{t} = \phi(t) + i\psi(t),$$

we see that  $\phi(t)$  and  $\psi(t)$  are real analytic. Further, we know that

$$(2.6) \quad r = te^{\phi(t)}$$

is exactly increasing in  $t \in [0, 1]$ , and its inverse function is  $t = t(r)$ .

Defining  $\theta(r)$  by

$$(2.7) \quad \theta(r) = \psi(t(r)),$$

we obtain the representation (2.4) of  $\Gamma$ .

Now we prove

**THEOREM 1.** If there exists a quasi-extremal curve

$$\Gamma = \{z: z = \alpha(r), 0 \leq r < R\}$$

of  $f(z)$  which is in the class  $A$  with modulus  $r$  such that  $\forall z_0 \in \Gamma$ ,  $g(z_0)$  is the maximum real value of  $g(z) = zf'(z)/f(z)$  on the circle  $|z| = |z_0|$ , then  $\Gamma$  is an extremal curve of  $f(z)$  and  $M(r)$  is real analytic on  $[0, R)$ .

**PROOF.** We may assume that  $f(z)$  is analytic on the closed disk  $|z| \leq R$ , other wise we consider a closed subdisk of  $|z| < R$ . Divide the disk  $|z| < R$  into several annular domain  $D_k$  given by

$$D_k = \{z: r_{k-1} < |z| < r_k; r_0 = 0; r_n = R; k = 1, 2, 3, \dots, n\}$$

such that  $f(z) \neq 0$  in every  $D_k$ . It is known that there are  $2^{n_k}$  quasi-extremal arcs

$$C_{kj} = \{z: z = \alpha_{kj}(r), j = 1, 2, 3, \dots, 2^{n_k}\}$$

of  $f(z)$  in  $D_k$ , which are analytic in  $(r_{k-1}, r_k)$ . Applying Lemma 1, we know that

$$(2.8) \quad g(\alpha(r)) = r \frac{d}{dr} \log |f(\alpha(r))|$$

and

$$(2.9) \quad g(\alpha_{k,j}(r)) = r \frac{d}{dr} \log |f(\alpha_{k,j}(r))|.$$

Then we have

$$(2.10) \quad \log |f(\alpha(r))| = \int_0^r g(\alpha(t)) \frac{dt}{t} \quad (0 \leq r < R)$$

and

$$(2.11) \quad \log |f(\alpha_{1j}(r))| = \int_0^r g(\alpha_{1j}(t)) \frac{dt}{t} \quad (0 \leq r < r_1),$$

which give

$$(2.12) \quad |f(\alpha_{1j}(r))| \leq |f(\alpha(r))| \quad (0 \leq r < r_1).$$

Also the inequality (2.12) is still true for  $r = r_1$ . Therefore we obtain

$$(2.13) \quad M(r) = |f(\alpha(r))| \quad (0 \leq r \leq r_1).$$

For  $r_1 < r' < r'' < r_2$ , it follows from (2.9) and (2.9) that

$$(2.14) \quad \frac{|f(\alpha_{2j}(r''))|}{|f(\alpha_{2j}(r'))|} = \exp \left\{ \int_{r'}^{r''} g(\alpha_{2j}(r)) \frac{dt}{t} \right\} \\ \leq \exp \left\{ \int_{r'}^{r''} g(\alpha(t)) \frac{dt}{t} \right\} \\ = \frac{|f(\alpha(r''))|}{|f(\alpha(r'))|},$$

so

$$(2.15) \quad \frac{|f(\alpha_{2j}(r''))|}{|f(\alpha(r''))|} \leq \frac{|f(\alpha_{2j}(r'))|}{|f(\alpha(r'))|}.$$

Letting  $r' \rightarrow r_1$  in (2.15), we see that

$$(2.16) \quad |f(\alpha_{2j}(r''))| \leq |f(\alpha(r''))|,$$

which proves that

$$(2.17) \quad M(r) = |f(\alpha(r))| \quad (0 \leq r \leq r_2).$$

To do the above process again and again, it follows that  $\Gamma$  is an extremal curve of  $f(z)$  and  $M(r)$  is real analytic on  $[0, R)$ .

Next, we derive

**THEOREM 2.** If  $g(z)$  is analytic and univalent in  $|z| < R$ , and if there exists a quasi-extremal curve of  $f(z) \in \Lambda$  which intersects every circle  $|z| = r$  ( $0 < r < R$ ) at only one point, then  $M(r)$  is real analytic.

**PROOF.** Because  $g(z)$  maps every quasi-extremal curve of  $f(z)$  onto

a half open interval on the real axis with the origin as an end point, the intersection of the real axis and the range of  $g(z)$  is an open interval  $L$  which is divided by the origin into two intervals  $L_1$  (left hand interval) and  $L_2$  (right hand interval). These inverse images  $\Gamma_1$  and  $\Gamma_2$  are two quasi-extremal curves of  $f(z)$  which are analytic curves.

Using Lemma 1 and Lemma 2, we have

$$(2.18) \quad \Gamma_j = \{z: z = re^{i\theta_j(r)}, j = 1, 2\},$$

where  $\theta_j(r)$  are real analytic and

$$(2.19) \quad g(re^{i\theta_j(r)}) = r \frac{d}{dr} \log |f(re^{i\theta_j(r)})|.$$

Then we have

$$(2.20) \quad |f(re^{i\theta_j(r)})| = \exp \left\{ \int_0^r g(te^{i\theta_j(t)}) \frac{dt}{t} \right\}.$$

Since

$$(2.21) \quad g(re^{i\theta_1(r)}) < 0 < g(re^{i\theta_2(r)}),$$

it follows that

$$(2.22) \quad M(r) = |f(re^{i\theta_2(r)})|$$

which is real analytic.

Further, we prove

**THEOREM 3.** Under the assumption in Theorem 2, there exists a certain open disk  $|z| < R_1$  such that  $M(r)$  is the maximum modulus of its complex analytic extension  $M(z)$  in  $|z| < R_1$ .

**PROOF.** Let  $D$  be the domain in which  $M(z)$  is analytic. Then the origin is contained in  $D$ , and there exists a open disk  $|z| < R_1$  in  $D$  such that, for any  $|z| < R_1$ ,

$$(2.23) \quad |ze^{i\theta_2(z)}| < R,$$

where  $\theta_2(z)$  is the complex analytic extension of  $\theta_2(r)$ . From (2.19), (2.21), and (2.22), we know that

$$(2.24) \quad \mu(r) = r \frac{M'(r)}{M(r)} = g(re^{i\theta_2(r)})$$

is the maximum real value of  $g(z)$  on the circle  $|z| = r$ , and  $\mu(r)$  is exactly increasing and real analytic. It follows from (2.24) that

$$(2.25) \quad \mu(z) = z \frac{M'(z)}{M(z)} = g(ze^{i\theta_2(z)})$$

and

$$(2.26) \quad \mu(\bar{z}) = \overline{\mu(z)}, \quad M(\bar{z}) = \overline{M(z)}, \quad \theta_2(\bar{z}) = \overline{\theta_2(z)}.$$

Suppose that  $M(r)$  is not the maximum modulus of  $M(z)$  in  $|z| < R_1$ , that is, there exists  $r_0$  ( $0 < r_0 < R_1$ ) such that  $M(r_0)$  is not the maximum modulus of  $M(z)$  on the circle  $|z| = r_0$ . Then there is a point  $z_0 = r_0 e^{i\theta_0}$  such that

$$(2.27) \quad \mu(\bar{z}_0) = \mu(z_0) > \mu(r_0)$$

by Theorem 1. This gives us that

$$(2.28) \quad g(\bar{z}_0 e^{i\theta_2(\bar{z}_0)}) = g(z_0 e^{i\theta_2(z_0)}) > \mu(r_0).$$

But, since

$$(2.29) \quad |\bar{z}_0 re^{i\theta_2(\bar{z}_0)}| |z_0 e^{i\theta_2(z_0)}| = r_0^2 |e^{i(\theta_2(\bar{z}_0) + \theta_2(z_0))}| = r_0^2,$$

(2.28) contradicts the fact that  $\mu(r)$  is the maximum real value on the circle  $|z| = r$  and exactly increasing. This completes the proof of Theorem 3.

**ACKNOWLEDGMENTS.** This work of the first author was supported, in part, by the Japanese Ministry of Education, Science and Culture under Grant-in-Aid for General Scientific Research.

## REFERENCES

- [1] M. Hadamard, Sur les fonctions entières, Bull. Soc. Math. France 24(1896), 186 - 187.
- [2] M. O. Bulmenthal, Sur le mode de croissance des fonctions entières, Bull. Soc. Math. France 35(1907), 213 - 232.
- [3] W. K. Hayman, A characterization of the maximum modulus of functions regular at the origin, J. Analyse Math. 1(1951), 135 - 154.
- [4] B. Anderson, On equivalent analytic functions, Arkiv Math. 1(1949), 77 - 92.

S. Owa: Department of Mathematics  
Kinki University  
Higashi-Osaka, Osaka 577  
Japan

W. Yang: Department of Mathematics  
Beijing Institute of Technology  
Beijing 100081  
People's Republic of China