

ON THE STRUCTURE OF CERTAIN CLASS OF VOLTERRA INTEGRAL OPERATORS AND ESTIMATES OF APPROXIMATION NUMBERS

V.D. Stepanov¹

*Computer Center of the Far-Eastern Branch of the Russian Academy of Sciences
Khabarovsk, Russia*

Introduction.

We study the integral operators of the form

$$Kf(x) = v(x) \int_0^x k(x, y)u(y)f(y) dy, \quad x > 0, \quad (1)$$

where the weight real functions $v(t)$ and $u(t)$ are locally integrable and the kernel $k(x, y) \geq 0$ satisfies the following condition: there exists a constant $D \geq 1$ such that

$$D^{-1}(k(x, y) + k(y, z)) \leq k(x, z) \leq D(k(x, y) + k(y, z)), \quad x > y > z \geq 0, \quad (2)$$

where D does not depend on x, y, z .

A few standard examples of the kernel $k(x, y) \geq 0$ satisfying (2) are

(1) $k(x, y) = (x - y)^\alpha$, $\alpha \geq 0$,

(2) $k(x, y) = \log^\beta(1 + x - y)$, $k(x, y) = \log^\beta\left(\frac{x}{y}\right)$; $\beta \geq 0$,

(3) $k(x, y) = \left(\int_y^x h(s) ds\right)^\alpha$, $\alpha \geq 0$, $h(s) \geq 0$,

as well as their various combinations. However, for instance, the kernel of the first kind with negative value of α does not satisfy (2).

The operators (1) with kernels satisfying (2) were intensively studied during the last decade and many authors made contributions in this topic, e.g. see the author's survey [St₁] with history and literature given there, and where the $L^p - L^q$ mapping properties of (1) were investigated.

Here, in Section 1, we give further extension of some characterization results of [St₁] on the Banach function spaces for the following problems:

(B) Boundedness,

(C) Compactness and measure of non-compactness.

In Section 2 we give the more detailing structural results for the operators (1), when they are compact in the Lebesgue spaces, namely

(S) Two-sided estimates of the Schatten-von Neumann ideal norms,

(N) Asymptotic behaviour and two-sided estimates of p -norms of approximation numbers.

1. Problems (B) and (C) in Banach function spaces.

Assume X and Y be two Banach spaces of measurable functions defined on $\mathbb{R}^+ = (0, \infty)$. First we consider the problem of the boundedness $K : X \rightarrow Y$ for the integral operator (1). This case was recently investigated by E. Berezhnoi [Ber], who, in particular, characterized the weak type estimates

¹The research work was partially supported by INTAS project 94-881.

for operator (1) with the kernel $k(x, y) \geq 0$ increasing with respect to the first variable and also the strong estimates, when $k(x, y) = 1$ and the spaces X and Y satisfy ℓ -condition (see Definition 3 below). E. Bereznoi [Ber] has also obtained some necessary and/or sufficient conditions for the boundedness of operators (1) with the restrictions on $k(x, y) \geq 0$, stronger than (2).

Definition 1 [BS]. A real normed linear space $X = \{f : \|f\|_X < \infty\}$ of Lebesgue-measurable functions on \mathbf{R}^+ is called a *Banach function space* (BFS), if in addition to the usual norm axioms $\|f\|_X$ satisfies the following properties:

1) $\|f\|_X$ is defined for every Lebesgue-measurable function f on \mathbf{R}^+ , and $f \in X$ if, and only if, $\|f\|_X < \infty$; and $\|f\|_X = 0$ if, and only if, $f = 0$ almost everywhere (a.e.);

2) $\|f\|_X = \||f|\|_X$ for all $f \in X$;

3) if $0 \leq f \leq g$ a.e., then $\|f\|_X \leq \|g\|_X$;

4) if $0 \leq f_n \uparrow f$ a.e., then $\|f_n\|_X \uparrow \|f\|_X$;

5) if $mes E < \infty$, then $\|\chi_E\|_X < \infty$;

6) if $mes E < \infty$, then $\int_E f(x) dx \leq C_E \|f\|_X$ for all $f \in X$.

Given BFS X , its *associate space* X' is defined by

$$X' = \left\{ g : \int_0^\infty |fg| < \infty \text{ for all } f \in X \right\}$$

and endowed with the associate norm

$$\|g\|_{X'} = \sup \left\{ \int_0^\infty |fg| : \|f\|_X \leq 1 \right\}.$$

X' is also the Banach function space satisfying axioms (1-6) and, moreover, X' is the norm fundamental subspace of the dual space X^* , that is the inequality

$$\|f\|_X = \sup \left\{ \int_0^\infty |fg| : \|g\|_{X'} \leq 1 \right\}$$

holds for all $f \in X$ [BS].

The spaces X, X' are the complete normed linear spaces and $X'' = X$ [BS].

X has *absolutely continuous norm* (AC norm), if for all $f \in X$, $\|f\chi_{E_n}\|_X \rightarrow 0$ for every sequence of sets $\{E_n\} \subset \mathbf{R}^+$ such that $\chi_{E_n} \rightarrow 0$ a.e. We assume throughout the paper that X' and Y have the AC-norms.

Let ℓ be a Banach sequence space (BSS), what means that axioms (1-6) are fulfilled with respect to the count measure and let $\{e_n\}$ denote the standard basis in ℓ .

Definition 2. Given BFS X and BSS ℓ , X is said to be ℓ -*concave*, if for any sequence of disjoint intervals $\{J_k\}$ such that $\bigcup J_k = \mathbf{R}^+$, and for all $f \in X$

$$\left\| \sum_k e_k \|\chi_{J_k} f\|_X \right\|_\ell \leq d_1 \|f\|_X,$$

where a finite positive constant d_1 independent on $f \in X$ and $\{J_k\}$. Analogously, BFS Y is said to be ℓ -*convex*, if for any sequence of disjoint intervals $\{I_k\}$ such that $\bigcup I_k = \mathbf{R}^+$, and for all $g \in Y$

$$\|g\|_Y \leq d_2 \left\| \sum_k e_k \|\chi_{I_k} g\|_Y \right\|_\ell$$

with a finite positive constant d_2 independent on $g \in Y$ and $\{I_k\}$.

Definition 3 [Ber]. We say, that Banach function spaces X, Y satisfy ℓ -condition, if there exist a Banach sequence space ℓ such that X is ℓ -concave and Y is ℓ -convex simultaneously.

Throughout the paper the uncertainties of the form $0 \cdot \infty$, $0/0$, ∞/∞ are taken equal to zero, the inequality $A \ll B$ means $A \leq cB$, where c depends only on D and, possibly, on d_1 and d_2 from Definition 2; however the relationship $A \approx B$ is interpreted as $A \ll B \ll A$ or $A = cB$. χ_E denotes the characteristic function of a set $E \subset \mathbf{R}^+$.

1.1. Boundedness. Put for all $t \geq 0$

$$A_0 = \sup_{t>0} A_0(t) = \sup_{t>0} \|\chi_{[t,\infty]} v\|_Y \|\chi_{[0,t]}(\cdot)k(t,\cdot)u(\cdot)\|_{X'}, \quad (3)$$

$$A_1 = \sup_{t>0} A_1(t) = \sup_{t>0} \|\chi_{[t,\infty]}(\cdot)k(\cdot,t)v(\cdot)\|_Y \|\chi_{[0,t]}u\|_{X'} \quad (4)$$

and let $A = A_0 + A_1$. Note, that $A_0 = A_1$, if $k(x, y) = 1$.

THEOREM 1.1. Let X and Y be BFS satisfying the ℓ -condition and let K be the integral operator of the form (1) with the kernel $k(x, y) \geq 0$ satisfying (2). Then $K : X \rightarrow Y$ is bounded, if and only if, A is finite. Moreover,

$$\|K\|_{X \rightarrow Y} \approx A. \quad (5)$$

Example 1. If $X = L^p$, $Y = L^q$, $1 \leq p, q \leq \infty$ are the Lebesgue spaces with the usual norms, then the ℓ -condition holds if, and only if, $p \leq q$ and quantities (3) and (4) transform into

$$A_0 = \sup_{t>0} A_0(t) = \sup_{t>0} \left(\int_t^\infty k^q(x, t) |v(x)|^q dx \right)^{1/q} \left(\int_0^t |u(y)|^{p'} dy \right)^{1/p'}$$

$$A_1 = \sup_{t>0} A_1(t) = \sup_{t>0} \left(\int_t^\infty |v(x)|^q dx \right)^{1/q} \left(\int_0^t k^{p'}(t, y) |u(y)|^{p'} dy \right)^{1/p'}$$

in this case, where $\frac{1}{p} + \frac{1}{p'} = 1$, $\frac{1}{q} + \frac{1}{q'} = 1$. For the case $1 < q < p$ in $L^p - L^q$ setting the following criterion is true [St₁]: $\|K\|_{L^p \rightarrow L^q} \approx B$, where $B = B_0 + B_1$ defined by

$$B_0 = \left\{ \int_0^\infty \left(\int_t^\infty k^q(x, t) |v(x)|^q dx \right)^{r/q} \left(\int_0^t |u(y)|^{p'} dy \right)^{r/q'} |u(t)|^{p'} dt \right\}^{1/r}$$

$$B_1 = \left\{ \int_0^\infty \left(\int_t^\infty |v(x)|^q dx \right)^{r/p} \left(\int_0^t k^{p'}(t, y) |u(y)|^{p'} dy \right)^{r/p'} |v(t)|^q dt \right\}^{1/r}$$

where $\frac{1}{r} = \frac{1}{q} - \frac{1}{p}$. If $k(x, y) = 1$, then $B_0 = \left(\frac{p'}{q}\right)^{1/r} B_1$ and the above criterion is valid for the range $0 < q < p$, $p \geq 1$ [S] with suitable modification, when $p = 1$ [SS].

Example 2. For $0 < r < \infty$, $0 < s \leq \infty$ and a locally integrable function $\varphi(x)$ on \mathbf{R}^+ , the Lorentz space $L_{\varphi}^{r,s} \equiv L_{\varphi}^{r,s}(\mathbf{R}^+)$ consists of all measurable functions f such that $\|f\|_{r,s,\varphi} < \infty$, where

$$\|f\|_{r,s,\varphi} = \left(\int_0^\infty \left(t^{1/r} f^{**}(t) \right)^s \frac{dt}{t} \right)^{1/s} \quad \text{for } 0 < s < \infty,$$

$$\|f\|_{r,s,\varphi} = \sup_{t>0} t^{1/r} f^{**}(t) \quad \text{for } s = \infty,$$

and

$$f^{**}(t) = \frac{1}{t} \int_0^t f^*(s) ds,$$

$$f^*(t) = \inf \left\{ x > 0 : \lambda_f(x) = \int_{\{y \in \mathbf{R}^+ : |f(y)| > x\}} \varphi(z) dz \leq t \right\}.$$

If $r = s$, then

$$\|f\|_{rr,\varphi} = \left(\int_0^\infty |f(x)|^r \varphi(x) dx \right)^{1/r}.$$

If $X = L_\varphi^r, Y = L_\psi^{pq}$ then the ℓ -condition holds if, and only if, $\max(r, s) \leq \min(p, q)$ and in this case the norm of $K : X \rightarrow Y$ is sandwiched by $A = A_0 + A_1$, where

$$A_0 = \sup_{t>0} A_0(t) = \left\| \chi_{[0,t]}(\cdot) k(t, \cdot) (u(\cdot)/\varphi(\cdot)) \right\|_{r',s',\varphi} \left\| \chi_{[t,\infty]} v \right\|_{pq,\psi},$$

$$A_1 = \sup_{t>0} A_1(t) = \left\| \chi_{[0,t]} (u/\varphi) \right\|_{r',s',\varphi} \left\| \chi_{[t,\infty]}(\cdot) k(\cdot, t) v(\cdot) \right\|_{pq,\psi}.$$

Remark 1.1. (i) If the ℓ -condition fails, then the lower bound in (5) is nevertheless true. However, there exist an operator, when (5) is valid for the spaces with no ℓ -condition. Indeed, if we take $k(x, y) = 1, v(x) = 1$, then in Lorentz space setting above the criterion (5) holds for $1 < r = s, q \geq r, 0 < p < \infty$ ([Sa], Theorem 2).

(ii) If $1 < r = s, 0 < q < r < \infty, 0 < p < \infty$, then the criterion for the boundedness of this operator is the following ([St₁], Theorem 2.2). Put $Uf(x) = \int_0^x f(y)u(y)dy$. Then

$$\sup_{f \neq 0} \frac{\|Uf\|_{pq,\psi}}{\|f\|_{rr,\varphi}} \approx \left(\int_0^\infty \left(\|\chi_{[t,\infty]}\|_{pq,\psi} \right)^\gamma d \left(\|\chi_{[0,t]}(u/\varphi)\|_{r',r',\varphi} \right)^\gamma \right)^{1/\gamma},$$

where $\frac{1}{\gamma} = \frac{1}{q} - \frac{1}{r}$.

1.2. Compactness and measure of non-compactness.

THEOREM 1.2. *Let the assumptions of Theorem 1 be fulfilled and the spaces X' and Y have the AC-norms. Then the operator $K : X \rightarrow Y$ is compact if, and only if, A is finite and*

$$\lim_{t \rightarrow a_i} A_i(t) = \lim_{t \rightarrow b_i} A_i(t) = 0, \quad i = 0, 1, \quad (6)$$

where

$$a_i = \inf\{t > 0 : A_i(t) > 0\}, \quad b_i = \sup\{t > 0 : A_i(t) > 0\} \quad i = 0, 1.$$

Remark 1.1. (i) In fact, it follows from the proof of Theorem 1.2, that $a_0 = a_1, b_0 = b_1$.

(ii) By many authors the condition (6) used to be formulated for the end-points, however it is easy to point out a formal counterexample, when A is finite and (6) is valid with $a_0 = a_1 = 0, b_0 = b_1 = \infty$, but K is non-compact. The matter is, that the condition (6) has to be formulated for the end-points of the real interval of non-zero action of K .

(iii) For the case $1 < q < p$ in $L^p - L^q$ setting the operator $K : L^p \rightarrow L^q$ is compact if, and only if, $B < \infty$ (see Example 1). It follows from the Ando theorem [A].

In the non-compact case we estimate the measure of non-compactness of K or, equivalently, the distance between K and the set of finite rank operators defined by

$$\alpha(K) = \inf \|K - P\|,$$

where the infimum is taken over all bounded linear maps $P : X \rightarrow Y$ of finite rank. To this end we need additional portion of notations. For $0 < a < z < b < \infty$ we put

$$J_L^0(a) = \sup_{0 < t < a} \|\chi_{[t,a]} v\|_Y \|\chi_{[0,t]}(\cdot) k(t, \cdot) u(\cdot)\|_{X'},$$

$$J_L^1(a) = \sup_{0 < t < a} \|\chi_{[t,a]}(\cdot) k(\cdot, t) v(\cdot)\|_Y \|\chi_{[0,t]} u\|_{X'},$$

$$J_L(z) = \max(J_L^0(z), J_L^1(a)), \quad J_L = \lim_{z \rightarrow a_0} J_L(z),$$

$$J_R^0(b) = \sup_{b < t < \infty} \|\chi_{[t,a]} v\|_Y \|\chi_{[0,t]}(\cdot) k(t, \cdot) u(\cdot)\|_{X'},$$

$$J_R^1(b) = \sup_{b < t < \infty} \|\chi_{[t,a]}(\cdot) k(\cdot, t) v(\cdot)\|_Y \|\chi_{[0,t]} u\|_{X'},$$

$$J_R(z) = \max(J_R^0(z), J_R^1(a)), \quad J_R = \lim_{z \rightarrow a_0} J_R(z),$$

$$J = \max(J_L, J_R).$$

THEOREM 1.3. *Let the assumptions of Theorem 2 be valid and $K : X \rightarrow Y$ be bounded. Then*

$$D^{-1}J \leq \alpha(K) \leq (d_1, d_2, D)J.$$

2. Problems (S) and (N) in Lebesgue spaces

2.1. Schatten-von Neumann ideal norms. Let H be a separable Hilbert space. Then the set of all linear bounded operators $T : H \rightarrow H$ forms the normed algebra B , where σ_∞ -the ideal of all compact operators. The theory of symmetrically normed (s.n.) ideals $\sigma_\Phi \subset \sigma_\infty$ was developed by using the s.n. functions Φ defined on the space of sequences with a finite number of non-zero terms ([GK], Chapter 3). If $T \in \sigma_\infty$, then $T^* \in \sigma_\infty$ and $(T^*T)^{1/2} \in \sigma_\infty$. To construct σ_Φ the sequences of singular numbers $s_j(T) = \lambda_j \left[(T^*T)^{1/2} \right]$ were used, with the eigenvalues $\lambda_j \geq 0$ taken according to their multiplicity and decrease. Formula $\|T\|_{\sigma_\Phi} = \Phi(s_j(T))$ defines the norm (quasinorm) in the s.n. ideal σ_Φ . The most well-known are the s.n. ideals σ_p related to the space of sequences l_p , $0 < p \leq \infty$. The norm (quasinorm) $\|T\|_{\sigma_p} = \left(\sum_j s_j^p(T) \right)^{1/p}$ is usually called by the Schatten-von Neumann norm (quasinorm). Thus, $\|T\|_{\sigma_2} = \|T\|$ and $\|T\|_{\sigma_2}$ is the Hilbert-Schmidt norm expressed for an integral operator $Tf(x) = \int T(x, y)f(y) dy$ by the formulae $\|T\|_{\sigma_2} = \left(\int \int |T(x, y)|^2 dx dy \right)^{1/2}$. It is known [BS], that in general the norm $\|T\|_{\sigma_p}$ of an integral operator substantially depends on the smoothness of its kernel, when $p < 2$, however for some particular operators of complex harmonic analysis the effective two-sided estimates of the Schatten-von Neumann norms are well known, e.g.

see [Pa], [P]. The aim of the section is to present a brief account of some results from [ES₂] and [St₂] about the Schatten-von Neumann ideal norms for the integral operators (1) with the condition (2) for their kernels.

Let $H = L^2(0, \infty)$ and

$$A_0^2 = \sup_{t>0} \int_t^\infty k^2(x, t) |v(x)|^2 dx \int_0^t |u(y)|^2 dy,$$

$$A_1^2 = \sup_{t>0} \int_t^\infty |v(x)|^2 dx \int_0^t k^2(t, y) |u(y)|^2 dy.$$

Theorem 1.1 and the Hilbert-Schmidt formula bring

$$\|K\|_{\sigma_\infty} \approx A_0 + A_1,$$

$$\|K\|_{\sigma_2} = \left(\int_0^\infty |v(x)|^2 dx \int_0^x k^2(t, y) |u(y)|^2 dy \right)^{1/2} = \left(\int_0^\infty |u(y)|^2 dy \int_y^\infty k^2(x, y) |v(x)|^2 dx \right)^{1/2}.$$

Using these formulas and applying the real method of interpolation we obtain the following

THEOREM 2.1. *Let K be an operator of the form (1) with the kernel satisfying (2) and $K \in \sigma_\infty$. Then*

$$\|K\|_{\sigma_p} \approx \left(\int_0^\infty \left[\left(\int_0^x k^2(x, y) |u(y)|^2 dy \right)^{p/2} \left(\int_x^\infty |v(y)|^2 dy \right)^{\frac{p}{2}-1} |v(x)|^2 + \left(\int_x^\infty k^2(y, x) |v(y)|^2 dy \right)^{p/2} \left(\int_0^x |u(y)|^2 dy \right)^{\frac{p}{2}-1} |u(x)|^2 \right] dx \right)^{1/p}, \quad 2 \leq p < \infty. \quad (7)$$

Remark 2.1. The upper bound of (7) is proved in [ES₂] and the lower one in [St₂]. In case $k(x, y) \equiv 1$ the formula (7) can be simplified and extended as follows. If

$$Hf(x) = v(x) \int_0^x f(y) u(y) dy, \quad (8)$$

then

$$\|H\|_{\sigma_p} \approx \left(\int_0^\infty \left(\int_0^x |u(y)|^2 dy \right)^{p/2} \left(\int_x^\infty |v(y)|^2 dy \right)^{\frac{p}{2}-1} |v(x)|^2 dx \right)^{1/p}, \quad 1 < p < \infty. \quad (9)$$

Remark 2.1. In alternate form the equivalence (9) for the case $u(y) = 1$ has been established in [N] and later this result has been widely extended in [NS] for the operator

$$I_\nu f(x) = \frac{v(x)}{x^\nu} \int_0^x (x-y)^{\nu-1} f(y) dy, \quad \nu > 1/2.$$

2.2. Approximation numbers. Recall that if $T : X \rightarrow Y$, then the n -th approximation number of T is defined by

$$a_n = \inf\{\|T - P\|, \text{rank } P < n\}, \quad n = 1, 2, \dots$$

The problem of asymptotic behaviour of the approximation numbers is well known and was treated in the monographs [EE], [K] and others and for the operator (8) in the papers [EEH₁], [EEH₂],

[ES₁], [LS₁], [LS₂]. Here we present the new result for the operator (8) easy comparable with formula (9).

THEOREM 2.2 *Let $1 < p, s < \infty$ and the integral operator $H : L^p(0, \infty) \rightarrow L^p(0, \infty)$ given by (8) be compact and $\{a_n\}$ is the sequence of the approximation numbers of H . Then*

$$\left(\sum_{n=1}^{\infty} a_n^s\right)^{1/s} \approx \left(\int_0^{\infty} \left(\int_0^x |u(y)|^{p'} dy\right)^{s/p'} \left(\int_0^x |v(y)|^p dy\right)^{\frac{s}{p}-1} |v(x)| dx\right)^{1/s},$$

$$\overline{\lim}_{n \rightarrow \infty} na_n \leq \gamma_p \int_0^{\infty} |uv| \leq 2 \lim_{n \rightarrow \infty} na_n, \quad p \neq 2,$$

$$\lim_{n \rightarrow \infty} na_n = \frac{1}{\pi} \int_0^{\infty} |uv|, \quad p = 2.$$

The last two formulas are proved in [EEH₂] and the proof of the first is based on the results of [EEH₂].

Remark 2.2. All the assertions of the paper have natural analogs for a finite interval instead of $(0, \infty)$ and for the dual operator K^* as well (see [St₁] for details).

Acknowledgement. The author wishes to thank Professor Takashi Yoshino for his kind invitation to give a talk on the workshop "Recent topics on the operator theory about the structure of operators" held in the Research Institute of Mathematical Science of Kyoto University, and his warm hospitality and the travel support.

References

- [A] T. Ando, On compactness of integral operators, *Indag. Math.*, 24 (1962) 235-239.
- [BS] C. Bennett and R. Sharpley, *Interpolation of operators*, Pure Appl. Math. 129, Academic Press, 1988.
- [Ber] E.I. Bereznoi, Sharp estimates of operators on the cones of ideal spaces, *Proc. Steklov Inst. Math.* (1994), Issue 3, 3-34.
- [BiS] M. Sh. Birman and M.Z. Solomjak, Estimates for the singular numbers of integral operators, *Russian Math. Surveys*, 32 (1977).
- [EE] D.E. Edmunds and W.D. Evans, *Spectral theory and differential operators*, Oxford: Univ. Press, 1987.
- [EEH₁] D.E. Edmunds, W.D. Evans and D.J. Harris, Approximation numbers of certain Volterra integral operators, *J. London Math. Soc.*, (2) 37 (1988), 471-489.
- [EEH₂] D.E. Edmunds, W.D. Evans and D.J. Harris, *Two-sided estimates of the approximation numbers of certain Volterra integral operators*, Centre for mathematical analysis and its applications, University of Sussex, Research Report No: 96/09, 1996, 25 pp.
- [ES₁] D.E. Edmunds and V.D. Stepanov, The measure of non-compactness and approximation numbers of certain Volterra integral operators. *Math. Ann.* 298 (1994) 41-66.
- [ES₂] D.E. Edmunds and V.D. Stepanov, On singular numbers of certain Volterra integral operators. *J. Funct. Anal.* 134 (1995) 222-246.
- [GK] I.C. Gohberg and M.G. Krein, *Introduction to the theory of linear non-selfadjoint operators*, AMS Tranl. Math. Monographs, 18, Providence, R. I., 1969.
- [K] H. König, *Eigenvalue distribution of compact operators*, Birkhäuser, Boston, 1986.

- [LS₁] E.N. Lomakina and V.D. Stepanov, On the compactness and approximation numbers of Hardy type integral operators in Lorentz spaces, *J. London Math. Soc.* (2) 53 (1996) 369-382.
- [LS₂] E.N. Lomakina and V.D. Stepanov, *On the Hardy type integral operators in Banach function spaces on semiaxis*, (in russian), Computer Center, Research Report No: 9-1996, Khabarovsk, 1996, 35 pp.
- [NS] J. Newman and M. Solomyak, Two-sided estimates on singular numbers for a class of integral operators on the semi-axis, *Integr. Equat. Oper. Th.*, 20 (1994) 335-349.
- [N] K. Nowak, Schatten ideal behaviour of a generalized Hardy operator, *Proc. Amer. Math. Soc.*, 118 (1993) 479-483.
- [Pa] O. Parfenov, Estimates of singular numbers of integral operators with holomorphic kernels, (in russian), *Bull. St.-Petersburg State Univ.*, Ser. 1, 8 (1992) 24-32.
- [P] V.V. Peller, A discription of Hankel operators of class S_p for $p > 0$, an investigation of rate of rational arroximation, and other applications, *USSR Math. Sbornik*, 122 (1985) 465-494.
- [Sa] E. Sawyer, Weighted Lebesgue and Lorentz norm inequalities for the Hardy operator, *Trans. Amer. Math. Soc.* 281 (1984) 329-337.
- [S] G. Sinnamon, Weighted Hardy and Opial type inequalities, *J. Math. Anal. Appl.*, 160 (1991) 433-445.
- [SS] G. Sinnamon and V. Stepanov, The weighted Hardy inequality: New proofs and the case $p = 1$, *J. London Math. Soc.*, (2) 54 (1996) 89-101.
- [St₁] V. Stepanov, *Weighted norm inequalities for integral operators and related topics*, Nonlinear analysis, function spaces and applications, vol. 5, Olimpia Press, Prague, 1994, 139-176.
- [St₂] V. Stepanov, On the lower bounds for Schatten-von Neumann norms of certain Volterra integral operators, *J. London Math. Soc.*, (to appear).

Vladimir Stepanov
 Computer Center
 Far-Eastern Branch
 Russian Academy of Sciences
 Shelest 118-205,
 Khabarovsk 680042, Russia
 E-mail: 1600@as.khabarovsk.su