

On the Dimension Formula for the Spaces of Jacobi Forms of Degree Two

明大理工 対馬 龍司 (Ryuji Tsushima)

§1. Result

Let $\mathfrak{S}_g = \{Z \in M_g(\mathbf{C}) \mid {}^t Z = Z, \operatorname{Im} Z > 0\}$ be the Siegel upper half plane of degree g and let $\Gamma_g = Sp(g, \mathbf{Z})$. If $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_g$, we denote $(AZ + B)(CZ + D)^{-1}$ by $M \langle Z \rangle$. Let $\mathbf{e}(z)$ denote $\exp(2\pi iz)$.

Definition 1.1. Let k, m be positive integers. A holomorphic function $f(Z, W)$ on $\mathfrak{S}_g \times \mathbf{C}^g$ is called a *Jacobi form of weight k and index m with respect to Γ_g* , if it satisfies the following transformation formulas and a regularity condition at infinity:

$$(1) \quad f(M \langle Z \rangle, {}^t(CZ + D)^{-1}W) = \det(CZ + D)^k \mathbf{e}(m {}^t W(CZ + D)^{-1} CW) f(Z, W),$$

for any $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_g$,

$$(2) \quad f(Z, W + Z\lambda + \mu) = \mathbf{e}(-m({}^t \lambda Z \lambda + 2 {}^t \lambda W)) f(Z, W), \quad \text{for any } \lambda, \mu \in \mathbf{Z}^g.$$

If f satisfies (1) and (2), f has a Fourier expansion of the form:

$$f(Z, W) = \sum_{N, r} c(N, r) \mathbf{e}(\operatorname{Tr}(NZ) + {}^t r W),$$

where N is over the symmetric half integral matrix of degree g and r is over the integral g -vector. The regularity condition at infinity is:

$$(3) \quad c(N, r) = 0 \text{ unless } 4mN - r {}^t r \text{ is semi-positive.}$$

Remark 1.2. If $g \geq 2$, then the condition (3) is superfluous ([Sh], [Y] and [Z]).

Definition 1.3. A Jacobi form f is called a *Jacobi cusp form* if $c(N, r) = 0$ unless $4mN - r {}^t r$ is positive definite in the Fourier expansion above.

Our main result is the following

Theorem 1.4. If $k \geq 4$, then the dimension of the space of Jacobi cusp forms $J_{k,m}^{cusp}(\Gamma_2)$ with respect to Γ_2 is given by the following Mathematica function:

```
JacobiTwo[k_, m_] := Block[{a, lk, x, y, m2, m3, m4, r, p, S1e, S1, S2e, S2, S3, SS, SSS},  
  mod[x_, y_] := Mod[x, y] + 1;
```

```

m2=Mod[m,2];
m3=Mod[m,3];
m4=Mod[m,4];

r=0;
While[EvenQ[m/2^r],r++];
p=m/2^r;

S1e=4*Sum[Mod[x^2,m],{x,1,m-1}];
S1=S1e+Sum[Mod[(2*x-1)^2,4*m],{x,1,m}];
S2e=16*Sum[Mod[x^2,m]^2,{x,1,m-1}];
S2=S2e+Sum[Mod[(2*x-1)^2,4*m]^2,{x,1,m}];
S3=Sum[Mod[x^2,4*m]^3,{x,1,2*m-1}];
SS=4*Sum[Mod[x^2,4*m]*Mod[x^2,m],{x,1,2*m-1}];
SSS=Sum[Mod[x^2,4*m]*Mod[y^2,4*m]*Mod[(x-y)^2,4*m],{x,1,2*m-1},{y,1,2*m-1}];

a=m^2*((2*k-3)*(2*k-4)*(2*k-5)/2^8/3^3/5-(2*k-4)/2^4/3^2+1/2^3/3);
a=a+(3*k-20)*S1/2^5/3^2+(-k+7)*S2/m/2^7/3+S3/m^2/2^8/3^2;
a=a+S1^2/m^2/2^7-S1*S2/m^3/2^8+SSS/m^3/2^8/3;

a=a+(2*k-3)*(2*k-4)*(2*k-5)/2^8/3^3/5+(2-k)/2^3/3^2+1/2^3/3;
a=a-m4^3/2^5/3^2+(10-k)*m4^2/2^7/3+(3*k-20)*m4/2^5/3^2;

lk={1,-1};
a=a+lk[[mod[k,2]]]*m*((2*k-3)*(2*k-5)/2^7/3^2-(k-2)/2^3/3+1/2^3);
a=a+lk[[mod[k,2]]]*(S1*(k-8)/m/2^6/3+S1*m4/m/2^7+m4*m*(k-8)/2^6/3);

lk={1,-1};
a=a+lk[[mod[k,2]]]*m*((2*k-3)*(2*k-5)/2^7/3+(5-3*k)/2^4/3+7/2^4/3);
a=a+lk[[mod[k,2]]]*(4*SS+(16-2*m4)*m*S1+4*(k-7+m4)*m*S1e-8*S2-S2e)/m^2/2^8;

a=a+(k-2)/2^4/3;
a=a+(m4-4)/2^5;

a=a+If[m2==0,1,0]*(k-2)/2^5;
a=a+If[m2==0,1,-1]*m4/2^5-If[m2==0,1,0]/2^3;

a=a+If[m3==0,3,1]*(k-2)/2^3/3;
a=a-If[m3==0,3,1]/2^2/3+If[m3==0,1,0]*Mod[m/3,3]/3^2;

a=a+(k-2)/2/3^3;
a=a-1/2^2/3;

lk={-1,-2*k+4,1,2*k-4};
a=a+If[m2==0,1,0]*lk[[mod[k,4]]]*m/2^5/3;
lk={0,-1,0,1};
a=a+If[m2==0,1,0]*lk[[mod[k,4]]]*(S1/m-4*m)/2^5;

lk={2*k-4,-1,-2*k+4,1};
a=a+If[m2==0,1,0]*lk[[mod[k,4]]]/2^5/3;

```

```

lk={1,0,-1,0};
a=a+If[m2==0,1,0]*lk[[mod[k,4]]]*(m4-4)/2^5;

lk={0,-1,0,1};
a=a+lk[[mod[k,4]]]*Sum[Mod[(2*y+m2)^2,8*m]-Mod[(2*y+m2)^2+4*m,8*m],
{y,0,m-1}]/m/2^6;

lk={1,0,-1,0};
a=a+If[m2==0,-1,0]*lk[[mod[k,4]]]/2^4;
a=a+lk[[mod[k+m,4]]]*(Mod[m,8]-Mod[m+4,8])/2^6;

If[m3==0,Goto[label1a],0];
lk={-2*k+3,-2*k+5,4*k-8};
a=a+If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]*lk[[mod[k,3]]]*m/2^4/3^3;
lk={-1,-1,2};
a=a+If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]*lk[[mod[k,3]]]*(S1/m-4*m)/2^4/3^2;
Goto[label2a];
Label[label1a];
lk={6*k-13,-6*k+11,2};
a=a+lk[[mod[k,3]]]*m/2^4/3^3;
lk={1,-1,0};
a=a+lk[[mod[k,3]]]*(S1/m-4*m)/2^4/3;
Label[label2a];

lk={2*k-3,2*k-5,-4*k+8};
a=a+lk[[mod[k,3]]]/2^4/3^3;
lk={1,1,-2};
a=a+lk[[mod[k,3]]]*(m4-4)/2^4/3^2;

lk={-2*k+3,-2*k+5,2,2*k-3,2*k-5,-2};
a=a+lk[[mod[k,6]]]*m/2^4/3^2;
lk={-1,-1,0,1,1,0};
a=a+lk[[mod[k,6]]]*(S1/m-4*m)/2^4/3;

If[m3==0,Goto[label1b],0];
lk={2*k-3,2*k-5,-2,-2*k+3,-2*k+5,2};
a=a+If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]*lk[[mod[k,6]]]/2^4/3^2;
lk={1,1,0,-1,-1,0};
a=a+If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]*lk[[mod[k,6]]]*(m4-4)/2^4/3;
Goto[label2b];
Label[label1b];
lk={2*k-7,-2*k+1,-4*k+8,-2*k+7,2*k-1,4*k-8};
a=a+lk[[mod[k,6]]]/2^4/3^2;
lk={1,-1,-2,-1,1,2};
a=a+lk[[mod[k,6]]]*(m4-4)/2^4/3;
Label[label2b];

```

```

If[m3==0,Goto[label1c],0];
lk={-1,-1,2};
a=a+If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]*Sum[If[Mod[y,3]==0,2,-1]*
(lk[[mod[k,3]]]*Mod[y^2,12*m]+lk[[mod[k+1,3]]]*Mod[y^2+4*m,12*m]+
lk[[mod[k+2,3]]]*Mod[y^2+8*m,12*m]),{y,0,2*m-1}]/m/2^4/3^3;
lk={1,-1,0};
a=a+Sum[If[Mod[y,3]==0,0,1]*(lk[[mod[k,3]]]*Mod[y^2,12*m]+lk[[mod[k+1,3]]]*Mod[y^2+4*m,12*m]+
lk[[mod[k+2,3]]]*Mod[y^2+8*m,12*m]),{y,0,2*m-1}]/m/
2^4/3^2;
Goto[label2c];
Label[label1c];
lk={1,-1,0};
a=a+Sum[lk[[mod[k,3]]]*Mod[9*y^2,12*m]+lk[[mod[k+1,3]]]*Mod[9*y^2+4*m,12*m]+
lk[[mod[k+2,3]]]*Mod[9*y^2+8*m,12*m],{y,0,2*m/3-1}]/m/2^3/3^2;
Label[label2c];

lk={-1,0,1};
a=a+lk[[mod[k,3]]]/2/3^2;
lk={1,1,-2};
a=a+(lk[[mod[k+m,3]]]*Mod[m,12]+lk[[mod[k+m+1,3]]]*Mod[m+4,12]+
lk[[mod[k+m+2,3]]]*Mod[m+8,12])/2^3/3^3;

a=a+0;

lk={0,1,0,-1};
a=a+If[m2==0,1,0]*lk[[mod[k,4]]]/2^3;

lk={-1,1,0};
a=a+lk[[mod[k,3]]]*If[m3==0,-3,1]/2^3/3^3;

lk={-1,1,0};
a=a+lk[[mod[k,3]]]/2^3/3^3;

If[m3==0,Goto[label2d],0];
lk={1,-1};
a=a+lk[[mod[k,2]]]*If[Mod[p,3]==1,1,-1]*If[EvenQ[r],1,-1]/2^2/3^3;
Label[label2d];

If[m3==0,Goto[label1e],0];
lk={1,-1,-2,-1,1,2};
a=a+If[Mod[p,3]==1,1,-1]*If[EvenQ[r],1,-1]*lk[[mod[k,6]]]/2^2/3^2;
Goto[label2e];
Label[label1e];
lk={1,1,0,-1,-1,0};
a=a+lk[[mod[k,6]]]/2^2/3;
Label[label2e];

```

```

If[m3==0,Goto[label1f],0];
lk={1,-1,-2,-1,1,2};
a=a+If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]*lk[[mod[k,6]]]/2^2/3^3;
Goto[label1f];
Label[label1f];
lk={1,1,0,-1,-1,0};
a=a+lk[[mod[k,6]]]/2^2/3^2;
Label[label2f];
lk={-1,1,0};
a=a+lk[[mod[k,3]]]/2^2/3;
If[m2==1,Goto[label2g],0];
If[m3==0,Goto[label1g],0];
lk={-1,-1,-2,-1,-1,0,1,1,2,1,1,0};
a=a+lk[[mod[k,12]]]*If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]/2^3/3;
Goto[label2g];
Label[label1g];
lk={3,1,0,-1,-3,-2,-3,-1,0,1,3,2};
a=a+lk[[mod[k,12]]]/2^3/3;
Label[label2g];
lk={1,1,0,-1,-1,-2,-1,-1,0,1,1,2};
a=a+lk[[mod[k,12]]]*If[m2==0,1,0]/2^3/3;
If[m3==0,Goto[label2h],0];
lk={1,-1};
a=a+lk[[mod[k,2]]]*If[Mod[p,3]==1,1,-1]*If[EvenQ[r],-1,1]/2^2/3^2;
Label[label2h];
a=a+0;
If[Mod[m,5]==0,Goto[label1i],0];
lk={0,-1,0,1,0};
a=a+lk[[mod[k,5]]]*If[Mod[p^2,5]==1,1,-1]*If[EvenQ[r],-1,1]/2/5;
Goto[label2i];
Label[label1i];
lk={2,1,0,-1,-2};
a=a+lk[[mod[k,5]]]/2/5;
Label[label2i];
lk={0,1,0,-1,0};
a=a+lk[[mod[k,5]]]/2/5;
Return[a];
]

```

§2. Methods

Definition 2.1. Let a and b be rational g -vectors. The theta function $\theta_{a,b}(Z, W)$ with characteristic (a, b) is a holomorphic function on $\mathfrak{S}_g \times \mathbf{C}^g$ defined by

$$\sum_{q \in \mathbf{Z}^g} \mathbf{e}\left((1/2)^t(q+a)Z(q+a) + {}^t(q+a)(W+b)\right).$$

For any integral g -vector r , we have

$$\theta_{a+r,b}(Z, W) = \theta_{a,b}(Z, W).$$

Hence $\theta_{a,0}(Z, W)$ depends only on $a \bmod \mathbf{Z}^g$. So we assume a is an element of $\mathbf{Q}^g/\mathbf{Z}^g$. If a runs a complete set of representatives of $\frac{1}{2m}\mathbf{Z}^g/\mathbf{Z}^g$, then $\theta_{a,0}(2mZ, 2mW)$ form a basis of theta function of degree $2m$. Therefore if f is a Jacobi form of index m , there exist uniquely determined holomorphic functions $f_r(Z)$ ($r \in \frac{1}{2m}\mathbf{Z}^g/\mathbf{Z}^g$) on \mathfrak{S}_g satisfying

$$f(Z, W) = \sum_r f_r(Z) \theta_{r,0}(2mZ, 2mW).$$

We define $(2m)^g$ -vectors:

$$F(Z) = (f_r(Z)), \quad \Theta(Z, W) = (\theta_{r,0}(2mZ, 2mW)).$$

Then by definition we have

$$(*) \quad {}^t F(M \langle Z \rangle) \Theta(M \langle Z \rangle, {}^t(CZ + D)^{-1}W) \\ = \det(CZ + D)^k \mathbf{e}(m^t W(CZ + D)^{-1} CW) {}^t F(Z) \Theta(Z, W).$$

We need the following transformation formula for the theta functions ([Si]).

Proposition 2.2. Let $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_g$. Then for any $r \in \frac{1}{2m}\mathbf{Z}^g/\mathbf{Z}^g$, we have

$$\theta_{r,0}(2mM \langle Z \rangle, 2m^t(CZ + D)^{-1}W) \\ = \det(CZ + D)^{1/2} \mathbf{e}(m^t W(CZ + D)^{-1} CW) \times \sum_s u_{rs}(M) \theta_{s,0}(2mZ, 2mW),$$

where s runs a complete set of representatives of $\frac{1}{2m}\mathbf{Z}^g/\mathbf{Z}^g$ and $(u_{rs}(M))_{r,s}$ is an unitary matrix of degree $(2m)^g$ depending only on M and the choice of $\det(CZ + D)^{1/2}$.

Let $u(M) = (u_{rs}(M))_{r,s}$. Then by the proposition we have

$$(**) \quad \Theta(M \langle Z \rangle, {}^t(CZ + D)^{-1}W) = \det(CZ + D)^{1/2} \mathbf{e}(m^t W(CZ + D)^{-1} CW) u(M) \Theta(Z, W).$$

From (*) and (**) we have

$$F(M \langle Z \rangle) = \det(CZ + D)^{k-1/2} \overline{u(M)} F(Z).$$

Namely, $F(Z)$ is a vector valued modular form with respect to the automorphic factor:

$$\det(CZ + D)^{k-1/2} \overline{u(M)}.$$

So we have the following proposition ([Sh]).

Proposition 2.3. *By the mapping: $f(Z, W) \mapsto F(Z)$, $J_{k,m}(\Gamma_g)$ is mapped isomorphically to the space of the vector valued modular forms with respect to the above automorphic factor and Γ_g .*

Let $\Theta(Z) = \sum_{\eta \in \mathbf{Z}^g} \mathbf{e}^{(t\eta Z\eta)}$ and let

$$\Gamma_0^g(4) = \left\{ \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_g \mid C \equiv O \pmod{4} \right\}.$$

If $M \in \Gamma_0^g(4)$, then $J(M, Z) := \Theta(M \langle Z \rangle)/\Theta(Z)$ is holomorphic on \mathfrak{S}_g and satisfies

$$J(M, Z)^2 = \det(CZ + D) \left(\frac{-1}{\det D} \right).$$

Let $\Gamma_g(N)$ be the principal congruence subgroup of level N of Γ_g . Namely,

$$\Gamma_g(N) = \{M \in \Gamma_g \mid M \equiv 1_{2g} \pmod{N}\}.$$

If $M \in \Gamma_g(4)$, we may assume that $\det(CZ + D)^{1/2} = J(M, Z)$. Then $u(M)$ becomes a representation of $\Gamma_g(4)$.

$\Gamma_g(N)$ is a normal subgroup of Γ_g . If $N \geq 3$, $\Gamma_g(N)$ acts on \mathfrak{S}_g without fixed points and the quotient space $X_g(N) := \Gamma_g(N) \backslash \mathfrak{S}_g$ is a (non-compact) manifold. $X_g(N)$ is a open subspace of a projective variety $\overline{X}_g(N)$ which was constructed by I. Satake ([Sa], Satake compactification). If $g \geq 2$, $\overline{X}_g(N)$ has singularities along its cusps: $\overline{X}_g(N) - X_g(N)$. Cusps of $\overline{X}_g(N)$ is (as a set) a disjoint union of copies of $X_{g'}(N)$'s ($0 \leq g' < g$). A desingularization $\tilde{X}_g(N)$ of $\overline{X}_g(N)$ was constructed by J.-I. Igusa ($g = 2, 3, 4$) ([Ig2]) and Y. Namikawa ([Nm]) and more generally by D. Mumford and others ([AMRT], Toroidal compactification).

Let \mathcal{V}_m be $\mathfrak{S}_g \times \mathbf{C}^{(2m)^g}$. $\Gamma_g(4N)$ acts on \mathcal{V}_m as follows:

$$M(Z, v) = (M \langle Z \rangle, \overline{u(M)}v).$$

$V_m := \Gamma_g(4N) \backslash \mathcal{V}_m$ is non-singular and is a vector bundle over $X_g(4N)$. Let \mathcal{H}_g be $\mathfrak{S}_g \times \mathbf{C}$. $\Gamma_g(4N)$ acts on \mathcal{H}_g as follows:

$$M(Z, v) = (M \langle Z \rangle, J(M, Z)v).$$

$H_g := \Gamma_g(4N) \backslash \mathcal{H}_g$ is a line bundle over $X_g(4N)$. H_g is extended to a line bundle \tilde{H}_g over $\tilde{X}_g(4N)$ and also to an ample line bundle \overline{H}_g over $\overline{X}_g(4N)$.

Now we have

Proposition 2.4. *If $m \mid N$, V_m is extendable to a vector bundle \tilde{V}_m over $\tilde{X}_g(4N)$. \tilde{V}_m is a flat vector bundle and the Chern class $c_i(\tilde{V}_m)$ ($i \geq 1$) is 0.*

Hence in the following we assume that the level is divisible by $4m$. Let $J_{k,m}(\Gamma_g(4mN))$ be the space of Jacobi forms with respect $\Gamma_g(4mN)$. This is canonically identified with the space

$$\Gamma(\tilde{X}_g(4mN), \mathcal{O}(\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-1)})),$$

which is the space of the global holomorphic sections of $\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-1)}$. Let $D := \tilde{X}_g(4mN) - X_g(4mN)$ be the divisor at infinity. D is a divisor with simple normal crossings. $J_{k,m}^{cusp}(\Gamma_g(4mN))$ is canonically identified with the space

$$S_{k,m}(\Gamma_g(4mN)) := \Gamma(\tilde{X}_g(4mN), \mathcal{O}(\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-1)} - D)).$$

$\mathcal{O}(\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-1)} - D)$ is the sheaf of germs of holomorphic sections which vanish along D and this is isomorphic to $\mathcal{O}(\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-1)} \otimes [D]^{\otimes(-1)})$, where $[D]$ is the line bundle associated with D . And this is isomorphic to $\mathcal{O}(\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-2g-3)} \otimes K_{\tilde{X}_g(4mN)})$, since $K_{\tilde{X}_g(4mN)} \otimes [D] \simeq \tilde{H}_g^{\otimes(2g+2)}$.

Since \tilde{V}_m is a flat vector bundle and \tilde{H}_g is positive on $X_g(4mN)$, we can prove the following theorem by the vanishing theorem of Kodaira-Nakano ([Kd], [Nk]).

Theorem 2.5. *If $k \geq g + 2$ and $p > 0$, then*

$$H^p(\tilde{X}_g(4mN), \mathcal{O}(\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-1)} - D)) \simeq \{0\}.$$

Since the Chern character $ch(\tilde{V}_m)$ of \tilde{V}_m is $(2m)^g$, from the above vanishing theorem and the theorem of Riemann-Roch-Hirzebruch we have

Theorem 2.6. *If $k \geq g + 2$, then*

$$\dim J_{k,m}^{cusp}(\Gamma_g(4mN)) = \dim S_{k,m}(\Gamma_g(4mN)) = (2m)^g \dim S_{k-1/2}(\Gamma_g(4mN)),$$

where $S_{k-1/2}(\Gamma_g(4mN))$ is the space of Siegel cusp forms of weight $k - 1/2$.

$M \in \Gamma_g$ acts on $S_{k,m}(\Gamma_g(4mN))$ as follows:

$$MF(M \langle Z \rangle) = \det(CZ + D)^{k-1/2} \overline{u(M)} F(Z).$$

Since $\Gamma_g(4mN)$ acts trivially, $\Gamma_g/\Gamma_g(4mN)$ acts on $S_{k,m}(\Gamma_g(4mN))$. Hence the dimension of $J_{k,m}^{cusp}(\Gamma_g) \simeq S_{k,m}(\Gamma_g)$ is calculated as an invariant subspace of $S_{k,m}(\Gamma_g(4mN))$ by using the holomorphic Lefschetz fixed point formula ([AS]).

We recall the holomorphic Lefschetz fixed point formula. Let X be a compact complex manifold and V a holomorphic vector bundle of rank n on X , and let G be a finite group of

automorphism of the pair (X, V) . For $g \in G$ let X^g be the set of fixed points of g . X^g is a disjoint union of submanifolds of X . Let

$$X^g = \sum_{\alpha} X_{\alpha}^g$$

be the irreducible decomposition of X^g , and let

$$N_{\alpha}^g = \sum_{\theta} N_{\alpha}^g(\theta)$$

denote the normal bundle of X_{α}^g decomposed according to the eigenvalues $e^{i\theta}$ of g . We put

$$\mathcal{U}^{\theta}(N_{\alpha}^g(\theta)) = \prod_{\beta} \left(\frac{1 - e^{-x_{\beta} - i\theta}}{1 - e^{-i\theta}} \right)^{-1},$$

where the Chern class of $N_{\alpha}^g(\theta)$ is

$$c(N_{\alpha}^g(\theta)) = \prod_{\beta} (1 + x_{\beta}).$$

Let $\mathcal{T}(X_{\alpha}^g)$ be the Todd class of X_{α}^g . Let $V|X_{\alpha}^g$ be the restriction of V to X_{α}^g and $ch(V|X_{\alpha}^g)(g)$ the Chern character of $V|X_{\alpha}^g$ with g -action (see below). Put

$$\tau(g, X_{\alpha}^g) = \left\{ \frac{ch(V|X_{\alpha}^g)(g) \cdot \prod_{\theta} \mathcal{U}^{\theta}(N_{\alpha}^g(\theta)) \cdot \mathcal{T}(X_{\alpha}^g)}{\det(1 - g|(N_{\alpha}^g)^*)} \right\} [X_{\alpha}^g]$$

and

$$\tau(g) = \sum_{\alpha} \tau(g, X_{\alpha}^g).$$

Note that in the definition of $\tau(g, X_{\alpha}^g)$ the terms except $ch(V|X_{\alpha}^g)(g)$ depend only on the base space X_{α}^g . We have

Theorem 2.7. ([AS])

$$\sum_{i \geq 0} (-1)^i \text{Tr}(g | H^i(X, \mathcal{O}(V))) = \tau(g).$$

To use the Lefschetz fixed point formula we have to classify the fixed points (sets). We classify (the irreducible components of) the fixed points of G in the following sense. Let Φ_1 and Φ_2 be the fixed points (sets). Φ_1 and Φ_2 is called *equivalent* if there is an element of G which maps Φ_1 biholomorphically to Φ_2 . Let Φ be one of fixed points (sets) and let

$$C(\Phi) = \{g \in G \mid g(x) = x \text{ for any } x \in \Phi\}.$$

Let $g \in C(\Phi)$ and $H = \langle g \rangle$ the subgroup of $C(\Phi)$ which is generated by g and let \widehat{H} be the character group of H . Let $\chi \in \widehat{H}$ and let V_{χ} be the subbundle of V consisting of the vectors

$v \in V|\Phi$ such that $g(v) = \chi(g)v$ for any $g \in C(\Phi)$. $V|\Phi$ is a direct sum of subbundles V_χ 's. We define

$$ch(V|\Phi)(g) := \sum_{\chi} \chi(g) ch(V_\chi).$$

Now we return to our case. The fixed points (sets) in the quotient space $X_2(4mN)$ were classified in [G]. The fixed points in the divisor at infinity are classified easily. In total there are 25 kinds of fixed points (sets). Let Φ be one of these fixed points (sets). We can prove

Lemma 2.8. *The direct summands \tilde{V}_χ of $\tilde{V}_m|\Phi$ are also flat vector bundles.*

Hence for $M \in C(\Phi)$ we have

$$ch(\tilde{V}_m|\Phi)(M) = \sum_{\chi} \chi(M) ch(\tilde{V}_\chi) = \sum_{\chi} \chi(M) \text{rank } \tilde{V}_\chi = \text{Tr } \overline{u(M)}$$

and

$$ch(\tilde{V}_m \otimes \tilde{H}_g^{\otimes(2k-1)} \otimes [D]^{\otimes(-1)}|\Phi)(M) = \text{Tr } \overline{u(M)} ch(\tilde{H}_g^{\otimes(2k-1)} \otimes [D]^{\otimes(-1)}|\Phi)(M).$$

Therefore we can apply the data when we computed the dimension of $S_k(\Gamma_2)$ by using the holomorphic Lefschetz fixed point formula ([Ts1]) and what we have to do is

- (a) to determine $\det(CZ + D)^{1/2} u(M)$ for $M \in C(\Phi)$,
 - (b) to evaluate the Gaussian sums which appear in $\text{Tr } \overline{u(M)}$,
- and
- (c) to execute a terrible computation.

§3. The case $m = 1$

The case of index one is very important concerning Saito-Kurokawa lifting.

Proposition 3.1.

$$\begin{aligned} \sum_{k=0}^{\infty} \dim J_{k,1}^{cusp}(\Gamma_2) t^k &= \sum_{k=0}^{\infty} \text{JacobiTwo}[k,1] t^k + t^3 \\ &= \frac{t^{10} + t^{12} + t^{14} + 2t^{16} + t^{18} + t^{21} - t^{26} + t^{27} - t^{28} + t^{29} + t^{35}}{(1-t^4)(1-t^6)(1-t^{10})(1-t^{12})}. \end{aligned}$$

Proof. Let φ_4 be the Eisenstein series of degree 2 and weight 4. Then if $f \in J_{k,1}^{cusp}(\Gamma_2)$, $\varphi_4 f \in J_{k+4,1}^{cusp}(\Gamma_2)$. Since $\dim J_{k,1}^{cusp}(\Gamma_2) = \text{JacobiTwo}[k,1] = 0$ for $k = 4, 5, 6, 7$, we have $\dim J_{k,1}^{cusp}(\Gamma_2) = 0$ for $k = 0, 1, 2, 3$. On the other hand we have $\text{JacobiTwo}[k,1] = 0$ for $k = 0, 1, 2$ and $\text{JacobiTwo}[3,1] = -1$. Hence the equality of the first line holds. \square

We use the following theorem concerning the surjectivity of Φ -operator ([D]).

Theorem 3.2. If k is even and $k \geq 6$, then

$$\dim J_{k,m}(\Gamma_2) = \dim J_{k,m}^{cusp}(\Gamma_2) + d_m^1 \dim J_{k,m}^{cusp}(\Gamma_1) + d_m^0 \dim J_{k,m}^{cusp}(\Gamma_0),$$

where $d_m^1, d_m^0 \geq 1$. We define that $J_{k,m}^{cusp}(\Gamma_0) = \mathbf{C}$.

Remark 3.3. If m is square free, then $d_m^1 = d_m^0 = 1$.

From above results we have

Corollary 3.4.

$$\sum_{k=0}^{\infty} \dim J_{k,1}(\Gamma_2) t^k = \frac{t^4 + t^6 + t^{10} + t^{12} + t^{21} + t^{27} + t^{29} + t^{35}}{(1-t^4)(1-t^6)(1-t^{10})(1-t^{12})}.$$

Proof. Let $M_k(\Gamma_g)$ be the space of Siegel modular forms of weight k with respect to Γ_g . We have $\dim M_4(\Gamma_3) = 1$ ([Ty]). Let α_4 the base of $M_4(\Gamma_3)$ and let $f_{4,1}$ be the coefficient of $e(Z_{33})$ in the Fourier-Jacobi expansion of α_4 . Since $f_{4,1}$ is not identically zero, we have $\dim J_{4,1}(\Gamma_2) \geq 1$. On the other hand we have $\dim J_{4,1}(\Gamma_2) \leq \dim J_{4,1}^{cusp}(\Gamma_2) + \dim J_{4,1}(\Gamma_1) = 0 + 1 = 1$. Hence we have $\dim J_{4,1}(\Gamma_2) = 1$ and Φ -operator is also surjective in case $k = 4$. Since $\sum_{k=0}^{\infty} \dim J_{k,1}(\Gamma_1) t^k = \frac{t^4 + t^6}{(1-t^4)(1-t^6)}$, we have

$$\sum_{k=0}^{\infty} \dim J_{k,1}(\Gamma_2) t^k = \sum_{k=0}^{\infty} \dim J_{k,1}^{cusp}(\Gamma_2) t^k + \frac{t^4 + t^6}{(1-t^4)(1-t^6)}.$$

□

Remark 3.5. $\bigoplus_{k=0}^{\infty} J_{k,1}(\Gamma_2)$ is a $\bigoplus_{k=0}^{\infty} M_k(\Gamma_2)$ -module. Because we have

$$\sum_{k=0}^{\infty} \dim M_k(\Gamma_2) t^k = \frac{1+t^{35}}{(1-t^4)(1-t^6)(1-t^{10})(1-t^{12})}$$

([Ig1]), $\bigoplus_{k=0}^{\infty} J_{k,1}(\Gamma_2)$ does not have a nice structure as a $\bigoplus_{k=0}^{\infty} M_k(\Gamma_2)$ -module but will be a free $\bigoplus_{k=0}^{\infty} M_{2k}(\Gamma_2)$ -module of rank 8.

Remark 3.6. Let $M_{2k-1/2}^+(\Gamma_0^2(4))$ be the plus space in $M_{2k-1/2}(\Gamma_0^2(4))$. Then there is an isomorphism between $J_{2k,1}(\Gamma_2)$ and $M_{2k-1/2}^+(\Gamma_0^2(4))$ ([Kh], [Ib1]). The dimension of $M_{k-1/2}(\Gamma_0^2(4))$ was computed in [Ts3] and the structure of $\bigoplus_{k=0}^{\infty} M_{k-1/2}(\Gamma_0^2(4))$ was determined in [Ib2]. Hence if one finds the generators of $\bigoplus_{k=0}^{\infty} M_{2k-1/2}^+(\Gamma_0^2(4))$, he can find the generators of $\bigoplus_{k=0}^{\infty} J_{2k,1}(\Gamma_2)$. According to T. Ibukiyama, he found the generators of $\bigoplus_{k=0}^{\infty} M_{2k-1/2}^+(\Gamma_0^2(4))$.

APPENDIX

Since we explained nothing about the detailed computation, we show the computation of the case of degree one here. Of course our result coincides with the result of Eichler-Zagier ([EZ] p.105 and p.121). In the computation we use Lemma A.2. We can also compute the dimension of $J_{k,m}^{cusp}(\Gamma)$ for any congruence subgroup Γ of $SL(2, \mathbf{Z})$ (cf. [Ts2] §1).

Theorem A.1. *If $k \geq 3$, then the dimension of the space of Jacobi cusp forms $J_{k,m}^{cusp}(\Gamma_1)$ is given by the following Mathematica function:*

```

JacobiOne[k_,m_]:=Block[{a,r,p,lk,x,y},
  mod[x_,y_]:=Mod[x,y]+1;
  r=0;
  While[EvenQ[m/2^r],r++];
  p=m/2^r;
  a=m*(2*k-15)/2^3/3+Sum[Mod[x^2,4*m],{x,1,2*m-1}]/m/2^3;
  a=a+If[Mod[k,2]==0,1,-1]*((2*k-15)/2^3/3+Mod[m,4]/2^3);
  lk={1,-1,-1,1};
  a=a+If[Mod[m,2]==0,1,0]*lk[[mod[k,4]]]/2^2;
  lk={0,-1,-1,0,1,1};
  a=a+lk[[mod[k,6]]]/2/3;
  If[Mod[m,3]==0,Goto[label1],0];
  lk={0,-1,1};
  a=a+If[Mod[p,3]==1,1,-1]*If[Mod[r,2]==0,-1,1]*lk[[mod[k,3]]]/2/3;
  Goto[label2];
Label[label1];
lk={2,-1,-1};
a=a+lk[[mod[k,3]]]/2/3;
Label[label2];
Return[a];
]

```

Proof. Let $M \in SL(2, \mathbf{Z})$ and let $u(M) = (u_{rs}(M))_{r,s}$. We denote $u_{rs}(M)$ by u_{ab} , where $r = \frac{a}{2m}$ and $s = \frac{b}{2m}$.

(a) Let $M = 1_2$. We choose $(cz+d)^{1/2} = 1^{1/2} = 1$. Then $u(M) = 1_{2m}$. Hence $\text{Tr}(\overline{u(M)}) = 2m$. So we have

$$\begin{aligned} \tau(1_2, \overline{X}_1(4mN)) &= (2m) \dim S_{k-1/2}(\Gamma_1(4mN)) \\ &= 2m \left(\left(k - \frac{3}{2} \right) \frac{(4mN)^3}{24} - \frac{(4mN)^2}{4} \right) \prod (1 - p^{-2}), \end{aligned}$$

where p is over the prime numbers which divide $4mN$. Therefore the contribution of 1_2 to the dimension of $J_{k,m}^{cusp}(\Gamma_1)$ is

$$\frac{\tau(1_2, \overline{X}_1(4mN))}{[\Gamma_1 : \Gamma_1(4mN)]} = 2m \left(\left(k - \frac{3}{2} \right) \frac{1}{24} - \frac{1}{16mN} \right).$$

(b) Let $M_r = \begin{pmatrix} 1 & r \\ 0 & 1 \end{pmatrix}$ ($1 \leq r \leq 4mN - 1$) and let $\zeta = \mathbf{e}(1/4mN)$. M_r fixes the cusp P_∞ of $\overline{X}_1(4mN)$. We choose $(cz+d)^{1/2} = 1^{1/2} = 1$. Then we have $u_{aa} = \mathbf{e}(a^2r/4m)$ and $u_{ab} = 0$, otherwise. Hence

$$\begin{aligned} \tau(M_r, P_\infty) &= \frac{ch(\tilde{V}_m \otimes \overline{H}_1^{\otimes(2k-1)} \otimes [D]^{\otimes(-1)}|P_\infty)(M_r)}{(1 - M_r|(N_\alpha^{M_r})^*)} \\ &= \frac{\text{Tr } \overline{u(M_r)} ch(\overline{H}_1^{\otimes(2k-1)} \otimes [D]^{\otimes(-1)}|P_\infty)(M_r)}{(1 - \zeta^{-r})} \\ &= \sum_{a=0}^{2m-1} \mathbf{e}(-a^2r/4m) \frac{\zeta^{-r}}{(1 - \zeta^{-r})}. \end{aligned}$$

For an integer a , we denote by $\text{Mod}[a, 4mN]$ the integer such that $a \equiv \text{Mod}[a, 4mN] \pmod{4mN}$ and $0 \leq \text{Mod}[a, 4mN] < 4mN$. Then similarly as [Ts1] Example (5.4), the sum of the contributions of M_r ($1 \leq r \leq 4mN - 1$) is equal to

$$\begin{aligned} \frac{1}{|C(P_\infty)|} \sum_{r=1}^{4mN-1} \sum_{a=0}^{2m-1} \frac{\mathbf{e}(-a^2r/4m)}{(\zeta^r - 1)} &= \frac{1}{8mN} \sum_{a=0}^{2m-1} \sum_{r=1}^{4mN-1} \frac{\zeta^{-a^2rN}}{(\zeta^r - 1)} \\ &= \frac{1}{8mN} \sum_{a=0}^{2m-1} \left(\frac{1 - 4mN}{2} + \text{Mod}[a^2, 4m] \cdot N \right) \\ &= -\frac{m}{2} + \frac{1}{8N} + \frac{1}{8m} \sum_{a=1}^{2m-1} \text{Mod}[a^2, 4m]. \end{aligned}$$

(c) Let $M = -1_2$. We choose $(cz+d)^{1/2} = (-1)^{1/2} = i$. Then $u_{-aa} = -i$ and $u_{ab} = 0$, otherwise. Hence $\text{Tr } (\overline{u(M)}) = \overline{u_{00} + u_{mm}} = 2i$. Therefore the contribution of -1_2 is

$$\frac{\tau(-1_2, \overline{X}_1(4mN))}{[\Gamma_1 : \Gamma_1(4mN)]} = 2(-1)^k \left(\left(k - \frac{3}{2} \right) \frac{1}{24} - \frac{1}{16mN} \right).$$

(d) Let $M = -M_r$. We choose $(cz+d)^{1/2} = (-1)^{1/2} = i$. Then $u_{-aa} = -i \cdot \mathbf{e}(a^2r/4m)$ and $u_{ab} = 0$, otherwise. Hence $\text{Tr } (\overline{u(M)}) = \overline{u_{00} + u_{mm}} = i \cdot (1 + \mathbf{e}(-m^2r/4m))$. Therefore the sum of the contributions of $-M_r$ ($1 \leq r \leq 4mN - 1$) is equal to

$$\begin{aligned} \frac{(-1)^k}{8mN} \sum_{r=1}^{4mN-1} \frac{1 + \zeta^{-m^2rN}}{(\zeta^r - 1)} &= \frac{(-1)^k}{8mN} ((1 - 4mN) + \text{Mod}[m^2, 4m] \cdot N) \\ &= (-1)^k \left(-\frac{1}{2} + \frac{1}{8mN} + \frac{\text{Mod}[m, 4]}{8} \right). \end{aligned}$$

(e) Let $M_i = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. M_i fixes $i = \sqrt{-1}$. We choose $(cz + d)^{1/2} = i^{1/2} = (1+i)/\sqrt{2}$. Then $u_{ab} = (1-i)\mathbf{e}(-ab/2m)/2\sqrt{m}$. Therefore the contribution of M_i is

$$\begin{aligned} \frac{1}{|C(i)|} \frac{1+i}{2\sqrt{m}} \sum_{a=0}^{2m-1} \mathbf{e}(a^2/2m) \frac{ch(\overline{H}_1^{\otimes(2k-1)}|i)(M_i)}{(1-M_i|(N_\alpha^{M_i})^*)} &= \frac{1+i}{8\sqrt{m}} \frac{((1+i)/\sqrt{2})^{2k-1}}{1+1} \sum_{a=0}^{2m-1} \mathbf{e}(a^2/2m) \\ &= \begin{cases} \frac{(i)^k(1+i)}{8}, & \text{if } m \text{ is even,} \\ 0, & \text{if } m \text{ is odd.} \end{cases} \end{aligned}$$

The contribution of M_i^{-1} is the complex conjugate of the contribution of M_i .

(f) Let $\rho = \mathbf{e}(1/3)$ and $M_\rho = \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$. M_ρ fixes ρ . We choose $(cz + d)^{1/2} = (-\rho)^{1/2} = i\rho^2$. Then $u_{ab} = (1+i)\mathbf{e}(a(2b-a)/4m)/2\sqrt{m}$. Therefore the contribution of M_ρ is

$$\begin{aligned} \frac{1}{|C(\rho)|} \frac{1-i}{2\sqrt{m}} \sum_{a=0}^{2m-1} \mathbf{e}(-a^2/4m) \frac{ch(\overline{H}_1^{\otimes(2k-1)}|\rho)(M_\rho)}{(1-M_\rho|(N_\alpha^{M_\rho})^*)} &= \frac{1-i}{12\sqrt{m}} \frac{(i\rho^2)^{2k-1}}{1-\rho^2} \sum_{a=0}^{2m-1} \mathbf{e}(-a^2/4m) \\ &= \frac{(-\rho)^{k+1}}{6(1-\rho^2)}. \end{aligned}$$

The contribution of M_ρ^{-1} is the complex conjugate of the contribution of M_ρ .

(g) Let $M = M_\rho^2 = \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$. M fixes ρ . We choose $(cz + d)^{1/2} = (\rho^2)^{1/2} = \rho$. Then $u_{ab} = (-1-i)\mathbf{e}(b(b+2a)/4m)/2\sqrt{m}$. Therefore the contribution of M is

$$\begin{aligned} \frac{1}{|C(\rho)|} \frac{-1+i}{2\sqrt{m}} \sum_{a=0}^{2m-1} \mathbf{e}(-3a^2/4m) \frac{ch(\overline{H}_1^{\otimes(2k-1)}|\rho)(M)}{(1-M|(N_\alpha^M)^*)} &= \frac{-1+i}{12\sqrt{m}} \frac{\rho^{2k-1}}{1-\rho} \sum_{a=0}^{2m-1} \mathbf{e}(-3a^2/4m) \\ &= \begin{cases} \frac{(\rho^2)^k}{6}, & \text{if } 3 \mid m, \\ \left(\frac{p}{3}\right) \cdot \frac{(\rho^2)^{k+1}(-1)^{r+1}}{6(1-\rho)}, & \text{if } 3 \nmid m, \end{cases} \end{aligned}$$

where $m = 2^r p$ and p is an odd integer. $\left(\frac{p}{3}\right)$ means the Legendre symbol. The contribution of M^{-1} is the complex conjugate of the contribution of M . \square

Lemma A.2. Let $\zeta = \mathbf{e}(1/4mN)$ and let $m = 2^r p$, where p is an odd integer. Then we have

$$(1) \quad \sum_{r=1}^{4mN-1} \frac{\zeta^{-ar}}{(1-\zeta^r)} = \frac{4mN-1}{2} - \text{Mod}[a, 4mN].$$

$$(2) \quad \sum_{a=0}^{2m-1} \mathbf{e}(a^2/2m) = \begin{cases} (1+i)\sqrt{2m}, & \text{if } m \text{ is even,} \\ 0, & \text{if } m \text{ is odd.} \end{cases}$$

$$(3) \quad \sum_{a=0}^{2m-1} \mathbf{e}(3a^2/4m) = \begin{cases} (1+i)\sqrt{3m}, & \text{if } 3 \mid m, \\ \left(\frac{p}{3}\right)(-1)^{r+1}(-1+i)\sqrt{m}, & \text{if } 3 \nmid m. \end{cases}$$

REFERENCES

- [AMRT] A. Ash, D. Mumford, M. Rapoport and Y. Tai, *Smooth Compactification of Locally Symmetric Varieties (Lie Groups: History, Frontiers and Applications, Vol 4.)*, Math. Sci. Press, Brookline MA, 1975.
- [AS] M. F. Atiyah and I. M. Singer, *The index of elliptic operator, III*, Ann. of Math. **87** (1968), 540–608.
- [D] J. Dulinsky, *A decomposition theorem for Jacobi forms*, Math. Ann. **303** (1995), 473–498.
- [EZ] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Progr. Math. **55** (1985), Birkhäuser, Boston.
- [G] E. Gottschling, *Die Uniformisierbarkeit der Fixpunkte eigentlich diskontinuierlicher Gruppen von biholomorphen Abbildungen*, Math. Ann. **169** (1967), 26–54.
- [Ib1] T. Ibukiyama, *On Jacobi forms and Siegel modular forms of half integral weights*, Comm. Math. Univ. St. Pauli **41** (1992), 109–124.
- [Ib2] ———, *On Siegel modular forms of half integral weight of $\Gamma_0(4)$ of degree two* (in preparation).
- [Ig1] J.-I. Igusa, *On Siegel modular forms of genus two, II*, Amer. J. Math. **86** (1964), 392–412.
- [Ig2] ———, *A desingularization problem in the theory of Siegel modular functions*, Math. Ann. **168** (1967), 228–260.
- [Kd] K. Kodaira, *On a differential geometric method in the theory of analytic stacks*, Proc. Nat. Acad. Sci. U. S. A. **39** (1953), 1268–1273.
- [Kh] W. Kohnen, *Modular forms of half integral weight on $\Gamma_0(4)$* , Math. Ann. **248** (1980), 249–266.
- [Nk] S. Nakano, *On complex analytic vector bundles*, J. Math. Soc. Japan **7** (1955), 1–12.
- [Nm] Y. Namikawa, *A new compactification of the Siegel space and degeneration of abelian varieties, I, II*, Math. Ann. **221** (1976), 97–141, 201–241.
- [Sa] I. Satake, *On the compactification of the Siegel space*, J. Indian Math. Soc. **20** (1956), 259–281.
- [Sh] G. Shimura, *On certain reciprocity-laws for theta functions and modular forms*, Acta Math. **141** (1978), 35–71.
- [Si] C. L. Siegel, *Moduln Abelscher Funktionen*, Nach. Akad. Wiss. Göttingen **25** (1960), 365–427.
- [Ts1] R. Tsushima, *On the spaces of Siegel cusp forms of degree two*, Amer. J. Math. **104** (1982), 843–885.
- [Ts2] ———, *Dimension formula for the spaces of Siegel cusp forms and a certain exponential sum*, Mem. Inst. Sci. Tech. Meiji Univ. **36** (1997), 1–56.
- [Ts3] ———, *Dimension formula for the spaces of Siegel cusp forms of half integral weight and degree two* (in preparation).
- [Ts4] ———, *Dimension formula for the spaces of Jacobi forms of degree two* (in preparation).
- [Ty] S. Tsuyumine, *On Siegel modular forms of degree three*, Amer. J. Math. **108** (1986), 755–862.
- [Y] T. Yamazaki, *Jacobi forms and a Maass relation for Eisenstein series*, J. Fac. Sci. Univ. Tokyo, Sect. IA **33** (1986), 295–310.
- [Z] C. Ziegler, *Jacobi forms of higher degree*, Abh. Math. Sem. Univ. Hamburg **59** (1989), 191–224.