

COMPUTATIONS OF CHOW RINGS AND THE MOD p MOTIVIC COHOMOLOGY OF CLASSIFYING SPACES

Nobuaki Yagita (柳田 伸顕)
Faculty of Education, Ibaragi University (茨城大学教育学部)

ABSTRACT. In this note, we explain how to compute mod p motivic cohomology over \mathbb{C} , the complex number field, by only using algebraic topology. Examples of algebraic spaces X are classifying spaces BG of algebraic groups.

1. CHOW RING, MILNOR K-THEORY, ÉTALE COHOMOLOGY

We use some category Spc of (algebraic) spaces, defined by Voevodsky, where schemes A , quotients A_1/A_2 and $colim(A_\alpha)$ are all contained ([Vo2],[Mo-Vo]). Here schemes are defined over a field k with $ch(k) = 0$. The motivic cohomology is the double indexed cohomology defined by Suslin and Voevodsky directly related with the Chow ring, Milnor K-theory and étale cohomology,


(CH) For a smooth scheme X , $H^{2n,n}(X) = CH^n(X)$: the classical Chow group.

(MK) $H^{n,n}(Spec(k)) \cong K_n^M(k)$, the Milnor K-group for the field k .

For a smooth variety X of $dim(X) = n$. The Chow ring is the sum $CH^*(X) = \bigoplus_i CH^i(X)$ where

$$CH^i(X) = \{(n-i)\text{cycles in } X\} / (\text{rational equivalence}).$$


Here the rational equivalence $a \equiv b$ is defined if there is a codimension i subvariety W in $X \times \mathbb{P}^1$ such that $a = p_* f^*(0)$ and $b = p_* f^*(1)$ where \mathbb{P}^1 is the projective line, p (resp. f) is the projection for the first (resp. second) factor.



The multiplications in $CH^*(X)$ is giving by intersections of cycles. Let $k = \mathbb{C}$. Let \mathbb{P}^n be the n -dimensional projective space. Then $CH^i(\mathbb{P}^n) \cong \mathbb{Z}\{L_{n-i}\}$ where $L_{n-i} \cong \mathbb{P}^{n-i}$ is an $n-i$ -dimensional subspace of \mathbb{P}^n . Hence the product is $L_{n-i} \cdot L_{n-j} = L_{n-i-j}$. This shows that

$$CH^*(\mathbb{P}^n) \cong \mathbb{Z}[y]/(y^{n+1}) \cong H^*(\mathbb{C}P^n)$$

identifying $y^i = L_{n-i}$.



1991 Mathematics Subject Classification. Primary 55P35, 57T25; Secondary 55R35, 57T05.
Key words and phrases. motivic cohomology, Chow ring.

Since Spc contains colimit , we can consider the infinite projective space $P^\infty = BG_m$ and the infinite Lens space $\text{colim}_n(A^n - \{0\}/Z/p) = L_p^\infty = BZ/p$. The Chow rings of BZ/p are given in [To 1]

$$(1.1) \quad CH^*(P^\infty) \cong H^{2*,*}(P^\infty) \cong Z[y], \quad CH^*(BZ/p) \cong H^{2*,*}(BZ/p) \cong Z[y]/(py)$$

with $\text{deg}(y) = (2, 1)$. For product of these spaces

$$(1.2) \quad CH^*(P^\infty \times \dots \times P^\infty) \cong Z[y_1, \dots, y_n]$$

$$(1.3) \quad CH^*(BZ/p \times \dots \times BZ/p) \cong Z[y_1, \dots, y_n]/(py_1, \dots, py_n).$$

Here note that $CH^*(X) \not\cong H^{\text{even}}(X(C))$ for the last case. Even if $H^*(X(C))$ is generated by even dimensional elements, there are cases that $CH^*(X) \not\cong H^*(X(C))$, e.g., the K3-surfaces have the cohomology $H^2(X(C)) \cong Z^{22}$ but there is a K3-surface such that $CH^1(X) \cong Z^i$ for each $1 \leq i \leq 20$.

The Milnor K-theory is the graded ring $\bigoplus_n K_n^M(k)$ defined by $K_n^M(k) = (k^*)^{\otimes n}/J$ where the ideal J is generated by elements $a \otimes (1-a)$ for $a \in k^*$. Hence $K_0^M(k) = Z$ and by definition $K_1^M(k)$ is just the multiplicative group k^* but written additively in the ring $K_*^M(k)$. Hilbert's theorem 90, which is essentially said that the Galois cohomology $H^1(G(k_s/k); k_s^*) = 0$, implies the isomorphism $K_1^M(k)/p \cong k^*/(k^*)^p \cong H^1(G(k_s/k); Z/p)$ for $1/p \in k$. Similarly we can define a map (the norm residue map) for any extension F of k of finite type

$$(BK) \quad K_n^M(F)/p \rightarrow H^n(G(F_s/F); \mu_p^{\otimes n})$$

where $\mu_p^{\otimes n}$ is the discrete $G(F_s/F)$ -module of n -th tensor power of the group of p -roots of 1.

The Bloch-Kato conjecture is that this map is an isomorphism for all field k and the Milnor conjecture is its $p = 2$ case. This conjecture is solved when $n = 2$ by Merkurjev-Susulin[Me-Su], and for $p = 2$ by Voevodsky [Vo1] by using the motivic cohomology.

Notice that $H^n(G(k_s/k); \mu_p^{\otimes n}) \cong H_{\text{et}}^n(\text{Spec}(k), \mu_p^{\otimes n})$ the étale cohomology of the point. The étale cohomology $H_{\text{et}}^*(X; Z/p)$ has the properties ;


(E.1) If k contains a primitive p -th root of 1, then there is the additive isomorphism

$$H_{\text{et}}^m(X, \mu_p^{\otimes n}) \cong H_{\text{et}}^m(X; Z/p).$$

(E.2) For smooth X over $k = C$,

$$H_{\text{et}}^m(X; Z/p^N) \cong H^m(X(C); Z/p^N) \quad \text{for all } N \geq 1.$$

The last cohomology is the usual mod p ordinary cohomology of C -rational point of X . Of course $H_{\text{et}}^*(\text{Spec}(C); Z/p) \cong Z/p$. It is known that $K_*^M(R)/2 \cong H_{\text{et}}^*(\text{Spec}(R); Z/2) \cong Z/2[\rho]$ with $\text{deg}(\rho) = 1$ for the real number field. Here $\rho = \{-1\} \in K_1^M(R) = R^*/R^2$. Let F_v be a local field with residue field k_v of $\text{ch}(k_v) \neq 2$. Then $K_*^M(F_v)/2 \cong H_{\text{et}}^*(\text{Spec}(F_v); Z/2) \cong \Lambda(\alpha, \beta)$ with $\text{deg}(\alpha) = \text{deg}(\beta) = 1$. Thus we know $\bigoplus_m H^{m,m}(pt; Z/2)$ for these cases.



2. THE REALIZATION MAP

In this section we consider the relation to the usual ordinary cohomology. Let R be \mathbb{Z} or \mathbb{Z}/p . The motivic cohomology has the following properties [Vo2].

(C1) $H^{*,*}(X; R)$ is a bigraded ring natural in X .

(C2) There are maps (realization maps)

$$t_c^{m,n} : H^{m,n}(X; R) \rightarrow H^m(X(C); R)$$

which sum up $t_c^{*,*} = \bigoplus_{m,n} t_c^{m,n}$ the natural ring homomorphism.

(C3) There are (the Bockstein, the reduced powers) operations

$$\beta : H^{*,*}(X; \mathbb{Z}/p) \rightarrow H^{*+1,*}(X; \mathbb{Z}/p)$$

$$P^i : H^{*,*}(X; \mathbb{Z}/p) \rightarrow H^{*+2(p-1)i, *+(p-1)i}(X; \mathbb{Z}/p)$$

which commutes with the realization map t_c .

(C4) For the projective space \mathbb{P}^n , there is an isomorphism

$$H^{*,*}(X \times \mathbb{P}^n/\mathbb{P}^{n-1}; R) \cong H^{*,*}(X; R)\{1, y'\}$$

with $\deg(y') = (2n, n)$ and $t_c(y') \neq 0$.

Here we consider some examples. Recall $H^*(CP^\infty = P^\infty(C); \mathbb{Z}/p) \cong \mathbb{Z}/p[y]$, $\deg(y) = 2$ and $H^*(B\mathbb{Z}/p(C) = B\mathbb{Z}/p; \mathbb{Z}/p) \cong \mathbb{Z}/p[y] \otimes \Lambda(x)$ with $\beta x = y$ (when $p = 2$, $y = x^2$). From the above properties (C1), (C2), we easily see that t_c is epic for $X = P^\infty$. Moreover there is $x' \in H^{1,1}(B\mathbb{Z}/p; \mathbb{Z}/p)$ such that $t_c(x') = x$ and from (C2), we also see t_c is epic for $X = B\mathbb{Z}/p$.

To see these facts hold for other spaces, we recall the Lichtenbaum motivic cohomology [Vo2]. Lichtenbaum defined the similar cohomology $H_L^{*,*}(X; R)$ by using the étale topology, while $H^{*,*}(X; R)$ is defined by using Nisnevich topology. Since Nisnevich covers are some restricted étale covers, there is the natural map $H^{*,*}(X; R) \rightarrow H_L^{*,*}(X; R)$. We say that the condition $B(n, p)$ holds if

$$B(n, p) : H^{m,n}(X; \mathbb{Z}(p)) \cong H_L^{m,n}(X; \mathbb{Z}(p)) \text{ for all } m \leq n + 1$$

and all smooth X . The Beilinson-Lichtenbaum conjecture is that $B(n, p)$ holds for all n, p . It is proved that the $B(n, p)$ condition is equivalent the Bloch-Kato conjecture (BK) for degree n and prime p . Hence $B(n, p)$ holds for $n \leq 2$ or $p = 2$. Moreover Suslin-Voevodsky proves



(L-E) If $1/p \in k$, then for all X ,

$$H_L^{m,n}(X; \mathbb{Z}/p) \cong H_{\text{ét}}^m(X; \mu_p^{\otimes n}).$$

Now we compute $H^{*,*}(pt = \text{Spec}(k); \mathbb{Z}/p)$. For a smooth X , it is known the following dimensional conditions:

(C5) For a smooth X , if $H^{m,n}(X; R) \neq 0$, then

$$m \leq n + \dim(X), m \leq 2n \text{ and } m \geq 0.$$



Hereafter this paper, we assume that k contains a primitive p -th root of 1 and $B(n, p)$ holds for all n but $X = \text{Spec}(k)$. Then

$$H^{m,n}(pt; Z/p) \cong H_{et}^m(pt; \mu_p^{\otimes n}) \cong H_{et}^m(pt; Z/p) \quad \text{if } m \leq n$$

and $H^{m,n}(pt; Z/p) \cong 0$ otherwise. Let $\tau \in H^{0,1}(pt; Z/p)$ be the element corresponding a generator of $H_{et}^0(\text{Spec}(k); \mu_p) \cong H_{et}^0(\text{Spec}(k); Z/p)$. Then we get the isomorphism

$$H^{*,*}(\text{Spec}(k); Z/p) \cong H_{et}^*(\text{Spec}(k); Z/p) \otimes Z/p[\tau]$$

since $\tau : H_{et}^m(pt; \mu_p^{\otimes n}) \cong H_{et}^m(pt; \mu_p^{\otimes(n+1)})$. Inparticular, for the real number field R and a local field F_v with the residue field k_v of $ch(k_v) \neq 2$

$$(2.1) \quad H^{*,*}(\text{Spec}(R); Z/2) \cong Z/2[\rho, \tau] \quad \text{with } deg(\rho) = (1, 1)$$

$$(2.2) \quad H^{*,*}(\text{Spec}(F_v); Z/2) \cong Z/2[\tau] \otimes \Lambda(\alpha, \beta) \quad \text{with } deg(\alpha) = deg(\beta) = (1, 1).$$

For $k = C$, $B(n, p)$ condition holds for $X = \text{Spec}(C)$, indeed $K_n^M(C) \cong 0$ for $n > 0$. Therefore

$$(2.3) \quad H^{*,*}(\text{Spec}(C); Z/p) \cong Z/p[\tau] \quad \text{with } deg(\tau) = (0, 1).$$

When $k = C$, if $B(n, p)$ condition holds for X , then it is immediate that

$$(2.4) \quad [\tau^{-1}]H^{*,*}(X; Z/p) \cong H^*(X(C); Z/p) \otimes Z/p[\tau, \tau^{-1}]$$

where the degree is defied by $deg(x) = (m, m)$ if $x \in H^m(X(C); Z/p)$.

Next we compute cohomology of P^∞ and BZ/p . For any (algebraic) map $f : X \rightarrow Y$ in the category Spc , we can construct the cofiber sequence

$$X \rightarrow Y \rightarrow cone(f) = Y/X$$

which induces the long exact sequence (Voevodsky [V2])

$$(2.5) \quad H^{*,*}(X; R) \leftarrow H^{*,*}(Y; R) \leftarrow H^{*,*}(Y/X; R) \leftarrow H^{*-1,*}(X; R).$$

In particular, we get the Mayer-Vietoris, Gysin and blow up long exact sequences.

By the cofiber sequence $P^{n-1} \rightarrow P^n \rightarrow P^n/P^{n-1}$ and (C4), we can inductively see that

$$(2.6) \quad H^{*,*}(P^n; Z/p) \cong H^{*,*}(pt; Z/p) \otimes Z/p[y]/(y^{n+1}) \quad \text{with } deg(y) = (2, 1)$$


Since $B(1, p)$ is always holds, $H^{1,1}(L_p^n; Z/p) \cong H^1(L_p^n; Z/p)$. Hence there is the element $x' \in H^{1,1}(L_p^n; Z/p)$ with $t_c(x') = x \in H^1(L_p^n; Z/p)$. The Lens space is identified with the sphere bundle associated with the line bundle

$$(A^n - \{0\}) \times_{(A - \{0\})} A \rightarrow (A^n - \{0\}) / (A - \{0\}) = P^n.$$


Where $(A^n - \{0\}) \times_{(A - \{0\})} A$ is the identification such that $(z_i, z) \sim (a^{-1}z_i, a^p z) \in (A^n - \{0\}) \times A$. Hence we get the cofibering $L_p^n \rightarrow P^n \xrightarrow{x_p} P^n$. Thus we get the additive isomorphism $H^{*,*}(L_p^n; Z/p) \cong H^{*,*}(P^n; Z/p)\{1, x\}$. This induces the ring isomorphism for $p = odd$

$$(2.7) \quad H^{*,*}(L_p^n; Z/p) \cong Z/p[y]/(y^{n+1}) \otimes \Lambda(x) \otimes H^{*,*}(pt; Z/p) \quad \text{with } deg(x) = (1, 1).$$

However note that when $p = 2$, we see $x^2 = y\tau + x\rho$ [Vo3] where $\rho \in H^{1,1}(pt; Z/p) \cong k^*/k^{2*}$ represents -1 . (hence $\rho = 0$ when $\sqrt{-1} \in k^*$.) This is proved by the wellknown facts $\{a, a\} = \{a, -1\} \in K_2^M(k)$.



$H^{*,*}(\text{Spec}(R); Z/2)$



$H^{*,*}(BZ/2; Z/2)$

$R = C$

Let us say that a space X satisfies the Kunneth formula for a space Y if $H^{*,*}(X \times Y; Z/p) \cong H^{*,*}(X; Z/p) \otimes_{H^{*,*}(pt; Z/p)} H^{*,*}(Y; Z/p)$.

By the above cofiber sequences, we can easily see that P^∞ and BZ/p satisfy the Kunneth formula for all spaces. In particular, we have the ring isomorphisms

$$(2.8) \quad H^{*,*}(P^\infty \times \dots \times P^\infty; Z/p) \cong Z/p[y_1, \dots, y_n] \otimes H^{*,*}(pt; Z/p)$$

$$(2.9) \quad H^{*,*}(BZ/p \times \dots \times BZ/p; Z/p) \cong Z/p[y_1, \dots, y_n] \otimes \Lambda(x_1, \dots, x_n) \otimes H^{*,*}(pt; Z/p)$$

(when $p = 2$, $x_i^2 = y_i \tau + x_i \rho$).

This fact is used to defined the reduced power operation P^i in (C3). Since the Sylow p subgroup of the symmetric group S_p of p -letters, is isomorphic to Z/p , we know the isomorphism

$$H^*(BS_p; Z/p) \cong H^*(BZ/p; Z/p)^{F_p} \cong Z/p[Y] \otimes \Lambda(x)$$

with identifying $Y = y^{p-1}$ and $X = xy^{p-2}$. If X is smooth (and suppose p is odd for easy of arguments), we can define the reduced powers (of Chow rings) as follows. Consider maps

$$H^{2*,*}(X; Z/p) \xrightarrow{i_1} H^{2p*,*}(X^p \times_{S_p} ES_p) \xrightarrow{\Delta^*} H^*(X; Z/p) \otimes_{H^{*,*}} H^{*,*}(BS_p; Z/p)$$

where i_1 is the Gysin map for p -th external power, and Δ is the diagonal map. For $deg(x) = (2n, n)$, the reduced powers are defined as

$$(2.10) \quad \Delta^* i_1(x) = \sum P^i(x) \otimes Y^{n-i} + \beta P^i(x) \otimes XY^{n-i-1}.$$

Hence note $deg(P^i) = deg(Y^i) = deg(y^{i(p-1)}) = (2i(p-1), i(p-1))$.

Voevodsky defined i_1 for non smooth X also and by using suspensions maps, he defined reduced powers for all degree elements in $H^{*,*}(X; Z/p)$ for all X [Vo 3].

Moreover we can see (Ho-Kriz [H-K])

$$(2.11) \quad H^{*,*}(BGL_n; Z/p) \cong Z/p[c_1, \dots, c_n] \otimes H^{*,*}(pt; Z/p)$$

where the Chern class c_i with $deg(c_i) = (2i, i)$ are identified with the elementary symmetric polynomial in $H^{*,*}(P^\infty \times \dots \times P^\infty; Z/p)$. So we can define the Chern class $\rho^*(c_i) \in H^{2*,*}(BG; Z/p)$ for each algebraic group G and for each representation $\rho: G \rightarrow GL_n$.

3. $H^{*,*}(X; Z/p)/Ker(t_c)$ AND OPERATION Q_i

In this section we always assume that X is smooth and $k = C$. Define a bidegree algebra by

$$(3.1) \quad h^{*,*}(X; Z/p) = \oplus_{m,n} H^{m,n}(X; Z/p)/Ker(t_c^{m,n}).$$

Suppose that $B(n, p)$ condition holds. By isomorphisms (B, p) , (L-E), (E1) and (E2), we have

$$H^{n,n}(X; Z/p) \cong H_L^{n,n}(X; Z/p) \cong H_{et}^n(X; \mu_p^{\otimes n}) \cong H_{et}^n(X; Z/p) \cong H^n(X(C); Z/p).$$

The realization map $t_c^{n,n}$ induces this isomorphism. Let $F_i = Im(t_c^{*,i})$. Then $\cup_i F_i = H^*(X(C); Z/p)$ and define the graded algebra $grH^*(X(C); Z/p) = \oplus F_{i+1}/F_i$. Thus we get the additive isomorphism

$$h^{*,*}(X; Z/p) \cong grH^*(X(C); Z/p) \otimes Z/p[\tau]$$

of bigraded rings. However the ring structures of both rings are different, in general. The cohomology $h^{*,*}(X; Z/p)$ is isomorphic to a $Z[\tau]$ -subalgebra B of $H^*(X(C); Z/p) \otimes Z/p[\tau, \tau^{-1}]$

with $\text{deg}(x) = (|x|, |x|)$ such that $B[\tau^{-1}] \cong H^*(X(\mathbb{C}); \mathbb{Z}/p) \otimes \mathbb{Z}/p[\tau, \tau^{-1}]$. Namely there is a \mathbb{Z}/p -basis $\{a_I\}$ of $H^*(X(\mathbb{C}); \mathbb{Z}/p)$ such that $B = \mathbb{Z}/p\{\tau^{-t_I} a_I\} \otimes \mathbb{Z}/p[\tau]$ for some $t_I \geq 0$.

Here we recall the Milnor primitive operation $Q_i = [Q_{i-1}, P^{p^{i-1}}]$

$$Q_i : H^{*,*}(X; \mathbb{Z}/p) \rightarrow H^{*+2p^{i-1}, *+p^{i-1}}(X; \mathbb{Z}/p)$$

which is derivative, $Q_i(xy) = Q_i(x)y + xQ_i(y)$. Note also $Q_i(\tau) = 0$ by dimensional reason of $H^{*,*}(pt; \mathbb{Z}/p) \cong \mathbb{Z}/p[\tau]$.


Lemma 3.1. *If $0 \neq Q_{i_1} \dots Q_{i_n} x \in H^{2*,*}(X; \mathbb{Z}/p)$, then x is a $\mathbb{Z}/p[\tau]$ -module generator.*

Proof. If $x = x'\tau$, then $\tau Q_{i_1} \dots Q_{i_n}(x') \neq 0$. But $Q_{i_1} \dots Q_{i_n}(x') = 0 \in H^{2*,* - 1}(X; \mathbb{Z}/p)$ since $H^{m,n}(X; \mathbb{Z}/p) = 0$ for $m > 2n$. \square

Define the weight by $w(x) = 2n - m$ for an element $x \in H^{m,n}(X; \mathbb{Z}/p)$ so that $w(x') = 0$ for $x' \in CH^*(X)/p$. Of course we get $w(xy) = w(x) + w(y)$, $w(P^i x) = w(x)$ and $w(Q_i(x)) = w(x) - 1$.

Corollary 3.2. *Suppose that $B(n, p)$ holds. If $x \in H^n(X(\mathbb{C}); \mathbb{Z}/p)$ and $Q_{i_1} \dots Q_{i_n}(x) \neq 0$, then there is a $\mathbb{Z}/p[\tau]$ -module generator $x' \in H^{n,n}(X; \mathbb{Z}/p)$ so that $t_c(x') = x$ and for each $0 \leq k \leq n$, $Q_{i_1} \dots Q_{i_k}(x')$ is also a $\mathbb{Z}/p[\tau]$ -module generator of $H^{*,*}(X; \mathbb{Z}/p)$.*

Proof. By $B(n, p)$ condition, $t_c^{n,n} : H^{n,n}(X; \mathbb{Z}/p) \cong H^n(X(\mathbb{C}); \mathbb{Z}/p)$. Hence there is an element $x' \in H^{n,n}(X; \mathbb{Z}/p)$ with $t_c(x') = x$. This means $w(x') = n$ and $w(Q_{i_1} \dots Q_{i_n}(x)) = 0$. From the above lemma, we get the corollary. \square



Now we consider the examples. The mod 2 cohomology of $BO(n)$ is $H^*(BO(n); \mathbb{Z}/2) \cong \mathbb{Z}/2[w_1, \dots, w_n]$ where the Stiefel-Whitney class w_i restricts the elementary symmetric polynomial in $H^*(B(\mathbb{Z}/2)^n; \mathbb{Z}/2) \cong \mathbb{Z}/2[x_1, \dots, x_n]$. Each element w_i^2 is represented by Chern class c_i of the induced representation $O(n) \subset U(n)$. Hence $c_i \in CH^*(BS(n); \mathbb{Z}/2) = H^{2*,*}(BO(n); \mathbb{Z}/2)$.

Proposition 3.3. $h^{*,*}(BO(n); \mathbb{Z}/p) \supset \mathbb{Z}/2[c_1, \dots, c_n] \otimes \Delta(w_1, \dots, w_n) \otimes \mathbb{Z}/2[\tau]$ where $\text{deg}(c_i) = (2i, i)$, $\text{deg}(w_i) = (i, i)$ and $w_i^2 = \tau^i c_i$.


Since $Q_{i-1} \dots Q_0(w_i) \neq 0$, each w_i is a $\mathbb{Z}/2[\tau]$ -module generator. However even $h^{*,*}(BO(n); \mathbb{Z}/2)$ seems very complicated. Consider the case $X = BO(3)$. The cohomology operations act by

$$\begin{array}{ccccccc} w_2 & \xrightarrow{Sq^1} & w_1 w_2 + w_3 & \xrightarrow{Sq^2} & w_2 w_1^2 + w_1^2 w_3 + w_1 w_2^2 + w_2 w_3 & \xrightarrow{Sq^1} & w_1^2 w_2^2 + w_3^2 \\ w_3 & \xrightarrow{Sq^1} & w_3 w_1 & \xrightarrow{Sq^2} & w_1 w_2 w_3 & & \end{array}$$

Theorem 3.4. *There is the isomorphism*

$$h^{*,*}(BO(3); \mathbb{Z}/2) \cong \mathbb{Z}/2[c_1, c_2, c_3] \{1, w_1, w_2, Q_0 w_2, Q_1 w_2, w_3, Q_0 w_3, Q_1 w_3\} \otimes \mathbb{Z}/2[\tau].$$

where $Q_0 w_2 = \tau^{-1}(w_1 w_2 + w_3)$, ...



W.S.Wilson ([W],[K-Y]) found a good $Q(i) = \Lambda(Q_0, \dots, Q_i)$ -module decomposition for $X = BO(n)$, namely,

$$(3.2) \quad H^*(X; Z/2) = \bigoplus_{-1} Q(i)G_i \quad \text{with} \quad Q_0 \dots Q_i G_i \in t_c(CH^*(X)).$$

Here G_{k-1} is quite complicated, namely, it is generated by symmetric functions

$$\sum x_1^{2i_1+1} \dots x_k^{2i_k+1} x_{k+1}^{2j_1} \dots x_{k+q}^{2j_q}, \quad k+q \leq n,$$

with $0 \leq i_1 \leq \dots \leq i_k$ and $0 \leq j_1 \leq \dots \leq j_q$; and if the number of j equal to j_u is odd, then there is some $s \leq k$ such that $2i_s + 2^s < 2j_u < 2i_s + 2^{s+1}$.

Then $w(G_i) \geq i$ in $h^{*,*}(X; Z/p)$, that means

Proposition 3.5. *Givenig the weight by $w(G_i) = i+1$, we have the incusion for $X = BO(n)$*

$$h^{*,*}(X; Z/2) \subset (\bigoplus_i Q(i)G_i) \otimes Z/2[\tau].$$

One problem is that the above inclusion is really isomorphism or not. The similar decomposition holds for $X = (BZ/p)^n$ and the above inclusion is an isomorphism. The case $X = BO(3)$ is also isomorphism. Since the direct decomposition of $BO(3)$ is complicated to write, we only write here that of $SO(3)$ since $O(3) \cong SO(3) \times Z/2$.

$$(3.3) \quad \begin{aligned} H^*(BSO(3); Z/2) &\cong Z/2[w_2, w_3] \cong Z/2[c_2, c_3]\{1, w_2, w_3 = Q_0 w_2, w_2 w_3 = Q_1 w_2\} \\ &\cong Z/2[c_2, c_3]\{w_2, Q_0 w_2, Q_1 w_2, c_3 = Q_0 Q_1 w_2\} \oplus Z/2[c_2] \\ &\cong Z/2[c_2, c_3]Q(1)\{w_2\} \oplus Z/2[c_2]. \end{aligned}$$

Since there is the isomorphism $O(2n+1) \cong SO(2n+1) \times Z/2$, the cohomology of $BSO(2n+1)$ is reduced from that of $BO(2n+1)$. However note that the situation for $BO(2n)$ is different.

The extraspecial 2-group 2_+^{1+2n} is the n -th central product of the dihedral group D_8 of order 8. It has a central extension

$$(3.4) \quad 0 \rightarrow Z/2 \rightarrow G \rightarrow V = \bigoplus^{2n} Z/2 \rightarrow 0$$

Let $H^*(BG; Z/2) \cong Z/2[x_1, \dots, x_{2n}]$. Then Quillen proved [Q2]

$$(3.5) \quad H^*(BG; Z/2) \cong Z/2[x_1, \dots, x_{2n}]/(f, Q_0 f, \dots, Q_{n-2} f) \otimes Z/2[w_{2^n}].$$

Here w_{2^n} is the Stiefel-Whitney class of the real 2^n dimensional irreducible representation restricting non zero on the center and $f = \sum_i x_{2i-1} x_{2i} \in H^2(BV; Z/2)$ represents the central extension (3.4).

Leting $y_i = x_i^2$ in $H^*(BG; Z/2)$, we can write $f^2 = \sum y_{2i-1} y_{2i}$,

$$(Q_{k-1} f)^2 = Q_0 Q_k f = \sum y_{2i-1}^2 y_{2i} - y_{2i-1} y_{2i}^2$$

NOBUAKI YAGITA

$$Q_{k-1}f = \sum y_{2i-1}^{2^{k-1}} x_{2i} - x_{2i-1} y_{2i}^{2^{k-1}}.$$

Now we consider in the motivic cohomology $H^{*,*}(BG; Z/2)$ and change $y_i = \tau^{-1}x_i^2$. Since $f = 0 \in H^{2,2}(BG; Z/2)$, we can see that $Q_{k-1}f = 0$ and $Q_k Q_0(f) = 0$ also in $H^{*,*}(BG; Z/2)$. However for general n , $\sum y_{2i} y_{2i-1} \neq 0$ in $H^{*,*}(BG; Z/2)$. Let

$$(3.6) \quad A = (Z/2[y_1, \dots, y_{2n}, c_{2^n}] / (Q_0 Q_k f, \dots, Q_0 Q_n f) \\ \otimes \Delta(x_1, \dots, x_2, w_{2^n}) / (f, Q_0 f, \dots, Q_{n-2} f) \otimes Z/2[\tau].$$

Lemma 3.6. For $G = 2_+^{1+2n}$, there is a map $A \rightarrow H^{*,*}(BG; Z/2)$ which induces the injection $A/(f^2) \subset h^{*,*}(BG; Z/2)$.

When $m = 0, 1, -1 \pmod 8$ and $m > 0$, we say that $Spin(m)$ is real type [Q2]. When $Spin(m)$ is real type, from Quillen, we know that $H^*(BSpin(m); Z/2) \subset H^*(BG; Z/2)$ where $G = 2_+^{2h+1}$, and h is the Hurwitz number (for details see [Q2]).

Corollary 3.7. Let $G = Spin(m)$ be real type and the Hurwitz number h , and let

$$A = (Z/2[c_2, c_3, \dots, c_m, c_{2^h}] / ((Q_1 Q_0 w_2), \dots, (Q_h Q_0 w_2)) \\ \otimes \Delta(w_2, \dots, w_m, w_{2^h}) / (w_2, Q_0 w_2, \dots, Q_{h-2} w_2) \otimes Z/2[\tau]$$

where $w_i, i \leq m$ (resp. w_{2^h}) is the Stiefel-Whitney class of the usual $SO(m)$ representation (resp. of the irreducible 2^h -dimensional spin representation). Then we have a map $A \rightarrow H^{*,*}(BG; Z/2)$ which induces the injection $A/(c_2) \subset h^{*,*}(BG; Z/2)$.

We study $Spin(7)$ and the exceptional Lie group G_2 . The cohomology of G_2 is given by $H^*(BG_2; Z/2) \cong Z/2[w_4, w_6, w_7]$ where w_i is the Siefel-Whitney class of the inclusion $G_2 \subset SO(7)$. The cohomology $H^*(BSpin(7); Z/2) \cong H^*(BG_2; Z/2) \otimes Z/2[w_8]$.

Corollary 3.8. Let $A = Z/2[c_2, c_4, c_6, c_7] \otimes \Delta(w_4, w_6, w_7) \otimes Z/2[\tau]$. Then there is the map $A \rightarrow H^{*,*}(BG_2; Z/2)$ which induces the injection $A/(c_2) \subset h^{*,*}(BG_2; Z/2)$. Similar facts hold for $BSpin(7)$ tensoring $Z/2[c_8]$.

The cohomology operations are given

$$w_4 \xrightarrow{Sq^2} w_6 \xrightarrow{Sq^1} w_7 \xrightarrow{Sq^4} w_4 w_7 \xrightarrow{Sq^2} w_7 w_6 \xrightarrow{Sq^1} w_7^2 \\ Q_1 Q_0(w_4 w_6) = w_7^2, \quad Q_2 Q_1 Q_0(w_4 w_6 w_7) = w_7^4.$$

Proposition 3.9. Let $w(w_4) = 2, w(w_{(4,6)}) = 2$ and $w(w_{(4,6,7)}) = 3$ with $t_c(w_{(i_1, \dots, i_n)}) = w_{i_1} \dots w_{i_n}$. Then we have the injection

$$h^{*,*}(BG_2; Z/2) \subset Z/2[c_4, c_6, c_7]$$

$$\otimes Z/2\{1, w_4, Sq^2 w_4, Q_1 w_4, Q_2 w_4, Sq^2 Q_2 w_4, w_{(4,6)}, w_{(4,6,7)}\} \otimes Z/p[\tau].$$

Remark. If $t_c^{4,3} \otimes Q$ is epic, then we can take $w_4 \in h^{4,3}(BG; Z/2)$, i.e., $w(w_4) = 2$. The kernel $Ker(t_c)^{2*,*}$ is not so big for $X = BG_2$. Indeed, it is known that

$$CH^*(BG_2) \cong Z_{(2)}[c_2, c_4, c_6, c_7] / (2^r(c_2^2 - 4c_4), 2c_7, c_2 c_7), \quad \text{for some } r \geq 0.$$

The cohomology operations are given in $H^*(BSO(7); Z/2)$

$$Q_1 Q_0 w_2 = w_3^2, \quad Q_2 Q_0 w_2 = w_5^2, \quad Q_3 Q_0 w_2 = w_7^2 w_2^2 + w_6^2 w_3^2 + w_5^2 w_4^2.$$

Hence we have $c_3 = 0, c_5 = 0, c_2 c_7 = 0$ in $CH^*(BG_2)$ but $c_2 \neq 0$.

From here we consider the case $p = \text{odd}$. One of the easiest examples is the case $G = PGL_3$ and $p = 3$. The mod 3 cohomology is given by ([K-Y],[Ve1])

$$(Z/3[y_2]\{y^2\} \oplus Z/3\{1, y_2, y_3, y_7\}[y_8]) \otimes Z/3[y_{12}]$$

It is known that y_2^2, y_2^3, y_8^2 and y_{12} are represented by Chern classes. Moreover $Q_1Q_0(y_2) = y_8$. Hence these elements are in the Chow ring, namely,

$$h^{2*,*}(BPGL_3; Z/3) \cong (Z/3[y_2]\{y_2^2\} \oplus Z/3[y_8]) \otimes Z/3[y_{12}].$$

The cohomology operations are given

$$y_2 \xrightarrow{\beta} y_3 \xrightarrow{P^1} y_7 \xrightarrow{\beta} y_8$$

Thus we get $h^{*,*}(PGL_3; Z/3)$ completely.

Theorem 3.10.

$$h^{*,*}(BPGL_3; Z/3) \cong (Z/3[y_2]\{y^2\} \oplus Z/3\{1\} \oplus Z/3[y_8] \otimes Q(1)\{y_2\}) \otimes Z/3[y_{12}] \otimes Z/3[\tau]$$

Next consider the extraspecial p -group $G = p_+^{1+2n}$. When $n > 2$, even the cohomology ring $H^*(BG(C); Z/p)$ are unknown, while it contains the subring

$$B = Z/p[y_1, \dots, y_{2n}, c_{p^n}]/(Q_1Q_0f, \dots, Q_nQ_0f).$$

where $f = \sum^n x_{2i-1}x_{2i}$ for $\beta x_i = y_i$ and $Q_kQ_0f = \sum y_{2i-1}y_{2i}^{p^k} - y_{2i-1}^{p^k}y_{2i}$. Since $f = 0 \in H^{2,2}(BG; Z/p)$, we have

Proposition 3.11. *Let $G = p_+^{1+2n}$ and $A = B \otimes Z/p[\tau]$. Then there is an injection $A \subset H^{*,*}(BG; Z/p)$*

We consider the case $n = 1$ here. Let us write $E = p_+^{1+2}$. The ordinary cohomology is known by Lewis [L], [Te-Y3], namely,

$$H^{even}(BE)/p \cong (Z/p[y_1, y_2]/(y_1^p y_2 - y_1 y_2^p) \oplus Z/p\{c_2, \dots, c_{p-1}\}) \otimes Z/p[c_p].$$

$$H^{odd}(BE) \cong Z/p[y_1, y_2, c_p]\{a_1, a_2\}/(y_1 a_2 - y_2 a_1, y_1^p a_2 - y_2^p a_1) \quad |a_i| = 3.$$

Theorem 3.12.

$$h^{*,*}(BE; Z/p) \cong (\{1, \partial^{-1}\}(H^*(BE)/p) - \{\partial^{-1}1\}) \otimes Z/p[\tau]$$

where $w(H^{even}(BE)/p) = 0, w(H^{odd}(BE)) = 1$ and ∂_p^{-1} ascends the weight one.

Proof. Since all elements in $H^{even}(BE)$ are generated by Chern classes, we have the isomorphism $h^{2*,*}(BG; Z/3) \cong H^{even}(BE)/p$. We know $H^{odd}(BE; Z/p)$ is generated as a $H^{even}(BE)/p$ -module by two elements a_1, a_2 such that $Q_1 a_i = y_i c_p$ [Te-Y3].

The mod p -cohomology is written additively $H^*(BE; Z/p) \cong \{1, \partial_p^{-1}\}H^*(BE)/p$. Here ∂_p is the (higher) Bockstein. All elements in $H^{odd}(BE)$ are just p -torsion and we can take $a'_i \in H^2(BE; Z/p)$ such that $\beta(a'_i) = a_i$. Thus we take $a'_i \in H^{2,2}(BE; Z/p)$ so that $a_i \in H^{3,2}(BE; Z/p)$.

Next consider elements $x = \partial_p^{-1}(y)$, $y \in H^{even}(BE)/p$. If $y \in (Ideal(y_1, y_2))$, then $\partial_p^{-1}(y) = \sum x_i b_i$ for $b_i \in H^{even}(BE)/p$, and hence we can take $w(\partial_p^{-1}(y)) = 1$. For other elements $y = c; c$ with $c \in Z/p[c_p]$, we can prove ([Ly]) that these elements are represented by transfer from a subgroup isomorphic to $Z/p \times Z/p$. Therefore we can also prove that $w(\partial_p^{-1}(y)) = 1$. Thus we complete the proof. \square

NOBUAKI YAGITA

REFERENCES

- [F] R.Field On the Chow ring of classifying space $BSO(2n, C)$ preprint (2000).
- [G-L] D. J. Green and I. J. Leary. Chern classes and extraspecial groups. *Manuscripta Math.* **88** (1995), 73-84.
- [H-K] Po Hu and I.Kriz. Some remarks on real and algebraic cobordim. preprint (2000).
- [I] K.Inoue The Brown-Peterson cohomology of $BSO(6)$. *J. Math. Kyoto Univ.* **32** (1992), 655-666.
- [K-Y] A. Kono and N. Yagita. Brown-Peterson and ordinary cohomology theories of classifying spaces for compact Lie groups. *Transactions of AMS.* **339** (1993), 781-798.
- [L] I. J. Leary. The integral cohomology rings of some p groups. *Math.Proc.Cambridge Philos.Soc.* **110** (1991), 245-255.
- [L-M] M. Levin and F. Morel. Cobordisme algébrique I,II. preprints (2001)
- [Me-Su] A.Merkurjev and A.Susulin. K -cohomology of Severi-Brauer varieties and the norm residue homomorphism. *Math. UssR Izvestiya* **21** (1983), 307-340.
- [Mo-Vo] F. Morel and V. Voevodsky. A^1 -homotopy theory of scheme. To appear (1998)
- [P] R. Pandharipande. Equivariant Chow rings of $O(k)$, $SO(2k+1)$, and $SO(4)$. *J. reine angew. Math.* **496** (1998) 131-148.
- [Q1] D. Quillen A cohomological criterion for p -nilpotence. *J.Pure and Applied Algebra* **1** (1971), 361-372.
- [Q2] D. Quillen. The mod 2 cohomology rings of extra-special 2-groups and the spinor groups. *Math. Ann.* **194** (1971), 197-212.
- [R-W-Y] D. C. Ravenel, W. S. Wilson and N. Yagita. Brown-Peterson cohomology from Morava K -theory. *K-theory* **15** (1998), 147-199.
- [Sc-Y] B. Schuster and N. Yagita. Transfer of chern classes in BP -cohomology and Cow rings. To appear in *Trans. AMS*.
- [Su-Vo] Motivic cohomology with finite coefficients and Bloch-Kato conjecture. preprint (1995).
- [Ta] H. Tamanoi The image of the BP Thom map for Eilenberg-MacLane spaces *Trans. AMS* **349** (1997) 1209-1237.
- [Te-Y1] M. Tezuka and N.Yagita. The varieties of the mod p cohomology rings of extraspecial p -groups for an odd prime p . *Math. Proc. Cambridge Phil. Soc.* **94** (1983) 449-459.
- [Te-Y2] M. Tezuka and N. Yagita Cohomology of finite groups and the Brown-Peterson cohomology. *Lecture Notes in Math.* **1370** (1989) 396-408.
- [Te-Y3] M. Tezuka and N. Yagita On odd prime components of cohomologies of sporadic simple groups and the ring of universal atable elements. *J.Algebra* **183** (1996) 483-513.
- [Tod] H.Toda Cohomology mod 3 of the classifying space BF_4 of the exceptional group F_4 . *J.Math.Kyoto Univ.* **13** (1973) 97-115.
- [To1] B.Totaro Torsion algebraic cycles and complex cobordism. *J. Amer. Math. Soc.* **10** (1997), 467-493.
- [To2] B.Totaro The Chow ring of classifying spaces. *Proc. of Symposia in Pure Math. "Algebraic K-theory"(1997:University of Washington,Seattle)* **67** (1999) 249-281.
- [Ve1] G.Vezzosi. On the Chow ring of the classifying space of $PGL_{3,C}$. *J.reine angew. Math.* **523** (2000), 1-54.
- [Ve2] G.Vezzosi. Brown-Peterson spectra in stable A^1 -homotopy theory. Preprint (2000).
- [Vo1] V. Voevodsky. The Milnor conjecture. Preprint (1996).
- [Vo2] V. Voevodsky (Noted by Weibel). Voevodsky's Seattle lectures : K -theory and motivic cohomology *Proc. of Symposia in Pure Math. "Algebraic K-theory"(1997:University of Washington,Seattle)* **67** (1999) 283-303.
- [Vo3] V. Voevodsky. Reduced power operations in motivic cohomology. preprint (2001).
- [Y1] N. Yagita. On relations between Brown-Peterson cohomology and the ordinary mod p cohomology theory. *Kodai Math.J* **7** (1984), 273-285.
- [Y2] N. Yagita. Chow ring of classifying spaces of extraspecial p groups , preprint (2000).

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, IBARAKI UNIVERSITY, MITO, IBARAKI, JAPAN
 E-mail address: yagita@mito.ipc.ibaraki.ac.jp