

# Univalence of certain integral operators

Virgil Pescar and Shigeyoshi Owa

## Abstract

Let  $\mathcal{A}_n$  be the class of functions  $f(z)$  which are analytic and  $n$ -fold symmetric in the open unit disk  $\mathbb{U}$ . The integral operator  $G_\alpha(z)$  for  $f(z) \in \mathcal{A}_n$  is considered. The object of the present paper is to derive univalence conditions of the integral operator  $G_\alpha(z)$  for  $f(z) \in \mathcal{A}_n$ .

## 1 Introduction

Let  $\mathcal{A}_n$  denote the class of functions  $f(z)$  of the form

$$f(z) = z + \sum_{k=1}^{\infty} a_{nk+1} z^{nk+1} \quad (n \in \mathbb{N} = \{1, 2, 3, \dots\})$$

which are analytic and  $n$ -fold symmetric in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$ . We denote by  $\mathcal{S}_n$  the subclass of  $\mathcal{A}_n$  consisting of functions  $f(z)$  which are univalent in  $\mathbb{U}$ . Many authors studied the problem of integral operators for functions  $f(z)$  in the class  $\mathcal{S}_1$ . In this sense, the following useful result is due to Pfaltzgraß [3].

**Theorem 1.1.** *If  $f(z)$  is univalent in  $\mathbb{U}$  and  $\alpha$  is complex number with  $|\alpha| \leq \frac{1}{4}$ , then the integral operator  $G_\alpha(z)$  given by*

$$G_\alpha(z) = \int_0^z (f'(t))^\alpha dt \tag{1}$$

*is also univalent in  $\mathbb{U}$ .*

Further, Pascu and Pescar [2] gave

**Theorem 1.2.** *If  $f(z) \in \mathcal{S}_1$  and  $\alpha$  is a complex number with  $|\alpha| \leq \frac{1}{4n}$ , then the integral operator  $G_{\alpha,n}(z)$  given by*

$$G_{\alpha,n}(z) = \int_0^z (f'(t))^\alpha dt$$

*is also in the class  $\mathcal{S}_1$  for all positive integer  $n$ .*

---

2000 *Mathematics Subject Classification:* Primary 30C45

*Key Words and Phrases:* Univalent,  $n$ -fold symmetric, integral operator.

## 2 Properties of integral operators

To discuss our problems for integral operators, we need to recall here the following lemma due to Becker [1].

**Lemma 2.1.** *If  $f(z) \in \mathcal{A}_1$  satisfies*

$$(1 - |z|^2) \left| \frac{zf''(z)}{f'(z)} \right| \leq 1 \quad (z \in \mathbb{U}), \quad (2)$$

then  $f(z) \in \mathcal{S}_1$ .

Applying the above lemma, we derive

**Theorem 2.1.** *If  $f(z) \in \mathcal{A}_1$  satisfies the inequality (2) for all  $z \in \mathbb{U}$ , then the integral operator  $G_\alpha(z)$  defined by (1) belongs to the class  $\mathcal{S}_1$  for all  $\alpha$  ( $|\alpha| \leq 1$ ).*

**Proof.** Note that  $G_\alpha(z) \in \mathcal{A}_1$  for  $f(z) \in \mathcal{A}_1$  and that

$$\frac{zf''(z)}{f'(z)} = \frac{1}{\alpha} \frac{zG_\alpha''(z)}{G_\alpha'(z)}.$$

It follows that

$$(1 - |z|^2) \left| \frac{zG_\alpha''(z)}{G_\alpha'(z)} \right| = |\alpha|(1 - |z|^2) \left| \frac{zf''(z)}{f'(z)} \right| \leq |\alpha| \leq 1$$

for  $z \in \mathbb{U}$ . Thus, using Lemma 2.1, we have  $G_\alpha(z) \in \mathcal{S}_1$ .

Next, we prove

**Corollary 2.1.** *If  $f(z) \in \mathcal{A}_1$  satisfies*

$$\left| \frac{f''(z)}{f'(z)} \right| \leq 1 \quad (z \in \mathbb{U}),$$

then the integral operator  $G_\alpha(z)$  defined by (1) is in the class  $\mathcal{S}_1$  with  $|\alpha| \leq \frac{3\sqrt{3}}{2}$ .

**Proof.** In view of the proof of Theorem 2.1, we see that

$$(1 - |z|^2) \left| \frac{zG_\alpha''(z)}{G_\alpha'(z)} \right| \leq |\alpha|(1 - |z|^2)|z| \leq 1,$$

because  $|\alpha| \leq \frac{3\sqrt{3}}{2}$  and

$$\max_{|z| \leq 1} (1 - |z|^2)|z| = \frac{2}{3\sqrt{3}}.$$

Thus, by Lemma 2.1, we prove that  $G_\alpha(z) \in \mathcal{S}_1$ .

Finally, we show

**Theorem 2.2.** *If  $f(z) \in \mathcal{A}_n$  satisfies*

$$\left| \frac{f''(z)}{f'(z)} \right| \leq |z|^{n-1} \quad (z \in \mathbb{U}),$$

*then the integral operator  $G_\alpha(z)$  defined by (1) belongs to the class  $\mathcal{S}_n$  with*

$$|\alpha| \leq \frac{(n+2)^{\frac{n+2}{2}}}{2n^{\frac{n}{2}}}.$$

**Proof.** Since

$$\frac{zf''(z)}{f'(z)} = \frac{1}{\alpha} \frac{zG_\alpha''(z)}{G_\alpha'(z)} = n(n+1)a_{n+1}z^n + \dots,$$

we have that

$$\begin{aligned} (1 - |z|^2) \left| \frac{zG_\alpha''(z)}{G_\alpha'(z)} \right| &= |\alpha|(1 - |z|^2) \left| \frac{zf''(z)}{f'(z)} \right| \\ &\leq |\alpha|(1 - |z|^2)|z|^n \quad (z \in \mathbb{U}). \end{aligned}$$

Note that

$$|\alpha| \leq \frac{(n+2)^{\frac{n+2}{2}}}{2n^{\frac{n}{2}}}$$

and

$$(1 - |z|^2)|z|^n \leq \frac{2n^{\frac{n}{2}}}{(n+2)^{\frac{n+2}{2}}} \quad (z \in \mathbb{U}).$$

This gives us that

$$(1 - |z|^2) \left| \frac{zG_\alpha''(z)}{G_\alpha'(z)} \right| \leq 1 \quad (z \in \mathbb{U}).$$

Further, it is easy to see that  $G_\alpha(z) \in \mathcal{A}_n$ . This completes the proof of the theorem.

**Remark.** For  $n = 1$ , Theorem 2.2 becomes Theorem 2.1.

## References

- [1] J. Becker, *Löwnersch Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen*, J. Reine Angew. Math. **255**(1972), 23 – 43.
- [2] N. N. Pascu and V. Pescar, *On the integral operators of Kim-Merkes and Pfaltzgraff*, Mathematica(Cluj) **32**(1990), 185 – 192.
- [3] J. A. Pfaltzgraff, *Univalence of the integral  $(f'(z))^\lambda$* , Bull. London Math. Soc. **7**(1975), 254

*V. Pescar  
Department of Mathematics  
Transilvania University of Brasov  
2200 Brasov  
Romania*

*Shigeyoshi Owa  
Department of Mathematics  
Kinki University  
Higashi-Osaka, Osaka, 577-8502  
Japan*