

## Parametrized Thue Equations — A Survey

Clemens Heuberger  
Institut für Mathematik B  
Technische Universität Graz  
8010 Graz  
Austria  
clemens.heuberger@tugraz.at\*

January 29, 2005

### Abstract

We consider families of parametrized Thue equations

$$F_a(X, Y) = \pm 1, \quad a \in \mathbf{N},$$

where  $F_a \in \mathbf{Z}[a][X, Y]$  is a binary irreducible form with coefficients which are polynomials in some parameter  $a$ .

We give a survey on known results.

## 1 Thue Equations

Let  $F \in \mathbf{Z}[X, Y]$  be a homogeneous, irreducible polynomial of degree  $n \geq 3$  and  $m$  be a nonzero integer. Then the Diophantine equation

$$F(X, Y) = m \tag{1}$$

is called a *Thue equation* in honour of A. Thue, who proved in 1909 [57]:

**Theorem 1 (Thue).** (1) has only a finite number of solutions  $(x, y) \in \mathbf{Z}^2$ .

Thue's proof is based on his approximation theorem: Let  $\alpha$  be an algebraic number of degree  $n \geq 2$  and  $\epsilon > 0$ . Then there exists a constant  $c_1(\alpha, \epsilon)$ , such that for all  $p \in \mathbf{Z}$  and  $q \in \mathbf{N}$

$$\left| \alpha - \frac{p}{q} \right| \geq \frac{c_1(\alpha, \epsilon)}{q^{n/2+1+\epsilon}}.$$

Since this approximation theorem is not effective, Thue's theorem is neither effective.

## 2 Number of Solutions

We call a solution  $(x, y)$  to  $F(x, y) = m$  primitive, if  $x$  and  $y$  are coprime integers. The problem of giving upper bounds (depending on  $m$  and the degree  $n$ ) for the number of primitive solutions goes back to Siegel. Such a bound has first been given by Evertse [14]. An improved version has been given by Bombieri and Schmidt [6]:

---

\*The author is supported by the grant S8307-MAT of the Austrian Science Fund.

**Theorem 2 (Bombieri-Schmidt [6]).** *There is an absolute constant  $c_2$  such that for all  $n \geq c_2$  the Diophantine equation  $F(X, Y) = m$  has at most  $215 \cdot n^{1+\omega(m)}$  primitive solutions, where  $\omega(m)$  denotes the number of prime factors of  $m$  and solutions  $(x, y)$  and  $(-x, -y)$  are regarded as the same.*

At least for  $m = \pm 1$ , this result is best possible (up to the constant 215), since the equation

$$X^n + (X - Y)(2X - Y) \dots (nX - Y) = \pm 1$$

has at least the  $n + 1$  solutions  $\pm\{(1, 1), \dots, (1, n), (0, 1)\}$ .

Sharper bounds have been obtained for special classes of Thue equations.

If only  $k$  coefficients of  $F(X, Y)$  are nonzero, the number of solutions depends on  $k$  and  $m$  only (and not on  $n$ ). For  $k = 3$ , this is proved by Mueller and Schmidt [41]: There are at most  $O(m^{2/n})$  solutions. The general case  $k \geq 3$  is proved in Mueller and Schmidt [42]: There are at most  $O(k^2 m^{2/n} (1 + \log m^{1/n}))$  solutions. Thomas [56] gives absolute upper bounds for the number of solutions for  $m = 1$  and  $k = 3$ : If  $n \geq 38$ , then there are at most 20 solutions  $(x, y)$  with  $|xy| \geq 2$ , where solutions  $(x, y)$  and  $(-x, -y)$  are only counted once. For smaller  $n$ , similar bounds are given.

If only 2 coefficients of  $F(X, Y)$  are nonzero, we arrive at the special case  $ax^n - by^n = \pm 1$  and we consider only the case  $ab \neq 0$ ,  $x > 0$ ,  $y > 0$ . This equation has been studied by many authors, starting with Delone [11] and Nagell [43], who proved that there is at most one solution for  $n = 3$ . Several authors have contributed to this question. Finally, Bennett [4] could prove that there is at most one solution  $(x, y)$ .

We now consider cubic Thue equations  $F(X, Y) = 1$ . If the discriminant of  $F$  is negative, there are at most 5 solutions, and the cases of 4 and 5 solutions can be listed explicitly. This has been shown independently by Delaunay [10] and Nagell [44] in the 1920's. If the discriminant is positive, there are at most 10 solutions, as it has been proved by Bennett [3]. Okazaki [47] proves that if the discriminant is at least  $5.65 \cdot 10^{65}$ , then there are at most 7 solutions. It is conjectured by Nagell [45], Pethő [48], and Lippok [35] that there are at most 5 solutions except for five equations (modulo equivalence) which have 6 or 9 solutions. We note that there are two families of cubic Thue equations which have exactly five solutions, cf. items 2 and 3 in the list in Section 4.1.

Okazaki [46] considers the analogous problem for quartic Thue equations  $F(X, Y) = \pm 1$ . If all roots of  $F(x, 1)$  are real and the discriminant is larger than a computable constant  $c_3$ , this equation has at most 14 solutions, where solutions  $(x, y)$  and  $(-x, -y)$  are counted once.

### 3 Algorithmic Solution of Single Thue Equations

Studying linear forms in logarithms of algebraic numbers, A. Baker could give an effective upper bound for the solutions of such a Thue equation in 1968 [1]:

**Theorem 3 (Baker).** *Let  $\kappa > n + 1$  and  $(x, y) \in \mathbb{Z}^2$  be a solution of (1). Then*

$$\max\{|x|, |y|\} < c_4 e^{\log^\kappa |m|},$$

where  $c_4 = c_4(n, \kappa, F)$  is an effectively computable number.

Since that time, these bounds have been improved; Bugeaud and Győry [7] give the following bound:

**Theorem 4 (Bugeaud-Győry).** *Let  $B \geq \max\{|m|, e\}$ ,  $\alpha$  be a root of  $F(X, 1)$ ,  $K := \mathbb{Q}(\alpha)$ ,  $R := R_K$  the regulator of  $K$  and  $r$  the unit rank of  $K$ . Let  $H \geq 3$  be an upper bound for the absolute values of the coefficients of  $F$ .*

*Then all solutions  $(x, y) \in \mathbb{Z}^2$  of (1) satisfy*

$$\max\{|x|, |y|\} < \exp\left(c_5 \cdot R \cdot \max\{\log R, 1\} \cdot (R + \log(HB))\right)$$

and

$$\max\{|x|, |y|\} < \exp\left(c_6 \cdot H^{2n-2} \cdot \log^{2n-1} H \cdot \log B\right),$$

with  $c_5 = 3^{r+27}(r+1)^{7r+19}n^{2n+6r+14}$  and  $c_6 = 3^{3(n+9)}n^{18(n+1)}$ .

The bounds for the solutions obtained by Baker's method are rather large, thus the solutions practically cannot be found by simple enumeration. For a similar problem Baker and Davenport [2] proposed a method to reduce drastically the bound by using continued fraction reduction. Pethő and Schulenberg [50] replaced the continued fraction reduction by the LLL-algorithm and gave a general method to solve (1) for the totally real case with  $m = 1$  and arbitrary  $n$ . Tzanakis and de Weger [61] describe the general case. Finally, Bilu and Hanrot [5] were able to replace the LLL-algorithm by the much faster continued fraction method and solve Thue equations up to degree 1000.

## 4 Families of Thue Equations

We study families of Thue equations

$$F_a(X, Y) = \pm 1, \quad a \in \mathbb{N} \quad (2)$$

where  $F_a \in \mathbb{Z}[a][X, Y]$  is an irreducible binary form of degree of at least 3 with coefficients which are integer polynomials in  $a$ . In the investigation of such families usually only two types of solutions appear: Firstly, there are *polynomial solutions*  $X(a), Y(a) \in \mathbb{Z}[a]$  which satisfy (2) in  $\mathbb{Z}[a]$ , and secondly, there occur (sometimes) single solutions for a few small values of the parameter  $a$ . However, Lettl [30] points out that the family  $X^6 - (a-1)Y^6 = a^2$  does not have any polynomial solution, but there are sporadic solutions for infinitely many values of the parameter  $a$ .

The first infinite parametrized families of Thue equations were considered by Thue [58] himself: He proved that the equation

$$(a+1)X^n - aY^n = 1, \quad X > 0, Y > 0 \quad (3)$$

has only the solution  $x = y = 1$  for  $a$  suitably large in relation to prime  $n \geq 3$ . For  $n = 3$ , the equation (3) has only this solution for  $a \geq 386$ . Of course, Bennett's result [4] cited in Section 2 implies that this is true for all  $n \geq 3$  and  $a \geq 1$ .

For a description of the techniques used to solve families of Thue equations, we refer to Heuberger [20]. Some automated procedures are presented in [26].

### 4.1 Families of Fixed Degree

In 1990, Thomas [53] investigated for the first time a parametrized family of cubic Thue equations of positive discriminant. Since 1990, the following particular families of Thue equations have been studied:

1.  $X^3 - (a-1)X^2Y - (a+2)XY^2 - Y^3 = 1$ .

Thomas [53] and Mignotte [36] proved that for  $a \geq 4$ , the only solutions are  $(0, -1)$ ,  $(1, 0)$  and  $(-1, +1)$ , while for the cases  $0 \leq a \leq 4$  there exist some nontrivial solutions, too, which are given explicitly in [53]. For the same form  $F_a(X, Y)$ , all solutions of the Thue inequality  $|F_a(X, Y)| \leq 2a + 1$  have been found by Mignotte, Pethő, and Lemmermeyer [39].

2.  $X^3 - aX^2Y - (a+1)XY^2 - Y^3 = X(X+Y)(X - (a+1)Y) - Y^3 = 1$ .

Lee [29] and independently Mignotte and Tzanakis [40] proved that for  $a \geq 3.33 \cdot 10^{23}$  there are only the solutions

$$(1, 0), (0, -1), (1, -1), (-a-1, -1), (1, -a).$$

Mignotte [37] could prove the same result for all  $a \geq 3$ .

3. Wakabayashi [66] proved that for  $a \geq 1.35 \cdot 10^{14}$ , the equation  $X^3 - a^2XY^2 + Y^3 = 1$  has exactly the five solutions  $(0, 1), (1, 0), (1, a^2), (\pm a, 1)$ .
4. Togbe [60] considered the equation  $X^3 - (n^3 - 2n^2 + 3n - 3)X^2Y - n^2XY^2 - Y^3 = \pm 1$ . If  $n \geq 1$ , the only solutions are  $(\pm 1, 0)$  and  $(0, \pm 1)$ .
5. Wakabayashi [64]:  $|X^3 + aXY^2 + bY^3| \leq a + |b| + 1$  for arbitrary  $b$  and  $a \geq 360b^4$  as well as for  $b \in \{1, 2\}$  and  $a \geq 1$ . He uses Padé approximations.
6. Thomas [55]: Let  $b, c$  be nonzero integers such that the discriminant of  $t^3 - bt^2 + ct - 1$  is negative,  $\Delta = 4c - b^2 > 0$ , and  $c \geq \min\{4.2 \times 10^{41} \times |b|^{2.32}, 3.6 \times 10^{41} \times \Delta^{1.1582}\}$ . Then the Thue equation  $X^3 - bX^2Y + cXY^2 - Y^3 = 1$  only has the trivial solutions  $(1, 0), (0, -1)$ .
7.  $X(X - a^{d_2}Y)(X - a^{d_3}Y) \pm Y^3 = 1$ .

This family was investigated by Thomas [54]. He proved that for  $0 < d_2 < d_3$  and

$$a \geq (2 \cdot 10^6 \cdot (d_2 + 2d_3))^{4.85/(d_3 - d_2)}$$

nontrivial solutions cannot exist. He also investigated this family with  $a^{d_1}$  and  $a^{d_2}$  replaced by monic polynomials in  $a$  of degrees  $d_1$  and  $d_2$ , respectively (see Theorem 5).

8.  $X^4 - aX^3Y - X^2Y^2 + aXY^3 + Y^4 = X(X - Y)(X + Y)(X - aY) + Y^4 = \pm 1$ .  
This quartic family was solved by Pethő [49] for large values of  $a$ ; Mignotte, Pethő, and Roth [38] solved it completely: The only solutions are  $\pm\{(0, 1), (1, 0), (1, 1), (1, -1), (a, 1), (1, -a)\}$  for  $|a| \notin \{2, 4\}$ . If  $|a| = 4$ , four more solutions exist. If  $|a| = 2$ , the family is reducible.
9.  $X^4 - aX^3Y - 3X^2Y^2 + aXY^3 + Y^4 = \pm 1$  has been solved for  $a \geq 9.9 \cdot 10^{27}$  by Pethő [49].
10.  $|bX^4 - aX^3Y - 6bX^2Y^2 + aXY^3 + bY^4| \leq N$ .  
For  $b = 1$  and  $N = 1$ , this equation has been solved completely by Lettl and Pethő [31]; Chen and Voutier [9] solved it independently by using the hypergeometric method. For the same form binary form  $F_{a,b}(X, Y)$ , Lettl, Pethő and Voutier [33] proved that  $|F_a(X, Y)| \leq 6a + 7$  has only trivial primitive solutions for  $a \geq 58$ , if  $b = 1$ . Furthermore,  $x^2 + y^2 \leq \max\{25a^2/(64b^2), 4N^2/a\}$  if  $a > 308b^4$ , cf. Yuan [67].
11. Togbé [59] gives all solutions to  $X^4 - a^2X^3Y - (a^3 + 2a^2 + 4a + 2)X^2Y^2 - a^2XY^3 + Y^4 = 1$  for  $a \geq 1.191 \cdot 10^{19}$  and  $a, a + 2, a^2 + 4$  squarefree.
12.  $|X^4 - a^2X^2Y^2 + Y^4| = |X^2(X - a)(X + a) + Y^4| \leq a^2 - 2$   
This family of Thue inequalities has only trivial solutions with  $|y| \leq 1$  for  $a \geq 8$  (Wakabayashi [62]).
13.  $|X^4 + 4aX^3Y + 6aX^2Y^2 + 4a^2XY^3 + a^2Y^4| \leq a^2$  has been solved for  $a \geq 205$  by Chen and Voutier [8].
14. Dujella and Jadrijević [12], [13] prove that  $|X^4 - 4cX^3Y + (6c + 2)X^2Y^2 + 4cXY^3 + Y^4| \leq 6c + 4$  has only trivial solutions for all  $c \geq 3$ .
15.  $X(X - Y)(X - aY)(X - bY) - Y^4 = \pm 1$ .  
All solutions of this two-parametric family are known for  $10^{2 \cdot 10^{28}} < a + 1 < b \leq a(1 + (\log a)^{-4})$ , cf. Pethő and Tichy [51]. The case of  $b = a + 1$  has been considered by Heuberger, Pethő and Tichy [23], where all solutions could be determined for all  $a \in \mathbb{Z}$ .
16. Jadrijević [27] proves that for every  $0.5 < s \leq 1$ , there is an effectively computable constant  $P(s)$  such that if  $a \neq 0$  and  $\max\{|a|, |b|\} \geq P(s)$  and  $\gcd(a, b) \geq \max\{|a|^s, |b|^s\}$ , then the equation  $X^4 - 2abX^3Y + 2(a^2 - b^2 + 1)X^2Y^2 + 2abXY^3 + Y^4 = 1$  only has trivial solutions. In particular,  $P(0.999) = 10^{27}$  and  $P(0.501) = 10^{36836}$ .

17. Wakabayashi [63] found all solutions of  $|X^4 - a^2X^2Y^2 - bY^4| \leq a^2 + b - 1$  for  $a \geq 5.3 \cdot 10^{10}b^{6.22}$ .
18.  $X(X^2 - Y^2)(X^2 - a^2Y^2) - Y^5 = \pm 1$ .  
For  $a > 3.6 \cdot 10^{19}$ , all solutions have been found by Heuberger [18].
19. Gaál and Lettl [15] investigated the family  $X^5 + (a-1)X^4Y - (2a^3 + 4a + 4)X^3Y^2 + (a^4 + a^3 + 2a^2 + 4a - 3)X^2Y^3 + (a^3 + a^2 + 5a + 3)XY^4 + Y^5 = \pm 1$  and found all solutions for  $|a| \geq 3.3 \cdot 10^{15}$ . The remaining cases have been solved in Gaál and Lettl [16].
20. Levesque and Mignotte [34] found all solutions of the equation  $X^5 + 2X^4Y + (a+3)X^3Y^2 + (2a+3)X^2Y^3 + (a+1)XY^4 - Y^5 = \pm 1$  for sufficiently large  $a$ .
21.  $X^6 - 2aX^5Y - (5a+15)X^4Y^2 - 20X^3Y^3 + 5aX^2Y^4 + (2a+6)XY^5 + Y^6 \in \{\pm 1, \pm 27\}$  was investigated by Lettl, Pethő, and Voutier. They found all solutions for  $a \geq 89$  by hypergeometric methods [33] and all solutions for  $a < 89$  by using Baker's method [32]. In [33], they also proved that  $|F_a(X, Y)| \leq 120a + 323$  (for the form  $F_a(X, Y)$  considered) has only trivial primitive solutions for  $a \geq 89$ .
22.  $X^8 - 8nX^7Y - 28X^6Y^2 + 56nX^5Y^3 + 70X^4Y^4 - 56nX^3Y^5 - 28nX^2Y^6 + 8nXY^7 + Y^8 = \pm 1$  has only trivial solutions for  $n \in \{a \in \mathbb{Z} : a + b\sqrt{2} = (1 + \sqrt{2})^{2k+1}, k \in \mathbb{N}\}$  with  $n \geq 6.71 \cdot 10^{32}$ . (Heuberger, Togbé and Ziegler [26]).

A more detailed survey on cubic families is contained in Wakabayashi [65].

## 4.2 Families of Relative Thue Equations

A few families of relative Thue equations have also been solved, i.e., families where the parameters and the solutions are elements of the same imaginary quadratic number field.

So let  $D > 0$  be an integer,  $k := \mathbb{Q}(\sqrt{-D})$ ,  $\mathfrak{o}_k$  its ring of algebraic integers, and  $\mu$  a root of unity in  $\mathfrak{o}_k$ .

1. For  $t \in \mathfrak{o}_k$  with  $|t| \geq 3.03 \cdot 10^9$ , the only solutions  $(x, y) \in \mathfrak{o}_k^2$  to  $X^3 - (t-1)X^2Y - (t+2)XY^2 - Y^3 = \mu$  satisfy  $\max\{|x|, |y|\} \leq 1$  and can be listed explicitly (Heuberger, Pethő, and Tichy [24]).
2. For  $t \in \mathfrak{o}_k$  with  $|t| > 2.88 \cdot 10^{33}$ , the only solutions  $(x, y) \in \mathfrak{o}_k^2$  to  $X^3 - tX^2Y - (t+1)XY^2 - Y^3 = \mu$  satisfy  $\min\{|x|, |y|\} \leq 1$  and can be listed explicitly (Ziegler [68]).
3. For  $s, t \in \mathfrak{o}_k$  with  $|t| \geq 5.3 \cdot 10^{10}|s|^{12.44}$  or  $s = 1$  and  $|t| > \sqrt{550}$ , all solutions  $(x, y) \in \mathfrak{o}_k^2$  to  $|X^4 - t^2X^2Y^2 + s^2Y^4| \leq |t|^2 - |s|^2 - 2$  are explicitly known (Ziegler [69]).

## 4.3 Families of Arbitrary Degree

Moreover, some general families of arbitrary degree have been considered. Apart from (3), the investigated general families are of the shape

$$F_a(X, Y) := \prod_{i=1}^n (X - p_i(a)Y) - Y^n = \pm 1, \quad (4)$$

where  $p_1, \dots, p_n \in \mathbb{Z}[a]$  are polynomials, which have been called *split families* by E. Thomas [54]. For  $i = 1, \dots, n$  it can easily be seen that  $(X, Y) \in \{\pm(p_i, 1), (\pm 1, 0)\}$  are solutions. Thomas conjectured that if

$$p_1 = 0, \quad \deg p_2 < \dots < \deg p_n$$

and the polynomials are monic, there are no further solutions for sufficiently large values of the parameter  $a$ . In [54] he proved this conjecture for  $n = 3$  under some technical hypothesis:

**Theorem 5.** Let  $u = \pm 1$ ,  $a(t), b(t) \in \mathbb{Z}[t]$  be monic polynomials and  $a := \deg a(t)$ ,  $b := \deg b(t)$  with  $0 < a < b$ . We write  $A(t) := a(t)/t^a - 1$  and  $B(t) := b(t)/t^b - 1$  and define for  $n \geq 1$

$$W(n) := \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j} (b \cdot A(n)^j - a \cdot B(n)^j),$$

which can be written in powers of  $1/n$  as  $W(n) = \sum_{j=1}^n w_j n^{-j}$ . Further we define  $J := \min\{j \in \mathbb{N} : w_j \neq 0\}$ .

If  $J \neq b - a$  or  $J = b - a \wedge 3w_J + 2b + a \neq 0 \wedge 3w_J - 2(b - a) \neq 0$ , then there is an effectively computable constant  $c_7$  depending on the coefficients of  $a(t)$  and  $b(t)$  such that for  $n \geq c_7$  the family of Thue equations

$$X(X - a(n)Y)(X - b(n)Y) + uY^3 = \pm 1$$

only has the solutions

$$\pm\{(1, 0), (0, u), (a(n)u, u), (b(n)u, u)\}.$$

Halter-Koch, Lettl, Pethő and Tichy [17] considered (4) for  $p_1 = 0, p_2 = d_2, \dots, p_{n-1} = d_{n-1}$  and  $p_n = a$ , where  $d_2, \dots, d_{n-1}$  are fixed distinct integers. They found all solutions for sufficiently large values of  $a$  assuming a conjecture of Lang and Waldschmidt [28]—which is a very sharp bound for linear forms in logarithms of algebraic numbers—:

**Theorem 6.** Let  $n \geq 3$ ,  $p_1 = 0, p_2 = d_2, \dots, p_{n-1} = d_{n-1}$  be distinct integers and  $p_n = a$ . Let  $\alpha = \alpha(a)$  be a zero of  $P(x) = \prod_{i=1}^n (x - p_i) - d$  with  $d = \pm 1$  and suppose that the index  $I$  of  $(\alpha - d_1, \dots, \alpha - d_{n-1})$  in  $\mathfrak{D}^\times$ , the group of units of  $\mathfrak{D} := \mathbb{Z}[\alpha]$ , is bounded by a constant  $J = J(d_1, \dots, d_{n-1}, n)$  for every  $a$  from some subset  $\Omega \in \mathbb{Z}$ . Assume further that the Lang-Waldschmidt conjecture is true. Then for all but finitely many values of  $a \in \Omega$  the Diophantine equation

$$\prod_{i=1}^n (x - p_i y) - dy^n = \pm 1$$

has only solutions  $(x, y) \in \mathbb{Z}^2$  with  $|y| \leq 1$ , except for the cases of  $n = 3$  and  $|d_2| = 1$  or  $n = 4$  and  $(d_2, d_3) \in \{(1, -1), (\pm 1, \pm 2)\}$ , where it has exactly one more solution for every value of  $a$ .

If  $\mathbb{Q}(\alpha)$  is primitive over  $\mathbb{Q}$  — especially if  $n$  is prime — then there exists a bound  $J = J(d_1, \dots, d_{n-1}, n)$  for the index  $I$  by lower bounds for the regulator of  $\mathfrak{D}$  (cf. Pohst and Zassenhaus [52], chapter 5.6, (6.22)). Applying the theory of Hilbertian fields and results on thin sets, primitivity is proved for almost all choices (in the sense of density) of the parameters, cf. [17].

The two exceptional families are those considered under 2 and 8 in the list in Section 4.1.

A similar family has been considered by Heuberger in [19], however, in this case, the result is unconditionally true:

**Theorem 7.** Let  $n \geq 4$  be an integer,  $d_2, \dots, d_{n-1}$  pairwise distinct integers and  $a$  an integral parameter. Furthermore we assume

$$d_2 + \dots + d_{n-1} \neq 0 \quad \text{or} \quad d_2 \cdots d_{n-1} \neq 0.$$

Let

$$F_a(X, Y) := (X + aY)(X - d_2Y)(X - d_3Y) \cdots (X - d_{n-1}Y)(X - aY) - Y^n.$$

Then there exists a (computable) constant  $c_8$  depending only on the degree  $n$  and  $d_2, \dots, d_{n-1}$ , such that for all  $a \geq c_8$ , the only solutions  $(x, y) \in \mathbb{Z}^2$  of the Diophantine equation

$$F_a(X, Y) = \pm 1$$

are  $\pm\{(1, 0), (-a, 1), (d_2, 1), (d_3, 1), \dots, (d_{n-1}, 1), (a, 1)\}$ .

In [25], Heuberger and Tichy considered a multivariate version of (4):

**Theorem 8.** Let  $n \geq 4$ ,  $r \geq 1$ ,  $p_i \in \mathbb{Z}[A_1, \dots, A_r]$  for  $1 \leq i \leq n$ . We make the following assumptions on the polynomials  $p_i$ :

$$\begin{aligned} \deg p_1 &< \dots < \deg p_{n-2} < \deg p_{n-1} = \deg p_n, \\ \text{LH}(p_n) &= \text{LH}(p_{n-1}), \text{ but } p_n \neq p_{n-1}. \end{aligned}$$

Furthermore we suppose that for  $p \in \{p_1, \dots, p_n, p_n - p_{n-1}\}$ , there exist positive constants  $t_p, c_p$  such that

$$|(\text{LH}(p))(a_1, \dots, a_r)| \geq c_p \cdot (\min_k a_k)^{\deg p} \quad \text{for } a_1, \dots, a_r \geq t_p.$$

Let

$$F_{a_1, \dots, a_r}(X, Y) := \prod_{i=1}^n (X - p_i(a_1, \dots, a_r)Y) - Y^n.$$

For every constant  $C > 1$  there is a constant  $t_0$  such that for all integers  $a_1, \dots, a_r$  satisfying  $t_0 \leq \min_k a_k$  and

$$\max_k a_k \leq C \cdot \min_k a_k,$$

the Diophantine equation

$$F_{a_1, \dots, a_r}(x, y) = \pm 1$$

considered for  $x, y \in \mathbb{Z}$  only has the solutions  $\{(\pm 1, 0)\} \cup \{\pm(p_i(a_1, \dots, a_r), 1) : 1 \leq i \leq n\}$ .

In Heuberger [21] Thomas' conjecture is proved under some technical hypothesis:

**Theorem 9.** Let  $n \in \mathbb{N}$ ,  $n \geq 3$  and  $p_i \in \mathbb{Z}[a]$  be monic polynomials for  $i = 1, \dots, n$ . We write

$$p_i(a) = a^{d_i} + k_i a^{d_i-1} + \text{terms of lower degree}, \quad i = 2, \dots, n,$$

allow  $p_1 = 0$  and assume

$$d_1 < d_2 < \dots < d_{n-1} < d_n \quad \text{and} \quad n + d_2 \geq 4.$$

Let

$$\delta_i := \begin{cases} 1 & \text{if } d_i - d_{i-1} = 1, \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad e := \sum_{i=2}^n d_i.$$

If  $\delta_4 = 1$  or

$$(e - d_2 + 2d_3)(k_2 - \delta_2) + (-e - 2d_2 + d_3)k_3 + (d_3 - d_2) \sum_{i=4}^n k_i \notin \{2\delta_3, -(e + d_3)\delta_3\}, \quad (5)$$

then there is a (computable) constant  $c_9 = c_9(p_1, \dots, p_n)$  depending on the coefficients of the polynomials  $p_i$  such that for all integers  $a \geq c_9$  the Diophantine equation

$$F_a(X, Y) := \prod_{i=1}^n (X - p_i(a)Y) - Y^n = \pm 1$$

only has the solutions

$$(\pm 1, 0) \text{ and } \pm(p_i(a), 1), 1 \leq i \leq n.$$

In [21], there is also a version with a stronger technical hypothesis than that in (5). For  $n = 3$ , that version improves Theorem 5.

Especially there are only trivial solutions if

$$\begin{aligned} \max(\deg p_1, 0) &< \deg p_2 < \deg p_3 < \dots < \deg p_n \\ \max(\deg p_1, 0) + \deg p_2 + \dots + \deg p_n &< 15. \end{aligned}$$

In Heuberger [22], an explicit constant  $c_9$  for Theorem 9 is given:

$$c_9 = \exp \left( 1.01(n+1)(n-1)!(n-1)^{n-2} \exp(1.04(n-2)(nd_n - n + 3)) \binom{nd_n - 1}{n-3} (2P+1)^{nd_n} \right),$$

where  $d_j = \deg p_j$  and  $P$  is an upper bound for the absolute values of the coefficients of the  $p_j$ ,  $j = 1, \dots, n$ .

## References

- [1] A. Baker, *Contribution to the theory of Diophantine equations. I. On the representation of integers by binary forms*, Philos. Trans. Roy. Soc. London Ser. A **263** (1968), 173–191.
- [2] A. Baker and H. Davenport, *The equations  $3x^2 - 2 = y^2$  and  $8x^2 - 7 = z^2$* , Quart. J. Math. Oxford **20** (1969), 129–137.
- [3] M. A. Bennett, *On the representation of unity by binary cubic forms*, Trans. Amer. Math. Soc. **353** (2001), 1507–1534.
- [4] ———, *Rational approximation to algebraic numbers of small height: the Diophantine equation  $|ax^n - by^n| = 1$* , J. Reine Angew. Math. **535** (2001), 1–49.
- [5] Yu. Bilu and G. Hanrot, *Solving Thue equations of high degree*, J. Number Theory **60** (1996), 373–392.
- [6] E. Bombieri and W. M. Schmidt, *On Thue's equation*, Invent. Math. **88** (1987), 69–81.
- [7] Y. Bugeaud and K. Györy, *Bounds for the solutions of Thue-Mahler equations and norm form equations*, Acta Arith. **74** (1996), 273–292.
- [8] J. H. Chen and P. M. Voutier, *The complete solution of  $aX^2 - bY^4 = 1$* , Preprint.
- [9] ———, *Complete solution of the Diophantine equation  $X^2 + 1 = dY^4$  and a related family of quartic Thue equations*, J. Number Theory **62** (1997), 71–99.
- [10] B. Delaunay, *Über die Darstellung der Zahlen durch die binären kubischen Formen von negativer Diskriminante*, Math. Z. **31** (1930), 1–26.
- [11] B. N. Delone, *Solution of the indeterminate equation  $x^3q + y^3 = 1$* , Izv. Akad. Nauk SSR (6) **16** (1922), 253–272.
- [12] A. Dujella and B. Jadrijević, *A parametric family of quartic Thue equations*, Acta Arith. **101** (2002), 159–170.
- [13] ———, *A family of quartic Thue inequalities*, Acta Arith. **111** (2004), 61–76.
- [14] J.-H. Evertse, *Upper bounds for the numbers of solutions of Diophantine equations*, Mathematisch Centrum, Amsterdam, 1983.
- [15] I. Gaál and G. Lettl, *A parametric family of quintic Thue equations*, Math. Comp. **69** (2000), 851–859.
- [16] ———, *A parametric family of quintic Thue equations II*, Monatsh. Math. **131** (2000), 29–35.
- [17] F. Halter-Koch, G. Lettl, A. Pethő, and R. F. Tichy, *Thue equations associated with Ankeny-Brauer-Chowla number fields*, J. London Math. Soc. (2) **60** (1999), 1–20.
- [18] C. Heuberger, *On a family of quintic Thue equations*, J. Symbolic Comput. **26** (1998), 173–185.
- [19] ———, *On families of parametrized Thue equations*, J. Number Theory **76** (1999), 45–61.
- [20] ———, *On general families of parametrized Thue equations*, Algebraic Number Theory and Diophantine Analysis. Proceedings of the International Conference held in Graz, Austria, August 30 to September 5, 1998 (F. Halter-Koch and R. F. Tichy, eds.), Walter de Gruyter, 2000, pp. 215–238.
- [21] ———, *On a conjecture of E. Thomas concerning parametrized Thue equations*, Acta Arith. **98** (2001), 375–394.



- [22] ———, *On explicit bounds for the solutions of a class of parametrized Thue equations of arbitrary degree*, *Monatsh. Math.* **132** (2001), 325–339.
- [23] C. Heuberger, A. Pethő, and R. F. Tichy, *Complete solution of parametrized Thue equations*, *Acta Math. Inform. Univ. Ostraviensis* **6** (1998), 93–113.
- [24] ———, *Thomas' family of Thue equations over imaginary quadratic fields*, *J. Symbolic Comput.* **34** (2002), 437–449.
- [25] C. Heuberger and R. F. Tichy, *Effective solution of families of Thue equations containing several parameters*, *Acta Arith.* **91** (1999), 147–163.
- [26] C. Heuberger, A. Togbé, and V. Ziegler, *Automatic solution of families of Thue equations and an example of degree 8*, *J. Symbolic Comput.* **38** (2004), 1145–1163.
- [27] B. Jadrijević, *A system of pellian equations and related two-parametric family of quartic Thue equations*, to appear in *Rocky Mountain J. Math.*
- [28] S. Lang, *Elliptic curves: Diophantine analysis*, *Grundlehren der Mathematischen Wissenschaften*, vol. 23, Springer, Berlin/New York, 1978.
- [29] E. Lee, *Studies on Diophantine equations*, Ph.D. thesis, Cambridge University, 1992.
- [30] G. Lettl, *Parametrized solutions of Diophantine equations*, to appear in *Math. Slovaca*.
- [31] G. Lettl and A. Pethő, *Complete solution of a family of quartic Thue equations*, *Abh. Math. Sem. Univ. Hamburg* **65** (1995), 365–383.
- [32] G. Lettl, A. Pethő, and P. Voutier, *On the arithmetic of simplest sextic fields and related Thue equations*, *Number Theory, Diophantine, Computational and Algebraic Aspects. Proceedings of the International Conference held in Eger, Hungary, July 29 – August 2, 1996* (K. Győry, A. Pethő, and V. T. Sós, eds.), de Gruyter, Berlin, 1998.
- [33] ———, *Simple families of Thue inequalities*, *Trans. Amer. Math. Soc.* **351** (1999), 1871–1894.
- [34] C. Levesque and M. Mignotte, Preprint.
- [35] F. Lippok, *On the representation of 1 by binary cubic forms of positive discriminant*, *J. Symbolic Comput.* **15** (1993), no. 3, 297–313.
- [36] M. Mignotte, *Verification of a conjecture of E. Thomas*, *J. Number Theory* **44** (1993), 172–177.
- [37] ———, *Pethő's cubics*, *Publ. Math. Debrecen* **56** (2000), 481–505.
- [38] M. Mignotte, A. Pethő, and R. Roth, *Complete solutions of quartic Thue and index form equations*, *Math. Comp.* **65** (1996), 341–354.
- [39] M. Mignotte, A. Pethő, and F. Lemmermeyer, *On the family of Thue equations  $x^3 - (n-1)x^2y - (n+2)xy^2 - y^3 = k$* , *Acta Arith.* **76** (1996), 245–269.
- [40] M. Mignotte and N. Tzanakis, *On a family of cubics*, *J. Number Theory* **39** (1991), 41–49.
- [41] J. Mueller and W. M. Schmidt, *Trinomial Thue equations and inequalities*, *J. Reine Angew. Math.* **379** (1987), 76–99.
- [42] ———, *Thue's equation and a conjecture of Siegel*, *Acta Math.* **160** (1988), 207–247.
- [43] T. Nagell, *Solution complète de quelques équations cubiques à deux indéterminées*, *J. de Math.* (9) **4** (1925), 209–270.

- [44] ———, *Darstellung ganzer Zahlen durch binäre kubische Formen mit negativer Diskriminante*, Math. Z. **28** (1928), 10–29.
- [45] ———, *Remarques sur une classe d'équations indéterminées*, Ark. Mat. **8** (1969), 199–214 (1969).
- [46] R. Okazaki, *Geometry of a quartic Thue equation*, Preprint available at <http://www1.doshisha.ac.jp/~rokazaki/>.
- [47] ———, *Geometry of a cubic Thue equation*, Publ. Math. Debrecen **61** (2002), 267–314.
- [48] A. Pethő, *On the representation of 1 by binary cubic forms with positive discriminant*, Number Theory, Lect. Notes Math., vol. 1380, Springer, 1987, pp. 185–196.
- [49] ———, *Complete solutions to families of quartic Thue equations*, Math. Comp. **57** (1991), 777–798.
- [50] A. Pethő and R. Schulenberg, *Effektives Lösen von Thue Gleichungen*, Publ. Math. Debrecen **34** (1987), 189–196.
- [51] A. Pethő and R. F. Tichy, *On two-parametric quartic families of Diophantine problems*, J. Symbolic Comput. **26** (1998), 151–171.
- [52] M. Pohst and H. Zassenhaus, *Algorithmic algebraic number theory*, Cambridge University Press, Cambridge etc., 1989.
- [53] E. Thomas, *Complete solutions to a family of cubic Diophantine equations*, J. Number Theory **34** (1990), 235–250.
- [54] ———, *Solutions to certain families of Thue equations*, J. Number Theory **43** (1993), 319–369.
- [55] ———, *Solutions to infinite families of complex cubic Thue equations*, J. Reine Angew. Math. **441** (1993), 17–32.
- [56] ———, *Counting solutions to trinomial Thue equations: A different approach*, Trans. Amer. Math. Soc. **352** (2000), 3595–3622.
- [57] A. Thue, *Über Annäherungswerte algebraischer Zahlen*, J. Reine Angew. Math. **135** (1909), 284–305.
- [58] ———, *Berechnung aller Lösungen gewisser Gleichungen von der Form*, Vid. Skrifter I Mat.-Naturv. Klasse (1918), 1–9.
- [59] A. Togbé, *On the solutions of a family of quartic Thue equations*, Math. Comp. **69** (2000), 839–849.
- [60] ———, *A parametric family of cubic Thue equations*, J. Number Theory **107** (2004), 63–79.
- [61] N. Tzanakis and B. M. M. de Weger, *On the practical solution of the Thue equation*, J. Number Theory **31** (1989), 99–132.
- [62] I. Wakabayashi, *On a family of quartic Thue inequalities I*, J. Number Theory **66** (1997), 70–84.
- [63] ———, *On a family of quartic Thue inequalities. II*, J. Number Theory **80** (2000), 60–88.
- [64] ———, *Cubic Thue inequalities with negative discriminant*, J. Number Theory **97** (2002), 222–251.
- [65] ———, *On families of cubic Thue equations*, Analytic Number Theory (K. Matsumoto and C. Jia, eds.), Developments in Mathematics, vol. 6, Kluwer Academic Publishers, 2002, pp. 359–377.
- [66] ———, *On a family of cubic Thue equations with 5 solutions*, Acta Arith. **109** (2003), 285–298.
- [67] P. Yuan, *On algebraic approximations of certain algebraic numbers*, J. Number Theory **102** (2003), 1–10.
- [68] V. Ziegler, *On a family of cubics over imaginary quadratic fields*, to appear in Period. Math. Hungar.
- [69] ———, *On a family of relative quartic Thue inequalities*, Preprint.