

Quadrature rule for Abel's equations: uniformly approximating fractional derivatives

Hiroshi Sugiura(Nanzan University)
 Takemitsu Hasegawa(University Fukui)

Abstract

An automatic quadrature method is presented for approximating fractional derivative $D^q f(x)$ of a given function $f(x)$, which is defined by an indefinite integral involving $f(x)$. The present method interpolates $f(x)$ in terms of the Chebyshev polynomials in the range $[0, 1]$ to approximate the fractional derivative $D^q f(x)$ uniformly for $0 \leq x \leq 1$, namely the error is bounded independently of x . Some numerical examples demonstrate the performance of the present automatic method.

1 Introduction

Fractional calculus (fractional integral and derivative) [6, 14, 17] has been often used recently in modeling many physical and engineering problems, see, say [8, 13, 16] and the references therein, see also [1] for the application in economics. For an interesting history (Leibniz, 30 September 1695) and scientific applications of fractional calculus, see a review due to Cafagna [4].

Let $f(s)$ be a sufficiently well-behaved function in $[0, 1]$. The Riemann-Liouville fractional integral $I^{1-q} f(s)$, where $0 < q < 1$, is defined by

$$I^{1-q} f(s) = \frac{1}{\Gamma(1-q)} \int_0^s f(t)(s-t)^{-q} dt, \quad 0 < s \leq 1,$$

where $\Gamma(1-q)$ is the gamma function [19]. On the other hand, The fractional derivative $D^q f(s)$ in the Riemann-Liouville version and the Caputo fractional derivative $D_*^q f(s)$ [4, 18] are defined by

$$D^q f(s) = \frac{d}{ds} [I^{1-q} f(s)] = \frac{d}{ds} \left[\frac{1}{\Gamma(1-q)} \int_0^s f(t)(s-t)^{-q} dt \right], \quad (1)$$

$$D_*^q f(s) = I^{1-q} \left[\frac{d}{ds} f(s) \right] = \frac{1}{\Gamma(1-q)} \int_0^s f'(t)(s-t)^{-q} dt, \quad 0 < s \leq 1, \quad (2)$$

respectively. Riemann-Liouville fractional derivative $D^q f(s)$ differs from $D_*^q f(s)$ as follows,

$$D^q f(s) = f(0)s^{-q}/\Gamma(1-q) + D_*^q f(s). \quad (3)$$

It is well known [2, p.134], [3, p.8], [5] that $D^q f(t)/\Gamma(q)$ gives the solution $y(t)$ of the generalized Abel equation [12, p.174],

$$\int_0^s y(t)(s-t)^{q-1} dt = f(s), \quad 0 < q < 1, \quad s > 0.$$

If $q-1$ is a positive non-integer, then $D^q f(s)$ is defined by

$$D^q f(s) = \frac{d^{m+1}}{ds^{m+1}} [I^{m+1-q} f(s)] = \frac{d^m}{ds^m} [D^{q-m} f(s)],$$

where m is the positive integer such that $m < q < m+1$.

The present method approximates the fractional derivatives $D^q f(s)$ uniformly for $0 < s \leq 1$, namely the errors of the approximations are bounded independently of s .

Let $J(s; f)$ be defined by

$$J(s; f) = \Gamma(1 - q) D_*^q f(s) = \int_0^s f'(t)(s - t)^{-q} dt, \quad (4)$$

then from (1), (2) and (3) we see that $D^q f(s)$ can be written by

$$D^q f(s) = \{f(0)s^{-q} + J(s; f)\}/\Gamma(1 - q).$$

Approximating $f(t)$, $0 \leq t \leq 1$, by a sum of the shifted Chebyshev polynomials $T_k(2t - 1)$,

$$f(t) \approx p_n(t) = \sum_{k=0}^n ' a_k T_k(2t - 1), \quad 0 \leq t \leq 1, \quad (5)$$

we have an approximation $J(s; p_n)$ to $J(s; f)$ as follows,

$$J(s; f) \approx J(s; p_n) = \int_0^s p_n'(t)(s - t)^{-q} dt. \quad (6)$$

In (5) the prime denotes the summation whose first term is halved. The Chebyshev coefficients a_k in (5) can be determined so that $p_n(t)$ may interpolate $f(t)$ at abscissae $t_j = \{1 + \cos(\pi j/n)\}/2$, $j = 0, \dots, n$, [20] as follows

$$a_k = \frac{2\delta_k}{n} \sum_{j=0}^n '' f(t_j) \cos \frac{\pi j k}{n},$$

where $\delta_k = 1$, ($k = 0, \dots, n - 1$), $\delta_n = 0.5$ and the double prime denotes the summation whose first and last terms are halved, and can be efficiently evaluated by using the FFT [11]. If $f(t)$ is a smooth function, then the sum of the Chebyshev polynomials (5) converges to $f(t)$ quickly as $n \rightarrow \infty$ [20].

To evaluate $J(s; p_n)$ given by (6) we use a polynomial $F_{n-1}(t)$ of degree $n - 1$ to write

$$\int_x^s \{p_n'(s) - p_n'(t)\}(s - t)^{-q} dt = \{F_{n-1}(s) - F_{n-1}(x)\}(s - x)^{1-q}. \quad (7)$$

From (7) we have

$$J(s; p_n) = \int_0^s p_n'(t)(s - t)^{-q} dt = \left[\frac{p_n'(s)}{1 - q} - F_{n-1}(s) + F_{n-1}(0) \right] s^{1-q}, \quad (8)$$

and the approximation $\tilde{D}_n^q f(s)$ to $D^q f(s)$ as follows

$$\tilde{D}_n^q f(s) = \{f(0)s^{-q} + J(s; p_n)\}/\Gamma(1 - q).$$

Functions $p_n'(t)$ and $F_{n-1}(t)$ of degree $n - 1$ are also expanded in terms of the shifted Chebyshev polynomials, see section 2.

This paper is organized as follows. In section 2 we express $F_{n-1}(t)$ in (7) by a sum of the Chebyshev polynomials and show the recurrence relation satisfied by the Chebyshev coefficients. In section 3 we estimate the error of the approximation to the fractional derivative, in particular to $J(s; f)$. In section 4 numerical examples are shown to demonstrate the performance of the present automatic quadrature method.

2 Evaluation of $F_{n-1}(t)$

Differentiating both sides of (7) with respect to x yields

$$\begin{aligned} & \{p'_n(s) - p'_n(x)\}(s-x)^{-q} \\ &= F'_{n-1}(x)(s-x)^{1-q} + \{F_{n-1}(s) - F_{n-1}(x)\}(1-q)(s-x)^{-q}, \end{aligned}$$

namely we have

$$p'_n(s) - p'_n(x) = F'_{n-1}(x)(s-x) + \{F_{n-1}(s) - F_{n-1}(x)\}(1-q). \quad (9)$$

To evaluate $F_{n-1}(s)$ in (8) we expand $F'_{n-1}(x)$ in terms of the shifted Chebyshev polynomials

$$F'_{n-1}(x) = \sum_{k=0}^{n-2} b_k T_k(2x-1), \quad 0 \leq x \leq 1, \quad (10)$$

where we have omitted the dependency of b_k on s . Integrating both sides of (10) gives

$$F_{n-1}(x) - F_{n-1}(s) = \sum_{k=1}^{n-1} \frac{b_{k-1} - b_{k+1}}{4k} \{T_k(2x-1) - T_k(2s-1)\}, \quad (11)$$

where we define $b_{n-1} = b_n = 0$. On the other hand, by using the relation $T_{k+1}(u) + T_{k-1}(u) = 2uT_k(u)$, $-1 \leq u \leq 1$, we have

$$\begin{aligned} (x-s)F'_{n-1}(x) &= F'_{n-1}(x)\{(2x-1) - (2s-1)\}/2 \\ &= \frac{1}{4} \sum_{k=0}^{n-1} \{b_{k+1} - 2(2s-1)b_k + b_{k-1}\}T_k(2x-1), \end{aligned} \quad (12)$$

where we set $b_{-1} = b_1$. Further by inserting $F_{n-1}(x) - F_{n-1}(s)$ and $(x-s)F'_{n-1}(x)$ given by (11) and (12), respectively and $p'_n(x)$ written by

$$p'_n(x) = \sum_{k=0}^{n-1} c_k T_k(2x-1), \quad (13)$$

into (9) we have

$$\left\{1 - \frac{1-q}{k}\right\}b_{k+1} - 2(2s-1)b_k + \left\{1 + \frac{1-q}{k}\right\}b_{k-1} = 4c_k, \quad 1 \leq k. \quad (14)$$

We can stably compute the recurrence relation (14) in the backward direction with starting values $b_n = b_{n-1} = 0$ to obtain b_k , $k = n-2, \dots, 0$. The Chebyshev coefficients c_k of $p'_n(x)$ given by (13) can be evaluated by the relation [15]

$$c_{k-1} = c_{k+1} + 4ka_k, \quad k = n, n-1, \dots, 1,$$

with starting values $c_n = c_{n+1} = 0$, where a_k are the Chebyshev coefficients of $p_n(x)$ in (5).

3 Error estimate

We estimate the error of the approximation to the fractional derivative $D^q f(s)$, particularly the error of $J(s; p_n)$. We shall use the notation that for $n \gg 1$, $a(n) \sim b(n)$ and $a(n) \lesssim b(n)$ mean that $\lim_{n \rightarrow \infty} a(n)/b(n) = 1$ and $\lim_{n \rightarrow \infty} a(n)/b(n) \leq 1$, respectively. Let $\omega_{n+1}(t)$ be defined by

$$\omega_{n+1}(t) = T_{n+1}(2t-1) - T_{n-1}(2t-1), \quad (15)$$

then $p_n(t)$ agrees with $f(t)$ at the zeros of $\omega_{n+1}(t)$, namely $\{1 + \cos(\pi j/n)\}/2$, $0 \leq j \leq n$. Let \mathcal{E}_ρ denote the ellipse in the complex plane $z = x + iy$,

$$\mathcal{E}_\rho : z = (w + w^{-1} + 2)/4, \quad w = \rho e^{i\theta}, \quad 0 \leq \theta \leq 2\pi, \quad (16)$$

with foci at $z = 0, 1$ and the sum of its major and minor axes equal to $\rho (> 1)$. We have the following theorem.

Theorem 3.1 *Suppose that $f(z)$ is single-valued and analytic inside and on \mathcal{E}_ρ defined by (16) and let $K = \max_{z \in \mathcal{E}_\rho} |f(z)|$. Then the approximation $J(s; p_n)$ given by (6) uniformly converges to $J(s; f)$ given by (4) as $n \rightarrow \infty$ as follows,*

$$|J(s; f) - J(s; p_n)| \leq \frac{16K \{n(\rho - 1)^2 + \rho\} \rho}{(1 - q)(\rho - 1)^4 (\rho^n - \rho^{-n})} = O(n\rho^{-n}), \quad \rho > 1. \quad (17)$$

Since our goal is to construct an automatic quadrature method, we wish to estimate the error of the approximation $J(s; p_n)$ (8) in terms of the available coefficients a_k of $p_n(t)$. Suppose that $f(z)$ is a meromorphic function which has only simple pole at the point $z = \alpha \equiv (\beta + \beta^{-1} + 2)/4$ in an ellipse \mathcal{E}_σ , $1 < \rho < \sigma$, where $1 < \rho < |\beta| < \sigma$. We have an estimated error $E_n(f)$ for the approximation $J(s; p_n)$

$$|J(s; f) - J(s; p_n)| \leq \frac{8nL_n + 2L'_n}{1 - q} \sim \frac{8nL_n}{1 - q} \lesssim \frac{8rn|a_n|}{(1 - q)(r - 1)^2} \equiv E_n(f), \quad (18)$$

where $r = |\beta|$.

Remark. The constant r may be estimated from the asymptotic behavior of $\{a_k\}$ [10].

Incidentally, an automatic quadrature of nonadaptive type is constructed from the sequence of the approximations $\{J(s; p_n)\}$ converging the integral $J(s; f)$, until a stopping criterion is satisfied. It is an usual and simple way to double the degree n of $p_n(t)$ (5) for generating the sequence $\{J(s; p_n)\}$ (8), see [9]. In order to make an automatic quadrature efficient, however, it is advantageous to have more chance of checking the stopping criterion than doubling n . To this end, as is shown in [11] we may generate the sequence of $\{p_n\}$, increasing the degree n more slowly as follows:

$$n = 6, 8, 10, \dots, 3 \times 2^i, 4 \times 2^i, 5 \times 2^i, \dots, \quad (i = 1, 2, 3, \dots)$$

and by using the FFT.

Stopping rule. We compute the sequence of $\{p_n(t)\}$ until $E_n(f)/\Gamma(1 - q)$, where $E_n(f)$ is given by (18), is less than or equal to the required tolerance ε for $D^q f(t)$.

4 Numerical examples

Table 1: Approximations of $D^{1/2}(s + 0.1)^{-0.5}$ with the required tolerance $\varepsilon = 10^{-6}$ for $0 \leq s \leq 1$. The number $n + 1$ of function evaluations required to satisfy ε is 41.

s	approximation	error
0.09	3.1300423140 508	0.35E-10
0.29	0.8494962341 923	0.63E-10
0.49	0.43199131159 67	0.48E-11
0.69	0.2718775997 025	0.41E-10
0.89	0.19102704860 19	0.84E-11

Examples in this section were computed in double precision: the machine precision is $2.22 \dots \times 10^{-16}$.

Table 2: Approximations for $D^q(s+a)^{q-1}$, $0 \leq s \leq 1$. The numbers $n+1$ of function evaluations required to satisfy the tolerances $\varepsilon = 10^{-5}$ and 10^{-9} are listed in the third and fifth columns, respectively. The actual maximum errors E_n in magnitude of approximations for $0 \leq s \leq 1$ are listed in the fourth and sixth columns.

q	a	$\varepsilon = 10^{-5}$		$\varepsilon = 10^{-9}$	
		$n+1$	E_n	$n+1$	E_n
0.1	0.01	129	6.8E-10	161	3.4E-12
	0.1	33	1.9E-8	49	2.6E-12
	1.0	13	8.8E-10	17	2.9E-13
0.5	0.01	97	6.4E-7	161	2.3E-12
	0.1	33	1.3E-8	49	1.7E-12
	1.0	13	7.2E-10	17	2.7E-13
0.9	0.01	81	2.7E-6	129	1.0E-10
	0.1	33	3.6E-9	49	3.6E-13
	1.0	13	3.3E-10	17	9.9E-14

Table 3: Approximations for $D^q e^{a(s-1)}$ with the required tolerances $\varepsilon = 10^{-5}$ and 10^{-9}

q	a	$\varepsilon = 10^{-5}$		$\varepsilon = 10^{-9}$	
		$n+1$	E_n	$n+1$	E_n
0.1	1	9	3.4E-11	13	7.8E-16
	6	17	8.9E-13	21	2.4E-15
	11	17	3.0E-9	25	5.9E-15
0.5	1	9	1.0E-10	13	1.3E-15
	6	17	3.6E-12	21	7.5E-15
	11	21	1.1E-11	25	1.2E-14
0.9	1	11	4.9E-13	13	1.1E-14
	6	17	2.5E-11	21	7.2E-14
	11	21	5.4E-11	25	9.0E-14

We compute $D^q f(s)$ for four types of $f(s)$, where (A) $f(s) = (s+a)^{q-1}$, (B) $f(s) = e^{a(s-1)}$, (C) $f(s) = \sin as$, and (D) $f(s) = s^{a/2} J_a(2\sqrt{s})$ and (E) $D^{1/2} f(s)$ where $f(s) = 1/(s^2+a^2)$. The exact values of the fractional derivatives of these functions are given, respectively, by

$$(A) \quad D^q(s+a)^{q-1} = \frac{1}{\Gamma(1-q)} \left(\frac{a}{s}\right)^q \frac{1}{s+a},$$

$$(B) \quad D^q e^{a(s-1)} = e^{-a} s^{-q} \sum_{k=0}^{\infty} \frac{(as)^k}{\Gamma(k-q+1)},$$

$$(C) \quad D^q \sin as = a s^{1-q} \sum_{k=0}^{\infty} \frac{(-1)^k (as)^{2k}}{\Gamma(2k+2-q)},$$

$$(D) \quad D^q s^{a/2} J_a(2\sqrt{s}) = s^{(a-q)/2} J_{a-q}(2\sqrt{s}),$$

$$(E) \quad D^{1/2} \left\{ \frac{1}{s^2+a^2} \right\} = \frac{1}{\sqrt{\pi s} (s^2+a^2)} - \frac{1}{\sqrt{\pi} a} \mathfrak{F} \left\{ \frac{\sin^{-1} \sqrt{s/(ia)}}{(ia-s)^{3/2}} \right\}.$$

Table 1 shows the approximations $\tilde{D}_n^{1/2}(s+0.1)^{-0.5}$ and actual errors $|D^{1/2}(s+0.1)^{-0.5} - \tilde{D}_n^{1/2}(s+0.1)^{-0.5}|$ with the required tolerance $\varepsilon = 10^{-6}$ for $s = 0.09 + i/5$, $i = 0, 1, \dots, 4$. The number $n+1$ of

Table 4: Approximations for $D^q \sin as$ with the required tolerances $\varepsilon = 10^{-5}$ and 10^{-9}

q	a	$\varepsilon = 10^{-5}$		$\varepsilon = 10^{-9}$	
		$n+1$	E_n	$n+1$	E_n
0.1	1	9	4.8E-11	13	4.4E-16
	8	17	1.1E-9	25	5.8E-14
	15	25	5.8E-11	33	3.5E-11
0.5	1	9	1.5E-10	13	8.9E-16
	8	17	4.6E-9	25	1.1E-13
	15	25	2.8E-10	33	1.2E-10
0.9	1	11	7.0E-13	13	4.4E-15
	8	21	6.2E-12	25	2.5E-13
	15	25	1.4E-9	33	4.4E-10

Table 5: Approximations for $D^q \sqrt{s}^a J_a(2\sqrt{s})$ with the required tolerances $\varepsilon = 10^{-5}$ and 10^{-9}

q	a	$\varepsilon = 10^{-5}$		$\varepsilon = 10^{-9}$	
		$n+1$	E_n	$n+1$	E_n
0.1	1.5	97	1.6E-7	1025	2.5E-11
	2.0	9	1.0E-14	11	2.2E-16
	2.5	25	5.2E-8	129	7.2E-12
0.5	1.5	129	1.1E-6	1025	1.1E-9*
	2.0	9	3.1E-14	11	3.3E-16
	2.5	33	8.6E-8	129	1.1E-10
0.9	1.5	161	1.6E-5*	1537	2.1E-8*
	2.0	9	1.5E-13	11	1.9E-15
	2.5	33	1.2E-6	161	1.2E-9*

Asterisk means the failure to satisfy the tolerances ε .

function evaluations required is 41. Table 2 also shows the result for the problem (A) with varied values of q and a , namely the numbers $n+1$ required to satisfy the tolerances $\varepsilon = 10^{-5}$ and 10^{-9} and the actual maximum errors E_n defined by

$$E_n = \max_{1 \leq j \leq m} |D^q f(s_j) - \tilde{D}_n^q f(s_j)|, \quad s_j = j/m, \quad j = 1, 2, \dots, m,$$

where we choose large m , say, $m = 1000$. Tables 3~6 show the results for the problems (B)~(E), respectively.

From Tables 2~6 we see that the present automatic method could approximate successfully the fractional derivatives (A)~(D) with varied values of q and a and (E) with varied values of a for $q = 1/2$ except for (D) with $a = 1.5$ and $q = 0.5, 0.9$ for $\varepsilon = 10^{-9}$, with $a = 1.5$ and $q = 0.9$ for $\varepsilon = 10^{-5}$ and with $a = 2.5$ and $q = 0.9$ for $\varepsilon = 10^{-9}$. The present method is not suitable for the functions with a singularity in $[0, 1]$ or singularities of higher-order derivatives. Indeed, as seen from Table 5, the present method sometimes fails to approximate the fractional derivatives of Bessel functions $s^{a/2} J_a(2\sqrt{s})$ of $a = 1.5$ and 2.5 since Bessel functions $J_a(s)$ of fractional orders a have discontinuous derivatives of higher order at $s = 0$.

Table 6: Approximations for $D^{1/2}\{1/(s^2 + a^2)\}$ with the required tolerances $\varepsilon = 10^{-5}$ and 10^{-9}

a	$\varepsilon = 10^{-5}$		$\varepsilon = 10^{-9}$	
	$n + 1$	E_n	$n + 1$	E_n
1	17	1.0E-10	21	1.8E-13
1/4	33	1.2E-8	49	1.3E-13
1/16	81	5.2E-9	97	1.8E-11

References

- [1] R. T. Baillie, Long memory processes and fractional integration in econometrics, *J. Econometrics*, 73, (1996) 5–59.
- [2] C. T. H. Baker, Numerical treatment of integral equations, Part A. Abel and Volterra equations, in *Numerical Algorithms* edited by J. L. Mohamed and J. E. Walsh, Clarendon Press, 1986.
- [3] H. Brunner, P. J. van der Houwen, *The Numerical Solution of Volterra Equations*, North-Holland, 1986.
- [4] D. Cafagna, Fractional calculus: A mathematical tool from the past for present engineers, *IEEE Industrial Electronics Magazine*, summer (2007) 35–40.
- [5] R. F. Cameron, S. McKee, The analysis of product integration methods for Abel's equation using fractional differentiation, *IMA J. Numer. Anal.*, 5, (1985) 339–353.
- [6] K. Diethelm, An algorithm for numerical solution of differential equations of fractional order, *Elec. Trans. Numer. Anal.*, 5, (1997) 1–6.
- [7] D. Elliott, Truncation errors in two Chebyshev series approximations, *Math. Comput.*, 19, (1965) 234–248.
- [8] N. Engheta, On the role of fractional calculus in electromagnetic theory, *IEEE Antennas and Propagation Magazine*, 39, (1997) 35–46.
- [9] W. M. Gentleman, Implementing Clenshaw-Curtis quadrature II. Computing the cosine transformation, *Comm. ACM*, 15, (1972) 343–346.
- [10] T. Hasegawa, T. Torii, I. Ninomiya, Generalized Chebyshev interpolation and its application to automatic quadrature, *Math. Comput.*, 41, (1983) 537–553.
- [11] T. Hasegawa, T. Torii, H. Sugiura, An algorithm based on the FFT for a generalized Chebyshev interpolation, *Math. Comput.*, 54, (1990) 195–210.
- [12] P. Henrici, *Applied and Computational Complex Analysis, Volume 3*, John Wiley & Sons, 1986.
- [13] F. Mainardi, Fractional relaxation–oscillation and fractional diffusion–wave phenomena, *Chaos, Solitons & Fractals*, 7, (1996) 1461–1477.
- [14] F. Mainardi, G. Pagnini, R. Gorenflo, Some aspects of fractional diffusion equations of single and distributed order, *Appl. Math. Comput.*, 187, (2007) 295–305.
- [15] J. C. Mason, D. C. Handscomb, *Chebyshev Polynomials*, Chapman & Hall, 2003.
- [16] R. Metzler, E. Barkai, J. Klafter, Anomalous diffusion and relaxation close to thermal equilibrium: A fractional Fokker–Planck equation approach, *Phys. Rev. Lett.*, 82, (1999) 3563–3567.

- [17] S. P. Mirevski, L. Boyadjiev, R. Scherer, On the Riemann–Liouville fractional calculus, g -Jacobi functions and F -Gauss functions, *Appl. Math. Comput.*, 187, (2007) 315–325.
- [18] Z. Odibat, Approximations of fractional integrals and Caputo fractional derivatives, *Appl. Math. Comput.*, 178, (2006) 527–533.
- [19] N. M. Temme, *Special Functions; An Introduction to the Classical Functions of Mathematical Physics*, John Wiley & Sons, 1996.
- [20] L. N. Trefethen, Is Gauss quadrature better than Clenshaw–Curtis?, *SIAM Review*, to appear.
- [21] J. Waldvogel, Fast construction of the Fejér and Clenshaw–Curtis quadrature rules, *BIT*, 46, (2006) 195–202.