

Singular diffusion equations – fundamentals, applications and open problems

東京大学・大学院数理科学研究科 儀我 美一 (Yoshikazu Giga)
Graduate School of Mathematical Sciences,
University of Tokyo

1 Introduction

This is an informal note on singular diffusion equations whose diffusion effect is so strong that the speed of evolution becomes a nonlocal quantity. Typical examples include the total variation flow as well as crystalline flow. Our goal in this paper shows current status of fields including open problems. There is already a review article [G1] on a similar topic so we mainly focus development after 2004.

2 Well-posedness

We begin with two examples. The first example is a total variation flow equation for $u = u(t, x)$, $x = (x_1, \dots, x_n)$ of the form

$$(1) \quad u_t = \operatorname{div} (\nabla u / |\nabla u|) + \lambda(f - u),$$

where f is a given function and $\lambda \in \mathbf{R}$. The second example is an anisotropic mean curvature flow equation for the graph of a function u of the form

$$(2) \quad u_t = \sqrt{1 + |\nabla u|^2} M(\vec{n})(\operatorname{div} ((DW) (\nabla u)) + \sigma(x, u)),$$

where W is a given convex function in \mathbf{R}^n which may not be C^1 and $M(> 0)$ is a given positive function; σ is also a given function; \vec{n} denotes the upward normal of $y = u(t, x)$, i.e., $\vec{n} = (-\nabla u / \sqrt{1 + |\nabla u|^2}, 1 / \sqrt{1 + |\nabla u|^2})$. Of course, (1) is a particular example of (2) by taking $M(\vec{n}) = 1 / \sqrt{1 + |\nabla u|^2}$, $W(p) = |p|$ and $\sigma(x, u) = \lambda(f(x) - u)$.

The total variation flow can be formally viewed as an L^2 -gradient flow of

$$E(u) = \int_{\Omega} (|\nabla u| + \frac{\lambda}{2}(u - f)^2) dx, \quad (\Omega : n - \text{dimensional manifold})$$

i.e., $u_t \in -\partial E(u)$, where ∂E denotes the L^2 -subdifferential of E . There is a nice abstract theory for the existence of solutions for the initial value

problem $\partial_t u \in -\partial E(u)$ when E is a lower semicontinuous function defined in a Hilbert space initiated by Y. Komura [Ko] and developed by H. Brezis and others in 1960s; see e.g. [B]. For our problem (1) it guarantees the unique solvability of the initial value problem for (1) with $\lambda \geq 0$ under, for example, periodic boundary condition (by taking $\Omega = \mathbf{T}^n$, a flat torus) with $f \in L^2(\Omega)$. Surprisingly, the speed (the right time derivative) is determined uniquely although the evolution law $u_t \in -\partial E(u)$ looks ambiguous.

Let us give a simple example to see that the speed is a nonlocal quantity. We take $\lambda = 0$ in (1) and consider one-dimensional problem to get

$$u_t = (\operatorname{sgn} u_x)_x.$$

If one considers $u_0(x) = \cos x$ as an initial data, it is expected that a flat part instantaneously develops from maximum points and minimum points. The other part is not expected to move. Heuristically, the place where slope equals zero feels very strong diffusion while other part stops. In fact, the solution is of the form

$$u(x, t) = \begin{cases} \cos x, & d(t) \leq |x| \leq \pi/2, \\ \cos d(t), & |x| \leq d(t) \end{cases}$$

and $d(t)$ is determined by

$$u_t(x, t) = 2/(\text{length of flat part}) (= 2/2d(t)), |x| < d(t)$$

with $d(0) = 0$. This fact is observed for example by T. Fukui and Y. Giga [FG] and R. Hardt and X. Zhou [HZ] a long time ago. It is clear that the speed is determined by a nonlocal quantity.

If we consider (2), it cannot be viewed as a gradient flow so general theory for $u_t \in -\partial E(u)$ does not apply. The notion of a solution itself should be studied. A viscosity like theory [CIL] developed by M.-H. Giga and Y. Giga [GG98Ar], [GG99], [GG01Ar] during 1998-2001 covers the case that the space dimension is one and σ equals a constant with some technical assumptions on W (which applies a piecewise linear convex function W). The underlining structure of their theory is that flat position (with slope corresponding to a jump of W) stays as flat so that the speed is constant. This hypotheses (which is verified by the subdifferential theory if it applies) is also fundamental to construct admissible crystalline flows by J. Taylor [T1], and independently by S. B. Angenent and M. E. Gurtin [AG]. Actually, the speed is determined

by solving an obstacle problem [KS] which may not be a constant when σ is not spatially constant (as observed in the work of [GG98DS]) or the space dimension $n \geq 3$ (as observed by G. Bellettini M. Novaga and M. Paolini [BNP]).

In fact, G. Bellettini and M. Novaga [BN] consider a surface evolution by singular energy of the form

$$V = \gamma \operatorname{div}_{\Gamma}(\nabla_p \gamma)(\vec{n}),$$

where γ is an interfacial energy and V is the normal velocity. Here γ is extended in \mathbf{R}^n as a positively 1-homogeneous function and $\nabla_p \gamma$ is its gradient. the quantity $\nabla_p \gamma(\vec{n})$ is often called the Cahn-Hoffman vector field. G. Bellettini and M. Novaga introduced a notion of solution and established a uniqueness result [BN]. However, the existence of solution is known only for convex initial data [BCCN]. Flat portion such that the speed is constant is called a calibrable set for E or a Cheeger set studied by people including B. Kawohl [KSch], [KL]. Its relation of the Rayleigh quotient of 1-Laplacian i.e., $\int |\nabla u| dx / \int |u| dx$ is well-studied [KSch], [KL]. It is interesting to characterize calibrable sets in a general setting [BNP1](cf. [BNP2]).

The case when σ is not constant is handled recently by M.-H. Giga and Y. Giga [GG09], where they study dependence of solutions of the obstacle problems with respect to the domain (flat portion) we consider. However, the theory is still for one-dimensional problem. However, this is a fundamental key to develop a level set theory [CGG], [ES], [G2] for all planer singular curvature flow including crystalline flow with nonuniform driving force term. There is another approach to construct an explicit solution by solving a free boundary problem ([GGR], [GR4], [GR5]). We also note that the speed agrees with the speed proposed by A. R. Roosen [R].

Well-posedness problem is widely open for 1-harmonic map flow:

$$u_t = \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right) + |\nabla u|u,$$

where u is a mapping to a unit sphere in \mathbf{R}^n . A global existence of weak solution is recently established by J. W. Barrett, X. Feng and A. Prohl [BFP]. However, its uniqueness is not known even locally (except one dimensional case see Y. Giga and R. Kobayashi [GK]). For surface diffusion like type problems unless problem is formulated like $u_t \in -\partial E(u)$ where subdifferential is taken in H^{-1} norm, the well-posedness is widely open (cf. Y. Kashima [Ka]).

3 Applications

Although there are several applications we just give here a rather unusual application. It is summarized as a application of vertical singular diffusion to have solutions with shocks.

We consider the graph Γ_t of solution u of the Burgers equation

$$(3) \quad u_t + uu_x = 0.$$

If we use upward normal velocity V of Γ_t in $\mathbf{R}^2 = \{(x, y) | x, y \in \mathbf{R}\}$ it is formally of the form.

$$(4) \quad V = y \text{ on } \Gamma_t$$

However, solution of (4) may overturn and may not be the graph of a function. For (3) we often consider an entropy solution with jump discontinuities instead of a multivalued (overturned) solution. What is a good equation instead of (4) to track an entropy solution? Singular diffusion equation gives an answer as rigorously discussed by M.-H. Giga-Y. Giga [GG03]. The main idea is to consider.

$$(5) \quad V = y + A \operatorname{div} (\nabla \gamma(\vec{n})),$$

where $\gamma(p_1, p_2) = |p_2|$ and $A > 0$ instead of (4). It is useful to calculate numerically (cf. T.-H. R. Tsai, Y. Giga and S. Osher [TGO]). The last term of (5) is called a vertical singular diffusion term. If $A > 0$ is taken sufficiently large with respect to size of jumps (but still finite), then the solution of (5) is the graph of entropy solutions. It is curious whether such singular diffusion can be used for sand pile problems studied by L. C. Evans, M. Feldman and R. F. Gariepy [EFG]. If the slope bound is 1 i.e., $|\nabla u| \leq 1$, a suggestive equation is

$$u_t = f(x) + A \int_{|\omega|=1} (\operatorname{sgn}(1 - \nabla u \cdot \omega))_x d\omega, \quad A > 0,$$

when the original equation is $u_t = f(x)$.

4 Approximation

It is important to develop several ways of approximation of solutions not only for construction of a solution but also to study properties of a solution. In fact, Bellettini and et al [BCCN] constructed crystalline surface evolution flow by using Almgren-Taylor-Wang [ATW] scheme for convex initial data.

General tendency is that various approximation is uniform with respect to mollification way of singular interfacial energy. This is actually proved for the anisotropic Allen-Cahn equation by Y. Giga, T. Ohtsuka and R. Schätzle [GOS] when the driving force term is spatially homogeneous but for arbitrary dimension. It is interesting to study recent differential game approximation by R. V. Kohn and S. Serfaty [KS1], [KS2] of a solution although approximation scheme for singular interfacial energy is not yet given. For example it is interesting to give a differential game interpretation for crystalline curvature flow.

5 Anisotropic version of constant mean curvature surface

It is well-known that a constant mean curvature embedded hypersurface in \mathbf{R}^n must be a sphere and is known as Alexandrov's theorem [A]. This result is recently extended for anisotropic case for smooth strictly convex interfacial energy by Y. He, H. Li, H. Ma, and J. Ge [HLMG] and by J. Zhai [Z]. The problem is that if the anisotropic curvature $\operatorname{div}_\Gamma \nabla_p \gamma(\vec{n})$ is constant everywhere on an embedded hypersurface, it is the boundary a Wulff shape

$$W_\gamma = \{x \in \mathbf{R}^n \mid x \cdot m \leq \gamma(m) \text{ for all } m \text{ such that } |m| = 1\}$$

up to dilation and translation. This problem (which is solved by [HLMG] for smooth strictly convex energy) was posed in the preprint version of [G2] and later by R. Morgan [M]. For singular interfacial energy it is known only for $n = 2$, the case of curves by P. Mucha and P. Rybka [MR]. For $n \geq 2$ it is widely open. For development of this subject for strictly convex smooth energy the reader is referred to [HLMG], [KP1], [KP2] and references cited there.

Even if the surface in \mathbf{R}^3 is immersed, if it is diffeomorphic to the sphere S^2 , the constant anisotropic curvature surface is the boundary of the Wulff shape up to dilation and translation. This is first proved by Y. Giga and J. Zhai [GZ] where interfacial energy is close to isotropic. A general result is proved by Y. He and H. Li [HL] recently for strictly convex smooth interfacial energy. However, it is an open problem for singular energy.

References

- [A] A. D. Alexandrov, Uniqueness theorems for surfaces in the large I, Vestnik Leningrad Univ., **11** (1956), 5-17.
- [ATW] F. Almgren, J. R. Taylor and L. Wang, Curvature-driven flows: a variational approach, SIAM J. Control Optim., **31** (1993), 387-438.
- [AG] S.B. Angenent and M.E. Gurtin, Multiphase thermomechanics with interfacial structure 2 Evolution of an isothermal interface, Arch. Rational Mech. Anal., **108** (1989), 323-391.
- [BFP] J. W. Barrett, X. Feng and A. Prohl, On p -harmonic map heat flows for $1 \leq p < \infty$ and their finite element approximations, SIAM J. Math. Anal., **40** (2008), 1471-1498.
- [BCCN] G. Bellettini, V. Caselles, A. Chambolle and M. Novaga, Crystalline mean curvature flow of convex sets, Arch. Ration. Mech. Anal., **179** (2006), 109-152.
- [BGN] G. Bellettini, R. Goglionone and M. Novaga, Approximation to driven motion by crystalline curvature in two dimensions, Adv. Math. Sci. Appl., **10** (2000), 467-493.
- [BN] G. Bellettini and M. Novaga, Approximation and comparison for non-smooth anisotropic motion by mean curvature in \mathbf{R}^N , Math. Models Methods Appl. Sci., **10** (2000), 1-10.
- [BNP] G. Bellettini, M. Novaga and M. Paolini, Facet-breaking for three-dimensional crystals evolving by mean curvature, Interfaces Free Bound., **1** (1999), 39-55.
- [BNP1] G. Bellettini, M. Novaga and M. Paolini, Characterization of facet breaking for nonsmooth mean curvature flow in the convex case, Interfaces and Free Boundaries, **3** (2001), 415-446.
- [BNP2] G. Bellettini, M. Novaga and M. Paolini, On a crystalline variational problem, part II: BV regularity and structure of minimizers on facets, Arch. Rational Mech. Anal., **157** (2001), 193-217.
- [B] H. Brezis, Operatures Maximaux Monotone, North-Holland, Amsterdam (1973).

- [CGG] Y.-G. Chen, Y. Giga and S. Goto, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations, *J. Differential Geom.*, **33** (1991), 749-786.
- [CIL] M. Crandall, H. Ishii and P.-L. Lions, User's guide to viscosity solutions of second order partial differential equations, *Bull. Amer. Math. Soc.*, **27** (1992), 1-67.
- [EFG] L.C. Evans, M. Feldman and R. F. Gariepy, Fast/slow diffusion and collapsing sandpiles, *J. Differential Equations*, **137** (1997), 166-209.
- [ES] L.C. Evans and J. Spruck, Motion of level sets by mean curvature, I, *J. Differential Geom.*, **33** (1991), 635-681.
- [FG] T. Fukui and Y. Giga, Motion of a graph by nonsmooth weighted curvature, In: *World Congress of Nonlinear Analysts '92* (ed. V. Lakshmikantham), Walter de Gruyter, Berlin I (1996), 47-56.
- [GG98Ar] M.-H. Giga and Y. Giga, Evolving graphs by singular weighted curvature, *Arch. Rational Mech. Anal.*, **141** (1998), 117-198.
- [GG98DS] M.-H. Giga and Y. Giga, A subdifferential interpretation of crystalline motion under nonuniform driving force, In: *Proc. of the International Conference in Dynamical Systems and Differential Equations*, Springfield Missouri, (1996), *Dynamical Systems and Differential Equations* (eds. W.-X. Chen and S.-C. Hu), Southwest Missouri Univ. **1** (1998), pp.276-287.
- [GG99] M.-H. Giga and Y. Giga, Stability for evolving graphs by nonlocal weighted curvature, *Commun in Partial Differential Equations*, **24** (1999), 109-184.
- [GG01Ar] M.-H. Giga and Y. Giga, Generalized motion by nonlocal curvature in the plane, *Arch. Rational Mech. Anal.*, **159** (2001), 295-333.
- [GG03] M.-H. Giga and Y. Giga, Minimal vertical singular diffusion preventing overturning for the Burgers equation, *Recent Advances in Scientific Computing and Partial Differential Equations*, *Contemp. Math.* **330** (2003), Amer. Math. Soc. Providence, pp.73-88.

- [GG09] M.-H. Giga and Y. Giga, A comparison principle for singular diffusion equations with spatially inhomogeneous driving force
- [G1] Y. Giga, Singular diffusivity – Facets, Shocks and more, Applied Math Entering the 21st Century, eds J. M. Hill and R. Moore, ICIAM 2003 Sydney (2004), pp.121-138, SIAM, Philadelphia.
- [G2] Y. Giga, Surface Evolution Equations – a level set approach, Birkhäuser (2006).
<http://eprints3.math.sci.hokudai.ac.jp/75/>, (2002), Preprint.
- [GGR] Y. Giga, P. Górka and P. Rybka, Nonlocal spatially inhomogeneous Hamilton-Jacobi equation with unusual free boundary, HUPSM # 933(2009).
- [GK] Y. Giga and R. Kobayashi, On constrained equations with singular diffusivity, Methods Appl. Anal., **10** (2003), 253-277.
- [GOS] Y. Giga, T. Ohtsuka and R. Schätzle, On a uniform approximation of motion by anisotropic curvature by the Allen-Cahn equations, Interfaces Free Bound., **8** (2006), 317-348.
- [GR4] Y. Giga and P. Rybka, Facet bending in the driven crystalline curvature flow in the plane, J. Geom. Anal., **18** (2008), 109-147.
- [GR5] Y. Giga and P. Rybka, Facet bending driven by the planar crystalline curvature with a generic nonuniform forcing term, J. Differential Equations, **246** (2009), 2264-2303.
- [GZ] Y. Giga and J. Zhai, Uniqueness of constant weakly anisotropic mean curvature immersion of sphere S^2 in \mathbf{R}^3 , Adv. Diff. Eq., **14** (2009), 601-619.
- [HZ] R. Hardt and X. Zhou, An evolution problem for linear growth functionals, Comm. Partial Differential Equations, **19** (1994), 1879-1907.
- [HL] Y. He and H. Li, Anisotropic version of a theorem of H. Hopf, Ann. Global Anal. Geom., **35** (2009), 243-247.
- [HLMG] Y. He, H. Li, H. Ma and J. Ge, Compact embedded hypersurfaces with constant higher order anisotropic mean curvatures, Indiana Univ. Math. J., **58** (2009), 853-868.

- [Ka] Y. Kashima, A subdifferential formulation of fourth order singular diffusion equations, *Adv. Math. Sci. Appl.*, **14** (2004), 49-74.
- [KL] B. Kawohl and Th. Lachand-Robert, Characterization of Cheeger sets for convex subsets of the plane, *Pacific J. Math.*, **225** (2006), 103-118.
- [KSch] B. Kawohl and F. Schuricht, Dirichlet problems for the 1-Laplace operator, including the eigenvalue problem, *Commun. Contemp. Math.*, **9** (2007), 515-543.
- [KS] D. Kinderlehrer and G. Stampacchia, *An Introduction to Variational Inequalities and their Applications*, Academic Press, New York (1980)
- [KS1] R. V. Kohn and S. Serfaty, A deterministic-control-based approach to motion by curvature, *Comm. Pure Appl. Math.*, **59** (2006), 344-407.
- [KS2] R. V. Kohn and S. Serfaty, A deterministic-control-based approach to fully nonlinear parabolic and elliptic equations, preprint.
- [KP1] M. Koiso and B. Palmer, Geometry and stability of surfaces with constant anisotropic mean curvature, *Indiana Univ. Math. J.*, **54** (2005), 1817-1852.
- [KP2] M. Koiso and B. Palmer, Stability of anisotropic capillary surfaces between two parallel planes, *Calc. Var. Partial Differential Equations*, **25** (2006), 275-298.
- [Ko] Y. Komura, Nonlinear semi-groups in Hilbert space, *J. Math. Soc. Japan*, **19** (1967) 493-507.
- [M] F. Morgan, Planar Wulff shape is unique equilibrium, *Proc. Amer. Math. Soc.*, **133** (2005), 809-813.
- [MR] P. B. Mucha and P. Rybka, A new look at equilibria in Stefan-type problems in the plane, *SIAM J. Math. Anal.*, **39** (2007), 1120-1134.
- [R] A. R. Roosen, Crystalline curvature and flat flow in a diffusion field, unpublished note (1994).
- [T1] J. Taylor, Constructions and conjectures in crystalline nondifferential geometry, In: *Differential geometry*, (eds. B. Lawson and K. Tenenblat), *Proceedings of the Conference on Differential Geometry*, Rio de

Janeiro, Pitman Monographs in Pure and Applied Math., **52** (1991), pp. 321-336, Pitman, London.

[TGO] Y.-H. R. Tsai, Y. Giga and S. Osher, A level set approach for computing discontinuous solutions of Hamilton-Jacobi equations, *Math. Comp.*, **72** (2003), 159-181.

[Z] J. Zhai, On the uniqueness conjecture of Wulff shape, preprint.