

WEAK AND STRONG CONVERGENCE THEOREMS FOR
UNIFORMLY ASYMPTOTICALLY REGULAR
NONEXPANSIVE SEMIGROUPS

山梨大学 厚芝幸子 (SACHIKO ATSUSHIBA)

1. INTRODUCTION

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$ and let C be a nonempty closed convex subset of H . Then, a mapping $T : C \rightarrow C$ is called nonexpansive if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in C$. We denote by $F(T)$ the set of fixed points of T . We know iteration procedures for finding a fixed point of a mapping T : Let x be an element of C and for each t with $0 < t < 1$, let x_t be a unique element of C satisfying $x_t = tx + (1 - t)Tx_t$. In 1967, Browder [7] proved the following strong convergence theorem.

Theorem 1.1. *Let H be a Hilbert space, let C be a nonempty bounded closed convex subset of H and let T be a nonexpansive mapping of C into itself. Let x be an element of C and for each t with $0 < t < 1$, let x_t be a unique element of C satisfying*

$$x_t = tx + (1 - t)Tx_t.$$

Then, $\{x_t\}$ converges strongly to the element of $F(T)$ nearest to x as $t \downarrow 0$.

Reich [13] and Takahashi and Ueda [24] extended Browder's result to those of a Banach space. Using the idea of Shimizu and Takahashi [14, 15] and the notion of sequence of means, Shioji and Takahashi [16] proved the strong convergence of Browder's type sequences for nonexpansive semigroups (see also [17, 18, 19]). Recently, Domingues Benavides, Acedo and Xu [9] proved Browder's type strong convergence theorems for uniformly asymptotically regular one-parameter nonexpansive semigroups. Acedo and Suzuki [1] generalized Domingues Benavides, Acedo and Xu's results concerning the conditions of the sequences in real numbers.

¹This research was supported by Grant-in-Aid for Scientific Research (C), Japan Society for the Promotion of Science.

2010 *Mathematics Subject Classification.* Primary 47H09, 47H10.

Key words and phrases. Fixed point, iteration, nonexpansive mapping, nonexpansive semigroup, strong convergence, weak convergence.

On the other hand, Xu and Ori [25] studied the following implicit iterative process for finite nonexpansive mappings T_1, T_2, \dots, T_r in a Hilbert space: $x_0 = x \in C$ and

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_n x_n \quad (1)$$

for every $n = 1, 2, \dots$, where α_n is a sequence in $(0, 1)$ and $T_n = T_{n+r}$. And they proved the weak convergence of the iterative process defined by (1) in a Hilbert space. Motivated by [25], author and Takahashi [6] introduced an implicit iterative process for a nonexpansive semigroup and then prove a weak convergence theorem for the nonexpansive semigroup by using the idea of mean (see also [2, 3, 4]).

In this paper, we study the implicit iterations (1) for one-parameter nonexpansive semigroups and prove a weak convergence theorem for a uniformly asymptotically regular one-parameter nonexpansive semigroup in a Hilbert space. We also prove a weak convergence theorem for a uniformly asymptotically regular nonexpansive semigroup (see also [22, 23]). Further, we study Browder's type iterations for nonexpansive semigroups. Then, we prove strong convergence theorems for uniformly asymptotically regular nonexpansive semigroups in Hilbert spaces by using the idea of [1, 7, 9, 22, 23]. And we give a strong convergence theorem for the nonexpansive semigroup by the viscosity approximation method.

2. PRELIMINARIES AND NOTATIONS

Throughout this paper, we denote by \mathbb{N} and \mathbb{R} the set of all positive integers and the set of all real numbers, respectively. We also denote by \mathbb{R}^+ the set of all nonnegative real numbers. Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$ and let C be a nonempty closed convex subset of H . Then, for every point $x \in H$, there exists a unique nearest point in C , denoted by $P_C x$, such that

$$\|x - P_C x\| \leq \|x - y\|$$

for all $y \in C$. P_C is called the metric projection of H onto C . It is characterized by

$$\langle P_C x - y, x - P_C x \rangle \geq 0$$

for all $y \in C$. See [23] for more details. The following result is well-known; see also [23].

Lemma 2.1. *Let C be a nonempty bounded closed convex subset of a Hilbert space H and let T be a nonexpansive mapping of C into itself. Then, $F(T) \neq \emptyset$.*

We write $x_n \rightarrow x$ (or $\lim_{n \rightarrow \infty} x_n = x$) to indicate that the sequence $\{x_n\}$ of vectors in H converges strongly to x . We also write $x_n \rightharpoonup x$ (or $w\text{-}\lim_{n \rightarrow \infty} x_n = x$) to indicate that

the sequence $\{x_n\}$ of vectors in H converges weakly to x . In a Hilbert space, it is well known that $x_n \rightharpoonup x$ and $\|x_n\| \rightarrow \|x\|$ imply $x_n \rightarrow x$.

Let S be a semitopological semigroup. A semitopological semigroup S is called right (resp. left) reversible if any two closed left (resp. right) ideals of S have nonvoid intersection. If S is right reversible, (S, \leq) is a directed system when the binary relation " \leq " on S is defined by $s \leq t$ if and only if $\{s\} \cup \overline{Ss} \supset \{t\} \cup \overline{S+t}$, $s, t \in S$, where \overline{A} is the closure of A . A commutative semigroup S is a directed system when the binary relation is defined by $s \leq t$ if and only if $\{s\} \cup (S+s) \supset \{t\} \cup (S+t)$.

Let C be a nonempty closed convex subset of a Hilbert space H . A family $\mathcal{S} = \{T(t) : t \in S\}$ of mappings of C into itself is said to be a nonexpansive semigroup on C if it satisfies the following conditions:

- (i) For each $t \in S$, $T(t)$ is nonexpansive;
- (ii) $T(ts) = T(t)T(s)$ for each $t, s \in S$.

We denote by $F(\mathcal{S})$ the set of common fixed points of \mathcal{S} , i.e., $F(\mathcal{S}) = \bigcap_{t \in S} F(T(t))$.

We say that a Banach space E satisfies *Opial's condition* [12] if for each sequence $\{x_n\}$ in E which converges weakly to x ,

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\| \quad (2)$$

for each $y \in E$ with $y \neq x$. In a reflexive Banach space, this condition is equivalent to the analogous condition for a bounded net which has been introduced in [10]. It is well known that this condition is equivalent to the analogous condition of $\overline{\lim}$ (see [5]). It is well known that Hilbert spaces satisfy Opial's condition (see [12, 23]).

Proposition 2.2 ([12]). *Let H be a Hilbert space. Let $\{x_n\}$ be a sequence in H converging weakly to $x \in H$. Then,*

$$\liminf_{n \rightarrow \infty} \|x_n - x\| < \liminf_{n \rightarrow \infty} \|x_n - y\| \quad (3)$$

for each $y \in E$ with $y \neq x$.

3. CONVERGENCE THEOREMS FOR ONE-PARAMETER NONEXPANSIVE SEMIGROUPS

In this section, we prove a weak convergence theorem for an asymptotically regular one-parameter nonexpansive semigroup by using the idea of [1, 9, 22, 23, 25]. Let C be a nonempty closed convex subset of a Hilbert space H . A family $\mathcal{S} = \{T(t) : t \in \mathbb{R}^+\}$ of mappings of C into itself satisfying the following conditions is said to be one-parameter nonexpansive semigroup on C :

- (i) for each $t \in \mathbb{R}^+$, $T(t)$ is nonexpansive;
- (ii) $T(0) = I$;

- (iii) $T(t+s) = T(t)T(s)$ for every $t, s \in \mathbb{R}^+$;
 (iv) for each $x \in C$, $t \mapsto T(t)x$ is continuous.

We say that one-parameter nonexpansive semigroup $\mathcal{S} = \{T(t) : t \in \mathbb{R}^+\}$ is asymptotically regular if

$$\lim_{s \rightarrow \infty} \|T(h+s)x - T(s)x\| = 0$$

for all $h \in \mathbb{R}^+$ and $x \in C$ (see also [22, 23]). The following lemma proved by Acedo and Suzuki ([1]).

Lemma 3.1 ([1]). *Let H be a Hilbert space and let C be a nonempty closed convex subset of H . Let $\mathcal{S} = \{T(s) : s \in \mathbb{R}^+\}$ be a one-parameter nonexpansive semigroup on C . Assume that $\mathcal{S} = \{T(s) : s \in \mathbb{R}^+\}$ is asymptotically regular, that is,*

$$\lim_{t \rightarrow \infty} \|T(h+t)x - T(t)x\| = 0$$

for all $h \in \mathbb{R}^+$ and $x \in C$. Then,

$$F(T(h)) = F(\mathcal{S})$$

for each $h \in \mathbb{R}^+$.

We say that one-parameter nonexpansive semigroup $\mathcal{S} = \{T(t) : t \in \mathbb{R}^+\}$ is uniformly asymptotically regular if for every $h \in \mathbb{R}^+$ and for every bounded subset K of C ,

$$\limsup_{s \in \mathbb{R}^+} \sup_{x \in K} \|T(h+s)x - T(s)x\| = 0.$$

holds.

We prove a weak convergence theorem for a uniformly asymptotically regular one-parameter nonexpansive semigroup (see [1, 9]).

Theorem 3.2. *Let H be a Hilbert space and let C be a nonempty closed convex subset of H . Let $\mathcal{S} = \{T(s) : s \in \mathbb{R}^+\}$ be a uniformly asymptotically regular one-parameter nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let $\{m_n\}$ be a sequence in \mathbb{N} such that $m_n \rightarrow \infty$ or $m_n \rightarrow N$ for some $N \in \mathbb{N}$. Let $\{\alpha_n\}$ be a sequence in \mathbb{R} such that $0 < \alpha_n < 1$, and $\alpha_n \rightarrow 0$. Let $u \in C$ and let $\{x_n\}$ be the sequence defined by*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n)(T(m_n))x_n$$

for each $n \in \mathbb{N}$. Then, $\{x_n\}$ converges weakly to a common fixed point of \mathcal{S} .

4. WEAK CONVERGENCE THEOREMS FOR NONEXPANSIVE SEMIGROUPS

In this section, we prove a weak convergence theorem for an asymptotically regular nonexpansive semigroup by using the idea of [1, 9, 22, 23, 25]. Let C be a nonempty closed convex subset of a Hilbert space H , let S be a commutative semigroup and let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C . We say that nonexpansive semigroup $\mathcal{S} = \{T(t) : t \in S\}$ is asymptotically regular if

$$\lim_{s \in S} \|T(h)T(s)x - T(s)x\| = 0$$

for all $h \in S$ and $x \in C$ (see also [22, 23]). The following lemma plays an important role in the proof of main theorem (see [1]).

Lemma 4.1. *Let H be a Hilbert space, let C be a nonempty closed convex subset of H , and let S be a commutative semigroup. Let $\mathcal{S} = \{T(t) : t \in S\}$ be a nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Assume that $\mathcal{S} = \{T(t) : t \in S\}$ is asymptotically regular, that is,*

$$\lim_{t \in S} \|T(h)T(t)x - T(t)x\| = 0$$

for all $h \in S$ and $x \in C$. Then,

$$F(T(h)) = F(\mathcal{S})$$

for each $h \in S$.

We say that nonexpansive semigroup $\mathcal{S} = \{T(t) : t \in S\}$ is uniformly asymptotically regular if for every $h \in S$ and for every bounded subset K of C ,

$$\limsup_{s \in S} \sup_{x \in K} \|T(h)T(s)x - T(s)x\| = 0.$$

holds.

We prove a weak convergence theorem for a uniformly asymptotically regular nonexpansive semigroup (see also [1, 9]).

Theorem 4.2. *Let H be a Hilbert space, let C be a nonempty closed convex subset of H , and let S be a commutative semigroup. Let $\mathcal{S} = \{T(t) : t \in S\}$ be a uniformly asymptotically regular nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let $\{m_n\}$ be a sequence in \mathbb{N} such that $m_n \rightarrow \infty$ or $m_n \rightarrow N$ for some $N \in \mathbb{N}$. Let $\{\alpha_n\}$ be a sequence in \mathbb{R} such that $0 < \alpha_n < 1$, and $\alpha_n \rightarrow 0$. Let $u \in C$, let $t \in S$, and let $\{x_n\}$ be the sequence defined by*

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n)(T(t))^{m_n} x_n$$

for each $n \in \mathbb{N}$. Then, $\{x_n\}$ converges weakly to a common fixed point of \mathcal{S} .

5. STRONG CONVERGENCE THEOREMS

Motivated by [1, 7, 9], we study Browder's type strong convergence theorems for uniformly asymptotically regular nonexpansive semigroups (see also [22, 23]).

Theorem 5.1. *Let H be a Hilbert space, let C be a nonempty closed convex subset of H , and let S be a commutative semigroup. Let $\mathcal{S} = \{T(t) : t \in S\}$ be a uniformly asymptotically regular nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let $\{m_n\}$ be a sequence in \mathbb{N} such that $m_n \rightarrow \infty$ or $m_n \rightarrow N$ for some $N \in \mathbb{N}$. Let $\{\alpha_n\}$ be a sequence in \mathbb{R} such that $0 < \alpha_n < 1$, and $\alpha_n \rightarrow 0$. Let $u \in C$, let $t \in S$, and let $\{x_n\}$ be the sequence defined by*

$$x_n = \alpha_n u + (1 - \alpha_n)(T(t))^{m_n} x_n$$

for each $n \in \mathbb{N}$. Then, $\{x_n\}$ converges strongly to Pu , where P is the metric projection from C onto $F(\mathcal{S})$.

We know that $f : C \rightarrow C$ is said to be a contraction on C if there exists $r \in (0, 1)$ such that

$$\|f(x) - f(y)\| \leq r\|x - y\|$$

for each $x, y \in C$. Using [21] and Theorem 5.1, we obtain the following strong convergence theorem by the viscosity approximation methods (see also [11]).

Theorem 5.2. *Let C be a nonempty closed convex subset of a Hilbert space H , let S be a commutative semigroup and let $\mathcal{S} = \{T(t) : T \in S\}$ be a uniformly asymptotically regular nonexpansive semigroup on C such that $F(\mathcal{S}) \neq \emptyset$. Let f be a contraction on C . Let $\{m_n\}$ be a sequence in \mathbb{N} such that $m_n \rightarrow \infty$ or $m_n \rightarrow N$ for some $N \in \mathbb{N}$. Let $\{\alpha_n\}$ be a sequence in \mathbb{R} such that $0 < \alpha_n < 1$, and $\alpha_n \rightarrow 0$. Let $u \in C$, let $t \in S$, and let $\{x_n\}$ be the sequence defined by*

$$x_n = \alpha_n f(x_n) + (1 - \alpha_n)(T(t))^{m_n} x_n$$

for each $n \in \mathbb{N}$. Then, $\{x_n\}$ converges strongly to Pu , where P is the metric projection from C onto $F(\mathcal{S})$.

REFERENCES

1. G. L. Acedo and T. Suzuki, *Browder's Convergence for Uniformly Asymptotically Regular Nonexpansive Semigroups in Hilbert Spaces*, Fixed Point Theory and Applications Volume 2010, Article ID 418030.
2. S. Atsushiba, *Strong convergence theorems for finite noneopansive mappings*, Comm. Appl. Non-linear Anal. **9** (2002), 57-68.

3. S. Atsushiba, *Strong convergence theorems for finite nonexpansive mappings in Banach spaces*, Proceedings of the Third International Conference on Nonlinear Analysis and Convex Analysis (W. Takahashi and T. Tanaka Eds.), pp.9-16, Yokohama Publishers, Yokohama, 2004.
4. S. Atsushiba, *Approximating zero points of accretive operators by an implicit iterative sequences*, Taiwanese J. Math., **11** (2007), 807–818.
5. S. Atsushiba and W. Takahashi, *Nonlinear Ergodic Theorems in a Banach Space Satisfying Opial's Condition*, Tokyo J. Math. **21** (1998), 61–81.
6. S. Atsushiba and W. Takahashi, *Weak and strong convergence theorem for nonexpansive semigroups in Banach spaces*, Fixed Point Theory and Applications, **2005** (2005), 343–353.
7. F. E. Browder, *Fixed-point theorems for noncompact mappings in Hilbert space*, Proc. Natio. Acad. Sci. Unit. Stat. Ameri., **53** (1965), 1272–1276.
8. F. E. Browder, *Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces*, Arch. Rational Mech. Anal. **24** (1967) 82–90.
9. T. Dominguez Benavides, G. L. Acedo, and H.-K. Xu, *Construction of sunny nonexpansive retractions in Banach spaces*, Bull. Austrl. Math. Soc., **66** (2002) 9–16.
10. A. T. Lau, *Semigroup of nonexpansive mappings on Hilbert space*, J. Math. Anal. Appl. **105** (1985), 514–522.
11. A. Moudafi, *Viscosity approximation methods for fixed-points problem*. J. Math. Anal. Appl., **241** (2000), 46–55.
12. Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73** (1967), 591–597.
13. S. Reich, *Strong convergence theorems for resolvents of accretive operators in Banach spaces*, J. Math. Anal. Appl. **75** (1980), 287–292.
14. T. Shimizu and W. Takahashi, *Strong convergence theorem for asymptotically nonexpansive mappings*, Nonlinear Anal. **26** (1996), 265–272.
15. T. Shimizu and W. Takahashi, *Strong convergence to common fixed points of families of nonexpansive mappings*, J. Math. Anal. Appl. **211** (1997), 71–83.
16. N. Shioji and W. Takahashi, *Strong convergence theorems for asymptotically nonexpansive semigroups in Hilbert spaces*, Nonlinear Anal. **34** (1998), 87–99.
17. N. Shioji and W. Takahashi, *Strong convergence of averaged approximants for asymptotically nonexpansive mappings in Banach spaces*, J. Approx. Theory **97** (1999), 53–64.
18. N. Shioji and W. Takahashi, *Strong convergence theorems for continuous semigroups in Banach spaces*, Math. Japon. **50** (1999), 57–66.
19. N. Shioji and W. Takahashi, *Strong convergence theorems for asymptotically nonexpansive semigroups in Banach spaces*, J. Nonlinear Convex Anal. **1** (2000), 73–87.
20. T. Suzuki, *Browder's type strong convergence theorems for infinite families of nonexpansive mappings in Banach spaces*, Fixed Point Theory Appl. **2006**, (2006), 1–16.
21. T. Suzuki, *Moudafi's viscosity approximations with Meir-Keeler contractions*, J. Math. Anal. Appl., **325** (2007), 342–352.
22. W. Takahashi, *The asymptotic behavior of nonlinear semigroups and invariant means*, J. Math. Anal. Appl., **109** (1985), 130–139.
23. W. Takahashi, *Nonlinear functional analysis*, Yokohama Publishers, Yokohama, 2000.

24. W. Takahashi and Y. Ueda, *On Reich's strong convergence theorems for resolvents of accretive operators*, J. Math. Anal. Appl. **104** (1984), 546–553.
25. H. K. Xu and R.G.Ori, *An implicit iteration process for nonexpansive mappings*, Numer. Funct. Anal. Optim. **22** g(2001), 767–773.

(S. Atsushiba) DEPARTMENT OF MATHEMATICS AND PHYSICS, INTERDISCIPLINARY SCIENCES COURSE, FACULTY OF EDUCATION AND HUMAN SCIENCES, UNIVERSITY OF YAMANASHI, 4-4-37, TAKEDA, KOFU, YAMANASHI 400-8510, JAPAN

E-mail address: `asachiko@yamanashi.ac.jp`