

# Discrete Geometry on 3 Colored Point sets in the Plane

Mikio Kano

Department of Computer and Information Sciences,  
Ibaraki University, Hitachi, Ibaraki, Japan  
kano@mx.ibaraki.ac.jp <http://gorogoro.cis.ibaraki.ac.jp>

## 1 3 colored point sets in the plane

Let  $R$ ,  $B$  and  $G$  denote disjoint sets of red points, blue points and green points in the plane, respectively. If no three points of  $R \cup B \cup G$  are collinear, we say that  $R$ ,  $B$  and  $G$  are *in general position* in the plane. We always assume that given sets of colored points are in general position.

We begin with the following well-known theorem on two colored point sets in the plane. Notice that a *geometric graph* is a graph drawn in the plane whose edges are straight line segments, and every edge of an *alternating matching* joins two points with distinct colors.

**Theorem 1** ([3]). *If  $|R| = |B|$ , then there exists an alternating non-crossing geometric perfect matching on  $R \cup B$  (see Figure 1).*




Figure 1: An alternating non-crossing geometric perfect matching on  $R \cup B$ .

We generalize the above theorem by considering 3 colored point sets. The standard proof of the following theorem is basically similar to that of the above Theorem 1, but more difficult.

**Corollary 2** (Kano, Suzuki, Uno [4]). *If  $|R \cup B \cup G| = 2n$ ,  $|R| \leq n$ ,  $|B| \leq n$  and  $|G| \leq n$ , then there exists an alternating non-crossing geometric perfect matching on  $R \cup B \cup G$ .*




Figure 2: An alternating non-crossing geometric perfect matching on  $R \cup B \cup G$ .

It is known as the discrete version of Ham-Sandwich theorem that if  $|R| = 2m$  and  $|B| = 2n$ , then there exists a bisector line  $l$  such that  $|left(l) \cap R| = m$  and  $|left(l) \cap B| = n$ . It is easy to see that there exist configurations of 3 colored points in the plane such that there exists no line  $l$  such that a half-plane determined by  $l$  contains the same number of each colored points. Thus the condition in the next theorem is necessary. For a set  $X$  of points in the plane, we denote the *convex hull* of  $X$  by  $conv(X)$ .

**Theorem 3** (Bereg and Kano [2]). *Assume that  $|R| = |B| = |G| = n$ , where  $n \geq 2$ . If all the vertices of  $conv(R \cup B \cup G)$  are red, then there exists a line  $l$  such that  $|right(l) \cap R| = |right(l) \cap B| = |right(l) \cap G| = k$  for some integer  $1 \leq k \leq n - 1$  (see Figure 3).*

We give one more result on three colored point sets in the plane, and explain a sketch of its proof.

**Theorem 4** (Bereg and etc. [1]). *Assume that  $n$  red points and  $n$  blue points and  $n$  green points lie on a circle in the plane. Then for every integer  $1 \leq k \leq n - 1$ , there exist two intervals  $I$  and  $J$  on the circle such that  $I \cup J$  contains exactly  $k$  red points,  $k$  blue points and  $k$  green points (see Figure 4).*

We give a sketch of its proof.

**Lemma 5.** *Let  $n \geq 2$  be an integer. Then every integer  $1 \leq k \leq n - 1$  can be obtained from  $n$  by applying the following functions  $f$  and  $g$  some times.*

$$f(x) = \lfloor x/2 \rfloor \quad \text{and} \quad g(x) = n - x$$




Figure 3: All the vertices of  $\text{conv}(R \cup B \cup G)$  are red; An line  $l$  such that  $\text{right}(l)$  contains exactly 3 red points, 3 blue points and 3 green points.




Figure 4: Two disjoint intervals  $I$  and  $J$  that contains exactly 3 red points, 3 blue points and 3 green points.

We show only one example, whose generalization gives us its proof. Suppose that  $n = 30$  and  $k = 2$ . Then  $\lfloor n/2 \rfloor = 15$ . We construct the following series of intervals as follows: if an interval  $[x, y]$  does not contain 15 and  $y < 15$ , then make an interval  $[2x, 2y + 1]$ . If  $[x, y]$  does not contain 15 and  $15 < x$ , then make an interval  $[30 - y, 30 - x]$ . If an interval  $[x, y]$  contains 15, then stop. Then we can obtain  $k = 2$  from  $\lfloor n/2 \rfloor = 15$  by applying the operations  $f(x)$  and  $g(x)$  as follows.

$$k = 2 \rightarrow [4, 5] \rightarrow [8, 11] \rightarrow [16, 23] \rightarrow [7, 14] \rightarrow [14, 29] \ni 15$$

$$2 \leftarrow 5 \leftarrow 11 \leftarrow 23 \leftarrow 7 \leftarrow 15$$

The next lemma follows immediately from Lemma 5

**Lemma 6.** *Let  $n \geq 2$  be an integer, and let  $X$  be a subset of  $\{0, 1, 2, \dots, n\}$ . Define two functions  $f$  and  $g$  as follows:*

$$f(x) = \lfloor x/2 \rfloor \quad \text{and} \quad g(x) = n - x$$

*If  $X$  has the following properties, then  $X = \{0, 1, 2, \dots, n\}$ .*

$$n \in X; \text{ and if } k \in X, \text{ then } g(k) \in X \text{ and } f(k) \in X.$$

*Sketch of the proof of Theorem 4.* Let us define

$$X = \{1 \leq x \leq n : \text{there exist two intervals } I \text{ and } J \text{ on the circle} \\ \text{such that } I \cup J \text{ contains exactly } x \text{ red points,} \\ x \text{ blue points and } x \text{ green points.}\}$$

It is easy to see that  $n \in X$ , and if  $k \in X$ , then the complement  $I \cup J$  on the circle contains exactly  $n - k$  red points,  $n - k$  blue points and  $n - k$  green points, which implies  $g(k) = n - k \in X$ . Moreover, we can show that if there exist intervals  $I$  and  $J$  on the circle such that  $I \cup J$  contains exactly  $k$  red points,  $k$  blue points and  $k$  green points, then there exist intervals  $I'$  and  $J'$  in  $I \cup J$  such that  $I' \cup J'$  contains exactly  $\lfloor k/2 \rfloor$  red points  $\lfloor k/2 \rfloor$  blue points and  $\lfloor k/2 \rfloor$  green points. Hence by Lemma 6,  $X = \{0, 1, 2, \dots, n\}$ , which implies that Theorem 4 holds.

## References

- [1] S. Bereg, F. Hurtado, M.Kano, M. Korman, D. Lara, C. Seara, R. Silveira, J. Urrutia, and K. Verbeek, Balanced partitions of 3-colored geometric sets in the plane. (in preparation).

- [2] S. Bereg and M.kano, Balanced line for a 3-colored point set in the plane, *The Electronic Journal of Combinatorics* **19** (2012) P33.
- [3] A. Kaneko and M. Kano, Discrete geometry on red and blue points in the plane - a survey, *Discrete and Computational Geometry Algorithms and Combinatorics* **25** (2003) 551-570.
- [4] M. Kano, K. Suzuki and M. Uno, Geometric graphs on three colored point sets in the plane, (in preparation)