

On some geometric constant and the extreme points of the unit ball

Hiroyasu Mizuguchi

Department of Mathematical Sciences,
Graduate School of Science and Technology, Niigata University

Abstract

Mitani and Saito introduced a geometric constant $\gamma_{X,\psi}$ by using the notion of ψ -direct sum. For $t \in [0, 1]$, the constant $\gamma_{X,\psi}$ is defined as a supremum taken over all elements of in the unit sphere. It is proved that for a Banach space X with a predual Banach space X_* , $\gamma_{X,\psi}$ can be calculated as the supremum can be taken over all extreme points of the unit ball.

1. Introduction

Let X be a Banach space with $\dim X \geq 2$. By S_X and B_X , we denote the unit sphere and the unit ball of X , respectively. The von Neumann-Jordan constant (shortly, NJ constant) $C_{NJ}(X)$ is defined as the smallest constant C for which

$$\frac{1}{C} \leq \frac{\|x + y\|^2 + \|x - y\|^2}{2(\|x\|^2 + \|y\|^2)} \leq C$$

holds for all $x, y \in X$, not both zero (Clarkson [2]). This constant has been considered in many papers. It is known that $1 \leq C_{NJ}(X) \leq 2$ for any Banach space X . From the parallelogram law it follows immediately that X is a Hilbert space if and only if $C_{NJ}(X) = 1$ ([3]). Recall that a Banach space X is uniformly non-square provided $C_{NJ}(X) < 2$ ([10]), where X is said to be uniformly non-square if there exists $\delta > 0$ such that $\|x + y\| \leq 2(1 - \delta)$ holds whenever $\|x - y\| \geq 2(1 - \delta)$, $x, y \in S_X$.

By the definition, the NJ constant is in the following form;

$$C_{NJ}(X) = \sup \left\{ \frac{\|x + y\|^2 + \|x - y\|^2}{2(\|x\|^2 + \|y\|^2)} : x, y \in X, (x, y) \neq (0, 0) \right\},$$

and it can be reformulated as

$$C_{NJ}(X) = \sup \left\{ \frac{\|x + ty\|^2 + \|x - ty\|^2}{2(1 + t^2)} : x, y \in S_X, 0 \leq t \leq 1 \right\}.$$

In 2006, the function γ_X was introduced by Yang and Wang [13];

$$\gamma_X(t) = \sup \left\{ \frac{\|x + ty\|^2 + \|x - ty\|^2}{2} : x, y \in S_X \right\}, \quad (t \in [0, 1]).$$

It is easy to see that the NJ constant $C_{NJ}(X)$ coincide with $\sup\{\gamma_X(t)/(1+t^2) : 0 \leq t \leq 1\}$. Thus, the function γ_X is useful to calculate the NJ constant $C_{NJ}(X)$ for some Banach spaces. In fact, they computed $C_{NJ}(X)$ for X being Day-James spaces $\ell_\infty\text{-}\ell_1$ and $\ell_2\text{-}\ell_1$ by using the function γ_X .

In the same paper [13], they noted that, for a finite dimensional Banach space the supremum can be taken over all extreme points of the unit ball. We obtained a generalization of this.

2. Preliminaries

Recall that a norm $\|\cdot\|$ on \mathbb{R}^2 is said to be *absolute* if $\|(x, y)\| = \||x|, |y|\|$ for all $(x, y) \in \mathbb{R}^2$, and *normalized* if $\|(1, 0)\| = \|(0, 1)\| = 1$. The ℓ_p -norms $\|\cdot\|_p$ ($1 \leq p \leq \infty$) are basic examples;

$$\|(x, y)\|_p = \begin{cases} (|x|^p + |y|^p)^{1/p} & \text{if } 1 \leq p < \infty, \\ \max\{|x|, |y|\} & \text{if } p = \infty. \end{cases}$$

The family of all absolute normalized norms on \mathbb{R}^2 is denoted by AN_2 . As in Bonsall and Duncan [1], AN_2 is in a one-to-one correspondence with the family Ψ_2 of all convex functions ψ on $[0, 1]$ with $\max\{1-t, t\} \leq \psi(t) \leq 1$ for all $0 \leq t \leq 1$. Indeed, for any $\|\cdot\| \in AN_2$ we put $\psi(t) = \|(1-t, t)\|$. Then $\psi \in \Psi_2$. Conversely, for all $\psi \in \Psi_2$ let

$$\|(x, y)\|_\psi = \begin{cases} (|x| + |y|)\psi\left(\frac{|y|}{|x| + |y|}\right) & \text{if } (x, y) \neq (0, 0), \\ 0 & \text{if } (x, y) = (0, 0). \end{cases}$$

Then $\|\cdot\|_\psi \in AN_2$, and $\psi(t) = \|(1-t, t)\|_\psi$ (cf. [8]). The functions corresponding to the ℓ_p -norms $\|\cdot\|_p$ on \mathbb{R}^2 are given by

$$\psi_p(t) = \begin{cases} \{(1-t)^p + t^p\}^{1/p} & \text{if } 1 \leq p < \infty, \\ \max\{1-t, t\} & \text{if } p = \infty. \end{cases}$$

In [11], the notion of ψ -direct sum of Banach spaces was introduced. Let X and Y be Banach spaces, and let $\psi \in \Psi_2$. The ψ -direct sum $X \oplus_\psi Y$ of X and Y is defined as the direct sum $X \oplus Y$ equipped with the norm

$$\|(x, y)\|_\psi = \|(\|x\|, \|y\|)\|_\psi \quad ((x, y) \in X \oplus Y).$$

This notion has been investigated by several authors.

In [5], Mitani and Saito introduced a geometrical constant $\gamma_{X,\psi}$ of a Banach space X , by using the notion of ψ -direct sum. For a Banach space X and $\psi \in \Psi_2$, the function $\gamma_{X,\psi}$ on $[0, 1]$ is defined by

$$\gamma_{X,\psi}(t) = \sup\{\|(x+ty, x-ty)\|_\psi : x, y \in S_X\}, \quad (t \in [0, 1]).$$

One can easily have $\gamma_X(t) = \gamma_{X,\psi_2}(t)^2/2$, where $\psi_2 \in \Psi_2$ is the function which corresponds to the ℓ_2 -norm $\|\cdot\|_2$.

Mitani and Saito showed that

Proposition 2.1. ([5])

(1) For any Banach space X , $\psi \in \Psi_2$ and $t \in [0, 1]$,

$$2\psi\left(\frac{1-t}{2}\right) \leq \gamma_{X,\psi}(t) \leq 2(1+t)\psi\left(\frac{1}{2}\right).$$

(2) For a Banach space X , $\psi \in \Psi_2$ and $t \in [0, 1]$,

$$\gamma_{X,\psi}(t) = \sup \{ \|(x + ty, x - ty)\|_\psi : x, y \in B_X \}.$$

(3) Let $\psi \in \Psi_2$ with $\psi \neq \psi_\infty$. Then a Banach space X is uniformly non-square if and only if $\gamma_{X,\psi}(t) < 2(1+t)\psi(1/2)$ for any (or some) t with $0 < t \leq 1$.

They obtained some other results on $\gamma_{X,\psi}$ (cf. [5]).

3. Results

An element $x \in S_X$ is called an extreme point of B_X if $y, z \in S_X$ and $x = (y + z)/2$ implies $x = y = z$. The set of all extreme points of B_X is denoted by $\text{ext}(B_X)$.

In [13], Yang and Wang noted that

Proposition 3.1. Let X be a finite dimensional Banach space. Then

$$\gamma_X(t) = \sup \left\{ \frac{\|x + ty\|^2 + \|x - ty\|^2}{2} : x, y \in \text{ext}(B_X) \right\}.$$

There exists some infinite-dimensional Banach spaces whose unit ball has no extreme point. However, from the Banach-Alaoglu Theorem and Krein-Milman Theorem, we have that for any Banach space, the unit ball of the dual space is the weakly* closed convex hull of its set of extreme points.

For $\psi \in \Psi_2$, the dual function ψ^* of ψ is defined by

$$\psi^*(s) = \sup_{t \in [0,1]} \frac{(1-s)(1-t) + st}{\psi(t)}$$

for $s \in [0, 1]$ ([4]). Then we have $\psi^* \in \Psi_2$ and that $\|\cdot\|_{\psi^*}$ is the dual norm of $\|\cdot\|_\psi$. It is easy to see that $\psi^{**} = \psi$.

Suppose that X is a Banach space with the predual Banach space X_* . Then the unit ball B_X is the weakly* closed convex hull of $\text{ext}(B_X)$, and the direct sum $X \oplus_\psi X$ is isomorphic to the dual of $X_* \oplus_{\psi^*} X_*$. Based on these facts, we obtain the following result.

Theorem 3.2. [6] Let X be a Banach space with the predual Banach space X_* . Then

$$\gamma_{X,\psi}(t) = \sup \{ \|(x + ty, x - ty)\|_\psi : x, y \in \text{ext}(B_X) \}$$

for any $\psi \in \Psi_2$ and any $t \in [0, 1]$.

In [9], Takahashi introduced the James and von Neumann-Jordan type constants of Banach spaces. For $p \in [-\infty, \infty)$ and $t \geq 0$, the James type constant is defined as

$$J_{X,p}(t) = \begin{cases} \sup \left\{ \left(\frac{\|x+ty\|^p + \|x-ty\|^p}{2} \right)^{1/p} : x, y \in S_X \right\} & \text{if } p \neq -\infty, \\ \sup \{ \min(\|x+ty\|, \|x-ty\|) : x, y \in S_X \} & \text{if } p = -\infty \end{cases}$$

(cf. [12, 14]). The von Neumann-Jordan type constant is defined as

$$C_p(X) = \sup \left\{ \frac{J_{X,p}(t)^2}{1+t^2} : 0 \leq t \leq 1 \right\}.$$

For $p \in [1, \infty)$ and $t \in [0, 1]$, it is easy to see that $J_{X,p}(t) = 2^{-1/p} \gamma_{X,\psi_p}(t)$. Thus we have the following results on the James and von Neumann-Jordan type constants.

Corollary 3.3. *Let X be a Banach space with the predual Banach space.*

(1) *For any $p \in [1, \infty)$ and any $t \in [0, 1]$,*

$$J_{X,p}(t) = \sup \left\{ \left(\frac{\|x+ty\|^p + \|x-ty\|^p}{2} \right)^{1/p} : x, y \in \text{ext}(B_X) \right\}$$

(2) *For any $p \in [1, \infty)$,*

$$C_p(X) = \sup \left\{ \frac{(\|x+ty\|^p + \|x-ty\|^p)^{2/p}}{2^{2/p}(1+t^2)} : x, y \in \text{ext}(B_X), 0 \leq t \leq 1 \right\}.$$

In particular, on the modulus of convexity and the NJ constant, one can easily have

$$\rho_X(t) = J_{X,1}(t) - 1 = \frac{\gamma_{X,\psi_1}(t)}{2} - 1$$

for any $t \in [0, 1]$, and

$$C_{NJ}(X) = C_2(X) = \sup \left\{ \frac{\gamma_{X,\psi_2}(t)^2}{2(1+t^2)} : 0 \leq t \leq 1 \right\}.$$

Hence we obtain

Corollary 3.4. *Let X be a Banach space with the predual Banach space. Then,*

$$\rho_X(t) = \sup \left\{ \frac{\|x+ty\| + \|x-ty\|}{2} - 1 : x, y \in \text{ext}(B_X) \right\}$$

for all $t \in [0, 1]$, and

$$C_{NJ}(X) = \sup \left\{ \frac{\|x+ty\|^2 + \|x-ty\|^2}{2(1+t^2)} : x, y \in \text{ext}(B_X), 0 \leq t \leq 1 \right\}.$$

4. Examples

For p, q with $1 \leq p, q \leq \infty$, the Day-James space $\ell_p\text{-}\ell_q$ is defined as the space \mathbb{R}^2 with the norm

$$\|(x_1, x_2)\|_{p,q} = \begin{cases} \|(x_1, x_2)\|_p & \text{if } x_1x_2 \geq 0, \\ \|(x_1, x_2)\|_q & \text{if } x_1x_2 \leq 0. \end{cases}$$

Yang and Wang [13] calculated the von NJ constant of the Day-James spaces $\ell_\infty\text{-}\ell_1$ and $\ell_2\text{-}\ell_1$ by using the notion of $\gamma_X(t)$.

Remark that $\ell_\infty\text{-}\ell_1$ and $\ell_2\text{-}\ell_1$ have the predual spaces $\ell_1\text{-}\ell_\infty$ and $\ell_2\text{-}\ell_\infty$, respectively (cf. [7]). Thus, from Theorem 3.2, we obtain

$$\gamma_{X,\psi}(t) = \sup\{\|(x + ty, x - ty)\|_\psi : x, y \in \text{ext}(B_X)\}$$

for X being $\ell_\infty\text{-}\ell_1$ or $\ell_2\text{-}\ell_1$.

Example 4.1. Let X be the Day-James space $\ell_\infty\text{-}\ell_1$, $\psi \in \Psi_2$ and $t \in [0, 1]$. Then

$$\text{ext}(B_X) = \{\pm(1, 1), (\pm 1, 0), (0, \pm 1)\},$$

and hence

$$\gamma_{X,\psi}(t) = (2 + t) \max \left\{ \psi \left(\frac{1}{2+t} \right), \psi \left(\frac{1+t}{2+t} \right) \right\}.$$

Example 4.2. Let X be the Day-James space $\ell_2\text{-}\ell_1$, $\psi \in \Psi_2$ and $t \in [0, 1]$. Then

$$\text{ext}(B_X) = \{(x_1, x_2) : x_1^2 + x_2^2 = 1, x_1x_2 \geq 0\},$$

and so

$$\begin{aligned} & \gamma_{X,\psi}(t) \\ &= (1 + t + \sqrt{1+t^2}) \max \left\{ \psi \left(\frac{1+t}{1+t+\sqrt{1+t^2}} \right), \psi \left(\frac{\sqrt{1+t^2}}{1+t+\sqrt{1+t^2}} \right) \right\}. \end{aligned}$$

We note that some geometric constants does not necessarily coincide with the supremum taken over all extreme points of the unit ball. The constant

$$C_Z(X) = \sup \left\{ \frac{\|x+y\| \|x-y\|}{\|x\|^2 + \|y\|^2} : x, y \in X, (x, y) \neq (0, 0) \right\}.$$

was introduced by Zbăganu [15]. As in the von Neumann-Jordan constant, this constant is reformulated as

$$C_Z(X) = \sup \left\{ \frac{\|x+ty\| \|x-ty\|}{1+t^2} : x, y \in S_X, 0 \leq t \leq 1 \right\}.$$

Example 4.3. Let X be the Day-James space $\ell_\infty\text{-}\ell_1$. Then

$$\sup \left\{ \frac{\|x+ty\| \|x-ty\|}{1+t^2} : x, y \in \text{ext}(B_X), 0 \leq t \leq 1 \right\} < C_Z(X).$$

From [9], the Zbăganu constant $C_Z(X)$ coincide with the von Neumann-Jordan type constant $C_0(X)$. Hence, for any $\psi \in \Psi_2$, the Zbăganu constant $C_Z(X)$ can not be expressed by the means of $\gamma_{X,\psi}$.

For any q less than 1, can we obtain a Banach space X in which the von Neumann-Jordan type constant $C_q(X)$ does not coincide with the supremum taken over all extreme points of the unit ball B_X ?

References

- [1] F. F. Bonsall and J. Duncan, *Numerical Ranges II*. London Math. Soc. Lecture Note Series, **10**, 1973.
- [2] J. A. Clarkson, *The von Neumann-Jordan constant for the Lebesgue spaces*. Ann. of Math., **38** (1937), 114–115.
- [3] P. Jordan and J. von Neumann, *On inner products in linear metric spaces*. Ann. of Math., **36** (1935), 719–723.
- [4] K.-I. Mitani, S. Oshiro and K.-S. Saito, *Smoothness of ψ -direct sums of Banach spaces*. Math. Inequal. Appl., **8** (2005), 147–157.
- [5] K.-I. Mitani and K.-S. Saito, *A new geometrical constant of Banach spaces and the uniform normal structure*. Comment. Math., **49** (2009), 3–13.
- [6] H. Mizuguchi, *Some geometric constant and the extreme points of the unit ball of Banach space, to appear in Rev. Roumaine Math. Pures Appl.*
- [7] W. Nilsrakoo and S. Saejung, *The James constant of normalized norms on \mathbb{R}^2* . J. Inequal. Appl., 2006, Art. ID 26265, 12pp.
- [8] K.-S. Saito, M. Kato and Y. Takahashi, *Von Neumann-Jordan constant of absolute normalized norms on \mathbb{C}^2* . J. Math. Anal. Appl., **244** (2000), 515–532.
- [9] Y. Takahashi, *Some geometric constants of Banach spaces—a unified approach*. Proc. of 2nd International Symposium on Banach and Function Spaces **II** (2008), 191–220.
- [10] Y. Takahashi and M. Kato, *von Neumann-Jordan constant and uniformly non-square Banach spaces*. Nihonkai. Math. J., **9** (1998), 155–169.
- [11] Y. Takahashi, M. Kato and K.-S. Saito, *Strict convexity of absolute norms on \mathbb{C}^2 and direct sums of Banach spaces*. J. Inequal. Appl., **7** (2002), 179–186.
- [12] C. Yang, *An inequality between the James type constant and the modulus of smoothness*. J. Math. Anal. Appl., **398** (2013), 622–629.
- [13] C. Yang and F. Wang, *On a new geometric constant related to the von Neumann-Jordan constant*. J. Math. Anal. Appl., **324** (2006), 555–565.

- [14] C. Yang and F. Wang, *Some properties of James type constant*. Appl. Math. Lett., **25** (2012), 538–544.
- [15] G. Zbăganu, *An inequality of M. Rădulescu and S. Rădulescu which characterizes the inner product spaces*. Rev. Roumaine Math. Pures Appl., **47** (2002), 253–257.