

Free product of generalized Gaussian processes, random matrices and positive definite functions on permutation groups

Marek Bożejko^{1*}; Wojciech Bożejko^{2†}

¹Institute of Mathematics
Polish Academy of Sciences
bozejko@gmail.com

²Department of Control Systems and Mechatronics
Wrocław University of Science and Technology
wojciech.bozejko@pwr.edu.pl

Dedicated to prof. Nobuaki Obata for his 60th birthday.

Abstract

The paper deals with the free product of generalized Gaussian process with function $t_b(V) = b^{H(V)}$, where $H(V) = n - h(V)$, $h(V)$ is the number of singletons in a pair-partition $V \in \mathcal{P}_2(2n)$. Some new combinatorial formulas are presented. Connections with free additive convolutions probability measure on \mathbb{R} are also done. Also new positive definite functions on permutations are presented and also it is proved that the function H is norm (on the group $S(\infty) = \bigcup S(n)$). Connection with random matrices and positive definite functions on permutations groups are also done.

1 Introduction

We present some new construction of generalized Gaussian processes and its relations with random matrices as well as with positive definite functions defined on permutations groups. The plan of the paper is following: first we present definitions and remarks on pair-partitions. Next, Markov random matrices and function h on pair-partitions are presented in the Section 3 – $\mathcal{P}_2(2n)$ as obtained by Bryc, Dembo, Jiang [B-D-J]. Generalized strong Gaussian processes (fields) $\{G(f) \text{ are showed in the Section 4, } f \in \mathcal{H}\}$ (GSGP), \mathcal{H} - real Hilbert space, as

*The work was partially supported by Opus grant no. DEC 2016/21/B/ST1/00628

†The work was partially supported by Opus grant no. DEC 2017/25/B/ST7/02181

well as the main and the new examples. The main theorem is placed in the Section 5. The free product of (GSGP) $G(f)$ and free Gaussian field $G_0(f)$ is again (GSGP) – this appears in this paper as the Theorem 3. In the Section 5 we also present the new interesting results on free convolutions of measures extending results of Bożejko and Speicher [B-Sp2]. Such negative-definite functions have many applications in the field of telecommunication, parallel and quantum computing as well as in operations research (see [B-Wo, Boz2]). We show that the function $H(\sigma) = n - h(\sigma)$, $\sigma \in S(n)$ is conditionally negative definite, i.e. for each $x > 0$, $\exp(-x H(\sigma))$ is positive functions on the permutation group $S(\infty) = \bigcup S(n)$. Also in the Theorem 6 (Section 6) it is shown that the function H defined as $d_H(\sigma, \tau) = H(\sigma^{-1}\tau)$ is left-invariant distance on $S(\infty)$.


2 Definitions and remarks on pair-partitions

Definition 1. Let γ_0 is the Wigner (semicircle) law with density $\frac{1}{\sqrt{2\pi}}\sqrt{4 - x^2}dx$. This is free Gaussian law. By γ_1 we denote the Normal law $N(0, 1)$ with density $\frac{1}{\sqrt{2\pi}}e^{-x^2/2}$. The moments (only even)

$$\int_{-\infty}^{\infty} x^{2n} d\gamma_1(x) = m_{2n}(\gamma_1) = \sum_{V \in \mathcal{P}_2(2n)} 1 = 1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1) = (2n-1)!! = p_{2n}, \tag{1}$$


where $\mathcal{P}_2(2n)$ is the set of all pair-partitions on $2n$ -elementary set $\{1, 2, \dots, 2n\}$.

The moments of the Wigner law $m_{2n}(\gamma_0) = \frac{1}{n+1} \binom{2n}{n}$ = the cardinality of all non-crossing pair-partitions of $\mathcal{P}_2(2n)$, where a partition $V \in \mathcal{P}_2(2n)$ has a crossing if there exists blocks $(i_1, j_1), (i_2, j_2) \in V$ such that $i_1 < i_2 < j_1 < j_2$:



Let $cr(V) = \#$ of all crossings of pair-partition V .

In contrary, the partition V is called *non-crossing*; $NC_2(2n)$ – denote the set of all non-crossing pair-partitions on $2n$ -elements set $\{1, 2, \dots, 2n\}$. Pictorially:



It is well known, that the cardinality of $NC_2(2n) = \frac{1}{n+1} \binom{2n}{n}$.

Definition 2. The block $B \in V \in \mathcal{P}_2(2n)$ is a **singleton**, if B has no crossing with other block $C \in V$.

Ex. $\overbrace{\overbrace{\overbrace{1 \ 2 \ 3 \ 4 \ 5 \ 6}}^{(3,4)=B}}$, the block $(3, 4) = B$ is a singleton.

Following [B-D-J], let us denote $h(V) = \#$ of singletons in the pair-partition $V \in \mathcal{P}_2(2n)$.

For example: if $V = \overbrace{\overbrace{1 \ 2 \ 3 \ 4 \ 5 \ 6}}$ then $h(V) = 1$.

Facts:

(α) If V is **non-crossing pair-partition** on $\{1, 2, \dots, 2n\}$, then $h(V) = n$.

The important fact for us is the following:

(β) If V is connected and $V \in \mathcal{P}_2(2n)$, $n > 1$, then $h(V) = 0$,

for example $h(\overbrace{\overbrace{1 \ 2 \ 3 \ 4 \ 5 \ 6}}) = 0$.

The pair-partition $V \in \mathcal{P}_2(2n)$ is *connected* if its graph is *connected* set. For example $\overbrace{1 \ 2}$ is connected. Also $\overbrace{\overbrace{1 \ 2 \ 3 \ 4 \ 5 \ 6}}$ is connected.

Let $cc(V) =$ the number of connected components of the graph of the partition $V \in \mathcal{P}_2(2n)$, and

let $c_{2n} = \#\{V \in \mathcal{P}_2(2n) : V \text{ is connected}\}$.

That sequence is *the free cumulant* of the classical Gaussian distribution $N(0, 1)$, i.e.:

$$c_2 = 1, \quad c_4 = 1, \quad c_6 = 4, \quad c_8 = 27, \quad c_{10} = 248, \dots$$

The following formula is due to Riordan [R], see also Belinschi, Bożejko, Lehner, Speicher [B-B-L-S]:

$$c_{2n+2} = n \sum_{i=1}^n c_{2i} c_{2(n-i-1)} \quad (2)$$

and that sequence is the moment sequence of some symmetric probability measure on real line, as it was proved in [B-B-L-S].

That fact is equivalent to the following result:

Theorem 1. [B-B-L-S] *Normal law γ_1 is infinitely divisible in free \boxplus -convolution (i.e.: $\gamma_1 \in ID(\boxplus)$).*

One of our aim of this work is to find different proof of that above result using different method and the function $h(V)$.

Let us first prove the following:

Proposition 1. *Let us define $T_{2n} = \sum_{V \in \mathcal{P}_2(2n)} h(V)$, and*

$$p_{2n} = (2n - 1)!! = \sum_{V \in \mathcal{P}_2(2n)} 1, \quad p_{2n} = 1, 3, 15, 105, \dots,$$


then

$$T_{2n+2} = (n + 1) \sum_{k=0}^n p_{2k} \cdot p_{(2n-2k)}. \quad (3)$$

That sequence T_{2n} is following: 1, 4, 21, 144, 1245, 13140, 164745, ... For more informations about this sequence see Sloane integer sequences database (<https://oeis.org/>) no. A233481.

The proof of the formula (3) is by a simple considerations, if we consider pair-partitions as lying on a circle.

For example ($n = 3$):



3 Markov random matrices and function h on pair-partitions $\mathcal{P}_2(2n)$

Let $\{X_{ij} : j \geq i \geq 1\}$ be an infinite upper triangular array of i.i.d. random variables and define $X_{ji} = X_{ij}$ for $j > i \geq 1$.

Let $X_n = [X_{ij}]_{1 \leq i, j \leq n}$, and

$$D_n = \text{diag}\left(\sum_{j=1}^n X_{ij}\right)$$

is a diagonal matrix . We define *Markov matrices* M_n as a random matrix given by

$$M_n = X_n - D_n,$$

so then each of rows (and columns) of M_n has a zero sum.

Here for a symmetric $n \times n$ matrix A , its empirical distribution is done as

$$\hat{\mu}(A) = \frac{1}{n} \sum_{j=1}^n \delta_{\lambda_j(A)},$$

$\lambda_j(A)$, $1 \leq i \leq n$, denote the eigenvalues of the matrix A and δ_s is the Dirac mass at the point $s \in \mathbb{R}$.

The Theorem of Bryc, Dembo, Jiang [B-D-J] is following:

Theorem 2 ([B-D-J]). *If X_{ij} are i.i.d. random variables with $\mathbb{E}X_{ij} = 0$ and $\mathbb{E}X_{ij}^2 = 1$, then $\hat{\mu}(\frac{M_n}{\sqrt{n}}$ converges weakly as $n \rightarrow \infty$ to the measure $\gamma_M = \gamma_0 \boxplus \gamma_1$, where \boxplus denotes the free additive convolution of probability measures.*

The even moments of the measure $\gamma_M = \gamma_0 \boxplus \gamma_1$ are following:

$$m_{2n}(\gamma_M) = \sum_{V \in \mathcal{P}_2(2n)} 2^{h(V)}, \text{ and } m_{2n+1}(\gamma_M) = 0,$$

γ_0 is the Wigner (semicircle) law done by density $\frac{1}{2\pi} \sqrt{4-x^2} \cdot \chi[-2, 2]$, γ_1 is the Normal law $N(0, 1)$.

4 Generalized Gaussian process (field) $G(f)$, $f \in \mathcal{H}$, \mathcal{H} - real Hilbert space. Main and new examples

Let in some probability system $(\mathcal{A}, \varepsilon)$, (\mathcal{A} - *-algebra with unit, and ε is state on \mathcal{A}). The family $G(f) = G(f)^* \in \mathcal{A}$, $f \in \mathcal{H}$ - real Hilbert space, is called *normalized generalized Gaussian process* (GGP), if for each orthogonal map $\mathcal{O} : \mathcal{H} \rightarrow \mathcal{H}$, for $f_j \in \mathcal{H}_{\mathbb{R}}$, we have

$$\begin{aligned} \varepsilon(G(f_1)G(f_2)\dots G(f_k)) &= \varepsilon(G(\mathcal{O}(f_1))G(\mathcal{O}(f_2))\dots G(\mathcal{O}(f_k))) = \\ &= \begin{cases} 0 & \text{for } k - \text{ odd,} \\ \sum_{V \in \mathcal{P}_2(2n)} t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle & \text{for } k = 2n, \end{cases} \end{aligned}$$

for some function $t : \mathcal{P}_2(\infty) \rightarrow \mathbb{C}$ which will be called positive definite on $\mathcal{P}_2(\infty)$ (see [B-Sp2], Guţă-Maassen [G-M1, G-M2], for further facts), with normalization $t(1 \ 2) = 1$.

The main examples of (GGP) are related to q-CCR relations [B-Sp1],[B-K-S]:

$$(q\text{-CCR}) \quad a(f)a^*(q) - q a^*(q)a(f) = \langle f, q \rangle I$$

$-1 \leq q \leq 1$, $a(f)\Omega = 0$, $f, q \in \mathcal{H}_{\mathbb{R}}$ and $\|\Omega\| = 1$, Ω - vacuum vector.

If we take $G_q(f) = a(f) + a^*(f)$, and as a state - vacuum state:

$\varepsilon(T) = \langle T\Omega | \Omega \rangle$, then we get q -Gaussian field and $t_q(V) = q^{cr(V)}$ for $V \in \mathcal{P}_2(2n)$, where $cr(V)$ is the number of crossings in a partition V .

Other examples were constructed by Bożejko-Speicher [B-Sp2] by the function:

$$t_s(V) = s^{n-cr(V)}, \quad a \leq s \leq 1.$$

That examples were important to prove that Normal law γ_1 is free infinitely divisible, i.e. $\gamma \in ID(\boxplus)$.

Namely, the following fact was proven in [B-Sp2]. For $s \geq 1$

$$m_{2n}(\gamma_1^{\boxplus s}) = \sum_{V \in \mathcal{P}_2(2n)} s^{cc(V)},$$

where \boxplus is the free additive convolution. Many other examples were done by Accardi-Bożejko [A-B], Guţă-Maassen [G-M1, G-M2], Bożejko-Yoshida [B-Y], Bożejko-Guţă [B-G], Bożejko [Boz1] and Bożejko-Wysoczański [B-W].

Our work presents among others the proof of the result of A. Buchholz [Buch], that it exists a explicite realization of generalized Gaussian process connected with the function of Bryc-Dembo-Jiang

$$t_b(V) = b^{n-h(V)} = b^{H(V)}$$

for $0 \leq b \leq 1$, $V \in \mathcal{P}_2(2n)$, and this is consequence of *our Main Theorem*.

We define generalized strongly Gaussian process $(G(f), t_G, \varepsilon)$, $f \in \mathcal{H}$, as generalized *Gaussian process*, such that the function $t_G = t$ on $\mathcal{P}_2(\infty)$ is *strongly multiplicative*, i.e.

$$t(V_1 \dot{\cup} V_2) = t(V_1) \cdot t(V_2),$$

for $V_1, V_2 \in \mathcal{P}_2(\infty)$, which are pair-partitions on disjoint sets.

The simple examples of strongly *multiplicative* functions on $\mathcal{P}_2(\infty)$ are following:

$$\begin{aligned} 1^\circ & q^{cr(V)}, \\ 2^\circ & s^{n-cc(V)}, \\ 3^\circ & b^{n-h(V)}, \end{aligned}$$

see also [B-G] for more strongly multiplicative examples, related to the Thoma characters on $S(\infty)$ group.

That classes of processes correspond to so called *pyramidal* independence, which has been considered by B. Kümmerer [K], see also [B-Sp2].

In all above examples: if $q = b = s = 1$, we get classical Gaussian process.

Important fact: if $\mathcal{H} = L^2(\mathbb{R}^+, dx)$ and $B_u = G(\chi_{[0,u]})$, where $\chi_{[0,u]}$ is the characteristic function of interval $[0, u]$, then B_u , for $u \geq 0$, is a realization of classical *Brownian motion*.

If we take $q = b = s = 0$, we get a construction of the *free Brownian motion* (see books of Voiculescu, Dykema, Nica [V-D-N], Nica, Speicher, [N-S] and Hiai-Petz [H-P]).

5 The main theorem

Theorem 3. *If $G(f)$ is normalized generalized strong Gaussian process, $f \in \mathcal{H}$, \mathcal{H} is a real Hilbert space, and $G_0(f)$ is the free Gaussian process, and operators*

$\{G(f) : f \in \mathcal{H}\}$ and $\{G_0(f) : f \in \mathcal{H}\}$ are free independent in some probability system $(\mathcal{A}, \varepsilon)$, then for each $0 \leq b \leq 1$, the process:

$$X_b(f) = \sqrt{b} G(f) + \sqrt{1-b} G_0(f), \quad f \in \mathcal{H}$$

is again generalized strong Gaussian process.

Moreover, if $G(f)$ corresponds to strongly multiplicative function

$$t_G : \bigcup_{A_n} \mathcal{P}_2(2n) \rightarrow \mathbb{C}$$

done by equation

$$\varepsilon(G(f_1) \dots G(f_k)) = \begin{cases} \sum_{V \in \mathcal{P}_2(2n)} t_G(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle, & \text{if } k = 2n, \\ 0, & \text{if } k \text{ odd.} \end{cases}$$

Then the corresponding strongly multiplicative function of the generalized Gaussian process X_b , $0 \leq b \leq 1$ is following:

$$t_{X_b}(V) = b^{H(V)} \cdot t_G(V), \quad \text{for } V \in \mathcal{P}_2(2n).$$

i.e.

$$\varepsilon(X_b(f_1) X_b(f_2) \dots X_b(f_{2n})) = \sum_{V \in \mathcal{P}_2(2n)} b^{H(V)} t_G(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle,$$

for $f_i \in \mathcal{H}$, and odd moments are zero.

In the proof of the Theorem 3 we will need the following Lemma.

Main Lemma. Let t be strongly multiplicative function corresponding to strongly generalized Gaussian process (field) $G(f) = G_t(f)$, $f \in \mathcal{H}_{\mathbb{R}}$ such that

$$\varepsilon(G(f_1) \dots G(f_k)) = \begin{cases} \sum_{V \in \mathcal{P}_2(2n)} t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle, & k = 2n, \\ 0, & \text{if } k = 2n + 1. \end{cases}$$


then the free cumulants are following:

$$r_k(G(f_1), G(f_2), \dots, G(f_k)) = \begin{cases} \sum_{\substack{V \in \mathcal{P}_2(2n) \\ cc(V) = 1}} t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle, & k = 2n, \\ 0, & k = 2n + 1. \end{cases} \quad (4)$$

Proof. Let $NC_e(2n)$ denotes the set of all even non-crossing partition \mathcal{V} of $2n$, which all blocks of \mathcal{V} are even.

As in the proof of Theorem 11 in [B-Y], we define the mapping $\Phi : \mathcal{P}_2(2n) \rightarrow NC_e(2n)$ as follows: given a pair-partition $V \in \mathcal{P}_2(2n)$, the connected components of V , will induce the even non-crossing partition $\Phi(V) = W$.

For example, if



(i.e. Φ in some sense forgets crossings of partitions).

Let us denote $G(f_j) = g_j$ in the proof. Since all odd moments of g_j vanish, hence also all odd free cumulants are vanish, i.e. $r_{2k+1}(g_{i_1}, \dots, g_{i_{2k+1}}) = 0$.

Therefore the free moment-cumulant formula is following:

$$\varepsilon(g_1 g_2 \dots g_{2n}) = \sum_{V \in NC_e(2k)} \prod_{\substack{B \in V \\ B = \{i_1, \dots, i_{2s}\}}} r_{2s}(g_{i_1}, \dots, g_{i_{2s}}). \quad (5)$$

By the assumption we have

$$\varepsilon(G(f_1) \dots G(f_{2n})) = \varepsilon(g_1 g_2 \dots g_{2n}) = \sum_{V \in \mathcal{P}_2(2n)} t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle \quad (6)$$

where t is strongly multiplicative.

Let us denote

$$\tilde{r}_{2k}(g_1, g_2, \dots, g_{2k}) = \sum_{\substack{\rho \in \mathcal{P}_2(2k) \\ cc(\rho) = 1}} t(\rho) \prod_{(i,j) \in \rho} \langle f_i | f_j \rangle.$$

We want to show that

$$\tilde{r}_{2k}(g_1, \dots, g_{2k}) = r_{2k}(g_1, \dots, g_{2k}).$$

By the strong multiplicativity of t , we have that the function

$$\tilde{t}(\pi) = t(\pi) \prod_{(i,j) \in \pi} \langle f_i | f_j \rangle$$

is also strong multiplicative on the $\mathcal{P}_2(2n)$.

Let us fix a non-crossing partition $V \in NC_e(2n)$. By the strong multiplicative property of \tilde{t} , we have

$$\sum_{\substack{\pi \in \mathcal{P}_2(2n) \\ \Phi(\pi) = V}} \tilde{t}(\pi) = \prod_{\substack{B \in V \\ B = \{i_1, i_2, \dots, i_{2s}\}}} \tilde{r}_{2s}(g_{i_1}, g_{i_2}, \dots, g_{i_{2s}}).$$

Therefore the formulas (4) and (5) implies that

$$\varepsilon(g_1 g_2 \dots g_{2n}) = \sum_{V \in NC_e(2n)} \prod_{\substack{B \in V \\ B = \{i_1, i_2, \dots, i_{2s}\}}} \tilde{r}_{2s}(g_{i_1}, g_{i_2}, \dots, g_{i_{2s}}).$$

Comparing with the formulas (4) and (5) and using Möbius inversion formula for the lattice of non-crossing partition (see Nica, Speicher book [N-S]) we get

$$\tilde{r}_{2s}(g_1, \dots, g_{2s}) = r_{2s}(g_1, \dots, g_{2s}).$$

■

Now we can start the proof of the Main Theorem using the Main Lemma.

Proof of the Main Theorem. By definition of the freeness of the families $\{G(f)\}_{f \in \mathcal{H}}$, $\{G_0(f)\}_{f \in \mathcal{H}}$ in the probability system $(\mathcal{A}, \varepsilon)$, we have that all mixed free cumulants

$$r_k(G_{\varepsilon_1}(f_1), G_{\varepsilon_2}(f_2), \dots, G_{\varepsilon_k}(f_k)) = 0, \text{ for all } k = 2, 3, \dots,$$

if the sequence $(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_k)$ is not constant ($\varepsilon_j \in \{0, 1\}$) and in the proof we denote $G_1(f) \stackrel{\text{def}}{=} G(f)$.

Therefore the free cumulants of $X_b(f) = \sqrt{b} G(f) + \sqrt{1-b} G_0(f)$ are following:

$$\begin{aligned} r_{2k}(X_b(f_1), \dots, X_b(f_{2k})) &= \\ &= b^k r_{2k}(G(f_1), \dots, G(f_{2k})) + (1-b)^k r_{2k}(G_0(f_1), \dots, G_0(f_{2k})) \end{aligned} \quad (7)$$

and all odd free cumulants of $X_b(f)$ are zero.

From the assumption $G_0(f)$ is the free normalized Gaussian process, i.e.

$$t_{G_0}(V) = \begin{cases} 1, & \text{if } V \in NC_2(2k), \\ 0, & \text{otherwise.} \end{cases}$$

and this is equivalent that

$$r_2(G_0(f_1), G_0(f_2)) = \langle f_1, f_2 \rangle$$

and

$$r_{2k}(G_0(f_1), \dots, G_0(f_{2k})) = 0,$$

for $k > 1$ and arbitrary $f_j \in \mathcal{H}_{\mathbb{R}}$.

That above facts follow at once from our Main Lemma, as we can see now:

Since by definition:

$$\varepsilon(G_0(f_1), \dots, G_0(f_{2k})) = \sum_{V \in NC_2(2k)} \prod_{(i,j) \in V} \langle f_i | f_j \rangle$$

hence by Main Lemma:

$$r_{2k}(G_0(f_1), \dots, G_0(f_{2k})) = \sum_{\substack{V \in NC_2(2k) \\ cc(V) = 1}} \prod_{(i,j) \in V} \langle f_i | f_j \rangle.$$

But if $V \in NC_2(2k)$, and $cc(V) = 1$, then we have that $k = 1$ and therefore for $k > 1$

$$r_{2k}(G_0(f_1), \dots, G_0(f_{2k})) = 0.$$

If now $V \in \mathcal{P}_2(2k)$ and $cc(V) = 1$, $k > 1$, then we have $H(V) = k - h(V) = k$, we get by (7) and Main Lemma

$$\begin{aligned} r_{2k}(X_b(f_1), \dots, X_b(f_{2k})) &= b^k r_{2k}(G(f_1), \dots, G(f_{2k})) = \\ &= \sum_{\substack{V \in \mathcal{P}_2(2k) \\ cc(V) = 1}} b^k t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle = \sum_{\substack{V \in \mathcal{P}_2(2k) \\ cc(V) = 1}} b^{H(V)} t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle. \end{aligned}$$

On the other hand, for $k = 1$ we have

$$r_2(X_b(f_1), X_b(f_2)) = b \langle f_1 | f_2 \rangle + (1 - b) \langle f_1 | f_2 \rangle = \langle f_1 | f_2 \rangle,$$

since $\{G(f)\}$ is normalized Gaussian process, i.e. $t(\square) = 1$ and since $H(\square) = 1 - h(\square) = 0$.

Therefore for all $k \geq 1$

$$r_{2k}(X_b(f_1), \dots, X_b(f_{2k})) = \sum_{\substack{V \in \mathcal{P}_2(2k) \\ cc(V) = 1}} b^{H(V)} t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle$$

Now using again the free moment-cumulant formula, our Main Lemma and the strong multiplicativity of the function $b^{H(V)}t(V)$, we get

$$\varepsilon(X_b(f_1), X_b(f_2), \dots, X_b(f_{2k})) = \sum_{V \in \mathcal{P}_2(2k)} b^{H(V)} t(V) \prod_{(i,j) \in V} \langle f_i | f_j \rangle. \quad \blacksquare$$

After that considerations a natural problem appears:

Problem 1. Is it true that if we have 2 strongly generalized Gaussian processes $\{G_1(f)\}_{f \in \mathcal{H}_{\mathbb{R}}}$, $\{G_2(f)\}_{f \in \mathcal{H}_{\mathbb{R}}}$ which are free, that the Gaussian process $Z(f) = G_1(f) + G_2(f)$ is again strongly Gaussian?

As a corollary from the Main Lemma we get well-known similar simple proposition for probability measures on real line (see [B-Sp2], [B-D-J], [Leh]):

Proposition 2. *If μ is symmetric measure on \mathbb{R} with all moments, and*

$$m_{2n}(\mu) = \sum_{V \in \mathcal{P}_2(2n)} t(V), \text{ where } t - \text{strongly multiplicative,}$$

then the free cumulants of the measure μ are of the form:

$$r_{2n}(\mu) = \sum_{\substack{V \in \mathcal{P}_2(2n) \\ cc(V) = 1}} t(V).$$

Now we show some special case of our results.

In particular case, let $f \in \mathcal{H}$, $\|f\| = 1$, and let the law of $G(f)$ is the probability measure μ on \mathbb{R} , $\mathcal{L}(G(f)) = \mu$, i.e.

$$\varepsilon(G(f)^k) = \int_{\mathbb{R}} \lambda^k d\mu(\lambda), \quad k = 0, 1, 2, \dots$$

and let γ_0 is the law of the Wigner-semicircle-free Gaussian law $G_0(f)$, then the law of the process $X_b(f)$:

$$\mathcal{L}(X_b(f)) = D_{\sqrt{b}}(\mu) \boxplus D_{\sqrt{1-b}}(\gamma_0) = \mu_b,$$

here \boxplus is the *free additive convolution*, and D_λ is the *dilation of the measure done by the formula*:

$$(D_\lambda \mu)(E) = \mu(\lambda^{-1}E), \text{ for Borel set } E \subset \mathbb{R}, \quad \lambda > 0.$$

Hence from the Main Theorem we get that the even moments of the measure μ_b are following:

$$m_{2n}(\mu_b) = \int \lambda^{2n} d\mu_b(\lambda) = \sum_{V \in \mathcal{P}_2(2n)} b^{H(V)} \cdot t_G(V)$$

and $m_{2n+1}(\mu_b) = 0$.

If we take the classical Gaussian process as $G(f)$, corresponding to $t(V) \equiv 1$, for all $V \in \mathcal{P}_2(2n)$, so we get as corollary the completely different proof of the theorems of A. Buchholz [Buch].


Corollary 1. *For all $0 \leq b \leq 1$, the function $t(V) = b^{H(V)}$ is strongly multiplicative, tricable and positive definite on the set of all pair-partitions $\mathcal{P}_2(\infty) = \bigcup_n \mathcal{P}_2(2n)$.*

Here the function $H(V) = n - h(V)$ is *tricable*, i.e.

$$H(V) = H(\vec{V}), \text{ where}$$

for a pair-partition $V = \{(i_1, j_1), (i_2, j_2), \dots, (i_n, j_n)\}$, $\vec{V} = \{(\bar{i}_1, \bar{j}_1), (\bar{i}_2, \bar{j}_2), \dots, (\bar{i}_n, \bar{j}_n)\}$, \vec{V} is the cyclic rotation of our partition V ($i_k \rightarrow 1 + i_k$ (modulo $2n$)).

For example:



Remark 1. If $(G(f), t, f \in \mathcal{H})$ is generalized Gaussian process and t is **tracable**, then ε is a trace on the algebra generated by $G(f), f \in \mathcal{H}$, i.e.

$$\varepsilon(G(f_1) \dots G(f_k)) = \varepsilon(G(f_k)G(f_1) \dots G(f_{k-1})),$$

in general $\varepsilon(XY) = \varepsilon(YX)$, for X, Y in $*$ -algebra generated by the field $G(f), f \in \mathcal{H}$.

6 Free convolutions of measures

As Corollary 1 from the *Main Theorem* we get the following generalization of the Theorem 6 from [B-Sp2].

Proposition 3. Let for $0 \leq b \leq 1, \rho_b = D_{\sqrt{b}}(\gamma_1) \boxplus D_{\sqrt{1-b}}(\gamma_0)$ then for $0 \leq c \leq 1$

$$\rho_{(bc)} = D_{\sqrt{c}} \rho_b \boxplus D_{\sqrt{1-c}} \gamma_0. \tag{8}$$

This is a simple case of the following reformulation of the Main Theorem:

Theorem 4. If μ is symmetric probability measure on \mathbb{R} with all moments, such that the even moments of the measure μ are following:

$$m_{2n}(\mu) = \sum_{V \in \mathcal{P}_2(2n)} t(V),$$

and t is normalized and strongly multiplicative, then for $0 \leq b \leq 1$, the moments of the measure $D_{\sqrt{b}}(\mu) \boxplus D_{\sqrt{1-b}} \gamma_0 = \mu_b$, are of the form:

$$m_{2n}(\mu_b) = \sum_{V \in \mathcal{P}_2(2n)} b^{H(V)} \cdot t(V).$$

and the free cumulants of the measure μ and μ_b are following (see [B-Sp1] and [B-D-J], page 96):

$$r_{2n}(\mu) = \sum_{\substack{V \in \mathcal{P}_2(2n) \\ cc(V) = 1}} t(V),$$

$$r_{2n}(\mu_b) = b^n \sum_{\substack{cc(V) = 1 \\ V \in \mathcal{P}_2(2n)}} t(V) = b^n r_{2n}(\mu), \quad \text{for } \underline{n \geq 1}, 0 \leq b \leq 1.$$

and $r_2(\mu_b) = r_2(\mu)$.

Remark 2. If a measure μ is free infinitely divisible (i.e. $\mu \in ID(\boxplus)$), then for $0 \leq b \leq 1$

$$\mu_b = D_{\sqrt{b}}(\mu) \boxplus D_{\sqrt{1-b}}(\gamma_0) \text{ is also infinitely divisible,}$$

since Wigner semicircle law $\gamma_0 \in ID(\boxplus)$.

And vice versa,
if $\mu_b \in ID(\boxplus)$, for all $0 < b < 1$, then $\mu \in ID(\boxplus)$.

Problem 2. If we take $t(V) \equiv 1$, i.e. $\mu = \gamma_1$, $0 \leq b \leq 1$, then

$$\sum_{V \in \mathcal{P}_2(2n)} b^{H(V)} = m_{2n}(\omega_b)$$

is the moment sequence of the probability measure $\omega_b = D_{\sqrt{b}}(\gamma_1) \boxplus D_{\sqrt{1-b}}(\gamma_0)$.

Is for $b > 1$, the sequence

$$m_{2n}(\mu) = \sum_{V \in \mathcal{P}_2(2n)} b^{H(V)}$$

the moment sequence of some symmetric probability measure?

See the paper [Boz1] on similar results for $q^{cr(V)}$, for $q > 1$.

7 Positive positive definite functions and „norm” on permutation group

In the paper [B-Sp2] we proved (Theorem 1) that if t is positive definite function on $\mathcal{P}_2(\infty) = \bigcup_{n=0}^{\infty} \mathcal{P}_2(2n)$, than for all natural n , the restriction of t to the permutation group $S(n)$ is also positive definite (in the usual sense), where the embedding $j : S(n) \rightarrow \mathcal{P}_2(2n)$ is done later.


We recall that $t : \mathcal{P}_2(\infty) \rightarrow \mathbb{C}$, is *positive definite function* on $\mathcal{P}_2(\infty)$, if there exists a *generalized Gaussian process (field)* $\{G_t(f), f \in \mathcal{H}\}$, \mathcal{H} – real Hilbert space, such that

$$\varepsilon[G_t(f_1)G_t(f_2)\dots G_t(f_k)] = \begin{cases} 0, & \text{if } k - \text{odd,} \\ \sum_{V \in \mathcal{P}_2(2n)} t(V) \prod_{(i,j) \in V} \langle f_i | j_j \rangle, & k = 2n \end{cases}$$

for some state ε on the $*$ -algebra generated by the $G_t(f)$, $f \in \mathcal{H}$ (see [G-M],[B-Sp2],[B-G],[Boz1],[B-Y], [B-W] for more examples of positive definite functions).

The our main theorem implies that the function $t_b(V) = b^{H(V)}$, for $0 \leq b \leq 1$, is positive definite on $\mathcal{P}_2(\infty)$, which gives another proof of Buchholtz theorem [Buch].

Now we define the **embedding** map $j : S(n) \rightarrow \mathcal{P}_2(2n)$ formulated as follows: for $\sigma \in S(n)$, $j(\sigma) = \{(k, 2n + 1 - \sigma(k)) : k = 1, 2, 3, \dots, n\} \in \mathcal{P}_2(2n)$. On the picture:



From that figure we can see that the number of *singletons* $h_n(\sigma) \stackrel{def}{=} h(j(\sigma))$, $\sigma \in S(n)$, is exactly the number of *fixed points* of the permutation σ , which are *isolated*.


The following Theorem is true:

Theorem 5. *The function h_{n+1} on $S(n + 1)$ is of the form*

$$h_{n+1} = \sum_{k=1}^{n+1} s_{k-1} \cdot \tilde{s}_{(n+1)-k},$$

and it is positive definite on the permutation group $S(n + 1)$.


Proof of the Theorem 5. Let $s_{k-1} = \chi_{s_{k-1}}$ is the characteristic function of the permutation group $S(k - 1)$ on $\{1, 2, \dots, k - 1\}$, and $\tilde{s}_{(n+1)-k}$ is the characteristic function of the symmetric group on the letters $\{k + 1, k + 2, \dots, n + 1\}$,



here $S(k - 1)$ is the group generated by inversions $\{\pi_1, \pi_2, \dots, \pi_{k-2}\}$, and the group $S(n + 1 - k)$ is generated by inversions $\{\pi_{k+1}, \pi_{k+1}, \dots, \pi_n\}$ (here $\pi_j = (j, j + 1)$ is the inversion (transposition) of $(j, j + 1)$). Hence $s_{k-1} \cdot \tilde{s}_{(n+1)-k}$ is the characteristic function of the group generated by $\{\pi_1, \dots, \pi_{k-2}, \pi_k, \pi_{k+1}, \dots, \pi_n\}$, so it is positive definite on the group $S(n + 1)$.

Let us define the function

$$h_{n+1}^{(k)}(\sigma) = \begin{cases} 1, & \text{if the singleton } (k, k) \text{ appears in the permutation } \sigma, \\ 0, & \text{otherwise.} \end{cases}$$



For the above picture one can see that: $h_{n+1}^{(k)} = s_{k-1} \cdot \tilde{s}_{(n+1)-k}$.

Since our function

$$h_{n+1} = \sum_{k=1}^{n+1} h_{n+1}^{(k)}, \text{ so we get}$$

$$h_{n+1} = \sum_{k=1}^n s_{k-1} \cdot \tilde{s}_{(n+1)-k}$$

and it is positive definite as finite sum of positive definite functions. \blacksquare

Corollary 2. For each $b \geq 1$ and $\sigma \in S(n)$

- (α) The function $S(n) \ni \sigma \rightarrow b^{h_n(\sigma)}$ is positive definite on $S(n)$.
- (β) The function $H_n(\sigma) = n - h_n(\sigma)$ is conditionally negative definite on $S(n)$ (i.e. $\exp(-x H_n(\sigma))$ is positive definite on $S(n)$ for all positive $x > 0$).
- (γ) The function $H_n(\sigma)$ is well defined on $S(\infty) = \bigcup S(n)$, $S(n) \subset S(n+1)$ (natural embedding) and $H_n = H_{n+1}|S(n)$, so we can define $H : S(\infty) \rightarrow \mathbb{R}$, as

$$H(\sigma) = H_n(\sigma) = n - h_n(\sigma), \text{ for } \sigma \in S(n).$$

Proof. The case (γ) can be easily checked, since the function $h_{n+1}^{(k)}$ is the characteristic function of the group generated by

$$\{\pi_1, \pi_2, \dots, \pi_{k-2}, \pi_{k+1}, \pi_{k+2}, \dots, \pi_n\},$$

so by the restriction of $h_{n+1}^{(k)}$ to the subgroup $\{\pi_1, \dots, \pi_{k-2}, \pi_{k+1}, \dots, \pi_{n-1}\}$, we get $h_{k+1}^{(k)} = h_n^{(k)} + 1$, so

$$H_n = H_{n+1}|S(n).$$

Both cases (α) and (β) follow from the theorems of I. Schur and I. Schoenberg (see [Boz0] or Berg, Christensen, Ressel [B-Ch-R]).

Now we can state:

Theorem 6. The function H is a „norm” on $S(n)$ and also on $S(\infty)$, i.e.

- (i) $H(e) = 0$
- (ii) $H(\sigma) = H(\sigma^{-1})$, $\sigma \in S(\infty)$
- (iii) $H(\sigma^{-1}\tau) \leq H(\sigma) + H(\tau)$, $\sigma, \tau \in S(\infty)$
- (iv) If we define a function $d(\sigma, \tau) = H(\sigma^{-1}\tau)$, $\sigma, \tau \in S(\infty)$, then d is a left-invariant metric on the group $S(\infty)$.

Proof. Let us see that for for $\sigma \in S_n$

$$H(\sigma) = n - \sum_{k=0}^{n-1} s_k(\sigma) \cdot \tilde{s}_{n-k}(\sigma) = \sum_{k=0}^{n-1} (1 - \chi_{s_k \times \tilde{s}_{n-k-1}}).$$

Let us denote $\Delta_k = 1 - \chi_{s_k \times \tilde{s}_{n-k-1}}$, $k = 0, 1, 2, \dots, n-1$, then Δ_k is conditionally negative definite and

$$\Delta_k(\sigma) = \begin{cases} 0, & \sigma \in S_k \times \tilde{S}_{n-1-k} \\ 1, & \text{otherwise,} \end{cases}$$

therefore $\sqrt{\Delta_k} = \Delta_k$.

By the well known property of the conditionally negative definite function (see [B-Ch-R], [Boz0]), we have

$$\Delta_k(\sigma\tau) = \sqrt{\Delta_k(\sigma\tau)} \leq \sqrt{\Delta_k(\sigma)} + \sqrt{\Delta_k(\tau)} = \Delta_k(\sigma) + \Delta_k(\tau).$$

Therefore our function

$$H = \sum_{k=0}^{n-1} \Delta_k$$

is also subadditive.

Other facts follow at once from the definition of the function H. ■

Remark 3. *More general considerations can be done for signed permutations group and arbitrary Coxeter groups using the result of the paper Bożejko et al. [B-G-M].*

8 Some questions and problems

Problem 3. If the assumption that the function $t : \mathcal{P}_2(\infty) \rightarrow \mathbb{R}$ is strongly multiplicative is necessary in our Main Theorem and Lemmas?

Problem 4. Let $\Gamma_b(\mathcal{H})$ is the von Neumann algebra generated by our b -Gaussian process

$$Y_b(f) = \sqrt{1-b}G_0(f) + \sqrt{b}G_1(f), \quad f \in \mathcal{H}_{\mathbb{R}},$$

where $G_0(f)$ is the free Gaussian process and $G_1(f)$ is the classical Gaussian process and $G_0(f)$ and $G_1(f)$ are free independent. As we can see in that von Neumann algebra $\Gamma_b(\mathcal{H})$, the vacuum is the trace, it is faithful and normal state.

Problem 5. Natural question is, if that von Neumann algebras $\Gamma_b(\mathcal{H})$ is a factor, $\Gamma_b(\mathcal{H}) = \{Y_b(f) : f \in \mathcal{H}_{\mathbb{R}}\}''$ (bicommutant), for $\dim \mathcal{H} \geq 2$.

Problem 6. If for $0 < b < 1$ that algebras $\Gamma_b(\mathcal{H})$ are isomorphic as von Neumann algebras?

Some others facts about our algebra $\Gamma_b(\mathcal{H})$ will be presented in the second part of our paper.

References

- [A-B] Accardi L., Bożejko M., Interacting Fock spaces and Gaussianization of probability measures, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* 1 (1998), 663–670.

- [B-B-L-S] Belinschi S.T., Bożejko M., Lehner F., Speicher R., The normal distribution is \boxplus -infinitely divisible, *Advances in Mathematics* 226 (2011), 3677–3698.
- [B-Ch-R] Berg Ch., Christiansen J., Rassel P., *Harmonic Analysis on Semigroups*, GTM 100, Springer (1984).
- [Boz0] Bożejko M., Positive and negative definite kernels on discrete groups, *Lectures at Heidelberg University* (1987).
- [Boz1] Bożejko M., Remarks on q -CCR relations for $|q| > 1$, in: *Banach Center Publications Vol. 78* (2007), 59–67.
- [B-G] Bożejko M., Guță M., Functors of white noise associated to characters of the infinite symmetric group, *Comm. Math. Phys.* 229 (2002), 209–227.
- [B-G-M] Bożejko M., Gal Ś., Młotkowski W., Positive definite functions on Coxeter groups with applications to operator spaces and noncommutative probability, *Comm. Math. Phys.* (2018), to appear.
- [B-Sp1] Bożejko M., Speicher R., An example of generalized Brownian motion, *Comm. Math. Phys.* 137(3) (1991) 519–531.
- [B-Sp2] Bożejko M., Speicher R., Interpolations between bosonic and fermionic relations given by generalized Brownian motions, *Math. Z.* 222 (1996), 135–160.
- [B-Y] Bożejko M., Yoshida H., Generalized q -Deformed Gaussian random variables, *Banach Center Publications*, Vol. 73 (2006), 127–140.
- [B-K-S] Bożejko M., Kümmerer B., Speicher R., q -Gaussian processes: Noncommutative and classical aspects, *Commun. Math. Phys.* 185 (1997), 129–154.
- [B-W] Bożejko M., Wysoczański J., New examples of convolution and noncommutative central limit theorems, *Banach Center Publ.* 43 (1998), 97–120.
- [Boz2] Bożejko W., Parallel path relinking method for the single machine total weighted tardiness problem with sequence-dependent setups, *Journal of Intelligent Manufacturing* 21(6), Springer (2010), 777–785.
- [B-Wo] Bożejko W., Wodecki M., On the theoretical properties of swap multi-moves, *Operations Research Letters*, Vol. 35 Issue 2, (2007), 227–231.
- [B-D-J] Bryc W., Dembo A., Jiang T., Spectral measure of large random Hankel, Markov and Toeplitz matrices, *Ann. of Prob.* 34 (2006), 1–38.
- [Buch] Buchholz A., New interpolations between classical and free Gaussian processes, Preprint, Wrocław (2008)..

- [G-M1] Guţă M., Maassen H., Symmetric Hilbert spaces arising from species of structures, *Math. Z.* 239 (2002), no. 3, 477-513.
- [G-M2] Guţă M., Maassen H., Generalised Brownian motion and second quantisation, *J. Funct. Anal.* 191 (2002), no. 2, 241-275.
- [H-P] Hiai F., Petz D., *The semicircle law, Free random variables and Entropy*, AMS (2000).
- [K] Kümmerner B., *Markov dilations and non-commutative Poisson processes*, Preprint, Tübingen (1987).
- [Leh] Lehner F., Cumulants in noncommutative probability theory, I-IV, *Math. Zeit.* 248 (2004), 67-100.
- [N-S] Nica A., Speicher R., *Lectures on the combinatorics of free probability* London Mathematical Society Lecture Note Series, Vol. 335 Cambridge University Press, 2006
- [R] Riordan J., The distribution of crossings of chords joining pairs of $2n$ points on a circle, *Math. Comp.* 29 (1975), 215-222.
- [V-D-N] Voiculescu D.V., Dykema K. J., Nica A., *Free random variables*, CRM Monograph Series, vol. 1, American Mathematical Society, Providence, RI, 1992.