

# Application of Resolvent CCR Algebras to Statistical Mechanics of Bosons on Lattices

Taku Matsui\*  
T.Kanda<sup>⊥</sup>

Graduate School of Mathematics, Kyushu University,  
744 Motoka, Nishi-ku, Fukuoka 819-0395, JAPAN

\* matsui@math.kyushu-u.ac.jp

<sup>⊥</sup> ma214013@math.kyushu-u.ac.jp

## 1 Introduction

In this note, we report application of the resolvent CCR algebras to statistical mechanics of Bosons on lattices. One of our motivation is originated from results on the quantum Ising model in a transversal magnetic field.

Let us recall known results of the quantum Ising model. First we consider the case of the model on one dimensional integer lattice . Let  $\mathfrak{A}$  be the UFH  $C^*$ -algebra which is an infinite tensor product of the algebra of 2 by 2 matrices where each component of the tensor is specified with a site in the integer lattice. We denote the sub-algebra of observables localized in  $\Lambda$  by  $\mathfrak{A}_\Lambda$  and we set  $\mathfrak{A}_{loc} = \cup_{\Lambda \subset \mathbf{Z}, |\Lambda| < \infty} \mathfrak{A}_\Lambda$ . For each natural number  $N$  the finite volume Hamiltonian  $H_N$  of the quantum Ising model on  $[-N, N]$  is defined by the following equation:

$$H_N = - \sum_{j=-N}^{N-1} \sigma_z^{(j)} \sigma_z^{(j+1)} + \lambda \sum_{j=-N}^N \sigma_x^{(j)}$$

where  $\sigma_{x,z}^{(j)}$  are Pauli spin matrices on the site  $j$  and  $\lambda$  is a real parameter. The limit

$$\alpha_t(Q) = \lim_{N \rightarrow \infty} e^{itH_N} Q e^{-itH_N}$$

exists in the norm topology and the Heisenberg time evolution  $\alpha_t$  gives rise to a  $C^*$ - dynamical system of  $\mathfrak{A}$ .

A state  $\omega$  of  $\mathfrak{A}$  is a  $\beta$ -KMS state if the time-dependent correlation function defined by  $F_{Q_1, Q_2}(t) \equiv \omega_\beta(\alpha_t(Q_1)Q_2)$  satisfy the following KMS condition:

$$F_{Q_1, Q_2}(t) = F_{Q_2, Q_1}(t + i\beta)$$

for any  $Q_1$  and  $Q_2$  which are entire analytic for  $\alpha_t$ . Note that  $Q$  is entire analytic if  $\alpha_t(Q)$  as a function of  $t$  has an analytic extension to the whole complex plain. The set of analytic elements is dense in  $\mathfrak{A}$ . As the thermal equilibrium state at the inverse temperature  $\beta$  satisfy this KMS condition, it is natural to regard  $\beta$ -KMS states as the equilibrium states for quantum systems with an infinite degree of freedom.

In the same spirit, we can introduce infinite volume ground states, namely, a state  $\omega$  of  $\mathfrak{A}$  is a ground state if the inequality

$$\lim_{N \rightarrow \infty} \omega(Q^*[H_N, Q]) \geq 0 \quad (Q \in \mathfrak{A}_{loc})$$

holds. The zero temperature limit of KMS states is a ground state defined in this manner.

In 1975, H.Araki proved uniqueness of  $\beta$ -KMS state for any one-dimensional quantum spin system with any short range interaction Hamiltonian. The ground state of the finite volume quantum Ising model is unique, and for the infinite volume ground states of the quantum Ising model, the following results are known.

**Theorem 1.1** (*H. Araki, Taku Matsui 1982CMP*)

- (i) *there exist precisely two infinite volume pure ground states if  $|\lambda| < 1$*
- (ii) *the  $\infty$  volume ground state is unique if  $|\lambda| \geq 1$*

For higher dimensional lattices  $\mathbf{Z}^d$  ( $2 \leq d$ ) the KMS state of quantum Ising model is unique at high temperature and at least two extremal low temperature KMS states exist. There exist two pure ground states when the transversal field is weak.

The problem we consider here is a Bosonic counterpart of quantum Ising models and in our opinion, a natural candidate is anharmonic crystals on  $\mathbf{Z}^d$ . The Hamiltonian of the quantum anharmonic crystal is written in the following form.

$$H = \sum_{k \in \mathbf{Z}^d} \{p_k^2 + V(x_k)\} + \sum_{k,l:|k-l|=1} \varphi(x_k - x_l).$$

where  $V(x)$  is a double well potential.

$$\lim_{x \rightarrow \infty} V(x) = \infty$$

with two local minima and  $\varphi$  represents interaction between adjacent particles.

One of mathematical difficulty for interacting Bose systems is to define the Heisenberg time evolution of quantum observables. In suitable setting, it is possible to show that the Heisenberg time evolution of the quantum anharmonic crystal does not exist as one-parameter group of automorphisms of the Weyl CCR algebra. Due to this fact, we employ the resolvent CCR algebra introduced by H.Grundling and D.Buchholz for their study of supersymmetric QFT.

Formally , the resolvent CCR algebra is generated by resolvent of linear combination of  $x_k$  and  $p_k$ . For example, in a quantum system with one degree of freedom, set

$$R_\lambda(s, t) = [\lambda i1 + (sx + tp)]^{-1} \quad \lambda, s, u \in \mathbf{R}$$

and the resolvent CCR algebra is the universal  $C^*$ -algebra generated by  $R_\lambda(s, t)$  and a unit.

More generally, let  $(V, \sigma)$  be a real symplectic space.  $R_{BG}$  is the universal  $C^*$ -algebra generated by a unit and  $\{R_\lambda(v) \mid v \in V\}$  satisfying relations For  $\lambda, \nu \neq 0, f, g \in V$

$$R_\lambda(0) = -\frac{i}{\lambda}, \quad R_\lambda(f)^* = R_{-\lambda}(f), \quad \nu R_\nu(\nu f) = R_\lambda(f)$$

$$R_\lambda(f) - R_\nu(f) = i(\nu - \lambda)R_\lambda(f)R_\nu(f)$$

$$[R_\lambda(f), R_\nu(g)] = i\sigma(f, g)R_\lambda(f)R_\nu(g)^2R_\lambda(f)$$

$$R_\lambda(f)R_\nu(g) = R_{\lambda+\nu}(f+g)\{R_\lambda(f) + R_\nu(g) + \sigma(f, g)R_\lambda(f)^2R_\nu(g)\}$$

Due to these relations, it is easy to see that there exists a trivial representation  $\pi$  of  $R_{BG}$  such that  $\pi(R_\lambda(f)) = 0$ . For representations and states of  $R_{BG}$  we require that field operators satisfying CCR (canonical commutation relations) can be reconstructed from  $R_\lambda(f)$ .

**Definition 1.2** (*H.Grundling and D.Buchholz*)

A representation  $\pi$  is regular if  $\pi(R_\lambda(f))$  is a resolvent of a closed operator for any  $f \in V$ ,

Regularity of a representation is equivalent to the condition  $\ker(R_\lambda(f)) = \{0\}$  for any  $f \in V$ . H.Grundling and D.Buchholz have shown that the CCR algebra can be reproduced for any regular representation of  $R_{BG}$ . They have shown the standard Fock representation is a regular faithful representation of  $R_{BG}$  as well.

## 2 Weakly Coupled Anharmonic Crystal

Now we consider the resolvent CCR algebra  $R_{BG}$  associated with weakly coupled anharmonic crystals. On each lattice site  $k$  in  $\mathbf{Z}^d$ , we have  $L_2(\mathbf{R}^d)$ . we set  $V = \mathbf{R}^{2\infty}$  which is a infinite direct sum of two dimensional symplectic space. Here we assume that all but finite components of the summand vanish.

We introduce the finite volume Hamiltonian of the weakly coupled anharmonic crystal as follows:

$$H_\Lambda = \sum_{k \in \Lambda} \{p_k^2 + w^2 x_k^2 + V(x_k)\} + \sum_{k, l \in \Lambda, |k-l|=1} \varphi(x_k - x_l)$$

Here we assume both  $V(x)$  and  $\varphi(x)$  are continuous functions, vanishing at infinity on  $\mathbf{R}^d$ .

The word "weakly coupled" is employed here as we presume  $\varphi(x)$  vanishes at infinity. We can show that

$$e^{itH_\Lambda} Q e^{-itH_\Lambda} \quad Q \in R_{BG}$$

is a well-defined automorphism of  $R_{BG}$ , but is not norm continuous in  $t$ . Thanks to Lieb-Robinson Bounds on the Fock representation, it is possible to prove existence of the infinite volume dynamics (the Heisenberg time evolution as automorphisms  $R_{BG}$ )

**Theorem 2.3** *The limit*

$$\lim_{\Lambda \rightarrow \mathbf{Z}^d} e^{itH_\Lambda} Q e^{-itH_\Lambda} = \alpha_t(Q)$$

*exists in the operator norm topology of the resolvent algebra, and  $\pi_F(\alpha_t(Q))$  is weakly continuous in  $t$  for the Fock representation  $\pi_F$ .*

As a function of  $t$ ,  $\alpha_t(Q)$  is not continuous in the norm topology of  $R_{BG}$ , nevertheless, we can define the  $\beta$ -KMS state.

We say a state  $\omega$  of  $R_{BG}$  is a  $\beta$ -KMS state if the following three conditions are valid:

- (i) The time dependent correlation function  $F_{Q_1, Q_2}(t) \equiv \omega_\beta(\alpha_t(Q_1)Q_2)$  is continuous for any  $t \in \mathbf{R}$
- (ii) For any  $Q_1$  and  $Q_2$  in  $R_{BG}$  there exists a complex function  $G_{Q_1, Q_2}(z)$  which is holomorphic in the strip  $\{z | 0 < \text{Im}z < \beta\}$ , and is bounded, continuous on the boundary of  $\{z | 0 < \text{Im}z < \beta\}$  such that  $G_{Q_1, Q_2}(t) = F_{Q_1, Q_2}(t)$ .
- (iii)  $G_{Q_1, Q_2}(t) = G_{Q_2, Q_1}(t)(t + i\beta)$

**Theorem 2.4** *For any  $\beta > 0$ , there exists a  $\beta$ -KMS state  $\omega_\beta$  of the resolvent algebra such that for any finite subset  $\Lambda \subset \mathbf{Z}^d$  the restriction of  $\omega_\beta$  to observables localized in  $\Lambda$  is normal to the Fock representation.*

This above normality is called locally normal to the Fock representation. If a KMS state  $\omega$  is locally normal to the Fock representation, we say  $\omega$  is a regular KMS state.

**Remark 2.5** *The following remarks are in order.*

- (i) *D. Buchholz considered a similar model in a slightly different setting. He restricted  $\alpha_t$  to a sub-algebra for which  $\alpha_t$  is continuous in  $t$ . The disadvantage of the approach is in the point that the sub-algebra does not contain functions of position operators and momentum operators.*
- (ii) *Non-regular KMS states exist, though, we cannot construct field operators in the GNS representation of non-regular KMS states, and we regard non-regular KMS states unphysical.*

By use of H.Araki's argument of relative entropy for quantum spin chains, and results of relative entropy of (not necessarily  $C^*$ -)  $*$ algebras due to A.Uhlmann, we can establish absence of phase Transition in one dimensional systems.

**Theorem 2.6** *If the dimension of the lattice is one, the regular KMS state of the weakly coupled anharmonic crystal is unique.*

Our weakly coupled anharmonic crystal is the first example of interacting quantum lattice models with unbounded spin in which uniqueness of KMS states established.

## References

- [1] S. Albeverio, Y. Kondratiev, Y. Kozitsky and M. Rockner, *The Statistical Mechanics of Quantum Lattice Systems: A Path Integral Approach*, (Ems Tracts in Mathematics), European Mathematical Society (2009)
- [2] H. Araki, *On uniqueness of KMS states of one-dimensional quantum lattice systems*, Comm. Math. Phys. **44**, no. 1, 1–7 (1975).
- [3] H. Araki, *Relative entropy of states of von Neumann algebras*, Publ. RIMS, Kyoto Univ. **13**, 173–192 (1977).
- [4] H. Araki, *Relative entropy for states of von Neumann algebras II*, Publ. RIMS, Kyoto Univ. **11**, 809–833 (1976).
- [5] O. Bratteli and D. Robinson, *Operator algebras and quantum statistical mechanics I*, 2nd edition (Springer, 1987).
- [6] O. Bratteli and D. Robinson, *Operator algebras and quantum statistical mechanics II*, 2nd edition (Springer, 1997).
- [7] D. Buchholz and H. Grundling, *Algebraic supersymmetry: A case study*, Commun. Math. Phys. **272**, 699–750 (2007)
- [8] D. Buchholz and H. Grundling, *The resolvent algebra: A new approach to canonical quantum systems*, J. Funct. Anal. **254**, 2725–2779 (2008)
- [9] D. Buchholz and H. Grundling, *Lie algebras of derivations and resolvent algebras*, Comm. Math. Phys. **320**, 455–467 (2013)
- [10] D. Buchholz, *The resolvent algebra for oscillating lattice systems: Dynamics, ground and equilibrium states* to appear in CMP.
- [11] M. Fannes and A. Verbeure, *On the Time Evolution Automorphisms of the CCR-Algebra for Quantum Mechanics*, Comm. Math. Phys. **35**, 257–264(1974)
- [12] M. B. Hastings. and T. Koma, *Spectral Gap and Exponential Decay of Correlations*, Comm. Math. Phys. **265**, 781–804 (2006).
- [13] F. Hiai, M. Ohya, M. Tsukada, *Sufficiency and relative entropy in \*-algebras with applications in quantum systems*, Pacific J. Math. **107**, no. 1, 117–140 (1983).
- [14] E. H. Lieb and D. W. Robinson, *The Finite Group Velocity of Quantum Spin Systems*, Comm. Math. Phys. **28**, 251–257 (1972).
- [15] R. Minlos, A. Verbeure and V. Zagrebnov, *A quantum crystal model in the light-mass limit: Gibbs states*, Rev. Math. Phys. **12** (7), 981–1032.(2000)
- [16] B. Nachtergaele and R. Sims, *Lieb-Robinson Bounds and the Exponential Clustering Theorem*, Comm. Math. Phys. **265**, 119–130 (2006).

- [17] B. Nachtergaele and R. Sims, *Recent Progress in Quantum Spin Systems*, Markov Processes Relat. Fields **13**, 315–329 (2007).
- [18] B. Nachtergaele and R. Sims, *Locality Estimates for Quantum Spin Systems*, arXiv:math-ph/0712.3318v1
- [19] B. Nachtergaele, H. Raz, B. Schlein and R. Sims, *Lieb-Robinson bounds for harmonic and anharmonic lattice systems*, Comm. Math. Phys. **286**, 1073–1098 (2009).
- [20] B. Nachtergaele and R. Sims, *Lieb-Robinson bounds in quantum many-body physics*, Entropy and the quantum, 141–176, Contemp. Math. **529**, Amer. Math. Soc. Providence, RI, 2010.
- [21] B. Nachtergaele and R. Sims, *On the dynamics of lattice systems with unbounded on-site terms in the Hamiltonian*, arXiv:math-ph/1410.8174
- [22] M. Ohya and D. Petz, *Quantum entropy and its use*, corrected Second printing, Texts and Monographs in Physics. Springer-Verlag, Berlin, 2004.
- [23] Powers, Robert T. and Sakai, Shoichiro, *Existence of ground states and KMS states for approximately inner dynamics*, Comm. Math. Phys. **39**, 273–288 (1975).
- [24] Uhlmann, A, *Relative entropy and the Wigner-Yanase-Dyson-Lieb concavity in an interpolation theory*, Comm. Math. Phys. **54**, no. 1, 21–32 (1977).